

Lipoprotein(a) and long-term structural valve degeneration of aortic bioprostheses

Marin Boute ^{1,2}, Paul Salembier^{1,2}, Anne-Catherine Pouleur ^{1,2},
 Agnès Pasquet ^{1,2}, Bernhard L. Gerber ^{1,2}, David De Azevedo ^{1,2},
 Damien Gruson^{2,3}, Laurent de Kerchove ^{2,4}, Joelle Kefer ^{1,2},
 Christophe Beauloye ^{1,2,5}, Sandrine Horman ², Frédéric Maes ^{1,2},
 Sophie Pierard ^{1,2,5}, and David Vancraeynest ^{1,2*}

¹Division of Cardiology, Cardiovascular Department, Cliniques Universitaires Saint-Luc, Avenue Hippocrate 10, Brussels 1200, Belgium; ²Pôle de Recherche Cardiovasculaire (CARD), Institut de Recherche Expérimentale et Clinique (IREC), Université Catholique de Louvain (UCLouvain), Avenue Hippocrate 55, Brussels 1200, Belgium; ³Department of Chemistry and Laboratory Medicine, Cliniques Universitaires Saint-Luc, Avenue Hippocrate 10, Brussels 1200, Belgium; ⁴Division of Cardiothoracic and Vascular Surgery, Cardiovascular Department, Cliniques Universitaires Saint-Luc, Avenue Hippocrate 10, Brussels 1200, Belgium; and ⁵Division of Cardiovascular Intensive Care, Cardiovascular Department, Cliniques Universitaires Saint-Luc, Avenue Hippocrate 10, Brussels 1200, Belgium

Received 31 August 2025; revised 13 October 2025; accepted 14 October 2025; online publish-ahead-of-print 17 November 2025

Aims

Structural valve degeneration (SVD) is the leading cause of late bioprosthetic valve failure. Lipoprotein(a) [Lp(a)] contributes to native aortic valve calcification, but its role in SVD is unclear. We investigated whether elevated Lp(a) is associated with SVD after bioprosthetic aortic valve replacement (AVR) and whether this differs between stenotic and regurgitant phenotypes.

Methods and results

We studied 174 bioprosthetic AVR patients with available Lp(a) levels over a median echocardiographic follow-up of 7.3 years (1372 studies). SVD was defined by VARC-3 criteria, and associations were analysed with Fine–Gray competing risk models. Lp(a) was evaluated categorically ($\leq$ or $>$ 125 nmol/L) and continuously using spline modelling. During follow-up, 40 patients developed SVD (22 stenotic, 9 mixed, and 9 regurgitant). The 15-year cumulative incidence was 51% with a median onset at 14.8 years. Elevated Lp(a) was associated with a higher risk of overall SVD (62% vs. 47%; SHR 2.06, 95% CI 1.09–3.91; $P = 0.026$) and specifically with stenotic/mixed phenotypes (SHR 2.57, 95% CI 1.26–5.23; $P = 0.009$). No association was observed with regurgitant phenotypes (SHR 0.85, 95% CI 0.19–3.92; $P = 0.84$). After multivariable adjustment, elevated Lp(a) remained an independent predictor of stenotic/mixed SVD (adjusted SHR 3.00, 95% CI 1.48–6.07; $P = 0.002$). Spline modelling showed a linear dose–response, with each 25 nmol/L increase in Lp(a) conferring 13% higher risk.

Conclusion


Elevated Lp(a) is independently associated with long-term risk of stenotic/mixed SVD. These findings highlight Lp(a) as a promising biomarker of prosthetic valve vulnerability and support investigation of emerging Lp(a)-lowering therapies to improve valve durability.

* Corresponding author. E-mail: david.vancraeynest@saintluc.uclouvain.be

© The Author(s) 2025. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.

Graphical Abstract



Elevated Lp(a) and phenotype-specific risk of bioprosthetic valve degeneration. In this study of bioprosthetic aortic valve replacement recipients with long-term transthoracic echocardiographic (TTE) follow-up, elevated lipoprotein(a) [Lp(a)] was independently associated with an increased risk of stenotic/mixed structural valve degeneration (SVD), with a linear dose–response but showed no association with isolated regurgitant degeneration. These findings highlight Lp(a) as a promising biomarker of bioprosthetic valve vulnerability, particularly for calcification-driven forms of degeneration

Keywords

prosthesis failure • echocardiography • biomarkers • competing risk analysis • spline modelling

Introduction

Bioprosthetic heart valves now represent the majority of aortic valve replacements (AVRs), with growing use in middle-aged patients expected to live for decades.^{1–4} As a result, long-term structural durability has become a critical concern. Structural valve degeneration (SVD), characterized by leaflet calcification, tearing, and hemodynamic deterioration, is the leading cause of late bioprosthetic failure.⁵ Established contributors include younger age, traditional cardiovascular risk factors, and prosthesis-related issues such as patient–prosthesis mismatch.⁶ Recent studies have explored circulating biomarkers, particularly lipid-related ones, as tools to refine SVD risk prediction.⁷

Among these, lipoprotein(a) [Lp(a)] is a biologically plausible candidate. It plays a causal role in native aortic valve stenosis via inflammation, oxidative stress, and mineralization—processes that also contribute to SVD.⁸ Yet, extrapolation is complicated by differences in timing and mechanism: SVD evolves over a shorter period and also involves bioprosthesis-specific factors such as structural fatigue and passive leaflet calcification.

Only a handful of studies have assessed the relationship between Lp(a) and SVD, with inconsistent results.^{9–13} These discrepancies likely reflect the methodological challenges of studying SVD, which emerges over a decade and is sometimes pre-empted by death or non-SVD re-intervention. Addressing this question requires extended follow-up and analytical methods that account for these competing risks.

Clarifying the role of Lp(a) in SVD is increasingly relevant, as Lp(a)-lowering agents such as pelacarsen and olpasiran enter late-stage cardiovascular trials and may reach clinical practice within a few years.¹⁴ Should a causal relationship be confirmed, Lp(a) could become a modifiable determinant of bioprosthetic valve durability.

In this study, we investigated whether elevated Lp(a) is independently associated with SVD following bioprosthetic AVR. We further hypothesized that Lp(a) might differentially influence stenotic vs. regurgitant SVD phenotypes, reflecting potentially distinct pathophysiological mechanisms.

Methods

Study design and population

This retrospective, single-centre study screened 3293 consecutive patients who underwent bioprosthetic AVR between 2001 and 2021. Of these, 174 patients had Lp(a) measured during routine follow-up and comprised the analytic cohort, after exclusion of three patients on chronic renal replacement therapy (see [Supplementary data online, Figure S1](#)). The primary objective was to evaluate whether elevated Lp(a) is independently associated with the risk of SVD after bioprosthetic AVR. The study was approved by the institutional review board.

Lp(a) assessment

Lp(a) was measured using an isoform-insensitive immunoturbidimetric assay [Roche Tina-quant® Lipoprotein(a) Gen.2, Roche Diagnostics,

Mannheim, Germany], which reports concentrations in molar units (nmol/L). This approach aligns with the current consensus recommending molar reporting to minimize bias from apo(a) size variation.⁸ Values were obtained as part of routine clinical care, typically for cardiovascular risk assessment, at the discretion of the treating physician. Lp(a) sampling occurred a median of 1 day before surgery [interquartile range (IQR) -64 to +3632 days]. Because ~90% of an individual's Lp(a) level is genetically determined and remains highly stable over decades, each measurement was considered representative of the patient's chronic Lp(a) exposure, independent of sampling date.^{15,16} For dichotomized analyses, a cutoff of 125 nmol/L was applied, in accordance with current guidelines.^{8,17}

Echocardiography and SVD definition

Transthoracic echocardiography (TTE) was performed using iE33 or EPIQ scanners (Philips Healthcare, Best, the Netherlands) by experienced operators. A post-operative TTE performed within 3 months of AVR served as the baseline reference for subsequent comparisons. Follow-up TTEs were obtained as part of routine clinical care, with frequency and timing determined by treating physicians. In total, 1372 TTEs were analysed (median 7 per patient; IQR 3–12). Measurements conformed to current guidelines and were obtained in sinus rhythm or averaged over three cardiac cycles in patients with atrial fibrillation. Parameters included mean transprosthetic gradient, effective orifice area (EOA), and Doppler velocity index (DVI); the latter two were measured when the mean gradient exceeded 20 mmHg. Prosthetic regurgitation was graded semi-quantitatively in accordance with guidelines.¹⁸ Prosthesis–patient mismatch was defined as recommended by current consensus, using BMI-adjusted thresholds and the predicted indexed effective orifice area (EOAi) rather than measured EOAi, to minimize variability from transient haemodynamics and operator dependence.¹⁹

SVD was defined according to Valve Academic Research Consortium 3 (VARC-3) criteria, for events of moderate or greater severity:⁵

- Stenotic phenotype: increase in mean gradient ≥ 10 mmHg to a value ≥ 20 mmHg with $\geq 25\%$ fall in EOA (≥ 0.3 cm²) or $\geq 20\%$ fall in DVI (≥ 0.10).
- Regurgitant phenotype: new or worsening intra-prosthetic regurgitation by ≥ 1 grade to at least moderate severity.
- Mixed phenotype: both criteria met.
- Overall SVD: first occurrence of stenotic and/or regurgitant phenotype.

The endpoint of the study was overall SVD. For phenotype-specific analyses, stenotic and regurgitant phenotypes were examined separately, with mixed phenotypes grouped with the stenotic category to avoid double counting of events. All cases meeting VARC-3 criteria were reviewed in consensus by two senior echocardiographers, and echocardiographic changes attributable to alternative causes (paravalvular leak, endocarditis, or leaflet thrombosis) were systematically excluded based on echocardiographic and clinical findings.

Clinical data and follow-up

Baseline clinical characteristics were recorded at the time of surgery. Follow-up information was retrieved from institutional electronic records and cross-checked with national health registries. Vital status was verified for every patient. Survivors without SVD were censored at the date of last echocardiographic contact. Death (all-cause) and redo AVR for other causes acted as competing events.

Statistical analysis

Continuous variables were summarized as mean $\pm$ SD or median [IQR] and categorical variables as counts and percentages. Between-group comparisons used Welch's *t*-test, Wilcoxon rank sum test, Pearson's chi-squared test, or Fisher's exact test as appropriate. Given the competing risk of death or redo AVR for non-SVD causes, cumulative incidence of SVD was

assessed with Fine–Gray subdistribution hazard models (*cmprsk* package v2.2), truncated at 15 years when $<10\%$ of patients remained at risk. Candidate predictors of valve failure were selected based on prior literature: age, sex, body surface area, hypertension, smoking, diabetes, LDL cholesterol, chronic kidney disease, and prosthesis–patient mismatch.⁶ All were included in a full multivariable model, with a parsimonious model derived by backward selection using the Akaike Information Criterion. To assess model stability, we performed bootstrap internal validation ($B = 2000$) of the full multivariable Fine–Gray model, applying the optimism-corrected calibration slope as a global shrinkage factor. We also explored penalized Fine–Gray regression (LASSO; *fastcmprsk* package v1.24.10) including the same covariates. To examine continuous associations, Lp(a) was modelled using a natural spline within a Fine–Gray competing risk model, with bootstrapped 95% confidence intervals. A Cox proportional hazards model using a restricted cubic spline (*rms* package v8.0) was also fitted for comparison. Finally, we performed a time-matched nested case–control analysis. Each SVD case was matched to up to two controls with similar follow-up duration (± 6 months) using nearest-neighbour matching based on Mahalanobis distance (*MatchIt* package v4.7.2). Analyses were also stratified by sex to explore potential effect modification. All analyses were conducted using R version 4.5.0. Statistical significance was set at a two-sided $P < 0.05$.


Results

Baseline characteristics and follow-up

We included 174 patients (mean age 72.6 ± 7.9 years; 36% female), of whom 133 (76%) had Lp(a) ≤ 125 nmol/L and 41 (24%) had Lp(a) > 125 nmol/L (Figure 1A). Baseline characteristics (Table 1) were well balanced between groups, except for a higher prevalence of coronary artery disease in the high Lp(a) group (51.2% vs. 33.1%; $P = 0.036$). Median echocardiographic follow-up was 7.3 years (IQR 4.5–10.9), with comparable durations in the low- and high-Lp(a) groups (7.3 vs. 7.2 years; $P = 0.840$). During follow-up, 40 patients developed SVD: 22 with a stenotic phenotype, 9 with a mixed (stenotic + regurgitant) phenotype, and 9 with an isolated regurgitant phenotype. The 15-year cumulative incidence of overall SVD was 51.0% (95% CI 38.1–62.5), with a median time to onset of 14.8 years (Figure 1B). Competing events (28 deaths and 3 redo AVRs for endocarditis) occurred in 31 patients, corresponding to a 15-year cumulative incidence of 22.7%, with no significant difference between Lp(a) groups ($P = 0.806$).

Association of Lp(a) by SVD phenotype

By 15 years, elevated Lp(a) was associated with a higher cumulative incidence of overall SVD [62% vs. 47% for lower Lp(a); Gray's test $P = 0.030$], with divergence between groups apparent by 10 years (37% vs. 18%; Figure 2A). Corresponding 10- and 15-year cumulative incidences for stenotic/mixed and regurgitant SVD phenotypes are shown in Figure 2B and C. In univariable Fine–Gray competing risk analysis, elevated Lp(a) (> 125 nmol/L) was associated with a twofold higher risk of overall SVD [subdistribution hazard ratio (SHR) 2.06, 95% CI 1.09–3.91; $P = 0.026$]. When phenotypes were analysed separately, this association was evident for stenotic/mixed SVD (SHR 2.57, 95% CI 1.26–5.23; $P = 0.009$) but no association was observed for regurgitant SVD (SHR 0.85, 95% CI 0.19–3.92; $P = 0.84$). Cumulative incidence curves were concordant, with clear separation between Lp(a) groups for stenotic/mixed SVD but not for regurgitant cases. In multivariable Fine–Gray models restricted to the stenotic/mixed SVD endpoint (Table 2), elevated Lp(a) remained an independent predictor (adjusted SHR 3.05, 95% CI 1.41–6.61; $P = 0.005$) in the full model including all predefined covariates. In the parsimonious model, only elevated Lp(a) (adjusted SHR 3.00, 95% CI 1.48–6.07; $P = 0.002$) and diabetes (adjusted SHR 2.68, 95% CI 1.30–5.55; $P = 0.008$) were retained as



When modelled as a continuous variable, each 25 nmol/L increment in Lp(a) conferred a 13% increase in the risk of stenotic/mixed SVD (SHR 1.13 per 25 nmol/L; 95% CI 1.05–1.22; $P = 0.002$). Results were consistent when Lp(a) was analysed on a log-transformed scale (SHR 1.39 per log-unit; $P = 0.024$). Spline analysis in the Fine–Gray model revealed an approximately linear dose–response, with no significant





Figure 2 Cumulative incidence of SVD by phenotype and Lp(a) group. Fine-Gray competing risk curves stratified by lipoprotein(a) [Lp(a)] group (≤ 125 nmol/L vs. >125 nmol/L) for (A) overall, (B) stenotic/mixed, and (C) regurgitant structural valve degeneration (SVD). Numbers at risk and competing events are displayed beneath each panel. Ten- and 15-year cumulative incidences are annotated on the curves. Gray's test *P*-values are indicated within each panel.



Study limitations

This study has several limitations. First, its retrospective, single-centre design may limit generalizability. However, prospective trials designed to capture long-term SVD outcomes face major logistical constraints due to the extended follow-up required. Second, Lp(a) testing was not performed systematically but at the discretion of treating physicians, raising the possibility of selection bias. Compared with the overall AVR population, the included patients were somewhat younger, with fewer comorbidities and more frequent implantation in later eras (see [Supplementary data online, Table S6](#)). While these limits representativeness, it is unlikely to explain the observed association between Lp(a) and SVD. Third, Lp(a) measurements were obtained at variable time points, but this is unlikely to have affected the results because Lp(a) concentrations are strongly genetically determined and highly stable over time. Notably, the association with stenotic/mixed SVD persisted when restricted to measurements obtained before or within 12 months after AVR, supporting the robustness of our conclusions. Finally, the analysis of isolated regurgitant SVD was limited by the small number of events, and sex-specific results remain underpowered given the smaller female subgroup. Both findings should therefore be considered exploratory. These limitations highlight the need for external validation in larger, multi-centre studies.

Conclusion

Elevated Lp(a) levels were independently associated with a higher long-term risk of stenotic and mixed SVD after bioprosthetic AVR, whereas no association was observed with isolated regurgitant degeneration. This phenotype-specific pattern suggests that Lp(a) may contribute predominantly to calcification-driven prosthetic valve failure. If confirmed in larger, multi-centre cohorts, a single Lp(a) measurement could refine SVD risk stratification and help tailor follow-up strategies. As the use of bioprosthetic valves continues to rise, dedicated studies represent a key next step to establish whether emerging Lp(a)-lowering therapies can translate into improved valve durability.

Supplementary data

Supplementary data are available at [European Heart Journal - Cardiovascular Imaging](#) online.

Author contributions

Marin Boute (Conceptualization [supporting]; Data curation [lead]; Formal analysis [lead]; Investigation [equal]; Methodology [equal]; Software [lead]; Visualization [lead]; Writing—original draft [lead]; Writing—review and editing [supporting]), Paul Salembier (Data curation [supporting]; Methodology [supporting]; Software [supporting]; Writing—review and editing [supporting]), Anne-Catherine Pouleur (Conceptualization [supporting]; Investigation [supporting]; Methodology [supporting]; Resources [equal]; Supervision [supporting]; Writing—review and editing [supporting]), Agnès Pasquet (Investigation [supporting]; Methodology [supporting]; Supervision [supporting]; Writing—review and editing [supporting]), Bernhard L Gerber (Investigation [supporting]; Methodology [supporting]; Resources [supporting]; Writing—review and editing [supporting]), David De Azevedo (Formal analysis [supporting]; Methodology [supporting]; Software [supporting]; Validation [supporting]; Writing—review and editing [supporting]), Damien Gruson (Data curation [supporting]; Resources [supporting]; Writing—review and editing [supporting]), Laurent de Kerchove (Data curation [supporting]; Resources [supporting]; Writing—review and editing [supporting]), Joelle Kefer (Data curation [supporting]; Resources [supporting];

Writing—review and editing [supporting]), Christophe Beauloye (Investigation [supporting]; Methodology [supporting]; Writing—review and editing [supporting]), Sandrine Horman (Conceptualization [supporting]; Investigation [supporting]; Methodology [supporting]; Writing—review and editing [supporting]), Frédéric Maes (Data curation [supporting]; Resources [supporting]; Writing—review and editing [supporting]), Sophie Pierard (Data curation [supporting]; Investigation [supporting]; Methodology [supporting]; Resources [supporting]; Writing—review and editing [supporting]), and David Vancraeynest (Conceptualization [lead]; Funding acquisition [supporting]; Investigation [supporting]; Methodology [supporting]; Project administration [supporting]; Supervision [equal]; Validation [supporting]; Writing—original draft [supporting]; Writing—review and editing [lead])

Funding

M.B. is a Research Fellow of the Fonds De La Recherche Scientifique—FNRS (Belgium).

Conflict of interest: Professor Bernhard Gerber is Editor-in-Chief of the *European Heart Journal—Cardiovascular Imaging*.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

- lung B, Delgado V, Rosenhek R, Price S, Prendergast B, Wendler O et al. Contemporary presentation and management of valvular heart disease: the euroobservational research programme valvular heart disease ii survey. *Circulation* 2019;**140**:1156–69.
- Kim KM, Arghami A, Habib R, Daneshmand MA, Parsons N, Elhalabi Z et al. The society of thoracic surgeons adult cardiac surgery database: 2022 update on outcomes and research. *Ann Thorac Surg* 2023;**115**:566–74.
- Sharaf OM, Beaver TM. Aortic valve 2024: which valve for which patient? *J Thorac Cardiovasc Surg* 2025;**170**:164–74.
- Isaacs AJ, Shuhaiber J, Salemi A, Isom OW, Sedrakyan A. National trends in utilization and in-hospital outcomes of mechanical versus bioprosthetic aortic valve replacements. *J Thorac Cardiovasc Surg* 2015;**149**:1262–9 e3.
- Généreux P, Piazza N, Alu MC, Nazif T, Hahn RT, Pibarot P et al. Valve academic research consortium 3: updated endpoint definitions for aortic valve clinical research. *Eur Heart J* 2021;**42**:1825–57.
- Rodriguez-Gabella T, Voisine P, Puri R, Pibarot P, Rodés-Cabau J. Aortic bioprosthetic valve durability: incidence, mechanisms, predictors, and management of surgical and transcatheter valve degeneration. *J Am Coll Cardiol* 2017;**70**:1013–28.
- Coinse A, Lancellotti P, Cote N, Ternacle J, Hecht S, Grapsa J et al. Blood biomarkers in left-sided valvular heart disease. *Eur Heart J* 2025;**46**:2289–305.
- Kronenberg F, Mora S, Stroes ESG, Ference BA, Arsenault BJ, Berglund L et al. Lipoprotein(a) in atherosclerotic cardiovascular disease and aortic stenosis: a European atherosclerosis society consensus statement. *Eur Heart J* 2022;**43**:3925–46.
- Salaun E, Mahjoub H, Dahou A, Mathieu P, Larose É, Després JP et al. Hemodynamic deterioration of surgically implanted bioprosthetic aortic valves. *J Am Coll Cardiol* 2018;**72**:241–51.
- Botezatu SB, Tzolos E, Kaiser Y, Cartledge TRG, Kwiecinski J, Barton AK et al. Serum lipoprotein(a) and bioprosthetic aortic valve degeneration. *Eur Heart J Cardiovasc Imaging* 2023;**24**:759–67.
- Farina JM, Chao CJ, Pereyra M, Roarke M, Said EF, Barry T et al. Role of lipoprotein(a) concentrations in bioprosthetic aortic valve degeneration. *Heart* 2024;**110**:299–305.
- Hu X, Wang C, Feng D, Li Z, Chen Y, Niu G et al. Association between lipoprotein(a) and long-term prognosis in patients receiving transcatheter aortic valve replacement. *J Clin Lipidol* 2025;**19**:544–53.
- Capoulade R, Croyal M, Senage T, Guimbretière G, Le May C, Rodrigues Oliveira A et al. Polymorphic size of lipoprotein(a) and structural bioprosthetic valve degeneration. *JACC Cardiovasc Imaging* 2025;**18**:1167–9.
- Malick WA, Goonewardena SN, Koenig W, Rosenson RS. Clinical trial design for lipoprotein(a)-lowering therapies: jacc focus seminar 2/3. *J Am Coll Cardiol* 2023;**81**:1633–45.
- Trinder M, Paruchuri K, Haidermota S, Bernardo R, Zekavat SM, Gilliland T et al. Repeat measures of lipoprotein(a) molar concentration and cardiovascular risk. *J Am Coll Cardiol* 2022;**79**:617–28.
- Ghouse J, Ahlberg G, Albertsen Rand S, Salling Olesen M, Vilhjalmsson B, Stender S et al. Within-person stability of lipoprotein(a) concentration. *Eur Heart J* 2025;**46**:1159–61.
- Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ et al. 2019 acc/aha guideline on the primary prevention of cardiovascular disease: executive

- summary: a report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines. *J Am Coll Cardiol* 2019;**74**:1376–414.
18. Zoghbi WA, Jone PN, Chamsi-Pasha MA, Chen T, Collins KA, Desai MY et al. Guidelines for the evaluation of prosthetic valve function with cardiovascular imaging: a report from the American society of echocardiography developed in collaboration with the society for cardiovascular magnetic resonance and the society of cardiovascular computed tomography. *J Am Soc Echocardiogr* 2024;**37**:2–63.
 19. Hahn RT, Pibarot P. Prosthesis-patient mismatch in transcatheter and surgical aortic valve replacement. *Ann Cardiothorac Surg* 2024;**13**:211–23.
 20. Lorusso R, Gelsomino S, Lucà F, De Cicco G, Billè G, Carella R et al. Type 2 diabetes Mellitus is associated with faster degeneration of bioprosthetic valve: results from a propensity score-matched Italian multicenter study. *Circulation* 2012;**125**:604–14.
 21. Liu F, Cai H. Diabetes and calcific aortic valve disease: implications of glucose-lowering medication as potential therapy. *Front Pharmacol* 2025;**16**:1583267.
 22. Thanassoulis G. Precision medicine for prosthetic valve deterioration. *J Am Coll Cardiol* 2018;**72**:252–4.
 23. Kostyunin AE, Yuzhalin AE, Rezvova MA, Ovcharenko EA, Glushkova TV, Kutikhin AG. Degeneration of bioprosthetic heart valves: update 2020. *J Am Heart Assoc* 2020;**9**:e018506.
 24. Iribarren AC, AlBadri A, Wei J, Nelson MD, Li D, Makkar R et al. Sex differences in aortic stenosis: identification of knowledge gaps for sex-specific personalized medicine. *Am Heart J Plus* 2022;**21**:100197.
 25. Enkhmaa B, Berglund L. Non-genetic influences on lipoprotein(a) concentrations. *Atherosclerosis* 2022;**349**:53–62.
 26. Bhatia HS, Ma GS, Taleb A, Wilkinson M, Kahn AM, Cotter B et al. Trends in testing and prevalence of elevated lp(a) among patients with aortic valve stenosis. *Atherosclerosis* 2022;**349**:144–50.
 27. Tsimikas S, Karwatowska-Prokopczuk E, Gouni-Berthold I, Tardif JC, Baum SJ, Steinhagen-Thiessen E et al. Lipoprotein(a) reduction in persons with cardiovascular disease. *N Engl J Med* 2020;**382**:244–55.
 28. O'Donoghue ML, Rosenson RS, Gencer B, Lopez JAG, Lepor NE, Baum SJ et al. Small interfering rna to reduce lipoprotein(a) in cardiovascular disease. *N Engl J Med* 2022;**387**:1855–64.