



Data Paper

New data on limno-terrestrial rotifers of the world

Dzmitry Lukashanets[‡], Nataliia Iakovenko^{§,¶,¶}, Diego Fontaneto[¶], Miloslav Devetter[¶], Karel Janko^{‡,¶}, Jolanta Ejsmont-Karabin[¶], Irena Bielańska-Grajner[^], Bernard Hallet[¶], Jerzy Smykla[‡], Iryna Kozeretka[¶], Vladlen Trokhymets[¶], Vítězslav Plášek^{¶,¶}, Oleksii Redchenko[‡], Murat Kaya[¶], Yasin Hazer[¶], Mariusz Wierzgoń[^]

[‡] Klaipėda University, Klaipėda, Lithuania

[§] Czech University of Life Sciences Prague, Prague, Czech Republic

[¶] Institute of Animal Physiology and Genetics AS ČR, Libečov, Czech Republic

[¶] University of Ostrava, Ostrava, Czech Republic

[#] Schmalhausen Institute of Zoology of NASU, Kyiv, Ukraine

[¶] National Research Council of Italy, Water Research Institute (CNR-IRSA), Verbania Pallanza, Italy

[¶] Biology Centre, Institute of Soil Biology AS ČR, České Budějovice, Czech Republic

[¶] Nencki Institute of Experimental Biology, Polish Academy of Sciences, Warsaw, Poland

[^] University of Silesia in Katowice, Katowice, Poland

[¶] Catholic University of Louvain, Louvain-la-Neuve, Belgium

[‡] Institute of Nature Conservation, Polish Academy of Sciences, Krakow, Poland

[¶] National Antarctic Scientific Center, Kyiv, Ukraine

[¶] Taras Shevchenko National University of Kyiv, Kyiv, Ukraine

[¶] University of Opole, Institute of Biology, Opole, Poland

[¶] Institute of Molecular Genetics CAS, Prague, Czech Republic

[¶] Istanbul Technical University, Istanbul, Türkiye

[¶] Bülent Ecevit University, Zonguldak, Türkiye

Corresponding author: Dzmitry Lukashanets (dzmitry.lukashanets@ku.lt)

Academic editor: Lyudmila Kamburska

Received: 12 Sep 2025 | Accepted: 22 Oct 2025 | Published: 30 Dec 2025

Citation: Lukashanets D, Iakovenko N, Fontaneto D, Devetter M, Janko K, Ejsmont-Karabin J, Bielańska-Grajner I, Hallet B, Smykla J, Kozeretka I, Trokhymets V, Plášek V, Redchenko O, Kaya M, Hazer Y, Wierzgoń M (2025) New data on limno-terrestrial rotifers of the world. Biodiversity Data Journal 13: e171929. <https://doi.org/10.3897/BDJ.13.e171929>

Abstract

Background

Limno-terrestrial rotifers, particularly those of the order Bdelloidea, inhabit bryophytes, lichens, soils and other periodically moist terrestrial habitats. Despite their high

abundance in all latitudinal zones, these rotifers remain poorly documented in biodiversity databases due to difficulties in preservation and morphological identification. As a result, the current knowledge of their global distribution is still highly fragmented and geographically biased, with the majority of species records concentrated in Europe (where the experts mostly collected material). Many regions like north-eastern Asia, North and South America, Africa and Australia remain under-represented in current knowledge on distribution of limno-terrestrial rotifers. Recent studies suggested that Bdelloidea exhibit distinct biogeographical patterns and potentially high levels of cryptic diversity and endemism, challenging the traditional view of the omnipresence of all microscopic taxa. Comprehensive, georeferenced occurrence data are essential to advance our understanding of bdelloid biodiversity and distribution, yet such data are still scarce in global platforms like the Global Biodiversity Information Facility (GBIF).

New information

The dataset provides new, georeferenced data on the occurrence of rotifer species inhabiting limno-terrestrial habitats worldwide. For the first time, occurrence records for 48 rotifer species (including nominal taxonomic 'subspecies') are published in GBIF.

In particular, the dataset significantly expands the information on the distribution ranges of the bdelloid rotifers (order Bdelloidea) in GBIF. We contributed 5,651 new occurrence records of bdelloids rotifers, increasing the number of records in GBIF by 47.5% (or by 61.6% when considering only previously georeferenced records).

Moreover, we add 394 new records to the faunas of 49 of 56 studied countries, in 17 of which limno-terrestrial rotifers were studied for the first time.

Additionally, for 19 countries, records of bdelloid rotifers are now available in GBIF for the first time.

Keywords

Bdelloidea, Rotifera, biogeography, bryophytes, lichens, soil, occurrence

Introduction

Limno-terrestrial microfauna represents a distinct ecological group of microscopic invertebrates inhabiting soil, litter, lichens, bryophytes and other environments that regularly undergo alternating periods of drying and humidity. This group includes various animal taxa such as rotifers (Rotifera), tardigrades (Tardigrada), nematodes (Nematoda) and others.

Within limno-terrestrial rotifers, bdelloids (Rotifera, Bdelloidea) are one of the most abundant and widely distributed groups. These obligate parthenogenetic organisms can

survive under extreme environmental conditions (complete freezing, desiccation etc.) by entering a state of cryptobiosis – a temporary pause of metabolism (Ricci and Caprioli 2005, Shmakova et al. 2021). They also possess remarkable capabilities for passive dispersal, transporting via wind, water and other vectors (Fontaneto 2019). As a result, bdelloids inhabit a broad range of limno-terrestrial habitats and are distributed globally (Ricci 1987, Fontaneto et al. 2006, Fontaneto et al. 2007, Fontaneto et al. 2008). In certain environments, particularly those lacking larger invertebrates, bdelloids dominate in microinvertebrate communities, reaching large densities and playing a key role in ecosystem processes (Zawierucha et al. 2021).

However, bdelloid rotifers remain a poorly-studied taxon in terms of both biology and distribution. These animals cannot be preserved using traditional fixative chemicals, they are difficult to identify morphologically and are often overlooked in studies of biodiversity, soil biology etc.

Our current knowledge of bdelloid fauna and distribution remains significantly biased in geographical aspect. Most of original descriptions of bdelloids originate from Central and Western Europe, where pioneer studies were conducted (Donner 1965). The majority of new European bdelloid species descriptions comes from Austria (Donner 1949, Donner 1950a, Donner 1950b, Donner 1952), former Czechoslovakia (Bartoš 1951, Bartoš 1959), Germany (Janson 1893, Wulfert 1942, Wulfert 1960), the Netherlands (De Koning 1947), Romania (Rudescu 1960), Sweden (Bērziņš 1982) and UK (Davis 1873, Murray 1905, Murray 1906a, Bryce 1915). Although there were surveys of other regions, including Africa (Murray 1908, Murray 1911e, Murray 1911f, Milne 1916, Koste 1996), Asia (Murray 1906b, Hauer 1938, Bartoš 1963, Sudzuki 1981, Koste and Zhuge 1996), North America (Murray 1911b, Harring and Myers 1922, Burger 1948, Bateman and Davis 1980), South America (Murray 1907, Murray 1913, Donner 1980, Koste and Böttger 1989), Australia, New Zealand and Pacific Islands (Murray 1911a, Murray 1911c, Murray 1911d, Haigh 1963, Haigh 1970, Ricci et al. 2003), Antarctica (Murray 1910, Dartnall and Hollowday 1985), the majority of the first descriptions still arises from Europe. In recent decades, research efforts have expanded into previously poorly-studied areas of Asia (Song and Min 2015, Song and Lee 2020, Zeng et al. 2020, Song and Lee 2022, Wang et al. 2023, Song and Lee 2024, Wang et al. 2025), including tropical regions (Fontaneto and Ricci 2004, Jaturapruerk and Maiphae 2025, Pokpongmongkol et al. 2025) and North America (Örstan 2018, Örstan 2021). Extensive research efforts were also recently made in Central and Eastern Europe (Kutikova 2005, Devetter 2007, Devetter 2009, Devetter 2010, Bielańska-Grajner et al. 2011, Iakovenko and Ovander 2011, Bielańska-Grajner et al. 2013, Lukashanets 2018).

The most complete identification key for bdelloid species is still that of Donner (1965), published in German and lacking more than 40 taxa described since (Örstan and Plewka 2017). Several more recent keys (e.g. Ricci and Melone (2000), Kutikova (2005), Bielańska-Grajner et al. (2013), Iakovenko et al. (2013)) are limited either to certain taxonomical ranks and groups or to regions (Poland, Russian Federation).

Many countries still have not been studied for their bdelloid fauna and taxonomy. For example, in the Arctic, bdelloid checklists are known mainly from Svalbard (Kaya et al. 2010, Coulson et al. 2024) with very little data from Greenland, Iceland, Russia and USA (Alaska) (De Ridder 1972). The main bulk of material in numerous surveys of Antarctica comes from Maritime Antarctic and Victoria Land (Iakovenko et al. 2015, Fontaneto et al. 2015), whilst data for East Antarctica are scarce (Lukashanets et al. 2021, Lukashanets et al. 2022). In Europe, the best studied continent in relation to bdelloid distribution, mostly fragmentary knowledge comes from Baltic States (Cimdiņš 1979, Zarubov and Cimdiņš 1990) and Scandinavia (Bērziņš 1982, Bērziņš 1987). Although the works of Fontaneto et al. (2006) and Fontaneto et al. (2007) greatly expanded the knowledge on mainland Italy, bdelloid fauna of Balkan countries, France, Portugal, Spain and Mediterranean Islands remains unknown for the most part. Other continents, especially central and north-eastern parts of Asia along with Africa and South America, still require extensive taxonomic and faunistic work.

Existing online repositories do not yet provide a comprehensive snapshot of bdelloid distribution. For example, as of July 2025, the Global Biodiversity Information Facility (GBIF) contains 11,894 occurrence records of bdelloids, from both aquatic and terrestrial ecosystems. From them, 2,590 (22%) are identified only to family or genus level and other 2,857 (or 24%) are noted as 'Bdelloidea sp.'. Some current identifications of bdelloids in GBIF (according to the provided photos) have taxa misidentified not only to species, but even to genus and family level (personal observations).

Research over the past 20 years revealed specific biogeographical patterns in bdelloids (Fontaneto et al. 2008, Velasco-Castrillón et al. 2014, Iakovenko et al. 2015), supporting the concept that microscopic animals do have biogeography (Martínez et al. 2025). Furthermore, it has been claimed that bdelloid diversity is significantly underestimated due to cryptic species and high regional endemism (Velasco-Castrillón et al. 2014, Iakovenko et al. 2015). The most recent published annotated checklist includes more than 460 species (Segers 2007), but the actual number is believed to be much higher (Fontaneto et al. 2010).

All this highlights the strong need to collect new data, especially from underexplored regions, to fill gaps in our understanding of limno-terrestrial rotifer diversity.

In this work, we provide a dataset of limno-terrestrial rotifers from diverse regions across the globe, including all continents, several islands and archipelagoes and both polar areas. For our survey, samples were collected from a wide range of non-aquatic, but periodically moistened habitats, such as bryophytes (mosses and liverworts), lichens, algal mats, soil, needle, leaf or mixed litter. All rotifers (not only bdelloids, although they made up the majority of records) were identified to the lowest possible taxonomic rank.

The dataset has been submitted to GBIF with complete accompanying metadata. This includes detailed information on the occurrence sites: habitat (the substrate from which rotifers were isolated), biotope (the environment where the substrate was collected),

locality (country and continent), elevation and exact coordinates. For each species, the density in the sample is provided.

The primary goal of this paper is to document and publish all verified global records of limno-terrestrial rotifers in GBIF. These data will also support future research on the local, regional and global distribution patterns of limno-terrestrial rotifers and will enable more in-depth taxonomic studies of this still poorly-understood taxon.

Project description

Title: ROTISFERA 'Global patterns of microinvertebrate distribution: does diversity decrease poleward in rotifers (Rotifera, Bdelloidea)?' NDICI-GEO-NEAR/2022/434-092-0068

Personnel: Dzmitry Lukashanets, Nataliia Iakovenko

Study area description: The project encompasses all major regions of the Earth.

Design description: To address the project's aims, we used samples collected starting from 1996 until the present in limno-terrestrial habitats throughout the world. More than 2,000 samples were analysed using conventional methods (analysis of rotifer morphology using light microscopy) resulting in creation of dataset with 5,729 rotifer species occurrences, 98.5% of which belong to bdelloids. The data cover all continents, including both polar zones and ranging from 0 to 4,610 m of altitude.

The following steps include estimating diversity patterns within defined geographical areas (alpha- and gamma-diversity, rarefaction curves, asymptotic diversity estimates) and globally – revealing the shape and prominence of the latitudinal diversity gradient using a hexagonal model and generalised additive model (GAM).

In parallel, the diversity of limno-terrestrial rotifers was assessed using a molecular approach. We applied metabarcoding of the COI mitochondrial gene for rotifers; however, this was not conducted at a global scale, but rather limited to Europe (from the Mediterranean to the Arctic). The results obtained from both methodological approaches were compared.

Funding: European Union

Sampling methods

Description: All samples used in the dataset were collected either within the framework of several research projects focused on the biodiversity and ecology of microfauna, or as a part of other field activities. The latter included fieldwork in remote and hard-to-reach areas (tropical rainforests, polar and alpine regions), incidentally in field trips, within national Antarctic expeditions, during landing at cruises of research vessels etc. The samples have been stored in the repositories of the dataset's main creators, Nataliia

Iakovenko, Diego Fontaneto and Dzmitry Lukashanets. For the short-term storage, we used gradual drying and stored samples at room temperature or at +8°C, while, for long-term preservation freezing at -20–25°C was used (samples from other than polar latitudes were first gradually dried at room temperature before freezing).

Sampling description: For the analysis of limno-terrestrial rotifers, sampling was performed mainly in forest ecosystems (taiga, boreal and mixed forests, subtropical forests, tropical rainforests), flatlands, urban areas and agricultural landscapes. In polar areas, we conducted sampling in moss tundra, at post-glacial moraine, in Antarctic oases and nunataks.

Samples of soil and litter were collected using 3.8 cm metal corer or metal scoops. Lichens, bryophytes and other vegetation were picked from the ground, tree barks and stones using tweezers or simply by hands (Fig. 2). The size of the samples varied from a few cm³ to several dozen cm³, depending on the amount of material available for collection. All samples were placed in paper or plastic bags and properly labelled. All sampling sites were georeferenced.




Figure 1. [doi](#)

All sampling sites included in the dataset.

Quality control: Specialists in taxonomy of Bdelloidea, Nataliia Iakovenko, Diego Fontaneto and Dzmitry Lukashanets, carried out all species identification while consulting other specialists (Aydin Örstan, Claudia Ricci). Primarily first descriptions together with the existing identification keys and reviews on the taxonomy of bdelloids were used. Georeferenced data were checked by placing latitudes and longitudes on a map.

Step description: Sample processing included the following steps:

(i) Microscopic animals were isolated from each sample. To do this, bryophytes, lichens, soil, litter, timber, fungi or pieces of vegetation were washed through a system of sieves, followed by flotation and centrifugation in a sugar solution (Freckman and Virginia 1993). For some samples, the method for extracting moss-dwelling rotifers described by Peters

et al. (1993) was used. For other samples, the substrates were washed repeatedly in distilled water.




Figure 2. [doi](#)

Examples of limno-terrestrial habitats: **A** moss on tree bark; **B** lichens on tree bark; **C** moss on ground; **D** leaf litter.

(ii) Rotifers were counted and sorted under a binocular microscope (Olympus SZ61, Olympus SZX10, NR.3 Nikon SMZ1000) at magnification ranging between 10x and 100x.

(iii) For species identification, live rotifer individuals were transferred to slides and examined using light microscopy (Nikon Eclipse Ts2R, NIB-100F inverted microscope, Olympus CX43) at magnifications ranging from 400x to 1000x. First descriptions of taxa and the existing identification keys were used: Bartoš (1951), Bartoš (1959), Donner (1965), Kutikova (2005). Body dimensions and proportions were measured following the protocol of Iakovenko et al. (2013) and Iakovenko et al. (2015).

(iv) Rotifer densities were equalised to 25 cm³ (measured as 25 ml standard sample of the dried substrate using a plastic tube). In case when the sample volume was smaller, it was measured and rotifer densities were counted per sample and then recalculated to 25 cm³. The standard sample volume was chosen empirically as a minimal volume to be considered as a feasible and adequate representation of rotifer biodiversity in a sample.

Geographic coverage

Description: The dataset includes records collected throughout the world, spanning latitudes from 77.9° S to 78.2° N and covering all continents (Fig. 1):

- (i) Europe – Austria, Belarus, Belgium, Bulgaria, Croatia, Czech Republic, Denmark, Estonia, Finland, France, Germany, Great Britain and Northern Ireland, Greece, Hungary, Italy, Latvia, Lithuania, Montenegro, Netherlands, North Macedonia, Norway, Poland, Portugal, Russian Federation (Caucasus and Karelia), Serbia, Slovakia, Spain, Sweden, Switzerland, Ukraine, also including the European Arctic (Iceland, Svalbard and Jan Mayen) and minor Mediterranean islands (Mallorca, Menorca, Malta, Pantelleria, Sardinia);
- (ii) Asia – Armenia, Brunei, China, Cyprus, Georgia, India, Japan, Kyrgyzstan, Malaysia, Myanmar, Nepal, Russian Federation (southern Siberia), Taiwan, Tajikistan, Turkey;
- (iii) Africa – Democratic Republic of Congo, Lesotho, Tunisia, also including Macaronesia (Madeira, Canary Islands);
- (iv) North America – western Greenland, USA (Alaska, Maryland and Virginia), also including the Caribbean (Martinique);
- (v) South America – Brazil, Chile;
- (vi) Australia and Oceania – Australia (mainland and Tasmania), New Zealand, Papua New Guinea;
- (vii) Antarctica – Maritime Antarctic (Antarctic Peninsula, Argentine Islands, James Ross Island, King George Island) and Continental Antarctica (Dronning Maud Land, Enderby Land, Victoria Land and Ross Sea Region).

Taxonomic coverage

Description: The dataset includes all records belonging to Phylum Rotifera, which we found in the samples of soil, bryophytes, lichens and other limno-terrestrial habitats. Monogonont rotifers (order Ploima) accounted for only 1.36% the records, whilst the majority of the records were bdelloids (order Bdelloidea). Species potentially new for science were indicated as *Genus* sp. or – in case of minor differences from the nominal description – as cf. It is noteworthy that 10 species amongst the ones that had been found were not typical for limno-terrestrial habitats, but rather occur in swamps and temporary water bodies (f. ex. *Philodina flaviceps*, *Synchaeta* sp.). Their encounter on terrestrial vegetation is most likely the result of the proximity to the waterbodies and/or high humidity of the habitat.

Taxa included:

Rank	Scientific Name	Common Name
phylum	Rotifera	rotifers

Temporal coverage

Notes: 1996–2025

Usage licence

Usage licence: Creative Commons Public Domain Waiver (CC-Zero)

Data resources

Data package title: New data on limno-terrestrial rotifers of the world

Resource link: <https://doi.org/10.15468/vymt96>

Alternative identifiers: https://cloud.gbif.org/eca/resource?r=limno-terrestrial_rotifera

Number of data sets: 1

Data set name: New data on limno-terrestrial rotifers of the world

Data format: Darwin Core Archive

Description: The dataset brings together all records of rotifers (Phylum Rotifera) found by N. Iakovenko, D. Lukashanets and D. Fontaneto worldwide in limno-terrestrial habitats – substrates that periodically undergo cycles of desiccation and hydration, but cannot be classified as temporary waterbodies, such as bryophytes, lichens, fungi, timber, upper layers of soil and litter (Lukashanets et al. 2025).

From 1996 to the present, we have collected and analysed more than 2,275 limno-terrestrial samples. Sampling covered all continents and over 50 countries, spanning from the Arctic to Antarctica, including most major biomes. The altitude of sampling sites ranged from 0 to 4,610 m above sea level. In total, the dataset includes 5,729 records of limno-terrestrial rotifers. We found 216 morphological species (including nominal ‘subspecies’ that, in bdelloids, are to be updated to species in future) belonging to 23 genera (nine of Monogononta and 14 of Bdelloidea). More than other 400 species-level entities were different in minor or major details from the original descriptions and potentially might be new taxa.

The submitted dataset is extensive and substantially contributes to the current knowledge of diversity and distribution of limno-terrestrial rotifers, which are still poorly known, especially outside Europe. We present it for the wider use in studies of microinvertebrate biodiversity and macroecology.

Column label	Column description
occurrenceID	An unique identifier for the record including the code of the country, year of sampling, ID of the sample and species name.

materialSampleID	An identifier of the sample, from which rotifers were isolated.
eventDate	Year of sampling.
decimalLatitude	Latitude in decimal degrees.
decimalLongitude	Longitude in decimal degrees.
verbatimElevation	Elevation above sea level in metres.
continent	The name of the continent where sampling occurred.
country	The name of the country where sampling occurred.
countryCode	ISO 3166-1 alpha-2 country code.
stateProvince	The name of the next smaller administrative region than country, where sampling occurred (used only for USA).
island	The name of the island where sampling occurred (if country includes different islands or mainland and islands).
islandGroup	The name of the archipelago where sampling occurred (if country includes different islands or mainland and islands).
locationRemarks	The name of biotope where sampling occurred.
habitat	A description of the type of the sample, from which rotifers were isolated.
basisOfRecord	The specific nature of the data record.
scientificName	The full scientific name, with authorship and date information.
individualCount	The number of individuals present at the time of the Occurrence.
identifiedBy	Name and surname of the person who identified the species.
identifiedByID	ORCID of the person who identified the species.
sampleSizeValue	A numeric value for a measurement of the size of the sample.
sampleSizeUnit	The unit of measurement of the size of the sample.
kingdom	The full scientific name of the kingdom in which the species is classified.
phylum	The full scientific name of the phylum in which the species is classified.
class	The full scientific name of the class in which the species is classified.
order	The full scientific name of the order in which the species is classified.
family	The full scientific name of the family in which the species is classified.
genus	The full scientific name of the genus in which the species is classified.
specificEpithet	The name of the species epithet of the scientificName.
infraspecificEpithet	The name of the infraspecies epithet of the scientificName.
taxon rank	The taxonomic rank of the record.

identificationQualifier	A controlled value to express the identifier's doubts about identification.
scientificNameAuthorship	The authorship information for the scientificName.
recordedBy	Name and surname of the person who collected the sample, from which rotifers were isolated.

Additional information

Limno-terrestrial rotifers: overview of taxonomy

The dataset includes a total of more than 600 taxa identified to species and below species level. The majority of the records belong to order Bdelloidea (representing over 98% of all species). Within Bdelloidea, the most diverse families are Habrotrichidae (particularly the genus *Habrotricha* with 264 species) and Philodinidae, notably the genera *Macrotrachela* (178), *Mniobia* (81) and *Philodina* (54 species). The genus *Adineta* (family Adinetidae) is represented with 43 species (Table 1).

Table 1.

Taxonomic ranks of rotifers (with the number of species) recorded in limno-terrestrial habitats.

Class	Order	Family	Genus	Number of species (including subspecies and species-level entities different from nominal description)
Eurotatoria	Bdelloidea	Adinetidae	<i>Adineta</i>	43
			<i>Bradyscela</i>	2
		Habrotrichidae	<i>Habrotricha</i>	264
			<i>Otostephanos</i>	11
			<i>Scepanotrocha</i>	11
		Philodinavidae	<i>Philodinavus</i> *	1
		Philodinidae	<i>Ceratotrocha</i>	6
			<i>Didymodactylos</i>	1
			<i>Dissotrocha</i>	2
			<i>Macrotrachela</i>	178
			<i>Mniobia</i>	81
			<i>Philodina</i>	54
			<i>Pleuretra</i>	12
			<i>Rotaria</i>	8
	Ploima	Dicranophoridae	<i>Encentrum</i>	4
			<i>Wierzejskiella</i>	1
		Lecanidae	<i>Lecane</i>	5
		Lepadellidae	<i>Colurella</i> *	1

Class	Order	Family	Genus	Number of species (including subspecies and species-level entities different from nominal description)
			<i>Lepadella</i>	2
			<i>Squatinella</i>	1
		Lindiidae	<i>Lindia</i> *	1
		Notommatidae	<i>Cephalodella</i>	3
		Synchaetidae	<i>Synchaeta</i> *	1

* only species different from nominal descriptions were recorded.

The dataset additionally includes records from the order Ploima, comprising 19 species in total. Amongst these, the most represented genera are *Lecane* (Lecanidae), *Encentrum* (Dicranophoridae) and *Cephalodella* (Notommatidae), along with several other genera with 1–2 species each (Table 1).

Overall, the taxonomic scope of the dataset reflects a comprehensive sampling of limno-terrestrial rotifers, with a focus on bdelloid taxa, which are typically dominant in these habitats.

The number of species recorded in different types of habitat varied from 205 (mosses) to seven (fungi); however, some bias related to different number of samples collected per habitat type is possible (Table 2).

Table 2.

Number of recorded taxa (families, genera, species) per type of habitat.

Habitat	Number of families	Number of genera	Number of species (including subspecies)
Bryophytes (excluding Sphagnum mosses)	7	19	205
Lichens	6	13	80
Sphagnum mosses	9	17	40
Algae / algal mats	5	8	33
Fungi	2	3	7
Grass	3	6	31
Leaf litter	6	13	67
Needle litter	5	13	63
Mixed litter	4	9	60
Soil	6	14	76
Sand / stones	3	7	26

New records

Our dataset contains 394 new records for 49 of 56 studied countries, in 17 of which limno-terrestrial rotifers were studied for the first time. These data are summarised in Table 3.

Table 3.

Total and new records of limno-terrestrial rotifers in the database.

Country/ Continent	Number of records	Number of species found (including subspecies and species designated as cf. or Genus sp.)	Species new for the country/continent
Antarctica	850	44	Data published
Armenia*	4	3	All new for the country
Australia (mainland)	6	6	–
Australia (Tasmania)	12	6	<i>Habrotrocha angusticollis angusticollis</i> , <i>H. aspera</i> , <i>H. placida</i>
Austria	13	8	–
Belarus	236	58	Data published
Belgium	17	10	<i>Macrotrachela ehrenbergii</i> , <i>M. nana nana</i> , <i>M. plicata plicata</i> , <i>Mniobia russeola</i> , <i>Mn. tetraodon</i> , <i>Philodina plena</i>
Brazil	1	1	–
Brunei*	7	5	All new for the country
Bulgaria*	66	29	<i>Adineta barbata</i> , <i>Bradyscela clauda</i> , <i>B. granulosa</i> , <i>Habrotrocha flaviformis</i> , <i>H. gracilis gracilis</i> , <i>H. mediocris</i> , <i>H. pavida</i> , <i>H. puella puella</i> , <i>H. pusilla pusilla</i> , <i>Macrotrachela concinna</i> , <i>M. ehrenbergii</i> , <i>M. libera libera</i> , <i>M. multispinosa multispinosa</i> , <i>M. papillosa</i> , <i>M. plicata hirundinella</i> , <i>M. punctata</i> , <i>M. quadricornifera quadricornifera</i> , <i>M. quadricornifera quadricorniferoides</i> , <i>Mniobia bredensis</i> , <i>Mn. russeola</i> , <i>Mn. scabrosa</i> , <i>Philodina plena</i> , <i>Ph. vorax</i> , <i>Pleuretra brycei</i> , <i>Rotaria sordida sordida</i>
Chile	81	25	<i>Mniobia scabrosa</i>
China	19	12	Data published
Croatia	6	6	<i>Didymodactylos carnosus</i> , <i>Macrotrachela quadricornifera quadricornifera</i> , <i>M. timida timida</i> , <i>Pleuretra lineata</i>
Cyprus*	34	17	All new for the country
Czech Republic	107	45	–
Congo*	15	10	All new for the country
Denmark (mainland)*	7	7	All new for the country

Country/ Continent	Number of records	Number of species found (including subspecies and species designated as cf. or Genus sp.)	Species new for the country/continent
Denmark (Greenland)	90	34	All new for the country
Estonia*	93	29	All new for the country
Finland	18	13	<i>Habrotrocha constricta</i> , <i>H. pavida</i> , <i>H. puella puella</i> , <i>Macrotrachela concinna</i> , <i>M. ehrenbergii</i> , <i>M. habita</i> , <i>M.</i> <i>multispinosa multispinosa</i> , <i>Mniobia russeola</i> , <i>Philodina plena</i> , <i>Rotaria sordida sordida</i>
France (mainland)	157	37	<i>Adineta steineri</i> , <i>A. vaga minor</i> , <i>Habrotrocha elegans</i> , <i>H.</i> <i>minuta</i> , <i>Macrotrachela nana nana</i> , <i>M. timida timida</i> , <i>Mniobia</i> <i>tentans</i>
France (Martinique)*	10	5	All new for the region
Georgia*	3	3	All new for the country
Germany	72	31	<i>Habrotrocha minima</i> , <i>H. novemdens</i> , <i>H. pulchra</i> , <i>H. pusilla</i> <i>pusilla</i>
Greece	2	2	<i>Macrotrachela timida timida</i>
Hungary	98	29	<i>Habrotrocha gracilis gracilis</i> , <i>H. pavida</i> , <i>H. puella puella</i> , <i>H.</i> <i>sylvestris</i> , <i>Mniobia edmondsoni</i> , <i>Philodina proterva</i>
Iceland	50	20	<i>Didymodactylos carnosus</i> , <i>Macrotrachela quadricornifera</i> <i>quadricorniferoides</i> , <i>M. quadricornifera scutellata</i> , <i>Mniobia</i> <i>magna</i> , <i>Mn. scarlatina</i>
India	2	2	–
Italy (mainland)	276	69	Data published
Italy (Mediterranean islands)	15	7	All new for the region
Japan	3	3	<i>Habrotrocha angusticollis reversa</i> , <i>Habrotrocha pusilla nuda</i>
Kyrgyzstan*	12	8	All new for the country
Latvia	94	34	<i>Adineta steineri</i> , <i>Ceratotrocha franzi</i> , <i>Didymodactylos carnosus</i> <i>Habrotrocha bidens</i> , <i>H. elusa vegeta</i> , <i>H. fusca</i> , <i>H.</i> <i>fuscochlaena</i> , <i>H. gracilis gracilis</i> , <i>H. rosa</i> , <i>H. tridens</i> , <i>Macrotrachela aculeata</i> , <i>M. habita</i> , <i>M. libera</i> , <i>M. multispinosa</i> <i>multispinosa</i> , <i>M. papillosa</i> , <i>M. plicata hirundinella</i> , <i>M.</i> <i>quadricornifera quadricornifera</i> , <i>Mniobia magna</i> , <i>Mn. montium</i> , <i>Mn. russeola</i> , <i>Mn. scarlatina</i> , <i>Philodina plena</i> , <i>Ph. vorax</i> , <i>Rotaria sordida</i> , <i>Scepanotrocha rubra</i>
Lesotho*	50	13	All new for the country
Lithuania	11	9	<i>Adineta steineri</i> , <i>Rotaria sordida sordida</i> , <i>R. sordida fimbriata</i>
Malaysia	9	5	All new for the country

Country/ Continent	Number of records	Number of species found (including subspecies and species designated as cf. or Genus sp.)	Species new for the country/continent
Malta*	3	3	All new for the country
Montenegro*	42	17	All new for the country
Myanmar	3	3	<i>Macrotrachela habita</i>
Nepal*	16	5	All new for the country
Netherlands	19	15	<i>Adineta steineri</i> , <i>H. gracilis gracilis</i> , <i>H. tripus</i> , <i>Macrotrachela quadricornifera scutellata</i> , <i>Mniobia tentans</i> , <i>Philodina intermedia</i>
New Zealand	38	11	<i>Ceratotrocha velata</i> , <i>Habrotrocha visa</i> , <i>Macrotrachela bilfingeri</i> , <i>M. multispinosa multispinosa</i>
North Macedonia*	24	13	All new for the country
Norway (mainland)*	11	9	All new for the country
Norway (Jan Mayen and Bear Island)*	19	13	All new for the region
Norway (Svalbard)	202	45	<i>Adineta elongata</i> , <i>Ceratotrocha velata</i> , <i>Habrotrocha pavidata</i> , <i>H. pusilla pusilla</i> , <i>Macrotrachela bilfingeri</i> , <i>M. crucicornis</i> , <i>M. decora</i> , <i>M. induta</i> , <i>M. intermedia</i> , <i>M. oblita</i> , <i>M. timida</i> , <i>Mniobia scarlatina</i>
Papua New Guinea*	2	2	All new for the country
Poland	518	86	Data published
Portugal (mainland)*	11	5	<i>Pleuretra lineata</i>
Portugal (Madeira)	23	15	—
Russian Federation	16	9	—
Serbia*	47	20	All new for the country
Slovakia	7	7	<i>Habrotrocha pusilla pusilla</i> , <i>Mniobia tentans</i>
Spain (mainland)	2	2	—
Spain (Mediterranean islands)	23	14	—
Spain (Canary islands)	2	2	<i>Habrotrocha gracilis gracilis</i>

Country/ Continent	Number of records	Number of species found (including subspecies and species designated as cf. or Genus sp.)	Species new for the country/continent
Sweden	60	33	<i>Habrotrocha flaviformis</i> , <i>H. minima</i> , <i>H. pavida</i> , <i>H. tripus</i> , <i>Macrotrachela insolita</i> , <i>M. plicata hirundinella</i> , <i>M.</i> <i>quadricornifera scutellata</i> , <i>Mniobia loxocorona</i> , <i>Scepanotrocha</i> <i>simplex</i>
Switzerland	16	12	<i>Habrotrocha minima</i> , <i>Macrotrachela libera libera</i> , <i>Pleuretra</i> <i>lineata</i>
Taiwan	2	1	–
Tajikistan*	7	5	All new for the country
Tunisia*	2	2	<i>Rotaria sordida sordida</i>
Turkey	77	32	<i>Habrotrocha flaviformis</i> , <i>H. pavida</i> , <i>H. puella puella</i> , <i>H. pulchra</i> <i>H. serpens</i> , <i>Macrotrachela inermis</i> , <i>M. multispinosa</i> <i>multispinosa</i>
Ukraine	1930	139	<i>Habrotrocha humilis</i>
UK (England)	41	27	<i>Habrotrocha minima</i> , <i>H. puella puella</i> , <i>Macrotrachela inermis</i> , <i>M. libera</i> , <i>M. minuta minuta</i>
UK (Northern Ireland)	7	6	<i>Habrotrocha elegans</i> , <i>H. filum</i>
USA	14	11	–

* the country or region was studied for the first time.

However, these new records include only species that are identified as corresponding to the existing descriptions. At the same time, more than 400 morphologically distinctive species in this database either had minor differences from these descriptions (designated as cf.) or exhibited new features (designated as Genus sp.) and, thus, might potentially be species new for science. The highest number of such entities were found in the least studied regions (Arctic, Antarctica, South America, Africa). This calls for increasing the efforts for not only faunistic surveys, but primarily for the thorough taxonomical work aiming at describing new species characteristic for such regions, combining both morphological and morphometrical analysis with DNA barcoding.

Acknowledgements

The dataset was created under the ROTISFERA project: ‘Global patterns of microinvertebrate distribution: does diversity decrease poleward in rotifers (Rotifera, Bdelloidea)?’ NDICI-GEO-NEAR/2022/434-092-0068 (funded by EU).

Sampling activities in some regions and analyses of the samples were implemented as part of the following projects:

SNOWLIFE 'Microfauna emerging in seasonal snow patch ecosystems – losers or winners?' P-MIP-23-374 (Europe – Baltic States);

'Taksonomia i ekologia wrotków (Rotifera) z rzędu pijawczaków (Bdelloidea) na terenie Polski' N304 0341 33 (Europe – Poland);

INTAS Young Scientist Fellowship 'Order Adinetida, Bdelloidea, Rotifera. An analysis of diversity and distribution of the taxon in Central and Southern Europe' 06-1000014-5639 (Central and Southern Europe);

Polish Ministry of Science and Higher Education within the programme Supporting International Mobility of Scientists and within grants 2P04F00127, NN304069033, NN305376438 and NN304014939 for years 2002–2015 and National Science Foundation within US Antarctic Program grants NSF BO-034-M and ANT 0739575 (Antarctica – Victoria Land);

Ukrainian State Special-Purpose Research Program in Antarctica (Maritime Antarctic).

'Unique community of ice-bound tardigrades and rotifers in evolutionary, physiological and ecological context' CSF 22-28778S (all regions);

Institute of Animal Physiology and Genetics Institutional Support RVO 67985904 (all regions).

Collection of data from western Greenland and Iceland was made possible through the participation of D. Lukashanets as a visiting scientist in the cruise 'Iceland and Greenland: edge of the Arctic' on board the National Geographic Explorer vessel (organised by Lindblad Expeditions and National Geographic, August 2022).

The authors gratefully acknowledge Prof. Claudia Ricci and Prof. Giulio Melone for the guidance, suggestions and help in initial mastering of the rotifer identification.

The authors would like to thank all other people who collected samples – Igor Balashov, Olena Bezukrovna, Jörg Bohlen, Oleg Borodin, Tatiana Chernyachovskaya, Lukáš Choleva, Maya Chumak, Zdenek Duriš, Igor Dykyy, Sevil Erdogan, Yury Giginiak, Olexiy Gumovsky, Mykhail Iakovenko, Vira Inshina, Mykola Ivanov, Jan Kavan, Šárka Kinzelová, Jan Kočí, Alexei Korniushev, Liudmila Kovalchuk, Anna Kravchenko, Olga Kuzmanenko, Tetyana Kuzmina, Vladislav Miamin, Branko Micevski, Krystína Muchová, Svitlana Nyporko, Aydin Örsan, Milan Paunovich, Xenya Pichko, Alexei Pindrus, Dmitry Polin, Klara Řeháková, Ekaterina Romanenko, Otto Strunecký, Anna Šobanová, Marina Tarashuk, Oxana Tishchenko, Robert Vargovich and Alena Zikmundová.

Author contributions

DL prepared dataset for publishing in GBIF and wrote the dataset description and the manuscript. NI participated in writing the manuscript, performed analysis of the new species records in different countries, made necessary corrections in the list of scientific

names and generally supervised the creation of the dataset. All species identifications have been performed by NI, DF and DL, with some consultations made by MD. Additionally, these authors conducted sampling: NI – mainly Ukraine, Southern and Central Europe; DF – mainly Italy, France and Switzerland; DL – Belarus, Baltic States, Iceland, Greenland, Svalbard and Antarctica. MD carried out sampling and organised fieldwork in Svalbard. JEK and IBG organised sampling in Poland and participated in collection, processing of samples and rotifer identification. All other authors provided suggestions for manuscript preparation and samples from remote areas and are listed in order of their contribution.

References

- Bartoš E (1951) The Czechoslovak Rotatoria of the order Bdelloidea. Vestník Československe Zoologicke Spolecnosti 15: 241-500.
- Bartoš E (1959) Virnici – Rotatoria. Vol. 15. Fauna CSR. 15. Fauna CSR. Nakladatelství Československe Akademie Ved, Prague. [In Czech].
- Bartoš E (1963) Die Bdelloideen der Moosproben aus China und Java. Věstník Československé Zoologické Společnosti 27: 31-42.
- Bateman LE, Davis CC (1980) The Rotifera of Hammock-Hollow formations in a poor (mesotrophic) fen in Newfoundland. Internationale Revue der gesamten Hydrobiologie und Hydrographie 127-153. <https://doi.org/10.1002/iroh.19800650108>
- Bērziņš B (1982) Die Unterarten von *Dissotrocha aculeata*. Limnologiska institutionen, Lunds universitet.
- Bērziņš B (1987) Rotifer occurrence in relation to pH. Hydrobiologia 147: 107-116. <https://doi.org/10.1007/BF00025733>
- Bielańska-Grajner I, Ejsmont-Karabin J, Yakovenko N (2011) Bdelloidea (Rotifera) from the Tatra National Park, with some species newly found in Poland. Fragmenta Faunistica 54: 103-111. <https://doi.org/10.3161/00159301FF2011.54.2.103>
- Bielańska-Grajner I, Ejsmont-Karabin J, Yakovenko N (2013) Wrotki (Rotifera, Bdelloidea). Fauna Sładowodna Polski. Wydawnictwo Uniwersytetu Łódzkiego.
- Bryce D (1915) On five new species of genus *Habrotrocha*. Journal of the Quekett Microscopical Club 2: 631-642. <https://doi.org/10.5962/bhl.part.29309>
- Burger A (1948) Studies on the moss dwelling bdelloids (Rotifera) of Eastern Massachusetts. Transactions of the American Microscopical Society 67: 111-142. <https://doi.org/10.2307/3223491>
- Cimdine P (1979) Rotifers as saprobity indicators. Hydrobiological Journal 15: 63-67. [In Russian].
- Coulson SJ, Bartlett J, Boström S, et al. (2024) On the terrestrial and freshwater invertebrate diversity of the High Arctic archipelago of Svalbard: a revised species inventory and synopsis of the community composition. Arctic Science <https://doi.org/10.1139/as-2024-0017>
- Dartnall HJ, Hollowday ED (1985) Antarctic Rotifers. British Antarctic Survey. Cambridge
- Davis H (1873) A new Callidina; with the results of experiments on the desiccation of rotifers. Monthly Microscopy Journal 9: 201-209. <https://doi.org/10.1111/j.1365-2818.1873.tb02268.x>

- De Koning P (1947) Nieuwe bdelloide rotatorien. *Biolog Jaarb* 14: 184-205.
- De Ridder M (1972) The zoology of Iceland, Volume II, Part 13, Rotatoria, The zoology of Iceland. Ejnar Munksgaard, Copenhagen and Reykjavik.
- Devetter M (2007) Soil rotifers (Rotifera) of the Kokořínsko Protected Landscape Area. *Biologia* 62: 222-224. <https://doi.org/10.2478/s11756-007-0034-3>
- Devetter M (2009) Spatiotemporal dynamics of soil rotifers in a South-Bohemian beech forest. *Pesquisa Agropecuária Brasileira* 44: 1027-1032. <https://doi.org/10.1590/S0100-204X2009000800035>
- Devetter M (2010) A method for efficient extraction of rotifers (Rotifera) from soils. *Pedobiologia* 53: 115-118. <https://doi.org/10.1016/j.pedobi.2009.07.001>
- Donner J (1949) Rotatorien der Humusböden. *Österreichische Zoologische Zeitschrift* 11 (1/2): 117-151. [In German].
- Donner J (1950a) Rädertiere des Bodens. *Mikrokosmos* 39[In German].
- Donner J (1950b) Rotatorien der Humusböden. Hüllen und Gehäuse bei bdelloiden Rädertieren, besonders bei Bodenbewohnern. *Österreichische Zoologische Zeitschrift* 11: 287-335.
- Donner J (1952) Bodenrotatorien im Winter. *Mikrokosmos* 42: 29-33.
- Donner J (1965) Ordnung Bdelloidea (Rotatoria, Rädertiere). Akademie Verlag, Berlin.
- Donner J (1980) Einige neue Forschungen über bdelloide Rotatorien, besonders in Boden. *Revue d'Écologie et de Biologie du Sol* 17: 125-143.
- Fontaneto D, Ricci C (2004) Rotifera: Bdelloidea. In: Yule CM, Sen YH (Eds) *Freshwater invertebrates of the Malaysian Region*. Academy of Science Malaysia, Kuala-Lumpur, 6 pp.
- Fontaneto D, Ficetola GF, Ambrosini R, Ricci C (2006) Patterns of diversity in microscopic animals: are they comparable to those in protists or in larger animals? *Global Ecology and Biogeography* 15: 153-162. <https://doi.org/10.1111/j.1466-822X.2006.00193.x>
- Fontaneto D, Herniou EA, Barraclough TG, Ricci C (2007) On the global distribution of microscopic animals: New worldwide data on bdelloid rotifers. *Zoological Studies* 46: 336-346.
- Fontaneto D, Barraclough TG, Chen K, Ricci C, Herniou EA (2008) Molecular evidence for broad-scale distributions in bdelloid rotifers: everything is not everywhere but most things are very widespread. *Molecular Ecology* 17 (13): 3136-3146. <https://doi.org/10.1111/j.1365-294X.2008.03806.x>
- Fontaneto D, Iakovenko N, Eyres I, Kaya M, Wyman M, Barraclough T (2010) Cryptic diversity in the genus *Adineta* Hudson & Gosse, 1886 (Rotifera: Bdelloidea: Adinetidae): a DNA taxonomy approach. *Hydrobiologia* 662 (1): 27-33. <https://doi.org/10.1007/s10750-010-0481-7>
- Fontaneto D, Iakovenko N, Smet WH (2015) Diversity gradients of rotifer species richness in Antarctica. *Hydrobiologia* 761: 235-248. <https://doi.org/10.1007/s10750-015-2258-5>
- Fontaneto D (2019) Long-distance passive dispersal in microscopic aquatic animals. *Movement Ecology* 7 (10). <https://doi.org/10.1186/s40462-019-0155-7>
- Freckman DW, Virginia RA (1993) Extraction of nematodes from Dry Valley Antarctic soils. *Polar Biology* 13: 483-487. <https://doi.org/10.1007/BF00233139>
- Haigh SB (1963) The Bdelloid Rotifers of New Zealand. Part 1. *Journal of the Quekett Microscopical Club* 29: 161-170.

- Haigh SB (1970) The Bdelloid Rotifers of New Zealand. Part 5. Microscopy 31: 326-330.
- Harring HK, Myers FJ (1922) The rotifer Fauna of Wisconsin. Transactions of the Wisconsin Academy of Sciences, Arts, and Letters 20: 553-662.
- Hauer J (1938) Die Rotatorien von Sumatra, Java und Bali nach den Ergebnissen der Deutschen Limnologischen Sunda Expedition. Archiv für Hydrobiologie 15: 296-384.
- Iakovenko NS, Ovander EM (2011) Annotated checklist of bdelloid rotifers (Rotifera: Eurotatoria, Bdelloidea) of Ukraine. Bulletin of the Kyiv Taras Shevchenko National University, Biology 58: 25-29. [In Ukrainian].
- Iakovenko NS, Kašparová E, Plewka M, Janko K (2013) Otostephanos (Rotifera, Bdelloidea, Habrotrochidae) with the description of two new species. Systematics and Biodiversity 11 (4): 477-494. <https://doi.org/10.1080/14772000.2013.857737>
- Iakovenko NS, Smykla J, Convey P, et al. (2015) Antarctic bdelloid rotifers: diversity, endemism and evolution. Hydrobiologia 761: 5-43. <https://doi.org/10.1007/s10750-015-2463-2>
- Janson FO (1893) Versuch einer Übersicht über die Rotatorien-Familie der Philodinaeen. Abb Naturwiss Ver Bremen 12: 1-85. [In German].
- Jaturapruek R, Maiphae S (2025) An updated checklist of freshwater rotifers in Thailand: Taxonomy, diversity and distribution. Biodivers Data J 11: 155363. <https://doi.org/10.3897/BDJ.13.e155363>
- Kaya M, Smet WH, Fontaneto D (2010) Survey of moss-dwelling bdelloid rotifers from middle Arctic Spitsbergen (Svalbard). Polar Biology 33: 833-842. <https://doi.org/10.1007/s00300-009-0761-8>
- Koste W, Böttger K (1989) Rotatorien aus Gewässern Ecuadors. Amazoniana 10: 407-438. [In German].
- Koste W (1996) On soil Rotatoria from a lithotelma near Halali Lodge in Etosha National Park in N-Namibia, South Africa. Internationale Revue der gesamten Hydrobiologie und Hydrographie 81: 353-365. [In German]. <https://doi.org/10.1002/iroh.19960810305>
- Koste W, Zhuge Y (1996) A preliminary report on the occurrence of Rotifera in Hainan. Quekett Journal of Microscopy 37: 666-683.
- Kutikova LA (2005) The bdelloid rotifers of the fauna of Russia. KMK Scientific Press Ltd, Moscow. [In Russian].
- Lukashanets D, Iakovenko N, Fontaneto D, Devetter M, Janko K, Ejsmont-Karabin J, Bielańska-Grajner I, Hallet B, Smykla J, Kozeretska I, Trokhymets V (2025) New data on limno-terrestrial rotifers of the world. Version 1.7. Marine Research Institute, Klaipeda University. Occurrence dataset. URL: <https://doi.org/10.15468/vymt96>
- Lukashanets DA (2018) Fauna and taxonomy of the subclass Bdelloidea Hudson (Class Eurotatoria De Ridder; phylum Rotifera Cuvier) in Belarus, the Eastern Europe region. Zoology and Ecology 28: 36-45. <https://doi.org/10.1080/21658005.2017.1417101>
- Lukashanets DA, Convey P, Ol B, et al. (2021) Eukarya biodiversity in the Thala Hills, East Antarctica. Antarctic Science 33 (6): 605-623. <https://doi.org/10.1017/S0954102021000328>
- Lukashanets DA, Hihiniak YH, Miamin VY (2022) Extremely high abundances of *Prasiola crista*-associated micrometazoans in East Antarctica. Polar Research 41 <https://doi.org/10.33265/polar.v41.7781>
- Martínez A, Bonaglia S, Di Domenico M, Fonseca G, Ingels J, Jörger K, Laumer C, Leasi F, Zeppilli D, Baldrighi E, Bik H, Cepeda D, Curini-Galletti M, Cutter A, dos Santos G, Fattorini S, Frisch D, Gollner S, Jondelius U, Kerbl A, Kocot K, Majdi N, Mammola S,

- Martín-Durán J, Menegotto A, Montagna P, Nascimento FA, Puillandre N, Rognant A, Sánchez N, Santos I, Schmidt-Rhaesa A, Schratzberger M, Semprucci F, Shimabukuro M, Sommerfield P, Struck T, Sørensen M, Wallberg A, Worsaae K, Yamasaki H, Fontaneto D (2025) Fundamental questions in meiofauna research highlight how small but ubiquitous animals can improve our understanding of Nature. *Communications Biology* 8 (1). <https://doi.org/10.1038/s42003-025-07888-1>
- Milne W (1916) On the bdelloid Rotifera of South Africa. *Journal of the Quekett Microscopical Club* 2 (13): 47-83. <https://doi.org/10.5962/bhl.part.9295>
 - Murray J (1905) On a new family and twelve new species of Rotifera of the order Bdelloidea, collected by the Lake Survey. *Transactions of the Royal Society of Edinburgh* 41: 367-386. <https://doi.org/10.1017/S0080456800034451>
 - Murray J (1906a) Bdelloid Rotifera of the Forth Area. *Proceedings of the Royal Physical Society of Edinburgh* 16: 215-229.
 - Murray J (1906b) Some Rotifera of the Sikkim Himalaya. *Journal of the Royal Microscopical Society London* 26 (6): 637-644. <https://doi.org/10.1111/j.1365-2818.1906.tb00619.x>
 - Murray J (1907) Some South American Rotifers. *Amer Natural Boston* 4: 97-101. <https://doi.org/10.1086/278718>
 - Murray J (1908) Some African rotifers. *Journal of the Royal Microscopical Society London* 28 (6): 665-670. <https://doi.org/10.1111/j.1365-2818.1908.tb00241.x>
 - Murray J (1910) Antarctic Rotifera. *British Antarctic Expedition, 1907–1909. Rep Sci Inv* 1: 41-65.
 - Murray J (1911a) Australian Rotifera: collected by the Shackleton Antarctic Expedition, 1909. *Journal of the Royal Microscopical Society*
 - Murray J (1911b) Canadian Rotifera: collected by the Shaekeleton Antarctic Expedition, 1909. *Journal of the Royal Microscopical Society*
 - Murray J (1911c) Rotifera of New Zealand: collected by the Shaekeleton Antarctic Expedition, 1907–1909. *Journal of the Royal Microscopical Society*
 - Murray J (1911d) Rotifera of some Pacific islands: collected by the Shackleton Antaretic Expedition, 1909. *Journal of the Royal Microscopical Society*
 - Murray J (1911e) Some African rotifers: Bdelloidea of tropical Africa. *Journal of the Royal Microscopical Society*
 - Murray J (1911f) South African rotifers: collected by the Shackleton Antarctic Expedition, 1907–1909. *Journal of the Royal Microscopical Society*
 - Murray J (1913) South American Rotifera. *Journal of the Royal Microscopical Society*. <https://doi.org/10.1111/j.1365-2818.1913.tb01020.x>
 - Örstán A, Plewka M (2017) Bdelloid rotifer taxa described since 1965 with the addition of a new species. *Quekett Journal of Microscopy* 43: 31-37.
 - Örstán A (2018) Taxonomic morphology of the genus *Adineta* (Rotifera: Bdelloidea: Adinetidae) with a new species from a suburban garden. *Zootaxa* 4524 <https://doi.org/10.11646/zootaxa.4524.2.3>
 - Örstán A (2021) An extraordinary new fluvial bdelloid rotifer, *Coronistomus impossibilis* gen. nov., sp. nov., with adaptations for turbulent flow (Rotifera: Bdelloidea: Coronistomidae fam. nov.). *Zootaxa* 4966 <https://doi.org/10.11646/zootaxa.4966.1.2>
 - Peters U, Koste W, Westheide W (1993) A quantitative method to extract moss-dwelling rotifers. *Hydrobiologia* 255: 339-341. <https://doi.org/10.1007/BF00025857>

- Pokpongmongkol P, Jaturapruek R, Saardrit P, Maiphae S (2025) Substrate preference and seasonal distribution of bdelloid rotifers in mosses in a primary forest in Thailand. *Diversity* 17 (3): 171. <https://doi.org/10.3390/d17030171>
- Ricci C, Melone G (2000) Key to the identification of the genera of bdelloid rotifers. *Hydrobiologia* 418: 73-80. <https://doi.org/10.1023/A:1003840216827>
- Ricci C, Shiel R, Fontaneto D, Melone G (2003) Bdelloid rotifers recorded from Australia with description of *Philodinavus aussiensis* n.sp. *Zoologischer Anzeiger* 242: 241-248. <https://doi.org/10.1078/0044-5231-00101>
- Ricci C, Caprioli M (2005) Anhydrobiosis in bdelloid species, populations and individuals. *Integrative and Comparative Biology* 45 (5): 759-763. <https://doi.org/10.1093/icb/45.5.759>
- Ricci CN (1987) Ecology of bdelloids: how to be successful. *Hydrobiologia* 147: 117-127. <https://doi.org/10.1007/BF00025734>
- Rudescu L (1960) Rotatoria / Fauna republicii populare Romine. Trochelmintes. Acad. Rep. Pop. Romine, Bucurest. [In Romanian].
- Segers H (2007) Annotated checklist of the rotifers (Phylum Rotifera), with notes on nomenclature, taxonomy and distribution. *Zootaxa* 1564 (1): 1-104.
- Shmakova L, Malavin S, Iakovenko N, Vishnivetskaya T, Shain D, Plewka M, Rivkina E (2021) A living bdelloid rotifer from 24,000-year-old Arctic permafrost. *Current Biology* 31 (11): 712-713. <https://doi.org/10.1016/j.cub.2021.04.077>
- Song MO, Min GS (2015) A new species and ten new records of bdelloid rotifers from Korea. *Zootaxa* 3964 (2): 211-227.
- Song MO, Lee C (2020) Descriptions of two new species of *Philodina* (Rotifera, Bdelloidea), with notes on seven rare bdelloids from Korea. *Zootaxa* 4772: 253-276. <https://doi.org/10.11646/zootaxa.4772.2.2>
- Song MO, Lee C (2022) Descriptions of *Philodinavus koreanus* n. sp. and two new species of *Philodina* (Rotifera, Bdelloidea) from Korea. *Zootaxa* 5493: 36-54. <https://doi.org/10.11646/zootaxa.5493.1.2>
- Song MO, Lee C (2024) Descriptions of three new species of *Philodinida* (Rotifera, Bdelloidea) from Korea. *Zootaxa* 5493: 36-54. <https://doi.org/10.11646/zootaxa.5493.1.2>
- Sudzuki M (1981) Faunistic and ecological studies of the sewage biota of Japan (metropolis Tokyo and its adjacent prefecture. *Verh Internat Ver Limnol* 21: 1094-1100.
- Velasco-Castrillón A, Page T, Gibson J, et al. (2014) Surprisingly high levels of biodiversity and endemism amongst Antarctic rotifers uncovered with mitochondrial DNA. *Biodiversity* 15 (2–3): 130-142. <https://doi.org/10.1080/14888386.2014.930717>
- Wang W, Yang Y, Cui Z (2023) High diversity and strong habitat preference of bdelloid rotifers in the moss and leaf litter from a small area of urban plain and adjacent hill in China. *Biodivers Conserv* 32: 2769-2789. <https://doi.org/10.1007/s10531-023-02630-x>
- Wang W, Yang Y, Örstan A, et al. (2025) Species diversity of bdelloid rotifer (Rotifera, Bdelloidea) in different areas in China, with a description of two new species. *ZooKeys* 1229: 1-23. <https://doi.org/10.3897/zookeys.1229.113385>
- Wulfert K (1942) Neue Rotatorienarten aus deutschen Mineralquellen. *Zoologischer Anzeiger* 197: 187-200.
- Wulfert K (1960) Die Radertiere saurer Gewässer der Dubener Heide. I. Die rotatorien des Zadlitzmoors und des Wildenhainer Bruchs. *Arch für Hydrobiol* 56: 261-298.
- Zarubov A, Cimdinş P (1990) Ecological aspects of distribution in rotifers of the Order Bdelloidea. *Conference Proceedings. Zoological Institute Publishing* [In Russian].

- Zawierucha K, Porazinska DL, Ficetola GF, et al. (2021) A hole in the nematosphere: tardigrades and rotifers dominate the cryoconite hole environment, whereas nematodes are missing. *Journal of Zoology* 313 (1): 18-36. <https://doi.org/10.1111/jzo.12832>
- Zeng Y, Nan W, Qing W, Iakovenko N, Ying L, Yufeng Y (2020) Bdelloid rotifers (Rotifera, Bdelloidea) of China: Diversity and new records. *ZooKeys* 941: 1-23. <https://doi.org/10.3897/zookeys.941.50465>