

Persistent Organic Pollutant Burden, Experimental POP Exposure, and Tissue Properties Affect Metabolic Profiles of Blubber from Gray Seal Pups

Kelly J. Robinson,^{*,†,‡} Ailsa J. Hall,[†] Cathy Debier,[‡] Gauthier Eppe,[§] Jean-Pierre Thomé,^{||} and Kimberley A. Bennett[⊥]

[†]Sea Mammal Research Unit, Scottish Oceans Institute, University of St Andrews, St Andrews, Fife KY16 8LB, United Kingdom of Great Britain and Northern Ireland

[‡]Louvain Institute of Biomolecular Science and Technology, Université Catholique de Louvain, Ottignies-Louvain-la-Neuve, Louvain-la-Neuve 1348, Belgium


[§]Center for Analytical Research and Technology (CART), B6c, Department of Chemistry, Université de Liège, Liège 4000, Belgium

^{||}Center for Analytical Research and Technology (CART), Laboratory of Animal Ecology and Ecotoxicology (LEAE), Université de Liège, Liège 4000, Belgium

[⊥]Division of Science, School of Science Engineering and Technology, Abertay University, Dundee DD1 1HG, United Kingdom of Great Britain and Northern Ireland

Supporting Information

ABSTRACT: Persistent organic pollutants (POPs) are toxic, ubiquitous, resist breakdown, bioaccumulate in living tissue, and biomagnify in food webs. POPs can also alter energy balance in humans and wildlife. Marine mammals experience high POP concentrations, but consequences for their tissue metabolic characteristics are unknown. We used blubber explants from wild, gray seal (*Halichoerus grypus*) pups to examine impacts of intrinsic tissue POP burden and acute experimental POP exposure on adipose metabolic characteristics. Glucose use, lactate production, and lipolytic rate differed between matched inner and outer blubber explants from the same individuals and between feeding and natural fasting. Glucose use decreased with blubber dioxin-like PCBs (DL-PCB) and increased with acute experimental POP exposure. Lactate production increased with DL-PCBs during feeding, but decreased with DL-PCBs during fasting. Lipolytic rate increased with blubber dichlorodiphenyltrichloroethane and its metabolites (DDX) in fasting animals, but declined with DDX when animals were feeding. Our data show that POP burdens are high enough in seal pups to alter adipose function early in life, when fat deposition and mobilization are vital. Such POP-induced alterations to adipose metabolic properties may significantly alter energy balance regulation in marine top predators, with the potential for long-term impacts on fitness and survival.



INTRODUCTION

The impact of environmental and organismal accumulation of man-made chemicals is a growing global challenge.^{1,2} Persistent organic pollutants (POPs) are toxic,³ accumulate in living tissues,⁴ and resist biological and chemical degradation.⁵ Widespread bans on their production initially lowered environmental POP levels,^{6–9} but reductions have slowed or ceased more recently.^{10–13} Legacy POPs are thus ubiquitous in the marine environment and remain a serious concern for top predators, in which biomagnification produces the highest burdens.^{4,11,14–16}

At high exposure levels POPs, such as polychlorinated biphenyls (PCBs), polybrominated diphenyl ethers (PBDEs) and organochlorine pesticides (OCPs), including dichlorodiphenyltrichloroethane (DDT), impair fertility, reproductive function, skeletal and neurological development, immunity and

thyroid function in humans and wildlife.^{17–24} Recently, several POPs have also been implicated as lipid disruptors in humans and rodents.^{25,26} In fish and zooplankton, POPs can increase fat deposition and lipogenic gene expression.^{27–31} Conversely, exposure to POPs, such as dioxins, lindane, and OCPs, can cause rapid energy reserve depletion, termed “wasting syndrome”.^{32–35} These apparently conflicting effects of different POPs on a variety of energetic processes highlight the need for further investigation of pollutant mixtures, doses and exposure routes, and their physiological consequences for organisms in natural environments.

Received: August 1, 2018

Revised: October 17, 2018

Accepted: October 19, 2018

Published: October 19, 2018



POP-induced alterations to energy balance may have significant individual and population level consequences because fatness is often positively associated with reproductive fitness and survival.³⁶ This is particularly true in marine mammals,^{37–39} which rely heavily on subcutaneous fat reserves for insulation^{40,41} and as a metabolic fuel when food is scarce. Many marine mammal species also undergo natural periods of fasting during moulting, breeding, migration, or the postweaning fast that occurs in many phocid pups.^{37,42–46}

Marine mammals typically experience exceptionally high POP burdens due to their high fat content and top trophic level, and are regarded as sentinels of marine pollution.^{14,16} POPs are transferred to suckling young as they undergo rapid fat accretion.⁴⁷ In fasting individuals, less hydrophobic POPs are released into the bloodstream as lipid is mobilized and more recalcitrant POPs concentrate in the remaining blubber.^{47–54} Phocid pups thus experience high POP exposure during both feeding and fasting, two fundamentally important periods for adipogenesis, fat mobilization, and maturation of energy balance pathways. Their reliance on adipose tissue to fuel metabolism during fasting or while searching for food in changing environments⁵⁵ places these top predators at high risk from adverse effects of POP exposure as a result of impaired fat cell function.

Marine mammals show structural changes in fatty acid (FA) content and thermal properties with blubber depth.^{54,56–61} Effects of blubber depth and nutritional state on metabolic function and sensitivity to POP exposure have not been explored. High POP levels are associated with altered adipose gene expression in seals,^{62–64} polar bears (*Ursus maritimus*),⁵⁵ and killer whales (*Orcinus orca*).⁶⁵ POP cocktails from polar bear fat can induce adipogenesis in mouse (*Mus musculus*) preadipocyte cell lines and primary polar bear preadipocytes.⁶⁶ However, the consequences of POP-induced gene expression changes for fat tissue function are unknown.

The major functions of fat tissue are to store lipid and release it for use as metabolic fuel. The balance between lipogenesis and lipolysis depends on the nutritional state of the organism and the hormonal milieu to which the tissue is exposed, and may be influenced by POPs. Glucose uptake, lactate production and glycerol release are important metabolic properties of adipose tissue that contribute to whole animal glucose homeostasis and energy balance. Adipose tissue synthesizes triacylglycerol (TAG) from glucose,^{67,68} such that glucose uptake rates can be used to investigate metabolic function of adipose tissue. Glucose uptake is altered by POPs at tissue and the whole animal level in other animals.^{69–87} Of glucose taken up by adipose, 20% is stored as TAG, and 70% is used to synthesize lactate.⁶⁷ Lactate production is thus an important route of glucose disposal and may be influenced by POP exposure. Glycerol is produced during lipolysis,⁸⁸ and as a byproduct of glycolysis,⁶⁸ and is thus a major secretory product of adipose tissue. Unlike FA, glycerol cannot be re-esterified by adipose in vitro,^{89,90} making its production a useful marker of lipolytic,⁸⁸ and, to some extent, glycolytic⁶⁸ activity.

We measured glucose uptake, lactate production and glycerol release as indices of metabolic activity in blubber by employing an explant approach we have developed using blubber biopsies from wild gray seals (*Halichoerus grypus*).⁹¹ Our approach allows experimental work to be performed in vitro in remote field locations to investigate fat tissue function,⁹¹ which, in marine mammals, was previously limited to measurement of enzyme activities.⁹² Here, we used blubber

explants from wild gray seal pups, which remain on land during both feeding (during suckling) and fasting (during the postweaning fast). We investigated whether metabolic properties of blubber differ by tissue depth, nutritional state, and intrinsic POP burden. We also investigated acute changes in blubber metabolic characteristics in response to exogenous experimental treatments with high, but biologically relevant, POP concentrations.

METHODS

Study Site and Animals. Field work was conducted on the Isle of May, Scotland (56°11'N, 02°33'W) from 7th November to 18th December 2016 under permit from Scottish Natural Heritage (SNH). Gray seal females that were part of a long-term study were observed from when they came ashore, and the date of birth of their pups was recorded. Pups were included if they were at an appropriate predicted mass for their age (number of days postpartum) late in the suckling period and if they did not have any obvious indications of ill health. We collected blubber biopsies for POP analysis and explant culture from 27 pups late in the suckling period (day 15). Pups were then left to wean naturally from their mothers, which occurred at approximately 18 days old. We then attempted to resample the same pups once during the postweaning fast (day 12 after weaning). As pups were still feeding for several days between the two sampling events, additional mass gain and increases in blubber lipid percentage were possible between the feeding/fasting sampling points. Four pups were removed from the study because they developed an unrelated infection. This gave a total of 27 pups (9 female and 18 male) sampled while feeding and 23 (7 female and 16 male) of the same animals sampled again during fasting.

Pups were defined as weaned based on daily observation to determine maternal departure date.⁴³ Pups were then transferred to a large (~600 m²) fenced off area of the island, away from the main colony, with access to a pool of fresh water, in accordance with U.K. Home Office guidelines and based on prior studies.^{43,93}

The sex of all pups was determined at early lactation (~day 5 of suckling). Pups were weighed at early and late lactation; when they were initially transferred into the pen (within 2 days after weaning); and every 5 days in the pen. Pups less than 30 kg at weaning were excluded from the study. To ensure pups were of a viable departure mass, early release criteria were set at loss of 25% of weaning mass or reaching 30 kg, whichever was the higher mass.⁴³ It was not necessary to release any of the study animals early. All pups were released into the colony with ready access to the sea.

All animal procedures were performed under the U.K. Home Office project license #70/7806 and conformed to the U.K. Animals (Scientific Procedures) Act, 1986. All research received prior ethical approval from Abertay University and the University of St Andrews Animal Welfare and Ethics Committee.

Biopsy Sampling. Biopsy sampling, transport, explant creation, and culture protocols were adapted from a previous study.⁹¹ Prior to sampling, pups were given a mass-specific intravenous dose of Zoletil and subcutaneous injections of Lignol at the biopsy sites. Three full depth blubber biopsies, one 6 mm and two 10 mm, were taken.⁹⁴ The 6 mm biopsy was immediately wrapped in foil and frozen at –20 °C for POP analysis.^{47,50} The two 10 mm biopsies were cut immediately, using sterile surgical scissors, into “inner” (blubber closest to

Table 1. Means and Ranges for Study Pup Masses and Fat Content of 6 mm Biopsies, with Median Concentrations and Ranges of POPs in Grey Seal Pup Blubber in Feeding and Fasting States

	feeding mean	feeding range	fasting mean	fasting range
pup mass (kg)	41.9	31.4–52.6	40	32.8–50.4
biopsy fat content (% lipid)	78.5	67.8–85.5	82.9	56.3–88.9
	feeding median	feeding range	fasting median	fasting range
$\sum$ NDL-PCB (ng g ⁻¹ lipid)	354.29	194.2–790.5	541.73	302.5–1253.8
$\sum$ DL-PCBs (ng g ⁻¹ lipid)	10.78	5.5–45.6	19.78	9.6–151.2
$\sum$ PBDE (ng g ⁻¹ lipid)	13.81	6.4–57.9	20.2	10.2–70.1
$\sum$ DDX (ng g ⁻¹ lipid)	169.09	98.5–420.8	209.56	128.4–345.7

muscle layer) and “outer” (blubber closest to the skin) tissue, and placed into separate 15 mL centrifuge tubes filled with warm (37 °C) sterile Krebs Ringer solution (pH 7.4; NaCl; 7.89 g/L; KCl; 0.373 g/L; MgSO₄·0.12 g/L; K₂HPO₄; 0.07 g/L; Glucose; 0.99 g/L; HEPES; 4.77 g/L; CaCl₂; 0.11 g/L⁹⁵ with 1% Antibiotic Antimycotic Solution (all chemicals supplied by Sigma-Aldrich) for transport back to the field laboratory. A single hidden suture (Ethicon Vicryl Suture (W9114) Violet, 3–0, 20 mm, 75 cm with VICRYL Suture 1/2 circle “Taper Point Plus” Needle) was used to close 10 mm biopsy sites, either vertically or horizontally through the skin.⁹⁶ Pups were observed remotely on a daily basis, and biopsy sites checked at each reweighing to document healing. The second sample was not taken until all the previous biopsy sites had healed.

Explant Culture and POP Exposure. All biopsies were returned to the field laboratory, processed, and placed in the incubator within 30 min of collection, which is within timeframes used in prior explant studies on livestock^{97,98} and wild seals.⁹¹ Upon arrival, tissue was washed with 1 mL sterile, warm Krebs Ringer buffer. Visible hair, muscle, or blood contaminated tissue was removed. Tissue was minced using a sterile scalpel into 5–10 mg pieces, and 100 mg portions of tissue were weighed out in a sterile Petri dish. The 100 mg explants were dispensed into preprepared 12 well plates warmed to 37 °C in a sterile, nonducted PCR hood. In total, 1500 μ L of either complete cell culture media (medium 199, Hanks’ Balanced Salts with 1% Antibiotic Antimycotic Solution, 1% FA supplement and 5% charcoal stripped fetal bovine serum) or complete media containing 150 ng/mL POP mix (PCB Aroclor standard mix 1 (Sigma-Aldrich), containing Aroclor 1016; 500 μ g/mL, Aroclor 1260; 500 μ g/mL, Decachlorobiphenyl (PCB 209); 50 μ g/mL and 2,4,5,6-Tetrachloro-*m*-xylene (an OCP); 50 μ g/mL in acetone:methanol solution (2:3)) was added per well to generate control and + POP experiments. Plates were placed in a humidified incubator maintained at 37 °C and 5% CO₂ (Thermo Scientific, Midi 40 CO₂ Incubator, model: 3404) for 24 h. All media was drawn off after 24 h and frozen at –20 °C. Explants were snap frozen in liquid nitrogen and stored at –80 °C.

Metabolite Measurement. Glucose, lactate, and glycerol concentrations were measured in media from explant experiments using Randox (County Antrim, U.K.) kits (glucose: GL364, lactate: LC2389, glycerol: GY105) and standards in an RX Monza (Randox) Clinical Chemistry analyzer (Model: 328-14-0914) as described previously.⁹¹ Internal quality control measurements lay within $\pm 15\%$. Intra and inter assay variability for sample analysis was $< 5\%$. Mean variation between replicates from the same individual and of the same nutritional state and tissue type for the three metabolites

analyzed were as follows; glucose 26.6% (range 0.6–144%), lactate 46.9% (range 3–164%), and glycerol 19.8% (range 0.5–106%). Rates of glucose removal and accumulation of lactate (indices of glycolysis) and glycerol (indices of lipolysis) were calculated per 100 mg tissue⁻¹ hour⁻¹.

POP Analysis. Extraction and detection of POPs took place at the Center for Analytical Research and Technology (CART), University of Liège, Belgium, using standard methods (Supporting Information SI 1). The 6 mm biopsies yielded 385 ± 7 (s.e.) mg blubber tissue per sample. Six non Dioxin-Like PCBs (NDL-PCBs) (28, 58, 101, 138, 153, 180), eight Dioxin-Like PCBs (DL-PCBs) (105, 114, 118, 123, 156, 157, 167, 189), and nine PBDEs (28, 47, 66, 85, 99, 100, 153, 154, 183) were analyzed at CART in the Department of Chemistry. Four OCPs (*o,p'*-DDT, *p,p'*-DDT, dichlorodiphenyldichloroethane (DDD), and dichlorodiphenyldichloroethylene (DDE)) were analyzed at CART in the Laboratory of Animal Ecology and Ecotoxicology (LEAE).

POP concentrations were measured in ng g⁻¹ lipid, summed into $\sum$ NDL-PCBs, $\sum$ DL-PCBs, $\sum$ PBDEs and $\sum$ DDXs (*o,p'*-DDT, *p,p'*-DDT, DDD, DDE) and log transformed prior to statistical analysis.

Statistical Analysis. Analyses were performed using the statistical package R 3.4.1.⁹⁹ *t* tests were used to compare log transformed POP concentrations between feeding and fasting samples. Pearson’s correlation coefficient was calculated between all four $\sum$ POP types (log transformed $\sum$ NDL-PCBs, $\sum$ DL-PCBs, $\sum$ PBDEs and $\sum$ DDXs). As all POP classes were highly significantly correlated, we explored the effects of each POP class separately. Generalized additive mixed models (GAMM)¹⁰⁰ were used to identify the variables that best explained glucose uptake, lactate production, and glycerol production. Explanatory variables explored in these models included log transformed $\sum$ NDL-PCBs, $\sum$ DL-PCBs, $\sum$ PBDEs, and $\sum$ DDXs in the blubber at time of sampling (each POP class tested separately in each model), the mass of the pup, the tissue depth (inner or outer), the nutritional state at sampling (feeding or fasting), the experimental conditions (control or + POP) and the pup’s sex. Interaction terms between nutritional state, tissue type, experimental conditions and POP concentrations were also investigated (see SI Table S1). The model was fitted using the multiple generalized cross validation library *mgcv*.¹⁰¹ The identities of individual pups were fitted as random effect smooths¹⁰² to account for the same individual generating multiple explants and any variation between replicates. The smoothing parameters were set by maximum likelihood to reduce the risk of overfitting.¹⁰³ The models were fitted with a Gaussian error distribution. Glucose data were log transformed and glycerol data were square root transformed. No transformation was needed for lactate data. Model selection was performed by backward stepwise




Figure 1. Scatter plot (left panel) and prediction line plot (right panel) showing the impact of intrinsic Σ DL-PCBs on glucose uptake in blubber from gray seal pups, comparing inner (black) /outer (blue) tissue and feeding (filled triangles or solid lines) /fasting (open circles or dashed lines) states ($n = 27$ feeding and $n = 23$ fasting pups). Individual was included as a random effect in the model.




Figure 2. Prediction line plots showing the impact of exposure to acute exogenous POPs on glucose uptake in blubber from gray seal pups, comparing inner (left panel) and outer (right panel) tissue and feeding/fasting states across the range of intrinsic Σ DL-PCBs in the blubber. Individual was included as a random effect in the model.

elimination through examination of R^2 values, Akaike's information criterion (AIC) values, dAIC, Akaike's weights, QQ, and residual plots

RESULTS AND DISCUSSION

POP Concentrations. Mean concentrations and ranges of the summed POPs in gray seal pup blubber are given in Table 1.

Higher POP levels when pups were fasting (Welch's two sample t-test, $p < 0.001$ for all log transformed POP classes) are consistent with previous work showing that lipid mobilization during fasting concentrates more lipophilic POPs in remaining fat tissue.^{50,51,53} Blubber may thus be more vulnerable to disruptive effects of contaminants during

fasting if the POPs can interact with cellular machinery to exert negative effects.

Glucose Uptake. Σ DL-PCBs, experimental conditions, tissue depth, and nutritional state best explained variation in glucose uptake, along with an interaction between tissue depth and nutritional state (GAMM: $R^2 = 0.36$, SI Table S2). Σ NDL-PCBs, Σ PBDEs and Σ DDX had no significant impact on glucose uptake (SI Table S1). Uptake was higher in inner compared to outer tissue ($p < 0.001$). Glucose uptake was lower during fasting than feeding in outer blubber ($p = 0.035$), but did not differ between nutritional states in inner tissue (Figure 1). In both inner and outer tissue, glucose uptake was inversely related to intrinsic Σ DL-PCBs ($p < 0.001$) (Figure 1) and the slope of the relationship did not depend on nutritional state. Acute exogenous POP exposure increased




Figure 3. Scatter plot (left panel) and prediction line plot (right panel) showing the impact of intrinsic Σ DL-PCBs on lactate generation/uptake in blubber from male gray seal pups ($n = 34$), comparing inner (black) /outer (blue) tissue and feeding (filled triangles or solid lines) /fasting (open circles or dashed lines) states. Individual was included as a random effect in the model.

glucose uptake compared to controls, (GAMM: $R^2 = 0.36$, $p < 0.001$, SI Table S2, Figure S2), and the increase was not affected by nutritional state or tissue depth (Figure 2). Doses of each PCB congener given in the explant POP exposure conditions, with the estimated percentage change for each PCB type measured in blubber are given in SI Table S3. A boxplot of glucose uptake data during different nutritional states and tissue depths is provided in SI Figure S1.

These data show for the first time that metabolic activity in phocid blubber is altered by the POPs accumulated in utero and during suckling. POPs have been postulated to be “obesogens” from epidemiological data in humans.²⁴ Experimental studies have demonstrated that POPs can alter adipocyte function in rodents and human cell lines.^{25,104} POPs are associated with altered adipose and liver expression of metabolic pathway genes in adult polar bears^{55,66} and ringed seals (*Pusa hispida*),^{63,64} but the functional consequences of such changes for fat tissue have not been identified. Recent experimental studies that replicate environmental or physiological POP concentrations and congener mixtures provide causal evidence of their immunological¹⁰⁵ and adipogenic effects.⁹¹ Our work builds on these findings by showing that existing POP concentrations in blubber tissue, generated in utero and through lactational transfer, can alter metabolic properties of adipose tissue in young, developing seals.

Our data are consistent with experimental studies showing PCB 77, 153 and dioxin reduce glucose and insulin tolerance in mice, guinea pigs (*Cavia porcellus*), rabbits (*Oryctolagus cuniculus*), and in human cell lines.^{69–77} PCBs are also associated with dysfunctional glucose homeostasis in humans.^{78–81} Here, only Σ DL-PCBs altered glucose uptake in blubber explants, consistent with an aryl hydrocarbon receptor (AhR)-dependent pathway in seals, as seen in other species.^{71,78} POP-induced reduced glucose use is accompanied by a fall in glucose transporter, GLUT4, mRNA expression and GLUT1 protein levels in other animals,⁷⁵ and may also occur in seals. The opposite effects of an exogenous POP exposure to the intrinsic Σ DL-PCB levels on glucose uptake may reflect the difference between short (overnight) in vitro and long-term (in

utero and over 15 days of suckling) exposure, which may produce opposing effects on membrane fluidity and transport properties.¹⁰⁶ Alternatively, either the DL-PCB dose in the acute exposure experiments may have been too low to affect glucose uptake (SI Table S3) or the exogenous POP cocktail produces different effects from the in vivo contaminant mixture. We cannot distinguish between these possibilities from our data.

Lower glucose uptake caused by DL-PCBs is likely to reduce fat deposition rate, which may be problematic for pups laying down fat while suckling. However, fat deposition may increase if acute exogenous POP exposure increases glucose uptake. Indeed, short-term exposure to a POP cocktail from polar bear blubber induced adipogenesis in a mouse preadipocyte cell line.⁶⁶ The timecourse of POP effects on both adipogenesis and lipogenesis needs to be explored.

NDL-PCBs, DDX, and PBDEs in blubber did not affect glucose uptake, despite evidence of their impacts on glucose homeostasis in other animals.^{82–87,107} Our data suggest that the negative relationship between PBDEs and first year survival of gray seal pups¹⁰⁴ is not driven by direct alterations to adipose metabolic properties. PBDEs may alter adipose function indirectly through endocrine disrupting effects, for example, by altering thyroid or insulin function,^{86,87,108,109} highlighting the need for improved understanding of energy balance regulation in seals, and its disruption by POPs.

Glucose uptake was higher in inner tissue under all conditions, suggesting deeper blubber has greater metabolic activity than superficial tissue. This finding is consistent with regional stratification in structure; the theorized role of smaller, more numerous adipocytes that are more metabolically active⁹² in deeper tissue; and the observation that deeper blubber contains higher glycolytic enzyme levels compared to superficial blubber in fin whales (*Balaenoptera physalis*⁹²).

Glucose uptake fell as the animals transitioned from feeding to fasting in outer blubber, but not in inner blubber. This likely reflects high levels of fat deposition throughout the blubber during feeding, followed by cessation of fat accumulation




Figure 4. Scatter plot (left panel) and prediction line plot (right panel) showing the impact of intrinsic Σ DDX on glycerol generation in blubber from gray seal pups, comparing inner (black) /outer (blue) tissue and feeding (filled triangles or solid lines) /fasting (open circles or dashed lines) states ($n = 27$ feeding and $n = 23$ fasting pups). Individual was included as a random effect in the model.

during fasting. Metabolic rate falls in fasting pups,^{42,110} and this is likely reflected in reduced demands by all tissues, including blubber. Lower sensitivity to residual insulin present in stripped FBS may explain lower glucose uptake and greater decrease in outer tissue from fasting animals, a state in which many animals become less sensitive to the effect of insulin on glucose uptake.^{111–113} The lack of decreased glucose uptake in inner blubber in fasting animals is surprising. It is, however, plausible that cell membrane expression of GLUT4 and/or insulin-independent GLUT1 are reduced to a greater extent in outer compared to inner blubber during fasting. Indeed, fasting northern elephant seals (*Mirounga angustirostris*) maintain expression of GLUT4 in adipose cell membranes.¹¹⁴

Previous studies of metabolic effects of POPs on marine mammals have focused on adults.^{55,63,64,66} Concentrations of POPs in adult seals are an order of magnitude greater than those found in pups, with adult seal blubber containing several thousand Σ PCBs (ng/g)¹¹⁵ compared to the levels detected in the pups here, which were in the hundreds of ng/g range (Table 1). Here, we have shown that pups have already accumulated high enough POP levels in their blubber to impact on adipose function at less than 3 weeks of age, when the ability to generate and regulate blubber is a key factor for survival.¹¹⁶ Growing and dividing adipocytes are more vulnerable to disruption by DL-PCBs because they have higher AhR levels than mature, differentiated adipocytes.^{117,118} The sensitivity to POP-induced alterations to adipose tissue metabolism is therefore likely to be greatest in young marine mammals, which are exposed to large quantities of POPs from their mothers^{47,50} and are likely to have a high proportion of proliferating preadipocytes as a result of rapid fat deposition.¹¹⁹ The energy balance and survival consequences of reduced glucose uptake in young pups now needs to be explored.

Lactate Production. Σ DL-PCBs, tissue depth, nutritional state, pup sex and interactions between Σ DL-PCBs and nutritional state, and tissue depth and nutritional state were retained in the model that best explained lactate production (SI Tables S1 and S2). Σ NDL-PCBs, Σ PBDEs, and Σ DDX had no significant impact on lactate production (SI Table S1).

Lactate production increased with DL-PCBs in feeding pups, but decreased with DL-PCBs when pups were fasting (GAMM: $R^2 = 0.29$, $p = 0.003$; SI Table S2, Figure S3). Lactate generation was higher in males than in females ($p = 0.05$) and higher in inner tissue than outer tissue for any given DL-PCB load in any given nutritional state ($p = 0.001$) (Figure 3). The difference between inner and outer tissue tended to be greater during feeding than during fasting ($p = 0.055$). Lactate generation was not impacted by acute exogenous experimental POP exposure. A boxplot of lactate generation/uptake data during different nutritional states and tissue depths is provided in SI Figure S2.

Lactate production is a major route of glucose disposal in fat tissue.⁶⁷ However, the inhibitory effect of DL-PCBs on glucose uptake is only consistent with lactate production during fasting. During feeding, lactate production increased with increased DL-PCBs in the face of reduced glucose uptake. One explanation is that the electron transfer chain experiences increased push force during suckling, in combination with inhibition by DL-PCBs,^{120,121} which could elevate lactate production through increased reliance on glycolysis. Suckling pups with higher DL-PCB burdens could thus experience greater risk of metabolic acidosis from higher adipose lactate production, with attendant risk of adipose dysfunction, greater demand on liver gluconeogenesis to process the lactate, and thus heightened energetic costs that could impact on efficiency of mass transfer and fat accretion during suckling.

Adipose lactate production is a direct consequence of lactate dehydrogenase (LDH) activity.¹²² LDH activity can fall during fasting,¹²³ and LDH gene expression and enzyme activity also differ substantially between adipose depots in rats.¹²² LDH activity can also be modulated by PCB exposure in rat and dog liver, and the effect is dose dependent.^{124,125} Shifts in LDH expression and activity may thus contribute to the differences in lactate production between depths and nutritional states seen here in response to DL-PCBs. Higher lactate generation in inner tissue and the greater difference between regions during feeding may be a result of greater glucose uptake and metabolic activity in inner blubber during suckling. Regional

differences in metabolic properties may also relate to adipocyte size.¹²⁶

Higher lactate production by adipose from males than females has not been reported, either in adult seals⁹¹ or in humans and rodents¹²² and may result from the small number of females here. However, since adult females accumulate fat faster than adult males¹²⁷ and female juvenile northern elephant seals spare fat more effectively than males,¹²⁸ it is possible that some metabolic differences are present in pups' blubber prior to overt size dimorphism. For example, females may use more lactate to synthesize FA¹²⁹ than males, releasing less net lactate.

Glycerol Production. Nutritional state, an interaction between tissue depth and nutritional state, and an interaction between nutritional state and intrinsic blubber DDX were significant terms in the best model to explain glycerol production (SI Table S1 and S2, Figure S4). In addition, extrinsic POP treatment, mass and sex were retained as non-significant variables. Unexpectedly, glycerol production, was higher during feeding than fasting (GAMM: $R^2 = 0.74$, $p < 0.001$ Figure 4). In addition, lipolysis was higher in outer than in inner tissue during feeding, whereas inner tissue had a higher lipolytic rate than outer tissue in the fasted state ($p = 0.0003$; SI Table S2). Finally, lipolytic rate increased with intrinsic blubber DDX in fasting animals, whereas the opposite relationship between DDX and lipolytic rate occurred in feeding pups ($p < 0.0001$, Figure 4). A boxplot of glycerol generation data from different nutritional states and tissue depths is provided in SI Figure S3.

POPs, such as dioxin and related compounds, and OCPs typically produce "wasting syndrome" at high doses,^{32–35} likely due to increased WAT lipolysis.¹³⁰ In contrast, some studies on rat adipose or rodent cell lines show no effect of DDT or DDE on lipolysis at lower doses.^{131,132} Our data show that lipolysis is increased in blubber of fasting gray seal pups in response to intrinsic tissue levels of DDT and its metabolites, but decreased in feeding pups. These data suggest that, as in bats, seal pups in negative energy balance may be most vulnerable to the presence of DDX in their tissues.³⁴ Interestingly, the increase in lipolysis is greater in outer tissue, which is proposed to be important for insulation,^{56–61} such that DDT may negatively affect both thermoregulatory and metabolic capability of blubber during fasting. A reduction in lipolytic rate with DDX in feeding animals could reflect inhibition of tissue turnover of FFA, increased lipogenesis or a shift toward lactate production rather than glycerol generation during glycolysis.⁶⁸

Lower glycerol production during fasting compared to feeding contrasts starkly with the typical fasting increase in lipolysis.^{133,134} Regional and nutritional differences may result from depth changes in metabolic activity and adipocyte size.¹³⁵ Lower glycerol production in fasting pups could also arise from a reduction in aldolase activity.¹³⁶ In addition, higher lipolytic rates in feeding animals may reflect their higher metabolic rate,^{42,110} faster rates of triglyceride turnover and/or greater production of 3 carbon (3C) intermediates to protect tissues against high fuel availability and provide fuel for actively growing tissue elsewhere.^{68,122} High rates of glycerol production in feeding animals could protect adipocytes from metabolic acidosis, minimize overproduction of oxidizing equivalents and maximize energy efficiency.^{68,137}

Value and Wider Application of the Explant Method. Wildlife species are facing increasing anthropogenic threats

that can limit their ability to find food and store fat as an energy reserve for lean times. Constrained, fragmented or reduced foraging ranges¹³⁸ and decreased prey availability^{139,140} limit opportunities for fat accumulation, and force individuals to rely on adipose stores more frequently to sustain them until the next feeding opportunity. Understanding the impact of natural and anthropogenic stressors on energy balance requires experimental interventions that are practically and ethically challenging to accomplish in marine mammals and other wildlife species. In vitro methods are increasingly being explored as alternatives to whole animal work and lethal sampling. Prior efforts have generated viable skin tissue samples for culture via dermal punches and biopsy darts in both marine^{141,142} and terrestrial¹⁴³ wildlife species. Given the importance of fat tissue to the health and survival of wildlife in rapidly changing environments, there is a critical need for methods to better understand their adipose physiology. We previously used blubber explants from wild seals, which were returned to tissue culture facilities in a mainland laboratory.⁹¹ Here, we collected and maintained viable blubber explants for 24 h in a remote field laboratory, with only core tissue culture equipment. The methodology outlined here could therefore be adapted for use in other wildlife species in remote field locations or where alternative in vitro methods are not available.

Outlook. Our explant method clearly shows promise in better understanding adipose tissue responses to contaminants, and can be expanded beyond the parameters measured here. Although establishment of cell lines would permit a wider range of exposures, including time courses, dose responses, and mixtures over longer duration experiments, such approaches have been hampered by the absence of typical proliferation and differentiation responses in primary cells from nonmodel species.⁵³ A focus on understanding the requirements of such cell types and generation of cell lines would greatly facilitate progress in toxicological studies on adipose from wildlife species.

In summary, we have shown that dioxin-like PCBs disrupt adipose glucose uptake and lactate production, and DDX disrupt lipolysis in gray seal pups, using an experimental explant approach. Our results are particularly important because they demonstrate POP-induced alterations to adipose tissue function in young animals in the first weeks of life, for which rapid fat accumulation during suckling and mobilization during fasting are vital for survival. They also highlight that different POPs may impact different metabolic characteristics, and that their effects are modified by nutritional state and tissue depth, producing complex effects on tissue function. Wider screening of metabolic properties of cells and tissue from nonmodel organisms is key to identify energy-balance disrupting effects of legacy and emerging contaminants. Our methods and findings also have wider application in understanding drivers of juvenile survival of other top marine predators, particularly in species that simultaneously experience high POP loads and rapid alterations in prey availability and habitat.

■ ASSOCIATED CONTENT

📄 Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.est.8b04240.

SI 1, Methodological details for POP detection; SI Table S1, candidate GAMM models exploring different summed POP types, potential interactions, and the top models in the selection process; SI Table S2, model output from all final GAMMs analyzing metabolic characteristics in explant experiments with their standard errors, estimates, and *p* values; predictor variables in parentheses indicate the tested group; SI Table S3, doses of each PCB type given in the explant PCB culture conditions, with the estimated percentage change for each PCB type measured in blubber tissue; Σ DL-PCB within total dose: 1.6 ng; Σ NDL-PCB within total dose: 202.6 ng; SI Figure S1, glucose uptake in inner and outer blubber tissue from feeding and fasting gray seal pups (*n* = 27 feeding and *n* = 23 fasting pups) with median, upper, and lower quartiles; 1.5 $\times$ interquartile range and outliers shown; significant differences between groups are detailed in the main text, as not all significant parameters and interactions influencing glucose uptake are plotted; individual was included as a random effect in the model; SI Figure S2, lactate production in inner and outer blubber tissue from feeding and fasting gray seal pups (*n* = 27 feeding and *n* = 23 fasting pups) with median, upper, and lower quartiles, 1.5 $\times$ interquartile range and outliers shown; significant differences between groups are detailed in the main text, as not all significant parameters and interactions influencing lactate generation/uptake are plotted; individual was included as a random effect in the model; SI Figure S3, glycerol generation in inner and outer blubber tissue from feeding and fasting gray seal pups (*n* = 27 feeding and *n* = 23 fasting pups) with median, upper, and lower quartiles, 1.5 $\times$ interquartile range and outliers shown; significant differences between groups are detailed in the main text, as not all significant parameters and interactions influencing glycerol generation are plotted; and individual was included as a random effect in the model (PDF)

AUTHOR INFORMATION

Corresponding Author

*E-mail: kjr33@st-andrews.ac.uk.

ORCID

Kelly J. Robinson: 0000-0002-6212-9710

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We would like to thank Simon Moss, Holly Armstrong, and everyone who assisted with colony observations and sample collection on the Isle of May during the 2016 season. We would like to thank Scottish National Heritage for the permit to work on the Isle of May and the skipper and crew of the Osprey and May Princess for help with gear and personnel transport. We would also like to thank Dr. Georges Scholl, Catherine Adam, and all of the staff at the University of Liège who provided training and assistance while K.J.R. was in the lab processing samples for POP detection, and Dr. James Turton at Abertay University for advice on adipocyte tissue cell culture techniques. The long term program of research on grey seals at the Isle of May was funded by the SMRU national capability grant (SMRU 1001). K.A.B. was funded by NERC

grant NE/M013723/1 and A.J.H. and K.J.R. were funded by NE/M01357X/1 for this project. The funding bodies had no role in the design of the study, the collection of samples, the analysis of samples or data, and the interpretation of the data.

REFERENCES

- (1) Hop, H.; Borgå, K.; Gabrielsen, G. W.; Kleivane, L.; Skaare, J. U. Food web magnification of persistent organic pollutants in poikilotherms and homeotherms from the Barents Sea. *Environ. Sci. Technol.* **2002**, *36*, 2589–2597.
- (2) Ashraf, M. A. Persistent organic pollutants; a global issue, a global challenge. *Environ. Sci. Pollut. Res.* **2017**, *24*, 4223–4227.
- (3) Van den Berg, M.; Birnbaum, L. S.; Denison, M.; De Vito, M.; Farland, W.; Feeley, M.; Fiedler, H.; Hakansson, H.; Hanberg, A.; Haws, L.; Rose, M.; et al. The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol. Sci.* **2006**, *93*, 223–241.
- (4) Johnson-Restrepo, B.; Kannan, K.; Rapaport, D. P.; Rodan, B. D. Polybrominated diphenyl ethers and polychlorinated biphenyls in human adipose tissue from New York. *Environ. Sci. Technol.* **2005**, *39*, 5177–5182.
- (5) Jones, K. C.; De Voogt, P. Persistent organic pollutants (POPs): state of the science. *Environ. Pollut.* **1999**, *100*, 209–221.
- (6) Fleet, R. R.; Plapp, F. W., Jr DDT residues in snakes decline since the DDT ban. *Bull. Environ. Contam. Toxicol.* **1978**, *19*, 383–388.
- (7) ATSDR (Agency for Toxic Substances and Disease Registry). Toxicological Profile for Polychlorinated Biphenyls. US Department of Health and Human Services, Public Health Services: Atlanta, GA, 2000; <https://www.atsdr.cdc.gov/toxprofiles/tp.asp?id=142&tid=26#bookmark12>.
- (8) Connell, D. W.; Miller, G.; Anderson, S. Chlorohydrocarbon pesticides in the Australian marine environment after banning in the period from the 1970s to 1980s. *Mar. Pollut. Bull.* **2002**, *45*, 78–83.
- (9) MacGregor, K.; Oliver, I. W.; Harris, L.; Ridgway, I. M. Persistent organic pollutants (PCB; DDT;HCH; HCB; and BDE) in eels (*Anguilla anguilla*) in Scotland: current levels and temporal trends. *Environ. Pollut.* **2010**, *158*, 2402–2411.
- (10) French, T. D.; Campbell, L. M.; Jackson, D. A.; Casselman, J. M.; Scheider, W. A.; Hayton, A. Long term changes in legacy trace organic contaminants and mercury in Lake Ontario salmon in relation to source controls trophodynamics and climatic variability. *Limnol. Oceanogr.* **2006**, *51*, 2794–2807.
- (11) Aguilar, A.; Borrell, A. DDT and PCB reduction in the western Mediterranean from 1987 to 2002, as shown by levels in striped dolphins (*Stenella coeruleoalba*). *Mar. Environ. Res.* **2005**, *59*, 391–404.
- (12) Mahmood, M.; Bhavsar, S. P.; Arhonditsis, G. B. Examination of temporal trends in Lake Erie fish communities using dynamic linear modelling. *J. Great Lakes Res.* **2013**, *39*, 437–448.
- (13) Sutton, R.; Sedlak, M. D.; Yee, D.; Davis, J. A.; Crane, D.; Grace, R.; Arsem, N. Declines in polybrominated di-ethyl ether contamination of San Francisco Bay following production phase-outs and bans. *Environ. Sci. Technol.* **2015**, *49*, 777–784.
- (14) Ross, P. S.; Ellis, G. M.; Ikononou, M. G.; Barrett-Lennard, L. G.; Addison, R. F. High PCB concentrations in free-ranging Pacific killer whales, *Orcinus orca*: effects of age, sex and dietary preference. *Mar. Pollut. Bull.* **2000**, *40*, 504–515.
- (15) Goerke, H.; Weber, K.; Bornemann, H.; Ramdohr, S.; Plötz, J. Increasing levels and biomagnification of persistent organic pollutants (POPs) in Antarctic biota. *Mar. Pollut. Bull.* **2004**, *48*, 295–302.
- (16) Desforges, J. P.; Hall, A.; McConnell, B.; Rosing-Asvid, A.; Barber, J. L.; Brownlow, A.; De Guise, S.; Eulaers, I.; Jepson, P. D.; Letcher, R. J.; Levin, M.; et al. Predicting global killer whale population collapse from PCB pollution. *Science* **2018**, *361*, 1373–1376.
- (17) Darnierud, P. O. Toxic effects of brominated flame retardants on man and wildlife. *Environ. Int.* **2003**, *29*, 841–853.

- (18) Toft, G.; Hagmar, L.; Giwercman, A.; Bonde, J. P. Epidemiological evidence of reproductive effects of persistent organochlorines in humans. *Reprod. Toxicol.* **2004**, *19*, 5–26.
- (19) Fisk, A. T.; de Wit, C. A.; Wayland, M.; Kuzyk, Z. Z.; Burgess, N.; Letcher, R.; Braune, B.; Norstrom, R.; Blum, S. P.; Sandau, C.; Lie, E.; Larsen, H. J. S.; Skaare, J. U.; Muir, D. C. G. An assessment of the toxicological significance of anthropogenic contaminants in Canadian arctic wildlife. *Sci. Total Environ.* **2005**, 351–352, 57–93.
- (20) Hammond, J. A.; Hall, A. J.; Dyrinda, E. A. Comparison of polychlorinated biphenyl (PCB) induced effects on innate immune functions in harbour and grey seals. *Aquat. Toxicol.* **2005**, *74*, 126–138.
- (21) Bonde, J. P.; Toft, G.; Rylander, L.; Rignell-Hydbom, A.; Giwercman, A.; Spano, M.; Manicardi, G. C.; Bizzaro, D.; Ludwicki, J. K.; Zvyezday, V.; Bonefeld-Jorgensen, E. C.; Pedersen, H. S.; Joensson, B. A. G.; Thulstrup, A. M. Fertility and markers of male reproductive function in Inuit and European populations spanning large contrasts in blood levels of persistent organochlorines. *Environ. Health Perspect.* **2007**, *116*, 269–277.
- (22) Chiu, A.; Chiu, N.; Beaubier, N. T.; Beaubier, J.; Nalesnik, R.; Singh, D.; Hill, W. R.; Lau, C.; Riebow, J. Effects and mechanisms of PCB ecotoxicity in food chains: Algae > fish > seal > polar bear. *J. Environ. Sci. Health C* **2000**, *18*, 127–152.
- (23) Kramer, S.; Hikel, S. M.; Adams, K.; Hinds, D.; Moon, K. Current status of epidemiological evidence linking polychlorinated biphenyls and non-hodgkin lymphoma, and the role of immune dysregulation. *Environ. Health Perspect.* **2012**, *120*, 1067–1075.
- (24) Czerska, M.; Zieliński, M.; Kaminska, J.; Ligocka, D. Effects of polybrominated diethyl ethers on thyroid hormones, neurodevelopment and fertility in humans and rodents. *Int. J. Occupat. Med. Environ. Health* **2013**, *26*, 498–510.
- (25) Grün, F.; Blumberg, B. Endocrine disruptors as obesogens. *Mol. Cell. Endocrinol.* **2009**, *304*, 19–29.
- (26) Regnier, S. M.; Sargis, R. M. Adipocytes under assault: environmental disruption of adipose physiology. *Biochim. Biophys. Acta, Mol. Basis Dis.* **2014**, *1842*, 520–533.
- (27) Yadetie, F.; Karlsen, O. A.; Eide, M.; Hogstrand, C.; Goksøyr, A. Liver transcriptome analysis of Atlantic cod (*Gadus morhua*) exposed to PCB 153 indicates effects on cell cycle regulation and lipid metabolism. *BMC Genomics* **2014**, *15*, 481.
- (28) Yadetie, F.; Oveland, E.; Doskeland, A.; Berven, F.; Goksøyr, A.; Karlsen, O. A. Quantitative proteomics analysis reveals perturbation of lipid metabolic pathways in the liver of Atlantic cod (*Gadus morhua*) treated with PCB 153. *Aquat. Toxicol.* **2017**, *185*, 19–28.
- (29) Speranza, E. D.; Colombo, M.; Tatone, L. M.; Cappelletti, N.; Migoya, M. C.; Colombo, J. C. Fatty acid alterations in the detritivorous *Prochilodus lineatus* promoted by opportunistic feeding on sewage discharges in the Rio de la Plata estuary. *J. Fish Biol.* **2016**, *89*, 2024–2037.
- (30) Lee, M. C.; Han, J.; Lee, S. H.; Kim, D. H.; Kang, H. M.; Won, E. J.; Hwang, D. S.; Park, J. C.; Om, A. S.; Lee, J. S. A brominated flame retardant 2, 2', 4, 4' tetrabrominated diphenyl ether (BDE-47) leads to lipogenesis in the copepod *Tigriopus japonicus*. *Aquat. Toxicol.* **2016**, *178*, 19–26.
- (31) Maisano, M.; Cappello, T.; Oliva, S.; Natalotto, A.; Giannetto, A.; Parrino, V.; Battaglia, P.; Romeo, T.; Salvo, A.; Spano, N.; Mauceri, A. PCB and OCP accumulation and evidence of hepatic alteration in the Atlantic bluefin tuna, *T. thynnus*, from the Mediterranean Sea. *Mar. Environ. Res.* **2016**, *121*, 40–48.
- (32) Schwetz, B. A.; Norris, J. M.; Sparschu, G. L.; Rowe, V. K.; Gehring, P. J.; Emerson, J. L.; Gerbig, C. G. Toxicology of chlorinated dibenzo-p-dioxins. *Environ. Health Perspect.* **1973**, *5*, 87–99.
- (33) Clark, D. R., Jr; Prouty, R. M. Experimental feeding of DDE and PCB to female big brown bats (*Eptesicus fuscus*). *J. Toxicol. Environ. Health* **1977**, *2*, 917–928.
- (34) Clark, D. R., Jr; Stafford, C. J. Effects of DDE and PCB (Aroclor 1260) on experimentally poisoned female little brown bats (*Myotis Lucifugus*): Lethal brain concentrations. *J. Toxicol. Environ. Health* **1981**, *7*, 925–934.
- (35) Swanepoel, R. M.; Racey, P. A.; Shore, R. F.; Speakman, J. R. Energetic effects of sublethal exposure to lindane on pipistrelle bats (*Pipistrellus pipistrellus*). *Environ. Pollut.* **1999**, *104*, 169–177.
- (36) Young, R. M. Fat, energy and mammalian survival. *Am. Zool.* **1976**, *16*, 699–710.
- (37) Pomeroy, P. P.; Fedak, M. A.; Rothery, P.; Anderson, S. Consequences of maternal size for reproductive expenditure and pupping success of grey seals at North Rona, Scotland. *J. Anim. Ecol.* **1999**, *68*, 235–253.
- (38) Hall, A. J.; McConnell, B. J.; Barker, R. J. The effect of total immunoglobulin levels, mass and condition on the first-year survival of grey seal pups. *Funct. Ecol.* **2002**, *16*, 462–474.
- (39) Lidgard, D. C.; Boness, D. J.; Bowen, W. D.; McMillan, J. I. State dependent male mating tactics in the grey seal: the importance of body size. *Behav. Ecol.* **2005**, *16*, 541–549.
- (40) Irving, L.; Hart, J. S. The metabolism and insulation of seals as bare-skinned mammals in cold water. *Can. J. Zool.* **1957**, *35*, 497–511.
- (41) Rosen, D. A.; Renouf, D. Seasonal changes in blubber distribution in Atlantic harbor seals: indications of thermodynamic considerations. *Mar. Mammal Sci.* **1997**, *13*, 229–240.
- (42) Reilly, J. J. Adaptions to prolonged fasting in free-living weaned gray seal pups. *Am. J. Physiol.* **1991**, *260*, R267–R272.
- (43) Bennett, K. A.; Speakman, J. R.; Moss, S. E.; Pomeroy, P.; Fedak, M. A. Effects of mass and body composition on fasting fuel utilisation in grey seal pups (*Halichoerus grypus fabricius*): an experimental study using supplementary feeding. *J. Exp. Biol.* **2007**, *210*, 3043–3053.
- (44) Waugh, C. A.; Nichols, P. D.; Noad, M. C.; Nash, S. B. Lipid and fatty acid profiles of migrating Southern Hemisphere humpback whales *Megaptera novaeangliae*. *Mar. Ecol.: Prog. Ser.* **2012**, *471*, 271–281.
- (45) Irvine, L. G.; Thums, M.; Hanson, C. E.; McMahon, C. R.; Hindell, M. A. Quantifying the energy stores of capital breeding humpback whales and income breeding sperm whales using historical whaling records. *R. Soc. Open Sci.* **2017**, *4*, 160290.
- (46) Oftedal, O. T. The adaptation of milk secretion to the constraints of fasting in bears, seals, and baleen whales. *J. Dairy Sci.* **1993**, *76*, 3234–3246.
- (47) Debier, C.; Pomeroy, P. P.; Dupont, C.; Joiris, C.; Comblin, V.; Le Boulengé, E.; Larondelle, Y.; Thomé, J. P. Quantitative dynamics of PCB transfer from mother to pup during lactation in UK grey seals *Halichoerus grypus*. *Mar. Ecol.: Prog. Ser.* **2003**, *247*, 237–248.
- (48) Hall, A. J.; Gulland, F. M. D.; Ylitalo, G. M.; Greig, D. J.; Lowenstine, L. Changes in blubber contaminant concentrations in California sea lions (*Zalophus californianus*) associated with weight loss and gain during rehabilitation. *Environ. Sci. Technol.* **2008**, *42*, 4181–4187.
- (49) Tartu, S.; Bourgeon, S.; Aars, J.; Andersen, M.; Polder, A.; Thiemann, G. W.; Welker, J. M.; Routti, H. Sea ice-associated decline in body condition leads to increased concentrations of lipophilic pollutants in polar bears (*Ursus maritimus*) from Svalbard, Norway. *Sci. Total Environ.* **2017**, *576*, 409–419.
- (50) Debier, C.; Pomeroy, P. P.; Dupont, C.; Joiris, C.; Comblin, V.; Le Boulengé, E.; Larondelle, Y.; Thomé, J. P. Dynamics of PCB transfer from mother to pup during lactation in UK grey seals *Halichoerus grypus*: differences in PCB profile between compartments of transfer and changes during the lactation period. *Mar. Ecol.: Prog. Ser.* **2003**, *247*, 249–256.
- (51) Debier, C.; Chalon, C.; Le Boeuf, B. J.; De Tillesse, T.; Larondelle, J. P.; Thome, J.-P. Mobilization of PCBs from blubber to blood in northern elephant seals (*Mirounga angustirostris*) during the post-weaning fast. *Aquat. Toxicol.* **2006**, *80*, 149–157.
- (52) Peterson, S. H.; Hassrick, J. L.; Lafontaine, A.; Thomé, J. P.; Crocker, D. E.; Costa, D. P.; Debier, C. Effects of age, adipose percent, and reproduction on PCB concentrations and profiles in an extreme fasting North Pacific marine mammal. *PLoS One* **2014**, *9*, No. e96191.

- (53) Louis, C.; Tift, M.; Crocker, D. E.; Alexander, D.; Smith, D. R.; Debier, C. Isolation of progenitor cells from the blubber of northern elephant seals (*Mirounga angustirostris*) in order to obtain an *in vitro* adipocyte model—preliminary results. *Mar. Mammal Sci.* **2015**, *31*, 764–773.
- (54) Louis, C.; Covaci, A.; Crocker, D. E.; Debier, C. Lipophilicity of PCBs and fatty acids determines their mobilisation from blubber of weaned northern elephant seal pups. *Sci. Total Environ.* **2016**, *541*, 599–602.
- (55) Tartu, S.; Lille-Langøy, R.; Størseth, T. R.; Bourgeon, S.; Brunsvik, A.; Aars, J.; Goksøyr, A.; Jenssen, B. M.; Polder, A.; Thiemann, G. W.; Torget, V.; Routti, H. Multiple-stressor effects in an apex predator: combined influence of pollutants and sea ice decline on lipid metabolism in polar bears. *Sci. Rep.* **2017**, *7*, 16487.
- (56) Aguilar, A.; Borrell, A. Patterns of lipid content and stratification in the blubber of fin whales (*Balaenoptera physalus*). *J. Mammal.* **1990**, *71*, 544–554.
- (57) Koopman, H. N.; Pabst, D. A.; Mclellan, W. A.; Dillaman, R. M.; Read, A. J. Changes in blubber distribution and morphology associated with starvation in the harbor porpoise (*Phocoena phocoena*): evidence for regional differences in blubber structure and function. *Physiol. Biochem. Zool.* **2002**, *75*, 498–512.
- (58) Best, N. J.; Bradshaw, C. J.; Hindell, M. A.; Nichols, P. D. Vertical stratification of fatty acids in the blubber of southern elephant seals (*Mirounga leonina*): implications for diet analysis. *Comp. Biochem. Physiol., Part B: Biochem. Mol. Biol.* **2003**, *134*, 253–263.
- (59) Koopman, H. N. Phylogenetic, ecological, and ontogenetic factors influencing the biochemical structure of the blubber of odontocetes. *Mar. Biol.* **2007**, *151*, 277–291.
- (60) Strandberg, U.; Käkälä, A.; Lydersen, C.; Kovacs, K. M.; Grahnl-Nielsen, O.; Hyvärinen, H.; Käkälä, R. Stratification, composition, and function of marine mammal blubber: the ecology of fatty acids in marine mammals. *Physiol. Biochem. Zool.* **2008**, *81*, 473–485.
- (61) Liwanag, H. E.; Berta, A.; Costa, D. P.; Budge, S. M.; Williams, T. M. Morphological and thermal properties of mammalian insulation: the evolutionary transition to blubber in pinnipeds. *Biol. J. Linn. Soc.* **2012**, *107*, 774–787.
- (62) Brown, T. M.; Ross, P. S.; Reimer, K. J.; Veldhoen, N.; Dangerfield, N. J.; Fisk, A. T.; Helbing, C. PCB related effects thresholds as derived through gene transcript profiles in locally contaminated ringed seals (*Pusa hispida*). *Environ. Sci. Technol.* **2014**, *48*, 12952–12961.
- (63) Brown, T. M.; Hammond, S. A.; Behsaz, B.; Veldhoen, N.; Birol, I.; Helbing, C. De novo assembly of the ringed seal (*Pusa hispida*) blubber transcriptome: a tool that enables identification of molecular health indicators associated with PCB exposure. *Aquat. Toxicol.* **2017**, *185*, 48–57.
- (64) Castelli, M. G.; Rusten, M.; Goksøyr, A.; Routti, H. mRNA expression of genes regulating lipid metabolism in ringed seals (*Pusa hispida*) from differently polluted areas. *Aquat. Toxicol.* **2014**, *146*, 239–246.
- (65) Buckman, A. H.; Veldhoen, N.; Ellis, G.; Ford, J. K. B.; Helbing, C. C.; Ross, P. S. PCB-associated changes in mRNA expression in killer whales (*Orcinus orca*) from the NE Pacific Ocean. *Environ. Sci. Technol.* **2011**, *45*, 10194–10202.
- (66) Routti, H.; Lille-Langøy, R.; Berg, M. K.; Fink, T.; Harju, M.; Kristiansen, K.; Rostkowski, P.; Rusten, M.; Sylte, I.; Øygarden, L.; Goksøyr, A. Environmental chemicals modulate polar bear (*Ursus maritimus*) peroxisome proliferator-activated receptor gamma (PPARG) and adipogenesis *in vitro*. *Environ. Sci. Technol.* **2016**, *50*, 10708–10720.
- (67) Marin, P.; Rebuffe-Scrive, M.; Smith, U.; Bjorntorp, P. Glucose uptake in human adipose tissue. *Metab., Clin. Exp.* **1987**, *36*, 1154–1160.
- (68) Rotondo, F.; Ho-Palma, A. C.; Remesar, X.; Fernández-López, J. A.; del Mar Romeero, M.; Alemany, M.; Romero, M. d. M. Glycerol is synthesized and secreted by adipocytes to dispose of excess glucose, via glycerogenesis and increased acyl-glycerol turnover. *Sci. Rep.* **2017**, *7*, 8983.
- (69) Baker, N. A.; Karounos, M.; English, V.; Fang, J.; Wei, Y.; Stromberg, A.; Sunkara, M.; Morris, A. J.; Swanson, H. I.; Cassis, L. A. Coplanar polychlorinated biphenyls impair glucose homeostasis in lean C57BL/6 mice and mitigate beneficial effects of weight loss on glucose homeostasis in obese mice. *Environ. Health Perspect.* **2013**, *121*, 105.
- (70) Baker, N. A.; English, V.; Sunkara, M.; Morris, A. J.; Pearson, K. J.; Cassis, L. A. Resveratrol protects against polychlorinated biphenyl-mediated impairment of glucose homeostasis in adipocytes. *J. Nutr. Biochem.* **2013**, *24*, 2168–2174.
- (71) Baker, N. A.; Shoemaker, R.; English, V.; Larian, N.; Sunkara, M.; Morris, A. J.; Walker, M.; Yiannikouris, F.; Cassis, L. A. Effects of adipocyte aryl hydrocarbon receptor deficiency on PCB-induced disruption of glucose homeostasis in lean and obese mice. *Environ. Health Perspect.* **2015**, *123*, 944.
- (72) Enan, E.; Matsumura, F. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)-induced changes in glucose transporting activity in guinea pigs, mice and rats *in vivo* and *in vitro*. *J. Biochem. Toxicol.* **1994**, *9*, 97–106.
- (73) Enan, E.; Lasley, B.; Stewart, D.; Overstreet, J.; Vandevort, C. A. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) modulates function of human luteinizing granulosa cells via cAMP signaling and early reduction of glucose transporting activity. *Reprod. Toxicol.* **1996**, *10*, 191–198.
- (74) Hsu, H. F.; Tsou, T. C.; Chao, H. R.; Kuo, Y. T.; Tsai, F. Y.; Yeh, S. V. Effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on adipogenic differentiation and insulin-induced glucose uptake in 3T3-L1 cells. *J. Hazard. Mater.* **2010**, *182*, 649–655.
- (75) Liu, P. C.; Matsumura, F. Differential effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the “adipose-type” and “brain-type” glucose transporters in mice. *Mol. Pharmacol.* **1995**, *47*, 65–73.
- (76) Kern, P. A.; Dicker-Brown, A.; Said, S. T.; Kennedy, R.; Fonseca, V. A. The stimulation of tumor necrosis factor and inhibition of glucose transport and lipoprotein lipase in adipose cells by 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Metab., Clin. Exp.* **2002**, *51*, 65–68.
- (77) Wu, H.; Yu, W.; Meng, F.; Mi, J.; Peng, J.; Liu, J.; Zhang, X.; Hai, C.; Wang, X. Polychlorinated biphenyls 153 induces metabolic dysfunction through activation of ROS/ NF- κ B signalling via downregulation of HNF1b. *Redox Biol.* **2017**, *12*, 300–310.
- (78) Arsenescu, V.; Arsenescu, R. I.; King, V.; Swanson, H.; Cassis, L. A. Polychlorinated biphenyl-77 induces adipocyte differentiation and proinflammatory adipokines and promotes obesity and atherosclerosis. *Environ. Health Perspect.* **2008**, *116*, 761.
- (79) Uemura, H.; Arisawa, K.; Hiyoshi, M.; Satoh, H.; Sumiyoshi, Y.; Morinaga, K.; Kodama, K.; Suzuki, T. I.; Nagai, M.; Suzuki, T. Associations of environmental exposure to dioxins with prevalent diabetes among general inhabitants in Japan. *Environ. Res.* **2008**, *108*, 63–68.
- (80) Uemura, H.; Arisawa, K.; Hiyoshi, M.; Kitayama, A.; Takami, H.; Sawachika, F.; Dakeshita, S.; Nii, K.; Satoh, H.; Sumiyoshi, Y.; Morinaga, K.; et al. Prevalence of metabolic syndrome associated with body burden levels of dioxin and related compounds among Japan’s general population. *Environ. Health Perspect.* **2009**, *117*, 568.
- (81) Everett, C. J.; Mainous, A. G., III; Frithsen, I. L.; Player, M. S.; Matheson, E. M. Association of polychlorinated biphenyls with hypertension in the 1999–2002 National Health and Nutrition Examination Survey. *Environ. Res.* **2008**, *108*, 94–97.
- (82) Camón, L.; Solà, C.; Martínez, E.; Sanfeliu, C.; Rodríguez-Farré, E. Cerebral glucose uptake in lindane-treated rats. *Toxicology* **1988**, *49*, 381–387.
- (83) Rivera, S.; Sanfeliu, C.; Rodríguez-Farré, E. Changes in regional brain 2[¹⁴C]deoxyglucose uptake induced in postnatal developing rats by single and repeated nonconvulsant doses of lindane. *Pestic. Biochem. Physiol.* **1992**, *43*, 241–252.
- (84) Mahmood, A.; Agarwal, N.; Sanyal, S.; Subrahmanyam, D. Effects of DDT (Chlorophenotane) Administration on Glucose Uptake and Brush Border Enzymes in Monkey Intestine. *Acta Pharmacol. Toxicol.* **1978**, *43*, 99–102.

- (85) Norberto, S.; Calhau, C.; Pestana, D.; Faria, A. Effects of Environmental Pollutants on MCF-7 Cells: A Metabolic Approach. *J. Cell. Biochem.* **2017**, *118*, 366–375.
- (86) Howell, G. E., III; Meek, E.; Kilic, J.; Mohns, M.; Mulligan, C.; Chambers, J. E. Exposure to p,p'-dichlorodiphenyldichloroethylene (DDE) induces fasting hyperglycemia without insulin resistance in male C57BL/6H mice. *Toxicology* **2014**, *320*, 6–14.
- (87) Ruzzin, J.; Petersen, R.; Meugnier, E.; Madsen, L.; Lock, E. J.; Lillefosse, H.; Ma, T.; Pesenti, S.; Sonne, S. B.; Marstrand, T. T.; Malde, M. K.; et al. Persistent organic pollutant exposure leads to insulin resistance syndrome. *Environ. Health Perspect.* **2010**, *118*, 465.
- (88) Langin, D. Control of fatty acid and glycerol release in adipose tissue lipolysis. *C. R. Biol.* **2006**, *329*, 598–607.
- (89) Vaughan, M. The production and release of glycerol by adipose tissue incubated *in vitro*. *J. Biol. Chem.* **1962**, *237*, 3354–3358.
- (90) Chakrabarty, K.; Bombeck, C. T.; Sigel, B.; Tauber, T. W.; Jeffay, H. Glycerokinase activity in human adipose-tissue as related to obesity. *Int. J. Obesity* **1984**, *8*, 609–622.
- (91) Bennett, K. A.; Robinson, K. J.; Moss, S. E.; Millward, S.; Hall, A. J. Using blubber explants to investigate adipose function in grey seals: glycolytic, lipolytic and gene expression responses to glucose and hydrocortisone. *Sci. Rep.* **2017**, *7*, 7731.
- (92) Pond, C. M.; Mattacks, C. A. The distribution, cellular structure, and metabolism of adipose tissue in the fin whale, *Balaenoptera physalus*. *Can. J. Zool.* **1988**, *66*, 534–537.
- (93) Robinson, K. J.; Twiss, S. D.; Hazon, N.; Moss, S.; Pomeroy, P. P. Positive social behaviours are induced and retained after oxytocin manipulations mimicking endogenous concentrations in a wild mammal. *Proc. R. Soc. London, Ser. B* **2017**, *284*, 20170554.
- (94) Bennett, K. A.; Hughes, J.; Stamatas, S.; Brand, S.; Foster, N. L.; Moss, S. E. W.; Pomeroy, P. P. Adiponectin and insulin in gray seals during suckling and fasting: relationship with nutritional state and body mass during nursing in mothers and pups. *Physiol. Biochem. Zool.* **2015**, *88*, 295–310.
- (95) Cold Spring Harbor Protocol. Krebs-Ringer Modified Buffer (KRB) **2014**, DOI: [10.1101/pdb.rec076406](https://doi.org/10.1101/pdb.rec076406).
- (96) Chaise, L. L.; Paterson, W.; Laske, T. G.; Gallon, S. L.; McCafferty, D. J.; Théry, M.; Ancel, A.; Gilbert, C. Implantation of subcutaneous heart rate data loggers in southern elephant seals (*Mirounga leonina*). *Polar Biol.* **2017**, *40*, 2307–2312.
- (97) Etherton, T. D.; Evock, C. M. Stimulation of lipogenesis in bovine adipose tissue by insulin and insulin-like growth factor. *J. Anim. Sci.* **1986**, *62*, 357–362.
- (98) May, S. G.; Savell, J. W.; Lunt, D. K.; Wilson, J. J.; Laurenz, J. C.; Smith, S. B. Evidence for preadipocyte proliferation during culture of subcutaneous and intramuscular adipose tissues from Angus and Wagyu crossbred steers. *J. Anim. Sci.* **1994**, *72*, 3110–3117.
- (99) R Development Core Team. R: A language and environment for statistical computing. R Foundation for Statistics Computing, Vienna, Austria. 2012, <http://www.R-project.org>.
- (100) Wood, S. *Generalized Additive Models: An Introduction with R*. Chapman and Hall/CRC: London, U.K., 2006.
- (101) Wood, S. mgcv: Mixed GAM Computation Vehicle with GCV/AIC/REML smoothness estimation. 2012, <http://cran.r-project.org/web/packages/mgcv/index.html>.
- (102) Wood, S. Low-rank scale-invariant tensor product smooths for generalized additive mixed models. *Biometrics* **2006**, *62*, 1025–1036.
- (103) Wood, S. Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *J. R. Stat. Soc. B* **2011**, *73*, 3–36.
- (104) Lee, D. H.; Porta, M.; Jacobs, D. R., Jr; Vandenberg, L. N. Chlorinated persistent organic pollutants, obesity, and type 2 diabetes. *Endocr. Rev.* **2014**, *35*, 557–601.
- (105) Desforges, J. P.; Levin, M.; Jasperse, L.; De Guise, S.; Eulaers, I.; Letcher, R. J.; Acquarone, M.; Nordøy, E.; Folkow, L. P.; Hammer Jensen, T.; Grøndahl, C.; et al. Effects of polar bear and killer whale derived contaminant cocktails on marine mammal immunity. *Environ. Sci. Technol.* **2017**, *51*, 11431–11439.
- (106) Totland, C.; Nerdal, W.; Steinkopf, S. Effects and location of coplanar and noncoplanar PCB in a lipid bilayer: a solid state NMR study. *Environ. Sci. Technol.* **2016**, *50*, 8290–8295.
- (107) Lim, J. S.; Lee, D. H.; Jacobs, D. R. Association of brominated flame retardants with diabetes and metabolic syndrome in the US population, 2003–2004. *Diabetes Care* **2008**, *31*, 1802–1807.
- (108) Hall, A. J.; Kalantzi, O. I.; Thomas, G. O. Polybrominated diphenyl ethers (PBDEs) in grey seals during their first year of life—are they thyroid hormone endocrine disruptors? *Environ. Pollut.* **2003**, *126*, 29–37.
- (109) Hall, A. J.; Thomas, G. O.; McConnell, B. J. Exposure to persistent organic pollutants and first-year survival probability in gray seal pups. *Environ. Sci. Technol.* **2009**, *43*, 6364–6369.
- (110) Nordøy, E. S.; Ingebretsen, O. C.; Blix, A. S. Depressed metabolism and low protein catabolism in fasting grey seal pups. *Acta Physiol. Scand.* **1990**, *139*, 361–369.
- (111) Bryant, N.; Govers, R.; James, D. Regulated transport of the glucose transporter GLUT4. *Nat. Rev. Mol. Cell Biol.* **2002**, *3*, 267–277.
- (112) Fowler, M. A.; Champagne, C. D.; Houser, D. S.; Crocker, D. E. Hormonal regulation of glucose clearance in lactating northern elephant seals (*Mirounga angustirostris*). *J. Exp. Biol.* **2008**, *211*, 2943–2949.
- (113) Houser, D. S.; Champagne, C. D.; Crocker, D. E. A non-traditional model of the metabolic syndrome: the adaptive significance of insulin resistance in fasting-adapted seals. *Front. Endocrinol.* **2013**, *4*, 164.
- (114) Viscarra, J. A.; Champagne, C. D.; Crocker, D. E.; Ortiz, R. M. 5' AMP-activated protein kinase activity is increased in adipose tissue of northern elephant seal pups during prolonged fasting-induced insulin resistance. *J. Endocrinol.* **2011**, *209*, 317–325.
- (115) Vanden Berghe, M.; Weijs, L.; Habran, S.; Das, K.; Bugli, C.; Rees, J. F.; Pomeroy, P.; Covaci, A.; Debier, C. Selective transfer of persistent organic pollutants and their metabolites in grey seals during lactation. *Environ. Int.* **2012**, *46*, 6–15.
- (116) Hall, A. J.; McConnell, B. J.; Barker, R. J. Factors affecting first-year survival in grey seals and their implications for life history strategy. *J. Animal. J. Anim. Ecol.* **2001**, *70*, 138–149.
- (117) Shimba, S.; Todoroki, K.; Aoyagi, T.; Tezuka, M. Depletion of aryl-hydrocarbon receptor during adipose differentiation in 3T3-L1 cells. *Biochem. Biophys. Res. Commun.* **1998**, *249*, 131–137.
- (118) Shimba, S.; Wada, T.; Tezuka, M. Arylhydrocarbon receptor (AhR) is involved in negative regulation of adipose differentiation in 3T3-L1 cells: AhR inhibits adipose differentiation independently of dioxin. *J. Cell. Sci.* **2001**, *114*, 2809–2817.
- (119) Sethi, J. K.; Vidal-Puig, A. J. Adipose Tissue Development, Structure and Function. In *Metabolic Basis of Obesity*; Ahima, R., Eds.; Springer, New York, NY, 2011.
- (120) Chesney, C. F.; Allen, J. R. Oxidative phosphorylation and respiration by liver mitochondria from polychlorinated biphenyl-intoxicated rats. *Biochem. Pharmacol.* **1974**, *23*, 1577–1582.
- (121) Nishihara, Y. Effects of polychlorinated biphenyls (Kanechlor 400) on isolated rat liver mitochondria. *Arch. Environ. Contam. Toxicol.* **1983**, *12*, 517–522.
- (122) Arriarán, S.; Agnelli, S.; Sabater, D.; Remesar, X.; Fernández-López, J. A.; Alemany, M. Evidences of basal lactate production in the main white adipose tissue sites of rats. Effects of sex and a cafeteria diet. *PLoS One* **2015**, *10*, e0119572.
- (123) Kolataj, A.; Rysińska, J.; Flak, P. The influence of selection on reaction to stress in mice. III. Influence of fasting, immobilization and exposure to cold on lactate dehydrogenase and aldolase activity in the liver and kidney. *J. Anim. Breed. Genet.* **1995**, *112*, 224–233.
- (124) Rao, C. V.; Banerji, S. A. Effect of feeding polychlorinated biphenyl (Aroclor 1260) on hepatic enzymes of rats. *Indian J. Exp. Biol.* **1990**, *28*, 149–151.
- (125) Sonne, C.; Leifsson, P. S.; Dietz, R.; Kirkegaard, M.; Jensen, A. L.; Shahmiri, S.; Letcher, R. J. Greenland sledge dogs (*Canis familiaris*) develop liver lesions when exposed to a chronic and dietary

low dose of an environmental organohalogen cocktail. *Environ. Res.* **2008**, *106*, 72–80.

(126) King, J. L.; DiGirolamo, M. Lactate production from glucose and response to insulin in perfused adipocytes from mesenteric and epididymal regions of lean and obese rats. *Obes. Res.* **1998**, *6*, 69–75.

(127) Beck, C. A.; Bowen, W. D.; Iverson, S. J. Sex differences in the seasonal patterns of energy storage and expenditure in a phocid seal. *J. Anim. Ecol.* **2003**, *72*, 280–291.

(128) Kelso, E. J.; Champagne, C. D.; Tift, M. S.; Houser, D. S.; Crocker, D. E. Sex differences in fuel use and metabolism during development in juvenile northern elephant seals. *J. Exp. Biol.* **2012**, *215*, 2637–2645.

(129) Katz, J.; Wals, P. A. Lipogenesis from lactate in rat adipose tissue. *Biochim. Biophys. Acta, Lipids Lipid Metab.* **1974**, *348*, 344–356.

(130) Roth, W. L.; Weber, L. W. D.; Stahl, B. U.; Rozman, K. A pharmacodynamic model of triglyceride transport and deposition during feed deprivation or following treatment with 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in the rat. *Toxicol. Appl. Pharmacol.* **1993**, *120*, 126–137.

(131) Rao, T. N.; Subrahmanyam, D. Effect of DDT on adipose tissue lipolysis in rats. *Toxicology* **1981**, *20*, 229–235.

(132) Howell, G. E., III; Mangum, L. Exposure to bioaccumulative organochlorine compounds alters adipogenesis, fatty acid uptake, and adipokine production in NIH3T3-L1 cells. *Toxicol. In Vitro* **2011**, *25*, 394–402.

(133) Sethi, J. K.; Vidal-Puig, A. J. Thematic review series: adipocyte biology. Adipose tissue function and plasticity orchestrate nutritional adaptation. *J. Lipid Res.* **2007**, *48*, 1253–1262.

(134) Duncan, R. E.; Ahmadian, M.; Jaworski, K.; Sarkadi-Nagy, E.; Sul, H. S. Regulation of lipolysis in adipocytes. *Annu. Rev. Nutr.* **2007**, *27*, 79–101.

(135) De Koster, J.; Van den Broeck, W.; Hulpio, L.; Claeys, E.; Van Eetvelde, M.; Hermans, K.; Hostens, M.; Fievez, V.; Opsomer, G. Influence of adipocyte size and adipose depot on the in vitro lipolytic activity and insulin sensitivity of adipose tissue in dairy cows at the end of the dry period. *J. Dairy Sci.* **2016**, *99*, 2319–2328.

(136) Orevi, M.; Gorin, E.; Shafrir, E. Adaptive changes in phosphofructokinase and aldolase in adipose tissue. *Eur. J. Biochem.* **1972**, *30*, 418–426.

(137) Romero, M. d. M.; Sabater, D.; Fernández-López, J. A.; Remesar, X.; Alemany, M. Glycerol production from glucose and fructose by 3T3-L1 cells: a mechanism of adipocyte defence from excess substrate. *PLoS One* **2015**, *10*, e0139502.

(138) Regehr, E. V.; Lunn, N. J.; Amstrup, S. C.; Stirling, I. A. N. Effects of earlier sea ice breakup on survival and population size of polar bears in western Hudson Bay. *J. Wildl. Manage.* **2007**, *71*, 2673–2683.

(139) Fa, J. E.; Ryan, S. F.; Bell, D. J. Hunting vulnerability, ecological characteristics and harvest rates of bushmeat species in afro-tropical forests. *Biol. Conserv.* **2005**, *121*, 167–176.

(140) Cheung, W. W.; Sarmiento, J. L.; Dunne, J.; Frölicher, T. L.; Lam, V. W.; Palomares, M. L. D.; Watson, R.; Pauly, D. Shrinking of fishes exacerbates impacts of global ocean changes on marine ecosystems. *Nat. Clim. Change* **2013**, *3*, 254–258.

(141) Mathews, E. A.; Keller, S.; Weiner, D. B. A method to collect and process skin biopsies for cell culture from free-ranging gray whales (*Eschrichtius robustus*). *Mar. Mammal Sci.* **1988**, *4*, 1–12.

(142) Marsili, L.; Fossi, M. C.; Neri, G.; Casini, S.; Gardi, C.; Palmeri, S.; Tarquini, E.; Panigada, S. Skin biopsies for cell cultures from Mediterranean free-ranging cetaceans. *Mar. Environ. Res.* **2000**, *50*, 523–526.

(143) Tovar, H.; Navarrete, F.; Rodríguez, L.; Skewes, O.; Castro, F. O. Cold storage of biopsies from wild endangered native Chilean species in field conditions and subsequent isolation of primary culture cell lines. *In Vitro Cell. Dev. Biol.: Anim.* **2008**, *44*, 309–320.