

Potential of engineering the myo-inositol oxidation pathway to increase stress resilience in plants

Anshu Alok^{1,2*}, Sadhana Singh², Pankaj Kumar³, Kaushal K. Bhati^{4*}

1. Department of Biotechnology, UIET, Punjab University, Chandigarh, India

2. Department of Plant Pathology, University of Minnesota, Saint Paul, MN 55108, USA

3. International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru, India

4. School of Agricultural Biotechnology, Punjab Agricultural University, Ludhiana, India

5. Louvain Institute of Biomolecular Sciences, Catholic University of Louvain, Louvain-la-Neuve, Belgium

*corresponding authors: Anshu Alok: anshualok2@gmail.com

Kaushal K. Bhati: Kaushal.bhati@uclouvain.be

Abstract

Myo-inositol is one of the most abundant form of inositol. The myo-inositol (MI) serves as substrates to diverse biosynthesis pathways and hence it is conserved across life forms. The biosynthesis of MI is well studied in animals. Beyond biosynthesis pathway, implications of MI pathway and enzymes hold potential implications in plant physiology and crop improvement. *Myo*-inositol oxygenase (MIOX) enzyme catabolized MI into D-glucuronic acid (D-GlcUA). The MIOX enzyme family is well studied across higher plants. More recently, the MI associated pathway's crosstalk with other important biosynthesis and stress responsive pathways in plants has drawn attention. The overall outcome from different plant species studied so far are very suggestive that MI derivatives and associated pathways could open new directions to explore stress responsive novel metabolic networks. There are evidences for upregulation of MI metabolic pathway genes, specially *MIOX* under different stress condition. We also found *MIOX* genes getting differentially expressed according to developmental and stress signals in

Arabidopsis and wheat. In this review we try to highlight the missing links and put forward a tailored view over myo-inositol oxidation pathway and MIOX proteins.

Introduction

Inositols are sugar-like carbohydrates that exist as seven isomeric forms in plant cells. Among these isomers, the most abundant *myo*-inositol (MI) could exist as free or bound form with phospholipids or inositol phosphate derivatives. The biosynthesis of MI generally starts with Glucose-6-phosphate; however, it could also synthesize from other inositol phosphates. Glucose-6-phosphate catalyzed into *myo*-inositol-1-phosphate by *myo*-inositol-1-phosphate synthase enzyme (MIPS), it is also first and rate-limiting step of biosynthesis of MI [1, 2, 3]. *Myo*-inositol is principally catabolized into D-glucuronic acid (D-GlcUA) by *myo*-inositol oxygenase (MIOX) enzyme (EC 1.13.99.1), which is the first step of this pathway [4]. There is various important functional role of D-GlcUA for MIOX is a unique monooxygenase encoded by *MIOX* gene, ubiquitous to all eukaryotic cells, including those in yeast, animals, humans and plants. Animal genome contains only single *MIOX* gene whereas it is a multigene family in plants. For instance, the genome of Thale cress *Arabidopsis thaliana* contains four *MIOX* genes (*AtMIOX 1, 2, 4, and 5*), numbered according to their chromosomal location [5]. Isoforms of *A. thaliana* *MIOX* are 66-84% identical at the amino acid level and the same catalytic domain is present in all four proteins. In wheat, three *MIOX* homologs are mapped on chromosomes 7A, 7B, and 7D [6]. Recently, five *MIOX* homologs were identified from tomato, on chromosomes 6, 10, 11, and 12 [7].

MI oxidation pathway is well interconnected with other important metabolic pathways. These include, the biosynthetic pathway of MI polyphosphates, pyrophosphates, and phytic acid (PA) [8], cell wall biosynthetic molecules such as galactinol, raffinose, and pinitol etc., Ascorbic acid (AsA) biosynthesis [7], and Phosphoinositides and Ins(1,4,5)P₃ [9] (Figure 1). Additionally, it is also linked with various pathways related to signal transduction and transportation of sugars and auxin [10]. MIOX oxidize MI and therefore directly influence the MI homeostasis. The triple mutant *miox1/2/4/5* in *A. thaliana* has increased MI and MI containing metabolites. MI is

the primary precursor for PA biosynthesis via lipid dependent and independent pathways. The PA content in seeds reduce in *CaMV35::AtMIOX4* over-expression line compared to wild type and *miox1/2/4/5* mutant [11].

During MI oxidation pathway, D-GlcUA is converted into uridine diphosphate (UDP) glucuronic acid by glucurono kinase enzyme. UDP-glucuronic acid (UDP-GlcUA) is an important sugar precursor for various nucleotide sugars need for synthesis of cell wall related polysaccharides. Though, the sugar nucleotide oxidation pathway is primarily used in biosynthesis of pectin (cell-wall polysaccharides), it is also important as substrate (UDP-GlcUA) for the production of cell-wall polysaccharides. Labeling ³H-*myo* inositol and inositol-2-(¹⁴C) in *AtMIOX1* and *AtMIOX2* mutants of *Arabidopsis* showed that approximately 70% of MI converted to D-GlcUA, which is further used in cell wall biosynthesis [12,13,14,5]. Thus, cell wall engineering and stress adaption by altering tissue specific expression of MIOX is an open niche to explore as genetic engineering and synthetic biology approaches.

Ascorbic acid (AsA) biosynthesis via MI oxidation pathway is well characterized and active among mammals. This process involves, D-GlcUA reduced into L-gulonic acid through the action of glucuronic acid reductase, which then finally oxidized to AsA [15,16]. However, whether AsA biosynthesis via MI oxidation pathway exists in plants is still debatable. Identification of D-galacturonate reductase and L-glucose in strawberry (*Fragaria ananassa*) indicate that MI may be an alternative precursor for AsA biosynthesis. Ectopic expression of *D-galacturonate reductase* gene in *Arabidopsis* improved vitamin C content by 2-3 folds [17,18]. Moreover, ectopic expression of the *AtMIOX4* gene in *Arabidopsis* increased AsA content in the transgenic line by two or three times, suggesting a correlation between MI oxidation and AsA biosynthesis [19]. Conversely, Endres and Tenhaken (2009) showed that *AtMIOX4* did not increase the AsA content but, instead control the MI level in *AtMIOX* over-expressed plants [20]. In tomato plants, the over-expression of *MIOX* gene increased AsA content in leaves and ripe fruits compared to control plants [7]. All these finding suggests that engineering myo-inositol pathway could be a good target to improve fruits with higher AsA.

To this end, multifaced role of MIOX and *myo*-inositol oxidation pathway proved to be significant influencers on multiple aspect of plant physiology. Our current work will revisit various research reports to bring a practical perspective on possible aspects of this pathway across broaden areas includes PA biosynthesis, cell wall biosynthesis, ascorbic acid biosynthesis, abiotic, and biotic stresses.

Diversity, gene structure and chromosomal locations

The plant kingdom in general have very discreet genomic diversity among *MIOX* like genes. In *Arabidopsis*, there are four homologs of the *MIOX* gene, labeled *AtMIOX1*, 2, 4, and 5 according to their chromosomal location. All members of this family have 11 exons and 10 introns [11]. The soybean (*Glycine soja*) genome contains five homologs (*GsMIOX1a*, 1*b*, 2*a*, 2*b*, and 4), located on chromosome 1, 7, and 8, respectively [21]. In wheat (*Triticum aestivum*), homeologous *TaMIOX* genes (*Ta7AMIOX*, *Ta7BMIOX*, and *Ta7DMIOX*) are situated on 7A, 7B, and 7D and have 11 exons and 10 introns [6]. In tomato (*Solanum lycopersicum*) five *MIOX* genes identified; *SIMIOX1*, 2, 3, 4, and 5, they are located on chromosomes 6, 10, 11, and 12 respectively. *SIMIOX1*, 2, 3, and 4, contain ten, one, seven and eight introns, respectively, while *SIMIOX5* contains none [7]. Recently, two genes, *MhMIOX1* & 2 were identified in Chinese crab apple (*Malus hupehensis* Rehd.). *MhMIOX1* contain six exons and five introns, whereas the *MhMIOX2* comprised nine exons and eight introns [22]. Diploid and tetraploid cotton contains 6 and 11 *MIOX* genes [23].

Transcriptional regulation of MIOX genes

MIOX promoters express differentially across plant tissues, developmental stages, and under abiotic stresses. Several drought associated transcription factors like MYB, MYB-like, MYC, and Myb-have putative binding sites on wheat *MIOX* promoter (*pro:TaMIOX*). Similar, MYB binding sites are also present in *Arabidopsis MIOX2* and *MIOX4* promoters. Drought, wound and water stress related transcription binding sites such as DRE2COREZMRAB17, T/GBOXATPIN2, and MYBCORE are present in wheat *MIOX* promoter. Dehydration, low temperature and ABA responsive transcription binding sites such as ABRELATERD1, DRECRTCOREAT, MYCATERD1, MYBCORE, ACGTATERD1, MYB1AT, MYCATRD22, LTRECOREATCOR15, CBFHV,

RYREPEATBNNAPA, MYCCONSENSUAT, and DRE2COREZMRAB17 were also found on *TaMIOXpro* [24].

Arabidopsis AtMIOX2 promoter (*AtMIOX2pro*) express constitutively whereas, *AtMIOX4pro* and *AtMIOX5pro* are specific to reproductive stages [5]. *Arabidopsis* transgenic lines expressing *uidA* gene under the control of *AtMIOX2pro*, showed its activity in the roots, cotyledons, developing seeds, anthers, stigma, and stipules. While *AtMIOX4pro::uidA* expression is reported in the primary root tip, sepals, young petals, anthers, top of the pistil, and stipules. The *AtMIOX5pro::uidA* construct, displayed strong GUS activity in pollen grains and the stigma while lower was in petals, sepals, and filament [25]. In another report, microarray data showed *AtMIOX4* and *AtMIOX5* were highly expressed in pollen of *Arabidopsis*, whereas *AtMIOX2* expressed in roots and seedlings in normal condition [26].

Expression of *AtMIOX* genes are also correlated with nutrient status and energy sensing [27]. *AtMIOX2pro* have strong activity in cotyledons, hypocotyl, and roots under nutrient deficient conditions (0.8% agar without Murishige and Skoog salts) while promoter was moderately active in low nutrient condition (half MS salts with 0.8% agar). However, in the absence (only 0.8% agar) of nutrients, expression of *uidA* gene under *AtMIOX4pro* was not detected. GUS activity of *AtMIOX4pro* under low nutrient condition was observed in cotyledons and roots. The strength of *AtMIOX2pro* and *AtMIOX4pro* were decreased in seedlings under optimal nutrient (half MS, 0.8% agar, and 3% glucose) [27]. The *AtMIOX* genes express differentially under biotic stress. When challenged with cyst nematode *Heterodera schachtii* for 10-15 days *proAtMIOX5::uidA* lines showed significant GUS activity in roots [28]. *In-silico* identification and mining of *cis*-regulatory elements present in all *AtMIOXpro*, positively correlates to expressions pattern related to abiotic and biotic stresses [29].

Rice *MIOX* promoter (*OsMIOXpro*) showed its activity in all tissues, including panicles, stems, leaves, and roots. *OsMIOXpro* was highly active under drought stress [30]. The promoter activity of *TaMIOXpro* was 5-7-fold higher in transgenic *Arabidopsis* leaf under abiotic stresses such as heat, cold, and drought as compared to controls. Apart

from this, *TaMIOXpro::uidA* lines showed lesser GUS activity during *myo*-inositol feeding assay of plants [24].

We identified that these *MIOX* genes express in very tissue specific manner. Though, the substantial level of *AtMIOX1* was higher in most tissues as compared to other isoforms in publicly available data (Figure 2A). Similar pattern was also observed for *TaMIOX* genes during developmental stages. Additionally, we tested that wheat *MIOX* genes could express differentially in response to specific environmental triggers (Figure 2B and C).

MIOX proteins and their homology

MIOX is a soluble protein and lacks any signal peptide and transmembrane helix [31,32,5]. Heterologous expression and purification of MIOX from yeast and *Arabidopsis*, showed MIOX enzymatic activity [32,19]. The yeast and *Arabidopsis* MIOX have 57% sequence similarity [5]. In wheat 77% of the protein sequence showed similarity with *AtMIOX1*, whereas this ranges between 68% to 94% with other plants [6]. Amino acids (aa) sequences of MIOX across all kingdom have 1-100 aa as variable region, 100-250 aa are highly conserved and 251-303 aa are less conserved on C-terminal. *TaMIOX* contain 3 substrate binding regions, two substrate binding sites, and six potential iron binding sites [6]. The occurrence of H₁₃₉ and D₁₄₀ aa in *TaMIOX* showed a strong affinity with metal ions and the HD-domain superfamily is well known for strong binding with metal ions [33]. The 3-D structure reveals that MIOX protein is a monomeric with a single-domain and helices. The 3-D crystal structure of MIOX was reported from animal source using crystallization X-ray diffraction. So far, the crystal structure of MIOX from plant has not been reported. In human MIOX proteins comprised eight α -helices and 2 antiparallel β -sheet [34]. Plant MIOX protein comprised nine α -helices, five forming the core and four the protein surface [33].

Interconnection of MI oxidation pathways to phytic acid biosynthesis

PA is the hexa-*kis*-phosphate ester of MI, which accumulates in seeds and is stored in the embryo, endosperm and seed coat. All seed crops contain PA, which may vary from one plant to another but it usually accounts for 65–85% of total phosphorus in seeds [35]. PA strongly binds with iron and other divalent minerals such as Fe, Zn and Mg, there by

limit their bioavailability. This correlation of PA and micronutrients loading to seed is well studied in crops like wheat [36]. There are high chances of possible link between MI and PA biosynthetic pathways as the *miox1/2/4/5* knock-out *Arabidopsis* showed higher PA content in vegetative tissues than wild type. While, overexpression of *AtMIOX4* also showed a decreased PA content in *Arabidopsis* seeds compared to wild type and non-functional mutants [11]. The *1-myo-inositol 1-P synthase (INO1)*, *Inositol monophosphatase (IMP)*, and *myo-inositol O-methyl transferase (IMT)* genes, which are linked with MI pathways, also demonstrated their potential for reducing PA content. Overexpression of *IMT* gene from ice plant (*Mesembryanthemum crystallinum*) under seed-specific promoters into *Brassica napus*, showed 19-35% phytate reduction in seeds [37]. RNAi mediated suppression of wheat *inositol-pentakisphosphate 2-kinase 1 (IPK1)* gene, leading to a 28–56% reduction in the phytic acid content of wheat seeds and an increase in iron and zinc content [38]. Similarly, in rice suppression of *myo-inositol 1-phosphate synthase* resulted in a decrease in PA content but a 1.3, 1.6 and 1.27-fold increase in calcium, iron and magnesium, respectively [39]. Overexpression of *MIOX* result in lower PA in different crops as the free pool of *myo-inositol* is greatly utilized in the conversion of D-Glucuronic acid (D-GlcUA). All these studied suggest that, *MIOX* overexpression might decrease PA content in seed and increase nutrient content in seeds.

Engineering downstream pathways: synthesis of cell wall precursors

The cell wall is a complex and dynamic structure that consists of four major components i.e. polysaccharides cellulose, non-cellulosic polysaccharides (hemicellulose and pectin) and proteins. Various glycosyltransferases promote the synthesis of hemicellulose and pectin in the golgi body. These glycosyltransferases transfer the sugar moiety from an activated nucleotide donor onto acceptors polysaccharide in the form of a UDP or GDP-sugar [40]. D-GlcUA is converted into D-GlcUA-1P through the action of Glucuronokinase (GlcAK), which is enzyme encoded by *Glucuronokinase* gene [3]. D-GlcUA-1P is further converted into UDP-GlcUA by UDP-sugar pyro-phosphorylase (USP). Conversion of sugars into nucleotide sugars is generally initiated by a substrate-specific sugar-1-kinase or USP (broad substrate specificity) [41]. UDP-GlcUA is a primary precursor to numerous

nucleotide sugars involved in biosynthesis of hemicellulose and pectin. These nucleotide sugars are UDP-galacturonic acid, UDP-D-xylose, UDP-arabinose, and UDP-apiose [42]. Cell wall polysaccharides that have nucleotide sugars originated from UDP-GlcUA account for approximately 50% of the cell wall biomass [43,40]. UDP-GlcUA is synthesized and utilized in cytoplasm that is further transported inside Golgi body for biosynthesis of UDP-galacturonic acid, UDP-D-xylose, UDP-arabinose, and UDP-apiose (Figure 3). UDP-Uronic Acid Transporter1 (UUAT1) is a Golgi-localized nucleotide sugar transporter that transports UDP-GlcUA [40]. In *Arabidopsis* and sea onion (*Ornithogalum caudatum*), the enzyme UDP-GlcA 4-epimerase (UGlcAE) is characterized to convert UDP-GlcUA into UDP-D-galacturonic acid [44]. The genes *AtUGlcAE1*, 2, and 3 that are present in the *Arabidopsis* genome encodes UGlcAE enzyme [44]. Decarboxylation of UDP-GlcUA leads to production of UDP-xylose with the help of UDP-glucuronic acid decarboxylase (UXS) enzyme. Further, the enzyme UDP-D-xylose epimerase encoded by *MUR4* gene converts UDP-D-xylose into UDP-L-arabinose [45, 46] The ectopic expression of *AtMIOX2* in tobacco (*Nicotiana tabacum*) and RNAi mediated down regulation of tobacco *MIOX* showed significant changes in arabinan and D-galacturonate content [47]. Recently, RNA sequencing and analysis of *AtMIOX4* over-expressing *Arabidopsis* line showed that the MIOX pathway is plays an important role in cell wall biosynthetic pathway [48].

Mysterious connection between MIOX and AsA biosynthetic pathways

L-ascorbic acid (AsA) is an antioxidant molecule crucial in various cellular functions across plants and animals. Because humans are unable to synthesize it, hence they must obtain it by eating different fruits and vegetables. In plants, however, AsA is synthesized by multiple pathways. Among them, L-galactose, D-galactouronate, and L-glucose are well known as initial precursors but the existence of a fourth pathway starting from MI is still questionable (Figure 4). An earlier study showed that the overexpression of the *AtMIOX4* gene in *Arabidopsis* resulted in a two to three times higher AsA content compared to the wild type [19]. However, other finding suggests that the AsA content does not increase in lines that overexpress *AtMIOX4* but does regulate the level of MI [20]. In rice *OsMIOX* regulates the MI content without affecting AsA content, with or

without stress [30]. The ectopic expression of *AtMIOX4*, *glucuronate reductase* or *L-gulono-1,4-lactone oxidase* genes increased AsA 1.5-3 times in transgenic *Arabidopsis* [49]. While, knockout of *AtGlcAK1*, next successive enzyme of MI oxidation pathway did not alter AsA content [50]. The overexpression of *SIMIOX4* in tomato showed an increased content of total AsA in leaves and red fruits relative to wild type. *SIMIOX4* overexpressing lines and the controls showed an improved AsA content in leaves when fed MI. In contrast, the AsA content in mature green fruit did not show any significant difference among these genotypes. These findings suggest that the MI pathway contributes directly to AsA biosynthesis and accumulation in tomato leaves [7]. Low-vitamin C (*vtc1-1* and *vtc2-1*) mutants in *Arabidopsis* are well characterized for producing less foliar AsA. Mutation in *GDP-d-mannose phosphorylase* and *GDP-L-galactose phosphorylase* resulted in *vtc1-1* and *vtc2-1* in *Arabidopsis* accumulates only 20-30% of the AsA content. Recently, a study showed that *AtMIOX4* is able to restore the AsA level of the *vtc* mutants as well as increasing biomass and growth rate relative to controls [51].

The role of MIOX in relieving cope of abiotic stress

Inositol and its derivatives are well known to involve in alleviating various abiotic stresses such as salt stress, drought, and UV-B induced oxidative stress [52,30,54]. MI and its derivatives are involved in combatting salt stress in plant in two key ways: (i) by protecting cellular and cytoplasmic structures from reactive oxygen species (ROS) i.e. and (ii) by controlling water pressure inside the plant cells [52]. Under salt stress, sodium is transported and accumulated in vacuoles whereas MI, D-ononitol, and D-pinitol accumulated in the cytoplasm of ice plant (*Mesembryanthemum crystallinum*) [54]. Rice *MIOX* (*OsMIOX*) was highly induced by drought, H₂O₂, salt, cold, and ABA (Abscisic acid). Overexpression of *OsMIOX* in rice significantly improved drought resistance [30]. Overexpression of *MIOX4* and *L-gulono-1,4-lactone oxidase* genes increased root and shoot biomass in *Arabidopsis*. Apart from this, phenotypic changes are reported as these lines had increased tolerance of cold, heat, salt, and pyrene [55,49]. *OsMIOX* is upregulated in transgenic rice lines that overexpress *PeaT1* gene during the drought treatment. Yeast two-hybrid assays suggested that *PeaT1* interacts with *OsMIOX*, indirect evidences indicate that *PeaT1* might promote the transcription of *OsMIOX* and also

participate in some yet to explore protein complexes to play a role in drought tolerance. In soybean, *GsMIOX1a* is highly expressed in flowers and under alkaline stress. *Arabidopsis* lines that overexpressed *GsMIOX1a* were more alkaline stress-tolerant than *atmiox1* mutant. *Vacuolar H⁺-pyrophosphatase*, *NADP-malic enzyme*, *KIN1*, and *RD29B*, like alkaline stress inducible markers genes were upregulated in *GsMIOX1a* overexpression plants [21]. *MIOX4* overexpressing tomato lines were tolerant to oxidative stress. The increased content of AsA in tomato might remove excess reactive oxygen species and protect tomato plants from oxidative damage [7]. Transcriptome analysis of *AtMIOX4* over-expressing lines upregulation of gene families that enhanced tolerance against cold, water, and heat stresses [48]. Apple (*Malus hupehensis* Rehd.) *MhMIOX1* and *MhMIOX2* were expressed under ABA and salt stress with NaCl treatment. Overexpression of *MhMIOX2* in *Arabidopsis* and poplar lines improved resistance to salt stress compared to the wild type. Overexpression of *MhMIOX2* in poplar lines under salt stress led to a decrease in H₂O₂ and MDA content and upregulated activity by POD, APX and CAT [22]. All aforementioned findings in *Arabidopsis*, soybean, tomato, rice and apple thus suggest that the MIOX pathway plays an important role in abiotic stress in plants.

The role of MIOX in combating biotic stresses

Transcriptome analysis of *Arabidopsis* and soybean syncytia induced by different nematodes showed up regulation of *MIOX* genes. One of 12 probe sets of *MIOX* genes was up-regulated by soybean root syncytia caused by *Heterodera glycines* [56]. In other study, probe set ID (Gma.17873.1.S1_s_at) corresponding to *AtMIOX2* gene was 20 times more expressed in syncytia five days post inoculation than it was in the control [56]. Total RNA sequencing showed that two *AtMIOX* genes were highly up-regulated in root syncytia caused due to *Heterodera schachtii* on 5 and 15 days of infection [57]. The real-time expression of *AtMIOX2* gene was eight times higher in root syncytia made by cyst nematode. In another report, the expression analysis confirmed that *AtMIOX4* and *AtMIOX5* were highly up-regulated in root syncytia [28]. Wild type *Arabidopsis* and the double mutants *miox1/2* and *miox4/5* were infected with *H. schachtii* to examine the possible role of MIOX. Authors reported smaller syncytia and females per plant were less

in the double mutant roots as compared to control root [28]. While, quadruple *miox1/2/4/5* mutant showed a substantial decrease in susceptibility to cyst nematode *H. schachtii*. Metabolite profiling of root syncytia of the quadruple *miox1/2/4/5* mutant showed that the MI and galactinol content was increased [58]. Galactinol is known to induce various defense-related genes, such as the antimicrobial *Thionin* gene in *Arabidopsis* [58]. Thus, the evidences we discussed so far strongly emphasize that MIOX plays an important role in growth and development of root syncytia and could be a potential target to reduce biotic stress impact caused by cyst nematodes. Transcriptome studies of the wild type and *AtMIOX4* over-expressing lines showed that various genes related to jasmonic acid biosynthesis were also up-regulated in the latter but not in the wild type. Genes related to parthenogenesis were less expressed in overexpressing lines whereas the opposite is true of genes such as o-methyl transferase that are related to the expression of phenolics in biosynthesis [48]. *AtMIOX4* over-expressing lines contain more glucoerucin and AsA than wild type, which suggests that overexpressing lines could be less prone to herbivore attack. It is interesting that relatively high amount of jasmonic acid in *Arabidopsis AtMIOX4* overexpressing lines indicates greater resistance to insects and fungi [59, 48].

Conclusive remarks

The MIOX pathways has great potential as novel player working at cellular level in response to environmental cues. The dispensable nature of pathway enzymes, while key roles of upstream and downstream regulators provide perfect environment to bio-engineer this pathway for stress resilience in plants. In present work, we tried to snapshot the cross talk of myo-Inositol oxidation pathway with some important biosynthetic pathways (Table 1). An important link we could recommend to explore further is Ins(1,4,5)P3 biosynthesis pathway. In near future, we hope to see some exciting outcomes from this crosstalk beyond just metabolic pathways. One such area could be tissue remodeling and developmental reprogramming in response to stress.

Deceleration

All authors read the draft and approved. Authors declares no conflict of interest.

KKB is supported by Belgian Funds for Scientific Research FRS-FNRS, Belgium with Chargé de recherches fellowship. Authors are thankful to Dr Margaret Avery, University of the Witwatersrand for useful suggestions to improve manuscript.

References

1. Chen IW and Charalampous FC (1966) Biochemical studies on inositol: IX. D-inositol 1-phosphate as intermediate in the biosynthesis of inositol from glucose 6-phosphate, and characteristics of two reactions in this biosynthesis. The Journal of Biological Chemistry 241; 2194-2199.
2. Loewus MW, FA Loewus (1980) The C-5 hydrogen isotope-effect in myo-inositol 1-phosphate synthase as evidence for the myo-inositol oxidation-pathway. Carbohydr Res 82: 333–342
3. Pieslinger AM, Hoepflinger MC and Tenhaken R (2010) Cloning of glucuronokinase from *Arabidopsis thaliana*, the last missing enzyme of the myo-inositol oxygenase pathway to nucleotide sugars. The Journal of Biological Chemistry 285; 2902-2910.
4. Valluru R, Van den Ende W (2011) Myo-inositol and beyond - Emerging networks under stress. Plant Science 181; 387–400.
5. Kanter U, Usadel B, Guerineau F, Li Y, Pauly, Tenhaken R (2005) The inositol oxygenase gene family of Arabidopsis is involved in the biosynthesis of nucleotide sugar precursors for cell-wall matrix polysaccharides. Planta 221; 243-254.
6. Alok A, Kaur H, Bhati KK, Kumar J, Pandey P, Upadhyay SK, Pandey A, Sharma NC, Pandey AK, Tiwari S (2015) Biochemical characterization and spatio-temporal expression of myo-inositol oxygenase (MIOX) from wheat (*Triticum aestivum* L.). Plant Gene 4; 10-19.
7. Munir S, Mumtaz M A, Ahiakpa J K, Liu G, Chen W, Zhou G, Zheng W, Ye Z and Zhang Y (2020) Genome-wide analysis of Myo-inositol oxygenase gene family in tomato reveals their involvement in ascorbic acid accumulation. BMC genomics 21, pp.1-15.

8. Coulibaly A, Kouakou B, Chen J (2011) Phytic acid in cereal grains: structure, healthy or harmful ways to reduce phytic acid in cereal grains and their effects on nutritional quality. *American Journal of Plant Nutrition and Fertilization Technology* 1; 1-22.
9. Williams SP, Gillaspay GE, Perera IY (2015) Biosynthesis and possible functions of inositol pyrophosphates in plants. *Frontiers in Plant Science* 6; 67.
10. Sparvoli F, Cominelli E (2015) Seed Biofortification and Phytic Acid Reduction: A Conflict of Interest for the Plant? *Plants* 4; 728-755.
11. Endres S and Tenhaken R (2011) Down-regulation of the myo-inositol oxygenase gene family has no effect on cell wall composition in *Arabidopsis*. *Planta* 234; 157-169.
12. Kroh M, Loewus F (1968) Biosynthesis of Pectic Substance in Germinating Pollen: Labeling with *Myo* inositol-2-14C. *Science* 160; 1352–1354.
13. Seitz B, Klos C, Wurm M, Tenhaken R (2000) Matrix polysaccharide precursors in *Arabidopsis* cell walls are synthesized by alternate pathways with organ-specific expression patterns. *The Plant Journal* 21; 537–546.
14. Loewus FA, Murthy PPN (2000) myo -Inositol metabolism in plants. *Plant Science* 150, 1–19.
15. Bánhegyi G, Braun L, Csala M, Puskás F, Mandl J (1997) Ascorbate Metabolism and Its Regulation in Animals. *Free Radical Biology and Medicine* 23; 793–803.
16. Smirnoff N, Conklin PL, Loewus FA (2001) Biosynthesis of ascorbic acid in plants: A Renaissance. *Annual Review of Plant Physiology and Plant Molecular Biology* 52; 437–467.
17. Agius F, Gonzalez-Lamothe R, Caballero JL, Munoz-Blanco J, Botella MA, Valpuesta V (2003) Engineering increased vitamin C levels in plants by overexpression of a D-galacturonic acid reductase. *Nature Biotechnology* 21; 177–181.
18. Wolucka BA, Van Montagu M (2003) GDP- Mannose 3', 5' Epimerase Forms GDP-L-gulose, a Putative Intermediate for the de Novo Biosynthesis of Vitamin C in Plants. *Journal of Biological Chemistry* 278; 47483–47490.

19. Lorence A (2004) *myo*-Inositol oxygenase offers a possible entry point into plant ascorbate biosynthesis. *Plant Physiology* 134; 1200-1205.
20. Endres S, Tenhaken R (2009) Myo-inositol oxygenase controls the level of myoinositol in *Arabidopsis*, but does not increase ascorbic acid. *Plant physiology*. 149; 1042-1049.
21. Chen C, Sun X, Duanmu H, Yu Y, Liu A, Xiao J, Zhu Y (2015) Ectopic expression of a *Glycine soja* myo-inositol oxygenase gene (*GsMIOX1a*) in *Arabidopsis* enhances tolerance to alkaline stress. *PLoS ONE* 10; 0129998.
22. Yang J, Yang J, Zhao L, Gu L, Wu F, Tian W, Sun Y, Zhang S, Su H and Wang L (2021) Ectopic expression of a *Malus hupehensis* Rehd. myo-inositol oxygenase gene (*MhMIOX2*) enhances tolerance to salt stress. *Scientia Horticulturae* 281, p.109898.
23. Li Z, Liu Z, Wei Y, Liu Y, Xing L, Liu M, Li P, Lu Q, Peng R (2021) Genome-wide identification of the MIOX gene family and their expression profile in cotton development and response to abiotic stress. *PLoS One* 9;16(7):e0254111.
24. Alok A, Kaur J, Tiwari S (2020) Functional characterization of wheat myo-inositol oxygenase promoter under different abiotic stress conditions in *Arabidopsis thaliana*. *Biotechnology Letters* 42(10), pp.2035-2047.
25. Alford SR (2009) Molecular characterization of two *myo*-inositol oxygenases in *Arabidopsis thaliana* (Doctoral dissertation, Virginia Tech) <http://hdl.handle.net/10919/26448>.
26. Zimmermann P, Hirsch-Hoffmann M, Hennig L, Gruissem W (2004) GENEVESTIGATOR *Arabidopsis* microarray database and analysis toolbox. *Plant Physiology* 136; 2621–2632.
27. Alford SR, Rangarajan P, Williams SP, Gillaspie GE. (2012) Myo-inositol oxygenase is required for responses to low energy conditions in *Arabidopsis thaliana*. *Frontiers in Plant Science* 3; 69.
28. Siddique S, Endres S, Atkins JM, Szakasits D, Wiecezorek K, Hoffman J, Bohlmann H (2009) Myo-inositol oxygenase genes are involved in the development of

syncytia induced by *Heterodera schachtii* in *Arabidopsis* roots. New Phytologist 184; 457–472.

29. Alok A, Sharma NC, Tiwari S (2016) Cis-element architectures of the promoter regions and expression pattern of *myo-inositol oxygenase (MIOX)* genes in *Arabidopsis thaliana*. International Journal of Agricultural Science and Research 6; 259-266.

30. Duan J, Zhang M, Zhang H, Xiong H, Liu P, Ali J, Li J, Li Z (2012) Plant Science OsMIOX , a myo -inositol oxygenase gene, improves drought tolerance through scavenging of reactive oxygen species in rice (*Oryza sativa* L). Plant Science 196; 143–151.

31. Arner RJ, Prabhu KS, Thompson JT, Hildenbrandt GR, Likens D, Reddy CC (2001) *myo*-Inositol oxygenase: molecular cloning and expression of a unique enzyme that oxidizes *myo*-inositol and *D-chiro*-inositol. The Biochemical journal 360; 313–20.

32. Kanter U, Becker M, Friauf E, Tenhaken R (2003) Purification, characterization and functional cloning of inositol oxygenase from *Cryptococcus*. Yeast 20; 1317–1329.

33. Brown PM, Caradoc-Davies TT, Dickson JMJ, Cooper GJS, Loomes KM, Baker EN (2006) Crystal structure of a substrate complex of myo-inositol oxygenase, a di-iron oxygenase with a key role in inositol metabolism. Proceedings of the National Academy of Sciences 103; 15032–15037.

34. Thorsell AG, Persson C, Voevodskaya N, Busam RD, Hammarström M, Gräslund S, Gräslund A, Hallberg BM. (2008) Structural and biophysical characterization of human myo-inositol oxygenase. The Journal of Biological Chemistry 283; 15209-16.

35. Perera I, Seneweera S and Hirotsu N (2018) Manipulating the Phytic Acid Content of Rice Grain Toward Improving Micronutrient Bioavailability. Rice 11; 4.

36. Bhati K K, Aggarwal S, Sharma S, Mantri S, Singh S P, Bhalla S, Kaur J, Tiwari S, Roy J K, Tuli R, Pandey AK (2014) Differential expression of structural genes for the late phase of phytic acid biosynthesis in developing seeds of wheat (*Triticum aestivum* L.). Plant Science, 224 74-85.

37. Dong J, Yan W, Bock C, Nokhrina K, Keller W and Georges F (2013) Perturbing the metabolic dynamics of myo-inositol in developing *Brassica napus* seeds through in vivo methylation impacts its utilization as phytate precursor and affects downstream metabolic pathways. *BMC plant biology* 13(1), pp.1-12.
38. Aggarwal S, Kumar A, Bhati KK, Kaur G, Shukla VK, Tiwari S, Pandey AK (2018) RNAi Mediated Down regulation of Inositol Penta kisphosphate Kinase (IPK1) in Wheat Grains Decreases Phytic Acid Levels and Increases Fe and Zn Accumulation. *Frontiers in Plant Science* 9; 259.
39. Ali N, Paul S, Gayen D, Sarkar SN, Datta SK, Datta K (2013) RNAi mediated down regulation of myo-inositol-3-phosphate synthase to generate low phytate rice. *Rice* 6(1):1-2.
40. Saez-Aguayo S, Rautengarten C, Temple H, Sanhueza D, Ejsmentewicz T, Sandoval-Ibañez O, Doñas D, Parra-Rojas JP, Ebert B, Lehner A and Mollet, JC (2017) UUAT1 is a Golgi-localized UDP-uronic acid transporter that modulates the polysaccharide composition of *Arabidopsis* seed mucilage. *The Plant Cell* 29(1), pp.129-143.
41. Geserick C and Tenhaken R (2013) UDP-sugar pyrophosphorylase is essential for arabinose and xylose recycling, and is required during vegetative and reproductive growth in *Arabidopsis*. *The Plant Journal* 74(2), pp.239-247.
42. Reboul R, Geserick C, Pabst M, Frey B, Wittmann D, Lütz-Meindl U, Léonard R and Tenhaken R (2011) Down-regulation of UDP-glucuronic acid biosynthesis leads to swollen plant cell walls and severe developmental defects associated with changes in pectic polysaccharides. *Journal of Biological Chemistry* 286(46), pp.39982-39992.
43. Zablackis E, Huang J, Muller B, Darvill AG and Albersheim P, 1995. Characterization of the cell-wall polysaccharides of *Arabidopsis thaliana* leaves. *Plant physiology* 107(4), pp.1129-1138.
44. Gu X, Bar-Peled M (2004) The biosynthesis of UDP-galacturonic acid in plants. Functional cloning and characterization of *Arabidopsis* UDP-D-glucuronic acid 4-epimerase. *Plant Physiology* 136(4):4256-64.

45. Burget EG, Verma R, Mølhøj M, Reiter WD (2003) The biosynthesis of L-arabinose in plants: molecular cloning and characterization of a Golgi-localized UDP-D-xylose 4-epimerase encoded by the MUR4 gene of *Arabidopsis*. *The Plant Cell* 15; 523-531.
46. Mølhøj M, Verma R and Reiter WD (2004) The biosynthesis of D-Galacturonate in plants. functional cloning and characterization of a membrane-anchored UDP-D-Glucuronate 4-epimerase from *Arabidopsis*. *Plant physiology* 135(3), pp.1221-1230.
47. do Nascimento DD, Conti G, Labate MT, Gutmanis G, Bertollo AL, de Andrade A, Bragatto J, Pagotto LO, Damin P, Moon DH, Labate CA (2012) Modulating Miox2 expression in *Nicotiana tabacum* and impacts on genes involved in cell wall biosynthesis. *Bioenergia em Revista: Diálogos* 2; 60-84.
48. Nepal N, Yactayo-Chang JP, Medina-Jiménez K, Acosta-Gamboa LM, González-Romero ME, Arteaga-Vázquez MA, Lorence A (2019) Mechanisms underlying the enhanced biomass and abiotic stress tolerance phenotype of an *Arabidopsis* MIOX overexpresser. *Plant Direct* 3; e00165.
49. Lisko KA, Torres R, Harris RS, Belisle M, Vaughan MM, Jullian B, Lorence A (2013) Elevating vitamin C content via overexpression of myo-inositol oxygenase and L-gulonolactone oxidase in *Arabidopsis* leads to enhanced biomass and tolerance. *In Vitro Cellular & Developmental Biology-Plant* 49; 643–655.
50. Ivanov Kavkova E, Blöchl C, Tenhaken R (2019) The *Myo*-inositol pathway does not contribute to ascorbic acid synthesis. *Plant Biology* 21; 95-102.
51. Acosta-Gamboa LM, Nepal N, Medina-Jimenez K, Campbell ZC, Cunningham SS, Lee JA and Lorence A (2021) Myo-inositol Oxygenase Overexpression Rescues Vitamin C Deficient *Arabidopsis* (vtc) Mutants. *bioRxiv*, doi.org/10.1101/2021.02.24.432757.

52. Nelson, DE, Rammesmayer G and Bohnert HJ (1998) Regulation of cell-specific inositol metabolism and transport in plant salinity tolerance. *The Plant Cell* 10(5), pp.753-764
53. Raynaud C, Yemets AI, Bergounioux C and Blume YB (2016) Involvement of Inositol Biosynthesis and Nitric Oxide in the Mediation of UV-B Induced Oxidative Stress. *Frontiers in Plant Science* (7), 1-12.
54. Nelson DE, Koukoumanos M and Bohnert HJ (1999) Myo-inositol-dependent sodium uptake in ice plant. *Plant physiology* 119(1), pp.165-172.
55. Tóth SZ, Nagy V, Puthur JT, Kovács L, Garab G (2011) The physiological role of ascorbate as photosystem II electron donor: protection against photoinactivation in heat-stressed leaves. *Plant Physiology* 156; 382-92.
56. Ithal N, Recknor J, Nettleton D, Hearne L, Maier T, Baum TJ, Mitchum MG (2007) Parallel genome-wide expression profiling of host and pathogen during soybean cyst nematode infection of soybean. *Molecular Plant–Microbe Interactions*. 20; 293–305.
57. Szakasits D, Heinen P, Wieczorek K, Hofmann J, Wagner F, Kreil DP, Sykacek P, Grundler FM and Bohlmann H (2009) The transcriptome of syncytia induced by the cyst nematode *Heterodera schachtii* in *Arabidopsis* roots. *The Plant Journal* 57(5), pp.771-784.
58. Siddique S, Endres S, Sobczak M, Radakovic ZS, Fragner L, Grundler FMW, Bohlmann H (2014) Myo-inositol oxygenase is important for the removal of excess myo-inositol from syncytia induced by *Heterodera schachtii* in *Arabidopsis* roots. *New Phytologist* 201; 476–485.
59. Suza WP, Staswick PE (2008) The role of JAR1 in jasmonoyl-L-isoleucine production during *Arabidopsis* wound response. *Planta* 227; 1221–1232.

537 **Table 1: Details of *MIOX* gene expression and their functional validation in different plants.**

Source plant	Gene name	Gene ID	Expression pattern	Function validation	References
<i>Arabidopsis</i>	<i>AtMIOX1</i>	At1g14520	Less in leaf & stem; high in flower		5, 47
	<i>AtMIOX2</i>	At2g19800	Root, stem & flower	Overexpression in tobacco changes arabinan and D-galacturonate	
	<i>AtMIOX4</i>	At4g26260	Higher in flower	Overexpression in tobacco changes arabinan and D galacturonate	
	<i>AtMIOX5</i>	At5g56640	Only in flower		
soybean	<i>GsMIOX1a</i>	Glyma07g01660	Higher in flower; In alkaline stress	Overexpression in <i>Arabidopsis</i> enhances tolerance to alkaline stress	21
	<i>GsMIOX1b</i>	Glyma08g21300	In alkaline stress		
	<i>GsMIOX2a</i>	Glyma01g00840	In alkaline stress		
	<i>GsMIOX2b</i>	Glyma07g15190	In alkaline stress		
	<i>GsMIOX4</i>	Glyma08g10690	In alkaline stress		
Wheat	<i>Ta7AMIOX</i>	TraesCS7A01G357800	High in spike & root; Under heat stress & <i>B. graminis</i> infection		24
	<i>Ta7BMIOX</i>	TraesCS7B01G269900	High in spike & root; Under heat stress & <i>B. graminis</i> infection		

	<i>Ta7DMIOX</i>	TraesCS7D01G364900	High in spike & root; Under heat stress & <i>B. graminis</i> infection		
Tomato	<i>SIMIOX1</i>	Solyc06g062430	Flower & fruit	NA	7
	<i>SIMIOX2</i>	Solyc10g005400	Leaf	NA	
	<i>SIMIOX3</i>	Solyc11g006570	Less in flower	NA	
	<i>SIMIOX4</i>	Solyc12g008650	High in flower & fruit, moderate in roots & leaves	Overexpression in tomato altered the expression of AsA related genes and AsA content in both leaves and fruits.	
	<i>SIMIOX5</i>	Solyc12g098120	Higher in flower		
Rice	<i>OsMIOX</i>	AK103977	Under drought and cold stresses	Overexpression in rice improves tolerance to drought stress	30
Apple	<i>MhMIOX1</i>	MDO0000161309	Under salt stress	NA	22
	<i>MhMIOX2</i>	MDO0000285285	Under salt stress	Overexpression in <i>Arabidopsis</i> leads to increased growth under salt stress.	
Cotton <i>Gossypium ranmondii</i>	<i>GrMIOX01</i>	XP_012472181.1			23
	<i>GrMIOX02</i>	XP_012435726.1			
	<i>GrMIOX03</i>	XP_012439315.1			
	<i>GrMIOX04</i>	XP_012447306.1			
	<i>GrMIOX05</i>	XP_012452593.1			
	<i>GrMIOX06</i>	XP_012455200.1			
Cotton	<i>GhMIOX1</i>	Ghir_A03G000510.1			
	<i>GhMIOX2</i>	Ghir_A05G013690.1	Under heat, cold, drought, and salt		

<i>Gossypium</i>	<i>GhMIOX3</i>	Ghir_A06G007270.1	Under cold		
<i>Hirsutum</i>	<i>GhMIOX4</i>	Ghir_A10G006080.1	Under heat, cold, drought, and salt		
	<i>GhMIOX5</i>	Ghir_A12G016630.1			
	<i>GhMIOX6</i>	Ghir_A12G022070.1			
	<i>GhMIOX7</i>	Ghir_D03G019230.1			
	<i>GhMIOX8</i>	Ghir_D05G013430.1	Under heat, cold, drought, and salt		
	<i>GhMIOX9</i>	Ghir_D06G007460.1	Under cold		
	<i>GhMIOX10</i>	Ghir_D10G006900.1	Under heat, cold, drought, and salt		
	<i>GhMIOX11</i>	Ghir_D12G016880.1			
	<i>GhMIOX12</i>	Ghir_D12G022100.1			

538

539

Figure legends:




Figure 1: Myo-inositol oxidation pathway and its interconnection to other pathways. The pathways towards L-ascorbic acid are represented with dot line, which is active in the animal system, whereas in plants it exists or not is in debate. The pathway towards cell wall biosynthesis is active in the plant system. The myo-inositol catalyzed into D-GlucA (D-glucuronic acid) by MIOX (myo-inositol oxygenase). D-GlucA converted into D-GlucA-1P (D-glucuronic acid- 1-phosphate) with the help of GLK (glucuronokinase). , D-GlucA-1P further catalyzed into UDP-D-GlcA by USP (UDP-sugar pyrophosphorylase). UDP-D-GlcA is an important precursor of numerous cell wall polysaccharides i.e. UDP-D-GaIA (UDP-D-galacturonic acid), UDP-D-xylose, UDP-D-apiose, and UDP-D-Arabinose. IMT - inositol methyltransferase. Myo-inositol kinase (MIK).




Figure 2: Differential transcript level of (A) Arabidopsis (B) wheat *MIOX* genes in different publicly available dataset from RNAseq experiments. Expression data for *AtMIOX* 1, 2, 4 and 5 genes in different stages of development in Embryo development (Bent cotyledon, early torpedo, Early heart, globular, 8/16 cell, late heart, mature green and late torpedo), Germination (zero hour to harvest stage), and Silique 3 to 12 DAP) were normalized for fold changes and hierarchical clustering was applied to group genes based on similarities in their expression, represented as a heat map. All the raw data taken from the <https://www.ebi.ac.uk/gxa/home>, in the foam of TPM (transcript per million read). Wheat *TaMIOX* gene expression was analyzed on <http://www.wheat-expression.com/>. (C) Different stress samples were selected and expression was tested using default settings. Each *TaMIOX* gene represent chromosome they are located.




Figure 3: UDP-D-GlcA biosynthesis and cellular transport. UDP-D-GlcA and UDP-D-Xylose are synthesized into cytoplasm. The NST (nucleotide sugar transporter) has been shown to transport UDP-GlcA over the Endoplasmic Reticulum (ER) membrane and to transport UDP-Xylose over the Golgi bodies (GB) membrane. The function NST transporter is still unclear. UXS1 (UDP-glucuronic acid decarboxylase) responsible for the conversion of UDP-D-GlcA to UDP-D-Xylose. UDP-D-Xylose is synthesized within the lumen of the ER or GB and further XylIT (xylosyltransferases) incorporates xylose in the glycans. GT: glucosyltransferase




Figure 4: Biosynthesis pathways of Ascorbic acid and its committed reactions in organelle. Galactose, D-Galacturonate, L-glucose, and MI are the initial precursors of AsA. Solid arrows symbolize the exist pathway in plants from its initial precursors. Dashed pink line represent the myo-inostiol oxidation pathway, which exist or not in plants are in debate. Dashed blue line symbolize the translocation of a metabolite from one cellular compartment to another. GalL, L-galactono-1,4-lactone; AsA, ascorbic acid; GLDH, L-galactono-1,4-lactone dehydrogenase; and DHA, dehydroascorbate.