



US 20070297984A1

(19) **United States**(12) **Patent Application Publication****Li et al.**(10) **Pub. No.: US 2007/0297984 A1**(43) **Pub. Date: Dec. 27, 2007**(54) **INHIBITION OF HIF-1 ACTIVATION FOR
ANTI-TUMOR AND ANTI-INFLAMMATORY
RESPONSES****Publication Classification**(51) **Int. Cl.***A61K* 38/43 (2006.01)*A61K* 49/00 (2006.01)*A61P* 43/00 (2006.01)*C12N* 15/11 (2006.01)*C12N* 5/06 (2006.01)*C12Q* 1/68 (2006.01)(52) **U.S. Cl.** **424/9.2**; 424/94.1; 435/325;
435/375; 435/6; 536/23.5; 800/13;
800/3(75) Inventors: **Chuan-Yuan Li**, Englewood, CO (US);
Fang Li, Greenwood Village, CO (US);
Pierre Sonveaux,
Ottignies-Louvain-la-Neuve (BE); **Mark**
W. Dewhirst, Durham, NC (US)

Correspondence Address:

JENKINS, WILSON, TAYLOR & HUNT, P. A.
3100 TOWER BLVD., Suite 1200
DURHAM, NC 27707 (US)


(57)

ABSTRACT

The presently disclosed subject matter generally relates to methods and compositions for inhibiting the expression and/or activation of hypoxia-inducible factor 1 (HIF-1) genes in a hypoxic cell. More particularly, the methods disclosed herein relate to inhibition of HIF-1 activation in a cell, increasing sensitivity of a tumor cell to radiation and/or chemotherapy, delaying tumor growth, inhibiting tumor blood vessel growth, inhibiting inflammatory responses in a cell through the use of compositions that prevent the nitrosylation of HIF-1, and methods for screening for new inhibitors of HIF-1 activation. Additionally, the compositions disclosed herein relate to compositions that can be employed in, and are identified by, the disclosed methods.

(73) Assignee: **Duke University**, Durham, NC(21) Appl. No.: **11/731,719**(22) Filed: **Mar. 30, 2007****Related U.S. Application Data**

(60) Provisional application No. 60/787,373, filed on Mar. 30, 2006.






Figure 1A




Figure 1B




Figure 1C




Figure 1D




Figure 2A




Figure 2B




Figure 2C




Figure 3A





Figure 3B

GSNO Control



HIF-1 α

β -actin

Figure 3C





Figure 3D

A



B




Figure 4




Figure 5A





Figure 5B

A

pnsps _{ey} cfyvdsdm	<i>Homo sapiens</i> (human)
pnsps _{ey} cfdvdsdm	<i>Mus musculus</i> (house mouse)
pnsps _{ey} cfdvdsdm	<i>Rattus norvegicus</i> (Norway rat)
pnsps _{ey} cfdvdsdm	<i>Spalax judaei</i> (subterranean mole rat)
pnsps _{ey} cfdvdsdm	<i>Bos grunniens</i> (domestic yak)
pnspm _{ey} cfqvdsdi	<i>Carassius carassius</i> (crucian carp)
epntpe _y cfdvdsdm	<i>Xenopus laevis</i> (frog)

B



C






Figure 6

D



E



F






Figure 6 (cont'd)

A**B****Figure 7**

C




Figure 7 (cont'd)

Spalax	RSNADPALNQEVALKLEPNNAESLELSFTMPQIQDQPASPSDGSSTRQSSPE	512
Eospalax	RSNADPALNQEVALKLEPNNAESLELSFTMPQIQDQPASPSDGSSTRQSSPE	513
Mus	RSSADPALNQEVALKLESPELGLSFTMPQIQDQPASPSDGSSTRQSSPERLLQENVNTP	521
Rattus	RSSADPALNQEVALKLESPELGLSFTMPQIQDQPASPSDGSSTRQSSPE	512
Microtus	RSSADPALNQEVALKLESPELGLSFTMPQIQDQPASPSDGSSTRQSSPE	512
Human	RSSADPALNQEVALKLEPNPESLELSFTMPQIQDQTPSPDGSSTRQSSPE	512
Pongo	RSSADPALNQEVALKLEPNPESLELSFTMPQIQDQPPSPDGSSTRQSSPE	513
Macaca	RSSADPALNQEVALKLEPNPESLELSFTMPQIQDQPPSPDGSSTRQSSPE	512
Spermaphilus	RSSADPALNQEVALKLEPNPESLELSFTMPQIQDQPASPSDGSSTRQSSPE	512
Bos	RSSADPALNQEVALKLEPNPESLELSFTMPQIQDQPASPSDGSSTRQSSPE	512
Pantholops	RSSADPALNQEVALKLEPNPESLELSFTMPQIQDQPASPSDGSSTRQSSPE	512
Canis	RSSADPALNQEVALKLEPNPESLELSFTMPQIQDQPASPSDGSSTRQSSPE	512
Oryctolagus	RSSADPALNQEVALKLEPNPESLELSFTMPQIQDQPASPSDGSSTRQSSPE	512
Gallus	RSNADPALNREVSVKLEPNTELELSFTMPQVQOPTSPSDASTSQSSPE	510
Danio	SSHATTAKSTLPCRRRHGCP	486
Xenopus	RSNVDPALNREVVIKMESNPRTTTCAIHHSTAIGARQPFYQFQSEPTEP	510
	*	
Spalax	-----PNSPSEYCFDVSMDMNVFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	568
Eospalax	-----PNSPSEYCFDVSMDMNVFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	569
Mus	NFSQPNSPSEYCFDVSMDMNVFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	581
Rattus	-----PNSPSEYCFDVSMDMNVFKLELVEKLFADTEAKNPFSAQDSDLLEMLAPYIPM	568
Microtus	-----PNSPSEYCFDVSMDMNVFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	568
Human	-----PNSPSEYCFYVDSMDMNEFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	568
Pongo	-----PNSPSEYCFYVDSMDMNEFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	569
Macaca	-----PNSPSEYCFYVDSMDMNEFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	568
Spermaphilus	-----PNSPSEYCFDVSMDMNVFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	568
Bos	-----PNSPSEYCFDVSMDMNVFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	568
Pantholops	-----PNSPSEYCFDVSMDMNVFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	568
Canis	-----PNSPSEYCFDVSMDMNVFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	568
Oryctolagus	-----PTSPSEYCFDVSMDMNVFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	568
Gallus	-----PSSPNDCYCFDVSMDMANEFKLELVEKLFADTEAKNPFSTQDSDLLEMLAPYIPM	566
Danio	-----LHPYT--	491
Xenopus	-----NTPEYCFDVSSEMASEFKDLVEKLFADTEAKAPFYYPGNDLLEMLAPYIPM	564

Figure 8 (cont'd)

Spalax	DDDFQLRLFDQLSPLESSSPQPGVNT---ATAFOOTQLQEPITSTTTT-TTT-----N	619
Eospalax	DDDFQLRSFDQLSPLESSSPNPPSVS-----TAFQOTQLQEPITTTTT-----E	614
Mus	DDDFQLRSFDQLSPLESNPPSPMSMT---VTGFOOTQLQKPIITATATTATT-----D	633
Rattus	DDDFQLRSFDQLSPLESNPPSPSVST---VTGFOOTQLQKPIITVATATATT-----D	620
Microtus	DDDFQLRSFDQLSPLESSSPSPSVST---VTGFOOTPLQKPIITAAATATTATTATTD	625
Human	DDDFQLRSFDQLSPLESSSPSPSVST---VTGFOOTPLQKPIITAAATATTATTATTD	622
Pongo	DDDFQLRSFDQLSPLESSSPSPSVST---VTGFOOTPLQKPIITAAATATTATTATTD	623
Macaca	DDDFQLRSFDQLSPLESSSPSPSVST---VTGFOOTPLQKPIITAAATATTATTATTD	622
Spermaphilus	DDDFQLRSFDQLSPLESSSPSPSVST---VTGFOOTPLQKPIITAAATATTATTATTD	617
Bos	DDDFQLRSFDQLSPLESSSPSPSVST---VTGFOOTPLQKPIITAAATATTATTATTD	619
Pantholops	DDDFQLRSFDQLSPLESSSPSPSVST---VTGFOOTPLQKPIITAAATATTATTATTD	619
Canis	DDDFQLRSFDQLSPLESSSPSPSVST---VTGFOOTPLQKPIITAAATATTATTATTD	619
Oryctolagus	DDDFQLRSFDQLSPLESSSPSPSVST---VTGFOOTPLQKPIITAAATATTATTATTD	614
Gallus	DDDFQLRSFDQLSPLESSSPSPSVST---VTGFOOTPLQKPIITAAATATTATTATTD	608
Danio	DDDFQLRSFDQLSPLESSSPSPSVST---VTGFOOTPLQKPIITAAATATTATTATTD	
Xenopus	DDDFQLRTFDQLSSLECDSSIPQTLGS--MTTLFHQSLSPSTS-----	605
Spalax	ELKTVTKDNIEDIKILIASPSSSTHTPKETTTATTSSPYSGTQSRRTASPNRAGQGVIEQTE	679
Eospalax	ELKTVTKDSTEDIKILITSPSSSTHTPKETTTATTSSPYSGTQSRRTASPNRAGQGVIEQTE	674
Mus	ESKTETKDNKEDIKILIASPSSSTHTPKETTTATTSSPYSGTQSRRTASPNRAGQGVIEQTE	692
Rattus	ESKAVTKDNIEDIKILIASPSTQVPQEMTTAKAS-AYSGTHSRTASPDRAKRVIEQTD	679
Microtus	ELKTVTKDNIEDIKILIASPSSSVHVPQETTAATAS-AYSAPHSRTASPDRAKRVIEQAE	684
Human	ELKTVTKDRMEDIKILIASPSTHTHKETTSATSSP-YRDTQSRRTASPNRAGKGVIEQTE	681
Pongo	ELKTVTKDRMEDIKILIASPSTHTHKETTSATSSP-YRDTQSRRTASPNRAGKGVIEQTE	682
Macaca	ELKTVTKDRMEDIKILIASPSTHTHKETTSATSSP-YRDTQSRRTASPNRAGKGVIEQTE	681
Spermaphilus	ELKTVTKDDMEDIKILIASWSPTHTAPKETTSAATSS-YNDTQSRRTASPNRAGKGVIEQTE	676
Bos	ELKTVTKDGMEDIKILIAFPSPHPVPEPPCATTSPP-YSDTQSRRTASPNRAGKGVIEQTE	678
Pantholops	ELKTVTKDGMEDIKILIAFPSPHPVPEPPCATTSPP-YSDTQSRRTASPNRAGKGVIEQTE	678
Canis	ELKTVTKDGEDIKILIAAPSPHTVPKVTTSATTSPP-YSDTQSRRTASPNRAGKGVIEQTE	678
Oryctolagus	TIKTSAKDSVEDMKVLIASPSPNHIKPKETASATTSPPYRDTQSRRTASPNRAGKGVIEQTE	674
Gallus	EIKPVAER-VDDVKALIVSPSPVHINDTSSAPASP-YSGNRSRTASPNRAGKGVIEQTE	666
Danio	-----CCRCACVHLSRSS-----	504
Xenopus	--DFKPEDAMSDLKTIIOQSP--VHMKESTSAVPSP-YNGNRSRTSSPVPAPAKAVVDKTE	660

Figure 8 (cont'd)

Spalax	EVNAPIQGSRNLLQGEELLRALDQVN	824
Eospalax	EVNAPIQGSRNLLQGEELLRALDQVN	819
Mus	EVNAPIQGSRNLLQGEELLRALDQVN	836
Rattus	EVNAPIQGSRNLLQGEELLRALDQVN	823
Microtus	EVNAPVQGSRNLLQGEDLLRALDQVN	828
Human	EVNAPIQGSRNLLQGEELLRALDQVN	826
Pongo	EVNAPIQGSRNLLQGEELLRALDQVN	827
Macaca	EVNAPIQGSRNLLQGEELLRALDQVN	826
Spermaphilus	EVNAPIQGSRNLLQGEELLRALDQVN	821
Bos	EVNAPIQGSRNLLQGEELLRALDQVN	823
Pantholops	EVNAPIQGSRNLLQGEELLRALDQVN	823
Canis	EVNAPIQGSRNLLQGEELLRALDQVN	823
Oryctolagus	EVNAPIQGSRNLLQGEELLRALDQVN	819
Gallus	EVNAPIQGNRNLLQGEELLRALDQVN	811
Danio	-----	
Xenopus	EVNAPVHGTRNLLQGEELLRALDQAN	805

Figure 8 (cont'd)




Figure 9A



Figure 9B

INHIBITION OF HIF-1 ACTIVATION FOR ANTI-TUMOR AND ANTI-INFLAMMATORY RESPONSES

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Patent Application Ser. No. 60/787,373, filed Mar. 30, 2006, the disclosure of which is incorporated herein by reference in its entirety.

GOVERNMENT INTEREST

[0002] The presently disclosed subject matter was made with U.S. Government support under Grant No. EB001882 from the U.S. National Institute of Bioimaging and Bioengineering, Grant No. CA81512 from the U.S. National Cancer Institute, and Grant No. DAMD17-02-0052 from the U.S. Department of Defense. Thus, the United States Government has certain rights in the presently disclosed subject matter.

TECHNICAL FIELD

[0003] The presently disclosed subject matter generally relates to methods and compositions for inhibiting the expression and/or activation of hypoxia-inducible factor 1 (HIF-1) gene products in a hypoxic cell. More particularly, the presently disclosed subject matter provides methods and compositions involved in inhibition of HIF-1 activation through the use of agents that prevent the nitrosylation of HIF-1.

BACKGROUND

[0004] In a typical clinical setting, radiation therapy and/or chemotherapy treatments are administered to the majority (>90%) of cancer patients. Therefore, along with surgery, radiation therapy and chemotherapy represent two of the three main modalities employed for cancer treatment. However, the therapeutic outcomes are still far from ideal for many types of tumors. The main problem associated with radiotherapy is the recurrence of tumors and/or the development of metastases at distant locations. For chemotherapy, the problem is the development of resistance. In both cases, new methods and compositions that can sensitize tumors to current treatments are highly desirable. Ideally, these methods and compositions should decrease local recurrences in patients treated with radiotherapy and/or should increase the efficacy of chemotherapeutic agents systemically. In addition, they should not have severe side effects.

[0005] What are needed, then, are new strategies and compositions for treating tumors and/or cancers via inhibition of HIF-1 activity and/or the upregulation of HIF-1 activity that results from radiotherapy and/or chemotherapy. The presently disclosed subject matter addresses this and other needs in the art.

SUMMARY

[0006] This Summary lists several embodiments of the presently disclosed subject matter, and in many cases lists variations and permutations of these embodiments. This Summary is merely exemplary of the numerous and varied embodiments. Mention of one or more representative features of a given embodiment is likewise exemplary. Such an

embodiment can typically exist with or without the feature(s) mentioned; likewise, those features can be applied to other embodiments of the presently disclosed subject matter, whether listed in this Summary or not. To avoid excessive repetition, this Summary does not list or suggest all possible combinations of such features.

[0007] The presently disclosed subject matter provides methods for increasing a sensitivity of a tumor in a subject to a treatment. In some embodiments, the methods comprise administering to the tumor a composition comprising an effective amount of an inhibitor of nitric oxide synthase, a nitric oxide scavenger, an inhibitor of HIF-1 nitrosylation, or a combination thereof, wherein (i) the tumor is resistant to radiation therapy, chemotherapy, or both radiation therapy and chemotherapy; and (ii) the administering increases the sensitivity of the tumor to the radiotherapy, the chemotherapy, or both the radiotherapy and the chemotherapy. In some embodiments, the inhibitor of nitric oxide synthase is selected from the group consisting of L-N(6)-(1-iminoethyl)lysine tetrazole-amide (SC-51); aminoguanidine (AG); guanidinoethyl disulfide; L-NG-nitroarginine methyl ester; mercaptoethylguanidine (MEG); N^ω-nitro-L-arginine methyl ester (L-NAME); N-(3-(aminomethyl)benzyl)acetamidine (1400W); N^G-monomethyl-L-arginine (L-NMMA); 7-nitroindazole (7-NI). In some embodiments, the nitric oxide scavenger is selected from the group consisting of hydroxocobalamin; 2-(4-carboxyphenyl)-4,4,5,5-tetramethylimidazoline-1-oxyl-3-oxide (carboxy-PTIO); diethyldithiocarbamate; AMD6221 (ruthenium[hydrogen(diethylenetriamino)]pentaacetato]chloride); and N-dithiocarboxy-sarcosine (DTCS). In some embodiments, the administering comprises administering a minimally therapeutic dose of an inhibitor of inducible nitric oxide synthase (iNOS). In some embodiments, the composition inhibits nitrosylation of Cys520 of SEQ ID NO: 6.

[0008] The presently disclosed subject matter also provides methods for delaying tumor growth in a subject. In some embodiments, the methods comprise (a) administering to the subject a composition comprising an effective amount of an inhibitor of nitric oxide synthase, a nitric oxide scavenger, an inhibitor of HIF-1 nitrosylation, or a combination thereof; and (b) treating a tumor that is resistant to radiation therapy, chemotherapy, or both radiation therapy and chemotherapy, with radiation therapy, chemotherapy, or both radiation therapy and chemotherapy, whereby tumor growth in the subject is delayed. In some embodiments, the composition inhibits nitrosylation of Cys520 of SEQ ID NO: 6. In some embodiments, the treating comprises treating the tumor with a subtherapeutic dose of ionizing radiation. In some embodiments, the treating comprises administering to the subject a therapeutically effective amount of cyclophosphamide. In some embodiments, the method further comprises promoting tumor regression.

[0009] The presently disclosed subject matter also provides methods for inhibiting tumor blood vessel growth in a subject. In some embodiments, the methods comprise (a) administering to the subject a composition comprising an effective amount of an inhibitor of nitric oxide synthase, a nitric oxide scavenger, an inhibitor of HIF-1 nitrosylation, or a combination thereof; and (b) treating a tumor that is resistant to radiation therapy, chemotherapy, or both radiation therapy and chemotherapy, with radiation therapy, chemotherapy, or both radiation therapy and chemotherapy,

whereby tumor blood vessel growth is inhibited. In some embodiments, the composition inhibits nitrosylation of Cys520 of SEQ ID NO: 6. In some embodiments, the methods further comprise delaying tumor growth in the subject. In some embodiments, the methods further comprise promoting tumor regression in the subject.

[0010] The presently disclosed subject matter also provides methods for inhibiting HIF-1 activity in a cell. In some embodiments, the methods comprise contacting the cell with a composition comprising an effective amount of an inhibitor of nitric oxide synthase, a nitric oxide scavenger, an inhibitor of HIF-1 nitrosylation, or a combination thereof, whereby HIF-1 activity in the cell is inhibited. In some embodiments, the cell is a tumor cell. In some embodiments, the tumor cell is present in a subject. In some embodiments, the subject is a mammal. In some embodiments, the mammal is a human. In some embodiments, the composition inhibits nitrosylation of Cys520 of SEQ ID NO: 6. In some embodiments, the methods further comprise exposing the tumor cell to a treatment selected from the group consisting of radiation therapy, chemotherapy, and combinations thereof.

[0011] The presently disclosed subject matter also provides methods for inhibiting an inflammatory response in a cell. In some embodiments, the methods comprise contacting the cell with a composition comprising an effective amount of an inhibitor of nitric oxide synthase, a nitric oxide scavenger, an inhibitor of HIF-1 nitrosylation, or a combination thereof, whereby an inflammatory response in the cell is inhibited. In some embodiments, the cell is present in a subject. In some embodiments, the subject is a mammal. In some embodiments, the mammal is a human. In some embodiments, the agent inhibits nitrosylation of Cys520 of SEQ ID NO: 6.

[0012] In some embodiments of the presently disclosed methods, the composition is provided to the subject in an implantable device. In some embodiments, the subject is a mammal, and in some embodiments the mammal is a human.

[0013] The presently disclosed subject matter also provides methods for identifying an inhibitor of nitrosylation of an HIF-1 polypeptide. In some embodiments, the methods comprise (a) providing a cell comprising a nucleic acid a nucleotide sequence comprising any of SEQ ID NOs: 18-21; (b) contacting the cell with a compound comprising a potential inhibitor of nitrosylation of an HIF-1 polypeptide; and (c) assaying nitrosylation of a cysteine residue present in the nucleic acid, whereby an inhibitor of nitrosylation of an HIF-1 polypeptide is identified. In some embodiments, the cell is present in a subject. In some embodiments, the subject is a mammal. In some embodiments, the mammal is a human. In some embodiments, the nucleic acid comprises an expression vector in which the nucleic acid is operably linked to a promoter that is active in the cell. In some embodiments, the expression vector is a transgene and the animal is a transgenic animal that expresses the nucleic acid. In some embodiments, the compound is administered to the transgenic animal via a route that results in the compound contacting the cell. In some embodiments, the methods further comprise comparing a level of nitrosylation of the cysteine residue present in the nucleic acid to a level of nitrosylation of the cysteine residue present in the nucleic

acid prior to the contacting step. In some embodiments, the cell is an in vitro cultured cell and the contacting is performed in vitro.

[0014] The presently disclosed subject matter also provides expression constructs comprising one or more of SEQ ID NOs: 18-21 operably linked to a promoter.

[0015] The presently disclosed subject matter also provides expression constructs comprising one or more of SEQ ID NOs: 18-21 operably linked to a promoter, with the proviso that all cysteine residues present within SEQ ID NOs: 18-21 have been replaced with a non-nitrosylatable amino acid. In some embodiments, the non-nitrosylatable amino acid is serine.

[0016] The presently disclosed subject matter also provides host cells comprising the disclosed expression constructs.

[0017] The presently disclosed subject matter also provides transgenic, non-human animals comprising the disclosed expression constructs.

[0018] The presently disclosed subject matter also provides for the use of inhibitors of nitric oxide synthases to prevent activation of HIF-1 activity in tumors by cancer therapy that include radiation and chemotherapy, the use of nitric oxide scavengers to prevent activation of HIF-1 activity in tumors by cancer therapy that include radiation and chemotherapy, the use of agents that can reduce the production of NO to sensitize tumors to radiotherapy and/or chemotherapy, the use of nitric oxide synthase or nitric oxide scavengers to inhibit inflammatory reaction through the inhibition of HIF-1 activation, the use of agents that can block the nitrosylation of HIF-1 α cysteine 520 for the purpose of enhancing cancer therapy, and the use of agents that can block the nitrosylation of HIF-1 α cysteine 520 for the purpose of inhibiting/attenuating inflammatory response.

[0019] This and other objects are achieved in whole or in part by the presently disclosed subject matter.

[0020] An object of the presently disclosed subject matter having been stated above, other objects and advantages of the presently disclosed subject matter will become apparent to those of ordinary skill in the art after a study of the following description and non-limiting Examples.

BRIEF DESCRIPTION OF THE DRAWINGS

[0021] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the United States Patent and Trademark Office upon request and payment of the necessary fee.

[0022] FIGS. 1A-1D depict the results of experiments to establish ODD-luc as a non-invasive reporter for HIF-1 α expression.

[0023] FIG. 1A depicts the domain structure of murine HIF-1 α and reporter proteins. The oxygen dependent degradation (ODD) domain is located between amino acids 401 to 613 (top construct) of murine HIF-1 α (SEQ ID NO: 3). The ODD-luc reporter coding sequence (middle construct) was obtained by inserting the ODD domain of murine HIF-1 α between an upstream cytomegalovirus (CMV) promoter and the downstream gene firefly luciferase coding

sequence (luc). The ODD and luc sequences were engineered to be in frame. A Kozak sequence and start codon (ATG) were inserted 5' of the ODD coding sequence. A luciferase coding sequence driven by the CMV promoter (lower construct) was used as a control.

[0024] FIG. 1B depicts the expression of ODD-luc in 4T1 cells determined from luc-dependent conversion of luciferin to luminescent oxoluciferin, and measurement of associated light emission. The top panel depicts representative photographs obtained through the Xenogen IVIS™ system of ODD-luc expression/activity in 4T1 cells cultured after activation of ODD-luc by CoCl₂ (240 μM for 12 hrs), exposure to a proteasome inhibitor (10 μM MG132 for 12 hrs), hypoxia (0.5% O₂ for 24 hrs), or anti-VHL siRNA transfection (VHL-KD). A color ladder is provided to the right of the panel. The average luc activities were calculated from triplicate experiments in each case. Significant differences were observed between control and treated cells ($p < 0.05$ in all cases, Student's *t* test). The bottom panel presents a graphical depiction of the activity levels presented in the photographic depictions in the top panel.

[0025] FIG. 1C depicts Western blot analysis showing down regulation of VHL protein expression after introducing an siRNA-expressing vector encoding an anti-VHL minigene into 4T1 cells as a stable, integrated construct. The sequence of the siRNA was AACATCACATTGCCAGTG-TAT (SEQ ID NO: 17). β-actin levels were used as loading controls.

[0026] FIG. 1D depicts Western blot analysis of wild type 4T1 cells treated as in FIG. 1B. Lysates of the cells were analyzed for endogenous HIF-1α protein expression using a rabbit anti-mouse HIF-1α polyclonal antiserum. VHL-KD: cells stably transduced with an siRNA gene against VHL. β-actin was used as the loading control.

[0027] FIGS. 2A-2C depict *in vivo* activation of HIF-1α by ionizing radiation in tumors.

[0028] FIG. 2A depicts luciferase activity in 4T1 tumors stably transduced with ODD-luc or CMV-luc reporter genes established in nude mice. Size-matched tumors were locally irradiated (6 Gy) at day 0. Luciferase activity in tumors was determined daily through non-invasive imaging. Fourteen animals were used in each group and the error bars indicate the standard error of the mean. The difference between the irradiated group and control was significant ($p < 0.05$ from day 3 to day 10 by two-way ANOVA).

[0029] FIG. 2B is a bar graph presenting the results of radiation-induced activation of endogenous HIF-1 binding activity to a hypoxia responsive element (HRE) measured by ELISA in tumors irradiated 5 days earlier. In each group, the average results from 4 tumor samples are shown ($p < 0.05$, Student's *t* test). Error bars represent standard deviation.

[0030] FIG. 2C is a bar graph presenting the results of a radiation-induced increase in intratumoral VEGF levels as measured by ELISA. In each group, the average results from 4 tumors are shown ($p < 0.05$, Student's *t* test). Error bars represent standard deviation.

[0031] FIGS. 3A-3D depict the results of experiments showing that nitric oxide is a key regulator of radiation-induced HIF-1α activation.

[0032] FIG. 3A presents the results of assays for luciferase activity in 4T1-ODD-luc or 4T1-luc transduced tumors established in the hind legs of nude mice and irradiated (at day 0) with or without the administration of L-NAME (at day-1). Luciferase levels were then monitored post irradiation. Tumors with the CMV-luc reporter were used as controls. Significant inhibition of ODD-luc expression were observed by the use of L-NAME. Eight animals were used in each group and the error bars indicate standard error of the mean. $p < 0.05$ from day 4 (two-way ANOVA). The top panel is a graph of the activities of the listed conditions from day 0 to day 10, and the bottom panel depicts representative photographs at various stages and under the listed conditions.

[0033] FIG. 3B is a bar graph depicting S-nitrosoglutathione-(GSNO) induced activation of ODD-luc cells *in vitro*. 4T1-ODD-luc cells were exposed to the NO donor GSNO at indicated dosage and monitored for ODD-luc expression. The data were normalized against cells that were not treated with GSNO. The error bars represent standard deviations. Each data point represents the average of triplicate experiments. Dose-dependent induction was observed. $p < 0.001$ (Student's *t* test)

[0034] FIG. 3C depicts Western blot analysis of endogenous HIF-1α protein levels after GSNO treatment (1 mM for 8 hours) in 4T1 cells. β-actin levels were used as loading control.

[0035] FIG. 3D is a bar graph depicting suppression of NO-mediated HIF-1α activation by a nitric oxide scavenger. 4T1-ODD-luc cells were exposed to GSNO (1 mM) in the presence or absence of a chemical NO scavenger—carboxy-PTIO (0.5 mM). The cells were monitored for luciferase expression 24 hours later. The error bars represent standard deviation and each data point represents the average of triplicate experiments. $p < 0.05$ (Student's *t* test).

[0036] FIGS. 4A and 4B are graphical representations demonstrating the role of the inducible form of nitric oxide synthase (iNOS) in radiation-induced HIF-1α activation.

[0037] FIG. 4A depicts the effect of an iNOS specific inhibitor. Subcutaneous tumors were established in the hind legs of nude mice through the use of 4T1-ODD-luc cells and irradiated with or without the administration of 1400W, an iNOS-specific inhibitor. ODD-luc level were then monitored daily post irradiation. Significant inhibition of radiation-induced HIF-1 activation was observed in the group treated with 1400W ($p < 0.001$ from day 4, two-way ANOVA).

[0038] FIG. 4B depicts the effect of a homozygous genetic disruption (i.e., knockout) of the iNOS gene on HIF-1α activation in a host animal. Tumors were established from B16F10-ODD-luc cells in syngeneic wild type or iNOS^{-/-} C57BL/6 mice. In some groups, mice received L-NAME one day before tumor irradiation (6 Gy) at day 0. Luciferase activities were determined every other day. Eight animals were used in each group and the error bars represent the standard errors of the mean. In wild type C57BL/6 mice (solid lines), the difference between L-NAME treated and non-treated groups was statistically significant ($p < 0.01$ on days 1, 3, and 5, two-way ANOVA test). In iNOS^{-/-} mice (broken lines), the difference between L-NAME treated and non-treated groups was not significant ($p > 0.05$ at all time points, two-way ANOVA).

[0039] FIGS. 5A and 5B depict the role of macrophages in radiation-induced HIF-1 α activation.

[0040] FIG. 5A depicts luciferase expression in tumors established from 4T1-ODD-luc cells in nude mice. In some mice, macrophages were depleted by injection of carrageenan. Selected groups of mice also received L-NAME one day before irradiation (6 Gy). Luciferase expression was determined every other day. Eight mice were included in each group. The error bars represent the standard errors of the mean.

[0041] FIG. 5B depicts immunohistochemistry analysis of HIF-1 α , iNOS, and macrophages in tumors. Mice with irradiated 4T1 tumors were sacrificed and their tumors excised 5 days after localized 6 Gy or sham irradiation of tumors. Shown in the left panel are representative results from co-staining of CD68 (a marker for macrophages (M ϕ) and iNOS. Co-staining of HIF-1 α and iNOS is shown on the right panel. In each case, merged pictures are provided. Orange color in both panels represents co-localization.

[0042] FIGS. 6A-6F depict normoxic prevention of HIF-1 α degradation through S-nitrosylation of cysteine 533.

[0043] FIG. 6A presents amino acid subsequence conservation across different species in the region of Cys 533 of murine HIF-1 α (GENBANK® Accession No. NP_034561 (SEQ ID NO: 3)). The subsequences presented include PNSPSEYCFYVDSMD (*Homo sapiens*; SEQ ID NO: 18); PNSPSEYCFDVDSDM (*Mus musculus*; SEQ ID NO: 19); PNSPSEYCFDVDSDM (*Rattus norvegicus*; SEQ ID NO: 19); PNSPSEYCFDVDSDM (*Spalax judaei*; SEQ ID NO: 19); PNSPSEYCFDVDSDM (*Bos grunniens*; SEQ ID NO: 19); PNSPMEYCFQVDSDI (*Carassius carassius*; SEQ ID NO: 20); and EPNTPEYCFDVDSEM (*Xenopus laevis*; SEQ ID NO: 21). See also FIG. 8.

[0044] FIG. 6B is a bar graph depicting the effects of various stimuli (0.5% hypoxia, proteasome inhibitor MG132, and CoCl₂) on the activation of wild type ODD-luc and C533S-ODD-luc in 4T1 cells. The experiments were carried out in the similar manner as those described in FIG. 1B. The data shown are the results of triplicate experiments. The error bars represent standard deviations. In all treatment groups, $p > 0.05$ between wild type and mutant ODD-luc expression levels (Student's t test).

[0045] FIG. 6C is a bar graph depicting luciferase activity in wild type ODD-luc or C533S ODD-luc transduced 4T1 cells treated with GSNO (1 mM). Significant attenuation of luc expression was observed in C533S-ODD-luc transduced cells ($p < 0.01$, Student's t test). Each data point is the results of triplicate experiments and the error bars represent standard errors. The unit for light output shown is p/sec/CM²/Sr.

[0046] FIG. 6D is a graph depicting luciferase activity in irradiated (6 Gy) tumors established from 4T1 cells transduced with wild type or C533S-ODD-luc. Luciferase expression was monitored every other day. Significant attenuation of luciferase expression was observed in C533S-ODD-luc-transduced 4T1 tumors ($p < 0.01$ from day 5, two-way ANOVA). Each group has five animals and the error bars represent standard errors of the mean.

[0047] FIG. 6E depicts the results of Western blot analysis of S-nitrosylation of C533 in the ODD domain. 4T1 cells transduced with wild type ODD or C533S-ODD (both with

a myc-tag at the 3' end for Western blot detection) were exposed to GSNO and then lysed. S-nitrosylation of ODD was determined through the biotin switch assay (Jaffrey & Snyder, 2001). A clear nitrosylation signal was observed for wild type ODD after GSNO treatment, but was not observed in C533S ODD with or without GSNO treatment.

[0048] FIG. 6F depicts the results of Western blot analysis demonstrating the absence of binding between nitrosylated ODD and VHL. 4T1 tumor cells were transduced with CMV-ODD-mycTag, CMV-C533S-ODD-mycTag, or CMV-HA-VHL. Where indicated, ODD-transfected cells were exposed to 1 mM GSNO for 8 hours. The lysate of ODD-transfected cells was admixed with lysate of cells expressing HA-VHL. Mixed lysates were immunoprecipitated with anti-HA antibody to pull down the VHL protein and any ODD bound thereto. The immunoprecipitate was then immunoblotted with antibody against mycTag to detect ODD bound to VHL. Total tagged ODD (Input ODD) and VHL (Input VHL) were detected by Western blot analysis with antibodies against the mycTag and the HA-tag, respectively.

[0049] FIGS. 7A-7C depict the enhanced anti-tumor efficacy of radiotherapy in combination with L-NAME. B16F10 and 4T1 tumors were established in syngeneic C57BL/6 and 4T1 mice, respectively and irradiated with 3 fractions of X-rays at 6 Gy/fraction (irradiation every other day). In some of the groups, L-NAME was administered in the drinking water one day before irradiation. Tumor sizes were monitored every other day. At least 6 animals were used in each treatment groups. Tumor sizes were then plotted against time for each tumor type. The error bars represent the standard errors of the mean.

[0050] FIG. 7A is a graph depicting 4T1 tumor growth delay.

[0051] FIG. 7B is a graph depicting B16F10 melanoma growth delay.

[0052] FIG. 7C is a bar graph depicting CD31⁺ cells (indicative of vasculature) in tumors excised from different groups on day 10. The tumors were excised, sectioned, and probed for the presence of vasculature by use of an antibody against CD31, which stained for endothelial cells. The average vascular length density of tumors was determined from five randomly chosen fields for each treatment type. The error bars represent the standard errors. The differences between the combined treatment group and the individual groups were significant ($p < 0.05$, one way ANOVA) in both tumor models.

[0053] FIG. 8 presents a maximized amino acid sequence alignment of HIF-1 α polypeptide sequences from the following organisms: *Spalax judaei* (GENBANK® Accession No. CAG29396; SEQ ID NO: 1); *Eospalax baileyi* (GENBANK® Accession No. ABB17537; SEQ ID NO: 2); *Mus musculus* (GENBANK® Accession No. NP_034561; SEQ ID NO: 3); *Rattus norvegicus* (GENBANK® Accession No. NP_077335; SEQ ID NO: 4); *Microtus oeconomus* (GENBANK® Accession No. AAY27087; SEQ ID NO: 5); *Homo sapiens* (GENBANK® Accession No. NP_001521; SEQ ID NO: 6); *Pongo pygmaeus* (GENBANK® Accession No. CAH93355; SEQ ID NO: 7); *Macaca fascicularis* (GENBANK® Accession No. BAE01417; SEQ ID NO: 8); *Spermophilus tridecemlineatus* (GENBANK® Accession No.

AAU14021; SEQ ID NO: 9); *Bos taurus* (GENBANK® Accession No. NP_776764; SEQ ID NO: 10); *Pantholops hodgsonii* (GENBANK® Accession No. AAX89137; SEQ ID NO: 11); *Canis familiaris* (GENBANK® Accession No. XP_852278; SEQ ID NO: 12); *Oryctolagus cuniculus* (GENBANK® Accession No. AAP43517; SEQ ID NO: 13); *Gallus gallus* (GENBANK® Accession No. NP_989628; SEQ ID NO: 14); *Danio rerio* (GENBANK® Accession No. NP_956527; SEQ ID NO: 15); and *Xenopus laevis* (GENBANK® Accession No. CAB96628; SEQ ID NO: 16).

[0054] FIGS. 9A and 9B depict ODD-LUC expression in 4T1 tumor cells that have been treated with cyclophosphamide.

[0055] FIG. 9A depicts a time course of ODD-luc change in 4T1 tumors treated with cyclophosphamide.

[0056] FIG. 9B depicts images of ODD-luc expression with (bottom 2 panels) or without (top 2 panels) cyclophosphamide exposure.

DETAILED DESCRIPTION

I. General Considerations

[0057] Radiotherapy and chemotherapy are two of the three main modalities of cancer therapy. However, for the majority of cancer patients, the therapeutic efficacy of chemotherapy or radiotherapy is not ideal. Many tumors are resistant to various chemotherapy and radiotherapy treatments. At the molecular level, the mechanisms involved in such resistance are not completely understood. However, recent studies indicate that the hypoxia-inducible factor 1 (HIF-1) factor might be involved. These studies have shown that radiation and chemotherapy can upregulate the level and activity of HIF-1 protein and this upregulation is related to increased tumor angiogenesis and tumor resistance to therapy. Furthermore, inhibition of HIF-1 activity can significantly increase the sensitivity of tumor cells to radiotherapy and chemotherapy.

[0058] Recent progress in the understanding of tumor physiology and the tumor microenvironment has yielded new targets that can be used to develop novel therapeutic agents. One such target is the hypoxia-inducible factor 1 (HIF-1). HIF-1 is a master transcriptional regulator that plays important roles in development, physiology, and many pathological processes (Semenza et al., 2000; Semenza, 2002; Semenza, 2003; Melillo, 2004). Originally identified as a transcription factor activated under conditions of abnormally low oxygen (Wang & Semenza, 1993a; Wang & Semenza, 1993b), HIF-1's potential roles in cancer biology are a topic of current interest. More than 60 genes have been identified as direct targets of HIF-1 activity (Semenza, 2003) including, but not limited to genes involved in angiogenesis, metabolic adaptation, apoptosis induction/resistance, and invasion/metastasis.

[0059] HIF-1 is a heterodimeric protein that consists of the constitutively expressed HIF-1 β subunit (also called aryl hydrocarbon receptor nuclear translocator; ARNT) and the highly regulated HIF-1 α subunit (Wang & Semenza, 1995). The overall activity of HIF-1 is determined by intracellular HIF-1 α level. In the past decade, certain insights related to HIF-1 α regulation have been realized. One significant advance has been the discovery of HIF-1 α regulation by oxygen tension, which is mainly mediated by the ubiquitin-

proteasome pathway. Under normoxic conditions, human HIF-1 α is hydroxylated by one or more prolyl hydroxylases (PHDs) at proline residues 402 and 564 in the oxygen dependent domain (ODD; Ivan et al., 2001; Jaakkola et al., 2001). This hydroxylation renders HIF-1 α susceptible to binding and ubiquitylation by E3 ubiquitin protein ligases, which contain the von Hippel-Lindau tumor suppressor protein (VHL; Pause et al., 1999; Maxwell et al., 1999; Maxwell et al., 2001). Ubiquitylated HIF-1 α is then rapidly degraded by the proteasome. Under hypoxic conditions, the enzymatic activities of PHDs are significantly reduced due to the oxygen-dependent nature of PHDs. As a result, HIF-1 α accumulates.

[0060] In addition to hydroxylation of the proline residues in the ODD domain by PHDs, hydroxylation of the asparagine at residue 803 in human HIF-1 α , which is located in the transactivation domain, by a polypeptide termed factor inhibiting HIF-1 protein (FIH-1) has been found to regulate the activity of HIF-1 α by preventing its interaction with two co-activators—p300 and CBP (Lando et al., 2002a; Lando et al., 2002b). Acetylation of a lysine residue (Lys 532 in human HIF-1 α) has also been shown to regulate HIF-1 by enhancing the binding of HIF-1 α to VHL and its subsequent degradation (Jeong et al., 2002). The ARD1 acetyl transferase has been shown to be responsible for this acetylation.

[0061] Still another recently identified mechanism of HIF-1 regulation is fumarate-dependent. Intracellular fumarate is regulated by fumarate hydratase, an enzyme in the tricarboxylic acid (TCA) cycle. Mutations in this gene, which occur in hereditary leiomyomatosis, were shown to cause increased levels of intracellular fumarate. The increased fumarate can act as a competitive inhibitor of prolyl hydroxylase, causing increased level of HIF-1 α to accumulate (Isaacs et al., 2005). A similar function has been identified for succinate dehydrogenase (SDH), which is another member of the TCA cycle. Mutations of SDH, a candidate tumor suppressor for renal cell carcinoma, leads to increased succinate level, which has been shown to inhibit PHD activity and to lead to increased HIF-1 α levels (Selak et al., 2005).

[0062] In solid tumors, HIF-1 α activity is regulated via several mechanisms. Hypoxia is a common feature of all solid tumor microenvironments by virtue of the rapid proliferation of tumor cells and the generally poor functionality of newly formed tumor vasculature. Therefore, in the majority of solid tumors, hypoxia plays an important role in upregulating HIF-1 α activity (Harris, 2002). In fact, hypoxia-induced HIF-1 activation and subsequent VEGF expression has been postulated to be a major driving force in tumor angiogenesis in solid tumors (Maltepe et al., 1997; Harris, 2002). As a result, a significant effort is now being devoted to the development of HIF-1 inhibitors as anti-cancer drugs (Giaccia et al., 2003; Sutphin et al., 2004).

[0063] In addition to hypoxia, many hypoxia-independent pathways of HIF-1 regulation have been identified. These are mainly genetic/epigenetic alterations that can upregulate the level and/or activity of the HIF-1 α polypeptide. Loss of VHL (Maxwell et al., 1999; Ohh et al., 2000) and/or p53 gene function (Ravi et al., 2000; Chen et al., 2003; Sanchez-Puig et al., 2005), which decreases the ubiquitylation and subsequent degradation of HIF-1 α protein, can significantly upregulate HIF-1 activity. In addition, mutations in the

PTEN tumor suppressor gene (Zundel et al., 2000; Zhong et al., 2000), which increase activity of the PI3K-AKT-mTOR signaling pathway (Laughner et al., 2001; Chan et al., 2002); ERBB2 gain of function mutations (Laughner et al., 2001); increased EGFR (Zhong et al., 2000), MEK-ERK (Fukuda et al., 2002), and/or IGF-1R signaling (Fukuda et al., 2003); and SRC gain of function mutations (Jiang et al., 1997) can all cause increased synthesis of the HIF-1 α protein and overall HIF-1 activation.

[0064] In addition to tumor microenvironmental conditions and genetic/epigenetic changes in host tumor cells, it has recently been shown that HIF-1 activity can also be modified by exposure to radiotherapy (Moeller et al., 2004; Moeller et al., 2005). Exposure to ionizing radiation appears to activate HIF-1 via a hypoxia-independent mechanism. This activation appears to be mediated by a post-transcriptional mechanism that involves the release of pre-stored HIF-1 α -encoding mRNAs in "stress granules" located in the cytoplasm (Moeller et al., 2004). The triggering signals were identified to be free radical species induced by exposure to ionizing radiation.

[0065] This important discovery indicates that tumors respond to radiotherapy by activating HIF-1, which mediates the expression of VEGF and other factors that protect tumor vasculature against cytotoxic therapy, thereby increasing overall tumor cell survival. Consistent with this hypothesis are data indicating that combining radiotherapy with HIF-1 inhibitors appears to synergize their anti-tumor effects (Moeller et al., 2004).

[0066] Accordingly, disclosed herein is the identification of nitric oxide as a major regulator of HIF-1 α activity during cancer treatment. Thus, NO inhibitors can be employed as sensitizers of cancer radiation and chemotherapy.

[0067] Also disclosed herein is the discovery that an important cysteine (Cys 520) residue in the human HIF-1 α protein is responsible NO-mediated activation of HIF-1 α during cancer therapy. This residue serves as the site for nitrosylation and subsequent activation of the HIF-1 α during cancer therapy. The absence of this residue abolishes the induction of HIF-1 α . Therefore, Cys 520 and corresponding residues in HIF-1 α polypeptides from other species are targets for drug development.

[0068] Also disclosed herein is the discovery that compositions that can inhibit the production of nitric oxide can significantly increase therapeutic efficacy of radiotherapy through the inhibition of radiation-induced HIF-1 α upregulation. Therefore, inhibitors of nitric oxide production can act as sensitizers of radiation and cancer treatment.

[0069] Also disclosed herein is the administration of agents that can inhibit the production of NO and subsequent nitrosylation and stabilization of the HIF-1 α protein in tumors before, during, or after radiation therapy or cytotoxic chemotherapy. One rationale is that these agents would be expected to decrease the level of NO in the tumor microenvironment and cause a concomitant reduction of the level of HIF-1 α that is induced by radiation or chemotherapy. Because HIF-1 α has been shown to be a key angiogenesis regulator and survival factor for tumors during cancer therapy, the lower level of HIF-1 α should allow for a better therapeutic outcome.

[0070] In addition, agents that can inhibit the nitrosylation and activation of HIF-1 α either directly or indirectly can

also serve as inhibitors of anti-inflammatory agents. To that end, disclosed herein is the discovery that treatment of macrophages with inflammation-causing agents can result in the stabilization and activation of HIF-1 α . As HIF-1 α has been shown to be important in mediating inflammatory response, agents that inhibit the nitrosylation and stabilization of HIF-1 α can also be used as anti-inflammatory agents.

II. Definitions

[0071] While the following terms are believed to be well understood by one of ordinary skill in the art, the following definitions are set forth to facilitate explanation of the presently disclosed subject matter.

[0072] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which the presently disclosed subject matter belongs. Although any methods, devices, and materials similar or equivalent to those described herein can be used in the practice or testing of the presently disclosed subject matter, representative methods, devices, and materials are now described.

[0073] Following long-standing patent law convention, the terms "a", "an", and "the" refer to "one or more" when used in this application, including the claims. Thus, for example, reference to "a tumor cell" includes a plurality of such tumor cells, and so forth.

[0074] As used herein, the term "about," when referring to a value or to an amount of mass, weight, time, volume, concentration, or percentage is meant to encompass variations of in some embodiments, $\pm 20\%$, in some embodiments $\pm 10\%$, in some embodiments $\pm 5\%$, in some embodiments $\pm 1\%$, and in some embodiments $\pm 0.1\%$ from the specified amount, as such variations are appropriate to perform the disclosed methods.

[0075] As used herein, "significance" or "significant" relates to a statistical analysis of the probability that there is a non-random association between two or more entities. To determine whether or not a relationship is "significant" or has "significance", statistical manipulations of the data can be performed to calculate a probability, expressed as a "p value". Those p values that fall below a user-defined cutoff point are regarded as significant. In some embodiments, a p value less than or equal to 0.05, in some embodiments less than 0.01, in some embodiments less than 0.005, and in some embodiments less than 0.001, are regarded as significant. Accordingly, a p value greater than or equal to 0.05 is considered not significant.

[0076] As used herein, the term "subject" refers to any organism for which application of the presently disclosed subject matter would be desirable. The subject treated in the presently disclosed subject matter in its many embodiments is desirably a human subject, although it is to be understood that the principles of the presently disclosed subject matter indicate that the presently disclosed subject matter is effective with respect to all vertebrate species, including mammals, which are intended to be included in the term "subject". Moreover, a mammal is understood to include any mammalian species in which treatment of a tumor and/or a cancer is desirable, particularly agricultural and domestic mammalian species.

[0077] More particularly provided is the treatment of mammals such as humans, as well as those mammals of

importance due to being endangered (such as Siberian tigers), of economic importance (animals raised on farms for consumption by humans) and/or social importance (animals kept as pets or in zoos) to humans, for instance, carnivores other than humans (such as cats and dogs), swine (pigs, hogs, and wild boars), ruminants (such as cattle, oxen, sheep, giraffes, deer, goats, bison, and camels), and horses. Also provided is the treatment of birds, including the treatment of those kinds of birds that are endangered, kept in zoos, as well as fowl, and more particularly domesticated fowl, i.e., poultry, such as turkeys, chickens, ducks, geese, guinea fowl, and the like, as they are also of economic importance to humans. Thus, contemplated is the treatment of livestock, including, but not limited to, domesticated swine (pigs and hogs), ruminants, horses, poultry, and the like.

[0078] The terms “small interfering RNA”, “short interfering RNA”, and “siRNA” are used interchangeably and refer to any nucleic acid molecule capable of mediating RNA interference (RNAi) or gene silencing. See e.g., Bass, 2001; Elbashir et al., 2001; and PCT International Publication Nos. WO 99/07409; WO 99/32619; WO 00/01846; WO 00/44895; WO 00/44914; WO 01/36646; WO 01/29058. A non-limiting example of an siRNA molecule of the presently disclosed subject matter is shown in SEQ ID NO: 17. In some embodiments, the siRNA comprises a double stranded polynucleotide molecule comprising complementary sense and antisense regions, wherein the antisense region comprises a sequence complementary to a region of a target nucleic acid molecule (for example, an mRNA encoding VHL). In some embodiments, the siRNA comprises a single stranded polynucleotide having self-complementary sense and antisense regions, wherein the antisense region comprises a sequence complementary to a region of a target nucleic acid molecule. In some embodiments, the siRNA comprises a single stranded polynucleotide having one or more loop structures and a stem comprising self complementary sense and antisense regions, wherein the antisense region comprises a sequence complementary to a region of a target nucleic acid molecule, and wherein the polynucleotide can be processed either in vivo or in vitro to generate an active siRNA capable of mediating RNAi. As used herein, siRNA molecules need not be limited to those molecules containing only RNA, but further encompass chemically modified nucleotides and non-nucleotides.

[0079] The term “gene expression” generally refers to the cellular processes by which a biologically active polypeptide is produced from a DNA sequence and exhibits a biological activity in a cell. As such, gene expression involves the processes of transcription and translation, but also involves post-transcriptional and post-translational processes that can influence a biological activity of a gene or gene product. These processes include, but are not limited to RNA syntheses, processing, and transport, as well as polypeptide synthesis, transport, and post-translational modification of polypeptides. Additionally, processes that affect protein-protein interactions within the cell (for example, the interaction between HIF-1 α and VHL) can also affect gene expression as defined herein.

[0080] As used herein, the term “modulate” refers to a change in the expression level of a gene, or a level of RNA molecule or equivalent RNA molecules encoding one or more proteins or protein subunits, or activity of one or more

proteins or protein subunits is upregulated or downregulated, such that expression, level, and/or activity is greater than or less than that observed in the absence of the modulator. For example, the term “modulate” can mean “inhibit” or “suppress”, but the use of the word “modulate” is not limited to this definition.

[0081] As used herein, the terms “inhibit”, “suppress”, “downregulate”, and grammatical variants thereof are used interchangeably and refer to an activity whereby gene expression (e.g., a level of an RNA encoding one or more gene products) is reduced below that observed in the absence of a composition of the presently disclosed subject matter. In some embodiments, inhibition results in a decrease in the steady state level of a target RNA. In some embodiments, inhibition results in an expression level of a gene product that is below that level observed in the absence of the modulator.

[0082] In some embodiments, the terms “inhibit”, “suppress”, “downregulate”, and grammatical variants thereof refer to a biological activity of a polypeptide or polypeptide complex that is lower in the presence of a modulator than that which occurs in the absence of the modulator. For example, a modulator can inhibit the ability of a polypeptide (e.g., an HIF-1 polypeptide) to interact with its target (e.g., VHL and/or a promoter sequence comprising a hypoxia response element (HRE)). This can be accomplished by any mechanism, including but not limited to enhancing its existence in an inactive form (e.g., enhancing the complexing of an HIF-1 with VHL and/or inhibiting the dissociation of an HIF-1 from VHL) and/or by enhancing the rate of degradation of an HIF-1.

[0083] As used herein, the terms “gene” and “target gene” refer to a nucleic acid that encodes an RNA, for example, nucleic acid sequences including, but not limited to, structural genes encoding a polypeptide. The target gene can be a gene derived from a cell, an endogenous gene, a transgene, etc. The cell containing the target gene can be derived from or contained in any organism, for example an animal. The term “gene” also refers broadly to any segment of DNA associated with a biological function. As such, the term “gene” encompasses sequences including but not limited to a coding sequence, a promoter region, a transcriptional regulatory sequence, a non-expressed DNA segment that is a specific recognition sequence for regulatory proteins, a non-expressed DNA segment that contributes to gene expression, a DNA segment designed to have desired parameters, or combinations thereof. A gene can be obtained by a variety of methods, including cloning from a biological sample, synthesis based on known or predicted sequence information, and recombinant derivation of an existing sequence.

[0084] In some embodiments, a gene is a hypoxia-inducible gene. As used herein, a “hypoxia-inducible gene” is a gene for which the expression level increases in response to hypoxia. In some embodiments, a hypoxia-inducible gene is a gene that is characterized by upregulated transcription in response to hypoxic conditions. Exemplary hypoxia-inducible genes thus include genes with hypoxia response elements (HREs) in their promoters. Under hypoxic conditions, transcription of these genes is induced as a result of activated HIF-1 binding to the HREs. Also as used herein, a hypoxia-inducible gene is a gene for which an activity of the gene

product changes in response to hypoxia. In these embodiments, a hypoxia-inducible gene is a gene for which the polypeptide encoded by the gene experiences a change in state in response to hypoxia. Such a change in state includes, but is not limited to a post-transcriptional modification or an interaction with another molecule (for example, a protein-protein interaction). Thus, as used herein, the term hypoxia-inducible gene includes, but is not limited to HIF-1 α and VHL, each of which undergoes a change in state (in this example, a dissociation one from the other) in response to hypoxia.

[0085] As is understood in the art, a gene comprises a coding strand and a non-coding strand. As used herein, the terms “coding strand” and “sense strand” are used interchangeably, and refer to a nucleic acid sequence that has the same sequence of nucleotides as an mRNA from which the gene product is translated. As is also understood in the art, when the coding strand and/or sense strand is used to refer to a DNA molecule, the coding/sense strand includes thymidine residues instead of the uridine residues found in the corresponding mRNA. Additionally, when used to refer to a DNA molecule, the coding/sense strand can also include additional elements not found in the mRNA including, but not limited to promoters, enhancers, and introns. Similarly, the terms “template strand” and “antisense strand” are used interchangeably and refer to a nucleic acid sequence that is complementary to the coding/sense strand.

[0086] As used herein, the terms “complementarity” and “complementary” refer to a nucleic acid that can form one or more hydrogen bonds with another nucleic acid sequence by either traditional Watson-Crick or other non-traditional types of interactions.

[0087] As used herein, the phrase “percent complementarity” refers to the percentage of contiguous residues in a nucleic acid molecule that can form hydrogen bonds (e.g., Watson-Crick base pairing) with a second nucleic acid sequence (e.g., 5, 6, 7, 8, 9, 10 out of 10 being 50%, 60%, 70%, 80%, 90%, and 100% complementary). The terms “100% complementary”, “fully complementary”, and “perfectly complementary” indicate that all of the contiguous residues of a nucleic acid sequence can hydrogen bond with the same number of contiguous residues in a second nucleic acid sequence.

[0088] As used herein, the term “cell” is used in its usual biological sense. In some embodiments, the cell is present in an organism, for example, mammals such as humans, cows, sheep, apes, monkeys, swine, dogs, cats, and rodents. In some embodiments, the cell is a eukaryotic cell (e.g., a mammalian cell, such as a human cell). The cell can be of somatic or germ line origin, totipotent or pluripotent, dividing or non-dividing. The cell can also be derived from or can comprise a gamete or embryo, a stem cell, or a fully differentiated cell.

[0089] As used herein, the term “RNA” refers to a molecule comprising at least one ribonucleotide residue. By “ribonucleotide” is meant a nucleotide with a hydroxyl group at the 2' position of a β -D-ribofuranose moiety. The terms encompass double stranded RNA, single stranded RNA, RNAs with both double stranded and single stranded regions, isolated RNA such as partially purified RNA, essentially pure RNA, synthetic RNA, recombinantly produced RNA, as well as altered RNA, or analog RNA, that

differs from naturally occurring RNA by the addition, deletion, substitution, and/or alteration of one or more nucleotides. Such alterations can include addition of non-nucleotide material. Nucleotides in the RNA molecules of the presently disclosed subject matter can also comprise non-standard nucleotides, such as non-naturally occurring nucleotides or chemically synthesized nucleotides or deoxynucleotides. These altered RNAs can be referred to as analogs or analogs of a naturally occurring RNA.

[0090] As used herein, the phrase “double stranded RNA” refers to an RNA molecule at least a part of which is in Watson-Crick base pairing forming a duplex. As such, the term is to be understood to encompass an RNA molecule that is either fully or only partially double stranded. Exemplary double stranded RNAs include, but are not limited to molecules comprising at least two distinct RNA strands that are either partially or fully duplexed by intermolecular hybridization. Additionally, the term is intended to include a single RNA molecule that by intramolecular hybridization can form a double stranded region (for example, a hairpin). Thus, as used herein the phrases “intermolecular hybridization” and “intramolecular hybridization” refer to double stranded molecules for which the nucleotides involved in the duplex formation are present on different molecules or the same molecule, respectively.

[0091] As used herein, the phrase “double stranded region” refers to any region of a nucleic acid molecule that is in a double stranded conformation via hydrogen bonding between the nucleotides including, but not limited to hydrogen bonding between cytosine and guanosine, adenosine and thymidine, adenosine and uracil, and any other nucleic acid duplex as would be understood by one of ordinary skill in the art. The length of the double stranded region can vary from about 15 consecutive basepairs to several thousand basepairs.

[0092] As used herein, the terms “corresponds to”, “corresponding to”, and grammatical variants thereof refer to a nucleotide sequence that is 100% identical to at least 19 contiguous nucleotides of a nucleic acid sequence of a hypoxia-inducible gene. Thus, a first nucleic acid sequence that “corresponds to” a coding strand of a hypoxia-inducible gene is a nucleic acid sequence that is 100% identical to at least 19 contiguous nucleotides of a hypoxia-inducible gene, including, but not limited to 5' untranslated sequences, exon sequences, intron sequences, and 3' untranslated sequences.

[0093] The term “tumor” as used herein encompasses both primary and metastasized solid tumors and carcinomas of any tissue in a subject, including, but not limited to breast; colon; rectum; lung; oropharynx; hypopharynx; esophagus; stomach; pancreas; liver; gallbladder; bile ducts; small intestine; urinary tract including kidney, bladder and urothelium; female genital tract including cervix, uterus, ovaries (e.g., choriocarcinoma and gestational trophoblastic disease); male genital tract including prostate, seminal vesicles, testes and germ cell tumors; endocrine glands including thyroid, adrenal, and pituitary; skin (e.g., hemangiomas and melanomas), bone or soft tissues; blood vessels (e.g., Kaposi's sarcoma); brain, nerves, eyes, and meninges (e.g., astrocytomas, gliomas, glioblastomas, retinoblastomas, neuromas, neuroblastomas, Schwannomas and meningiomas). The term “tumor” also encompasses solid tumors arising from hematopoietic malignancies such as leukemias, including

chloromas, plasmacytomas, plaques and tumors of mycosis fungoides and cutaneous T-cell lymphoma/leukemia, and lymphomas including both Hodgkin's and non-Hodgkin's lymphomas. The term "tumor" also encompasses radioresistant and/or chemoresistant tumors, including, but not limited to radioresistant and/or chemoresistant variants of the any of the tumor listed above.

[0094] The terms "radiosensitivity" and "radiosensitive", as used herein to describe a tumor, refer to a quality of susceptibility to treatment using ionizing radiation. Thus, radiotherapy can be used to delay growth of a radiosensitive tumor. Radiosensitivity can be quantified by determining a minimal amount of ionizing radiation that can be used to delay tumor growth. Thus, the term "radiosensitivity" refers to a quantitative range of radiation susceptibility.

[0095] The terms "sensitivity to chemotherapy", "chemosensitivity", and "chemosensitive", as used herein to describe a tumor, refer to a quality of susceptibility to treatment using chemotherapy. Thus, chemotherapy can be used to delay growth of a tumor sensitive to chemotherapy. Sensitivity of chemotherapy can be quantified by determining a minimal dosage of chemotherapy that can be used to delay tumor growth. Thus, the phrase "sensitivity to chemotherapy" refers to a quantitative range of chemotherapy susceptibility.

[0096] The terms "radiation resistant tumor" and "radioresistant tumor" each generally refer to a tumor that is substantially unresponsive to radiotherapy when compared to other tumors. Representative radiation resistant tumor models include glioblastoma multiforme and melanoma. Similarly, the terms "chemotherapy resistant tumor" and "chemoresistant tumor" generally refer to a tumor that is substantially unresponsive to chemotherapy when compared to other tumors.

[0097] The term "delaying tumor growth" refers to a decrease in duration of time required for a tumor to grow a specified amount. For example, treatment with the compositions and/or methods disclosed herein can delay the time required for a tumor to increase in volume 3-fold relative to an initial day of measurement (day 0) or the time required to grow to 1 cm³.

[0098] The term "increase," as used herein to refer to a change in radiosensitivity and/or sensitivity to chemotherapy of a tumor, refers to change that renders a tumor more susceptible to destruction by ionizing radiation and/or chemotherapy. Alternatively stated, an increase in radiosensitivity and/or chemosensitivity refers to a decrease in the minimal amount of ionizing radiation and/or chemotherapy that effectively delays tumor growth. An increase in radiosensitivity and/or chemosensitivity can also comprise delayed tumor growth when a composition of the presently disclosed subject matter is administered with radiation and/or chemotherapy as compared to a same dose of radiation and/or chemotherapy alone. In some embodiments, an increase in radiosensitivity and/or chemosensitivity refers to an increase of at least about 2-fold, in some embodiments an increase of at least about 5-fold, and in some embodiments an increase of at least 10-fold. In some embodiments of the presently disclosed subject matter, an increase in radiosensitivity and/or chemosensitivity comprises a transformation of a radioresistant and/or chemoresistant tumor to a radiosensitive and/or chemosensitive tumor.

[0099] The term "tumor regression" generally refers to any one of a number of indices that suggest change within the tumor to a less developed form. Such indices include, but are not limited to a destruction of tumor vasculature (for example, a decrease in vascular length density or a decrease in blood flow), a decrease in tumor cell survival, a decrease in tumor volume, and/or a decrease in tumor growth rate. Methods for assessing tumor growth delay and tumor regression are known to the skilled artisan.

[0100] The term "nucleic acid molecule" refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. Unless specifically limited, the term encompasses nucleic acids containing known analogues of natural nucleotides that have similar properties as the reference natural nucleic acid. Unless otherwise indicated, a particular nucleotide sequence also implicitly encompasses complementary sequences, subsequences, elongated sequences, as well as the sequence explicitly indicated. The terms "nucleic acid molecule" or "nucleotide sequence" can also be used in place of "gene", "DNA", "cDNA", "RNA", or "mRNA". Nucleic acids can be derived from any source, including any organism.

[0101] The term "isolated", as used in the context of a nucleic acid molecule or polypeptide, indicates that the nucleic acid molecule or polypeptide exists apart from its native environment and is not a product of nature. An isolated nucleic acid molecule or polypeptide can exist in a purified form or can exist in a non-native environment such as a host cell.

[0102] The terms "identical" or percent "identity" in the context of two or more nucleotide or polypeptide sequences refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same, when compared and aligned for maximum correspondence, as measured using one of the sequence comparison algorithms disclosed herein or by visual inspection.

[0103] The term "substantially identical", in the context of two nucleotide sequences, refers to two or more sequences or subsequences that have in some embodiments at least 60%, in some embodiments about 70%, in some embodiments about 80%, in some embodiments about 90%, in some embodiments about 95%, in some embodiments, about 97%, and in some embodiments about 99% nucleotide identity, when compared and aligned for maximum correspondence, as measured using one of the following sequence comparison algorithms (described herein below) or by visual inspection. In some embodiments, the substantial identity exists in nucleotide sequences of at least 50 residues, in some embodiments in nucleotide sequence of at least about 100 residues, in some embodiments in nucleotide sequences of at least about 150 residues, and in some embodiments in nucleotide sequences comprising complete coding sequences.

[0104] In one aspect, polymorphic sequences can be substantially identical sequences. The terms "polymorphic", "polymorphism", and "polymorphic variants" refer to the occurrence of two or more genetically determined alternative sequences or alleles in a population. An allelic difference can be as small as one base pair. As used herein in regards to a nucleotide or polypeptide sequence, the term "substantially identical" also refers to a particular sequence

that varies from another sequence by one or more deletions, substitutions, or additions, the net effect of which is to retain biological activity of a gene, gene product, or sequence of interest.

[0105] For sequence comparison, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer program, subsequence coordinates are designated if necessary, and sequence algorithm program parameters are selected. The sequence comparison algorithm then calculates the percent sequence identity for the designated test sequence(s) relative to the reference sequence, based on the selected program parameters.

[0106] Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, 1981, by the homology alignment algorithm of Needleman & Wunsch, 1970, by the search for similarity method for Pearson & Lipman, 1988, by computerized implementations of these algorithms (GAP, BEST-FIT, FASTA, and TFASTA, in the Wisconsin Genetics Software Package, available from Accelrys Inc., San Diego, Calif., United States of America), or by visual inspection. See generally, Ausubel, 1995.

[0107] In some embodiments, an algorithm for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described by Altschul et al., 1990. Software for performing BLAST analyses is publicly available through the website of the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold. These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when the cumulative alignment score falls off by the quantity X from its maximum achieved value, the cumulative score goes to zero or below due to the accumulation of one or more negative-scoring residue alignments, or the end of either sequence is reached. The BLAST algorithm parameters W , T , and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength $W=11$, an expectation $E=10$, a cutoff of 100, $M=5$, $N=-4$, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix. See Henikoff & Henikoff, 1992.

[0108] In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences. See e.g., Karlin & Altschul, 1993. One measure of similarity provided by the

BLAST algorithm is the smallest sum probability ($P(N)$), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a test nucleic acid sequence is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid sequence to the reference nucleic acid sequence is in some embodiments less than about 0.1, in some embodiments less than about 0.01, and in some embodiments less than about 0.001.

[0109] Another indication that two nucleotide sequences are substantially identical is that the two molecules specifically or substantially hybridize to each other under stringent conditions. In the context of nucleic acid hybridization, two nucleic acid sequences being compared can be designated a “probe” and a “target”. A “probe” is a reference nucleic acid molecule, and a “target” is a test nucleic acid molecule, often found within a heterogeneous population of nucleic acid molecules. A “target sequence” is synonymous with a “test sequence”.

[0110] The phrase “hybridizing substantially to” refers to complementary hybridization between a probe nucleic acid molecule and a target nucleic acid molecule and embraces minor mismatches that can be accommodated by reducing the stringency of the hybridization media to achieve the desired hybridization.

[0111] “Stringent hybridization conditions” and “stringent hybridization wash conditions” in the context of nucleic acid hybridization experiments such as Southern and Northern blot analysis are both sequence- and environment-dependent. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, 1993. Generally, highly stringent hybridization and wash conditions are selected to be about 5° C. lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. Typically, under “stringent conditions” a probe will hybridize specifically to its target subsequence, but to no other sequences.

[0112] The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Very stringent conditions are selected to be equal to the T_m for a particular probe. An example of highly stringent hybridization conditions for Southern or Northern Blot analysis of complementary nucleic acids having more than about 100 complementary residues is overnight hybridization in 50% formamide with 1 mg of heparin at 42° C. An example of highly stringent wash conditions is 15 minutes in 0.1× standard saline citrate (SSC), 0.1% (w/v) SDS at 65° C. Another example of highly stringent wash conditions is 15 minutes in 0.2×SSC buffer at 65° C. (see Sambrook & Russell, 2001 for a description of SSC buffer and other stringency conditions). Often, a high stringency wash is preceded by a lower stringency wash to remove background probe signal. An example of medium stringency wash conditions for a duplex of more than about 100 nucleotides is 15 minutes in 1×SSC at 45° C. Another example of medium stringency wash for a duplex of more than about 100 nucleotides is 15 minutes in 4-6×SSC at 40° C. For short probes (e.g., about 10 to 50 nucleotides), stringent conditions typically involve salt concentrations of less than about 1M Na^+ ion, typically about

0.01 to 1M Na⁺ ion concentration (or other salts) at pH 7.0-8.3, and the temperature is typically at least about 30° C. Stringent conditions can also be achieved with the addition of destabilizing agents such as formamide. In general, a signal to noise ratio of 2-fold or higher than that observed for an unrelated probe in the particular hybridization assay indicates detection of a specific hybridization.

[0113] The following are examples of hybridization and wash conditions that can be used to clone homologous nucleotide sequences that are substantially identical to reference nucleotide sequences of the presently disclosed subject matter: a probe nucleotide sequence hybridizes in one example to a target nucleotide sequence in 7% sodium dodecyl sulfate (SDS), 0.5M NaPO₄, 1 mM EDTA at 50° C. followed by washing in 2×SSC, 0.1% SDS at 50° C.; in another example, a probe and target sequence hybridize in 7% sodium dodecyl sulfate (SDS), 0.5M NaPO₄, 1 mM EDTA at 50° C. followed by washing in 1×SSC, 0.1% SDS at 50° C.; in another example, a probe and target sequence hybridize in 7% sodium dodecyl sulfate (SDS), 0.5M NaPO₄, 1 mM EDTA at 50° C. followed by washing in 0.5×SSC, 0.1% SDS at 50° C.; in another example, a probe and target sequence hybridize in 7% sodium dodecyl sulfate (SDS), 0.5M NaPO₄, 1 mM EDTA at 50° C. followed by washing in 0.1×SSC, 0.1% SDS at 50° C.; in yet another example, a probe and target sequence hybridize in 7% sodium dodecyl sulfate (SDS), 0.5M NaPO₄, 1 mM EDTA at 50° C. followed by washing in 0.1×SSC, 0.1% SDS at 65° C.

[0114] The term “subsequence” refers to a sequence of a nucleic acid or polypeptide that comprises a part of a longer nucleic acid or polypeptide sequence.

[0115] The term “elongated sequence” refers to an addition of nucleotides (or other analogous molecules) or amino acid residues incorporated into the nucleic acid or polypeptide. For example, a polymerase (e.g., a DNA polymerase) can add sequences at the 3' terminus of the nucleic acid molecule. In addition, the nucleotide sequence can be combined with other DNA sequences, such as promoters, promoter regions, enhancers, polyadenylation signals, intronic sequences, additional restriction enzyme sites, multiple cloning sites, and other coding segments.

[0116] The terms “operatively linked” and “operably linked”, as used herein, refer to a nucleic acid molecule in which a promoter region is connected to a nucleotide sequence in such a way that the transcription of that nucleotide sequence is controlled and regulated by the promoter region. Similarly, a nucleotide sequence is said to be under the “transcriptional control” of a promoter to which it is operably linked. Techniques for operatively linking a promoter region to a nucleotide sequence are known in the art.

[0117] The terms “heterologous gene”, “heterologous DNA sequence”, “heterologous nucleotide sequence”, “exogenous nucleic acid molecule”, or “exogenous DNA segment”, as used herein, each refer to a sequence that originates from a source foreign to an intended host cell and/or, if from the same source, is modified from its original form. Thus, a heterologous gene in a host cell includes a gene that is endogenous to the particular host cell but has been modified, for example by mutagenesis and/or by isolation from native transcriptional regulatory sequences. The terms also include non-naturally occurring multiple copies

of a naturally occurring nucleotide sequence. Thus, the terms refer in some embodiments to a DNA segment that is foreign or heterologous to the cell, or is homologous to the cell but in a position within the host cell nucleic acid wherein the element is not ordinarily found.

[0118] The term “expression vector” as used herein refers to a nucleotide sequence capable of directing expression of a particular nucleotide sequence in an appropriate host cell, comprising a promoter operatively linked to the nucleotide sequence of interest which is operatively linked to termination signals. It also typically comprises sequences required for proper translation of the nucleotide sequence. The construct comprising the nucleotide sequence of interest can be chimeric. The construct can also be one that is naturally occurring but has been obtained in a recombinant form useful for heterologous expression.

[0119] The term “promoter” or “promoter region” each refers to a nucleotide sequence within a gene that is positioned 5' to a coding sequence and functions to direct transcription of the coding sequence. The promoter region comprises a transcriptional start site, and can additionally include one or more transcriptional regulatory elements. In some embodiments, a method for the presently disclosed subject matter employs a hypoxia inducible promoter.

[0120] A “minimal promoter” is a nucleotide sequence that has the minimal elements required to enable basal level transcription to occur. As such, minimal promoters are not complete promoters but rather are subsequences of promoters that are capable of directing a basal level of transcription of a reporter construct in an experimental system. Minimal promoters include but are not limited to the CMV minimal promoter, the HSV-tk minimal promoter, the simian virus 40 (SV40) minimal promoter, the human β -actin minimal promoter, the human EF2 minimal promoter, the adenovirus E1B minimal promoter, and the heat shock protein (hsp) 70 minimal promoter. Minimal promoters are often augmented with one or more transcriptional regulatory elements to influence the transcription of an operably linked gene. For example, cell-type-specific or tissue-specific transcriptional regulatory elements can be added to minimal promoters to create recombinant promoters that direct transcription of an operably linked nucleotide sequence in a cell-type-specific or tissue-specific manner.

[0121] Different promoters have different combinations of transcriptional regulatory elements. Whether or not a gene is expressed in a cell is dependent on a combination of the particular transcriptional regulatory elements that make up the gene's promoter and the different transcription factors that are present within the nucleus of the cell. As such, promoters are often classified as “constitutive”, “tissue-specific”, “cell-type-specific”, or “inducible”, depending on their functional activities in vivo or in vitro. For example, a constitutive promoter is one that is capable of directing transcription of a gene in a variety of cell types. Exemplary constitutive promoters include the promoters for the following genes which encode certain constitutive or “housekeeping” functions: hypoxanthine phosphoribosyl transferase (HPRT), dihydrofolate reductase (DHFR); (Scharfmann et al., 1991), adenosine deaminase, phosphoglycerate kinase (PGK), pyruvate kinase, phosphoglycerate mutase, the β -actin promoter (see e.g., Williams et al., 1993), and other constitutive promoters known to those of skill in the art.

“Tissue-specific” or “cell-type-specific” promoters, on the other hand, direct transcription in some tissues and cell types but are inactive in others. Exemplary tissue-specific promoters include the PSA promoter (Yu et al., 1999; Lee et al., 2000), the probasin promoter (Greenberg et al., 1994; Yu et al., 1999), and the MUC1 promoter (Kurihara et al., 2000) as discussed above, as well as other tissue-specific and cell-type specific promoters known to those of skill in the art.

[0122] The term “transcriptional regulatory sequence” or “transcriptional regulatory element”, as used herein, each refers to a nucleotide sequence within the promoter region that enables responsiveness to a regulatory transcription factor. Responsiveness can encompass a decrease or an increase in transcriptional output and is mediated by binding of the transcription factor to the DNA molecule comprising the transcriptional regulatory element.

[0123] The term “transcription factor” generally refers to a protein that modulates gene expression by interaction with the transcriptional regulatory element and cellular components for transcription, including RNA polymerase, Transcription Associated Factors (TAFs), chromatin-remodeling proteins, and any other relevant protein that impacts gene transcription.

[0124] The terms “reporter gene” or “marker gene” or “selectable marker” each refer to a heterologous gene encoding a product that is readily observed and/or quantitated. A reporter gene is heterologous in that it originates from a source foreign to an intended host cell or, if from the same source, is modified from its original form. Non-limiting examples of detectable reporter genes that can be operatively linked to a transcriptional regulatory region can be found in Alam & Cook, 1990 and PCT International Publication No. WO 97/47763. Exemplary reporter genes for transcriptional analyses include the lacZ gene (see e.g., Rose & Botstein, 1983), Green Fluorescent Protein (GFP; Cubitt et al., 1995), luciferase, and chloramphenicol acetyl transferase (CAT). Reporter genes for methods to produce transgenic animals include but are not limited to antibiotic resistance genes, for example the antibiotic resistance gene confers neomycin resistance. Any suitable reporter and detection method can be used, and it will be appreciated by one of skill in the art that no particular choice is essential to or a limitation of the presently disclosed subject matter.

[0125] An amount of reporter gene can be assayed by any method for qualitatively or quantitatively determining presence or activity of the reporter gene product. The amount of reporter gene expression directed by each test promoter region fragment is compared to an amount of reporter gene expression to a control construct comprising the reporter gene in the absence of a promoter region fragment. A promoter region fragment is identified as having promoter activity when there is significant increase in an amount of reporter gene expression in a test construct as compared to a control construct. The term “significant increase”, as used herein, refers to a quantified change in a measurable quality that is larger than the margin of error inherent in the measurement technique, in one example an increase by about 2-fold or greater relative to a control measurement, in another example an increase by about 5-fold or greater, and in yet another example an increase by about 10-fold or greater.

[0126] Nucleic acids of the presently disclosed subject matter can be cloned, synthesized, recombinantly altered, mutagenized, or combinations thereof. Standard recombinant DNA and molecular cloning techniques used to isolate nucleic acids are known in the art. Exemplary, non-limiting methods are described by Silhavy et al., 1984; Ausubel et al., 1992; Glover & Hames, 1995; and Sambrook & Russell, 2001). Site-specific mutagenesis to create base pair changes, deletions, or small insertions is also known in the art as exemplified by publications (see e.g., Adelman et al., 1983; Sambrook & Russell, 2001).

III. Compositions

[0127] The compositions disclosed herein can be employed in vitro and/or in vivo in order to perform the disclosed methods. In some embodiments, the compositions described herein comprise an agent selected from the group consisting of an inhibitor of nitric oxide synthase, a nitric oxide scavenger, an inhibitor of NIF-1 nitrosylation, and combinations thereof.

[0128] III.A. Inhibitors of Nitric Oxide Synthase(s)

[0129] In some embodiments, an agent comprises an inhibitor of nitric oxide synthase (NOS). As is known in the art, there are several nitric oxide synthases, including but not limited to neural/neuronal NOS (nNOS; also referred to as NOS1), inducible NOS (iNOS; also referred to as NOS2), and endothelial NOS (eNOS; also referred to as NOS3). Nucleic acid and amino acid sequences for each of these NOS gene products are present in the GENBANK® database, each of which is expressly incorporated by reference herein in its entirety. For example, human NOS sequences present in the GENBANK® database include GENBANK® Accession Nos. U17327 and AAA62405 (nNOS nucleic acid and amino acid sequences, respectively), NM_000625 and NP_000616 (iNOS nucleic acid and amino acid sequences, respectively), and BC069465 and AAH69465 (eNOS nucleic acid and amino acid sequences, respectively).

[0130] Various inhibitors of NOS have been identified, some of which are selective and other of which are non-selective for one or more specific NOS type. As used herein, a “selective” or “specific” NOS inhibitor demonstrates markedly greater specificity for one of the NOS types (e.g., iNOS) than it does for the other two (e.g., eNOS and nNOS), while “non-selective” or “non-specific” NOS inhibitors demonstrate approximately equivalent inhibition of two or more of the NOS types. Both non-selective and selective NOS inhibitors are appropriate for use in the methods and compositions for the presently disclosed subject matter. Representative NOS inhibitors thus include, but are not limited to L-N(6)-(1-iminoethyl)lysine tetrazole-amide (SC-51); aminoguanidine (AG); guanidinoethylthiol; L-NG-nitroarginine methyl ester; mercaptoethylguanidine (MEG); N^o-nitro-L-arginine methyl ester (L-NAME); N-(3-(aminomethyl)benzyl)acetamide (1400W); N^G-monomethyl-L-arginine (L-NMMA); 7-nitroindazole (7-NI), L-NIL(N⁶-(1-iminoethyl)-lysine (L-NIL); and N⁵-(1-iminoethyl)-L-ornithine (L-NIO); as well as their pharmaceutically acceptable salts and other derivatives.

[0131] III.B. Nitric Oxide Scavengers

[0132] In some embodiments, a composition of the presently disclosed subject matter comprises a nitric oxide (NO)

scavenger. As used herein, the phrase “nitric oxide scavenger” refers to a molecule that binds to nitric oxide or otherwise makes the nitric oxide less available to take part in a biochemical process within a cell. Nitric oxide scavengers are also known, and include, but are not limited to vitamin B12, particularly in the hydroxocobalamin form; 2-(4-carboxyphenyl)-4,4,5,5-tetramethylimidazoline-1-oxyl-3-oxide (carboxy-PTIO); diethyldithiocarbamate; AMD6221 (ruthenium[hydrogen(diethylenetri-nitrilo)pentaacetato]chloride); and N-dithiocarboxy-sarcosine (DTCS), as well as pharmaceutically acceptable salts and other derivatives thereof.

[0133] III.C. Inhibitors of HIF-1 Nitrosylation

[0134] In some embodiments, a composition of the presently disclosed subject matter comprises an inhibitor of HIF-1 nitrosylation. As used herein, the phrase “inhibitor of HIF-1 nitrosylation” refers to any molecule that inhibits, either completely or partially, nitrosylation of an HIF-1 polypeptide. As such, this phrase encompasses NOS inhibitors, NO scavengers, and any other molecule that can inhibit the nitrosylation of an HIF-1 polypeptide. Representative other molecules include, but are not limited to peptides, peptide mimetics, proteins, antibodies or fragments thereof, small molecules, nucleic acids, and combinations thereof. The term “small molecule” as used herein refers to a compound, for example an organic compound, with a molecular weight in one example of less than about 1,000 daltons, in another example less than about 750 daltons, in another example less than about 600 daltons, and in yet another example less than about 500 daltons. A small molecule also has in one example a computed log octanol-water partition coefficient in the range of about -4 to about +14, more preferably in the range of about -2 to about +7.5.

[0135] As is known in the art, polypeptides such as HIF-1 can be S-nitrosylated on cysteine residues. As disclosed herein, nitrosylation of C533 of murine HIF-1 α (which corresponds to C520 of human HIF-1 α) interferes with the interaction between HIF-1 α and VHL, which in turn inhibits the ubiquitylation and subsequent degradation of HIF-1 α by the proteasome. S-nitrosylation of HIF-1 α at this highly conserved cysteine (see also SEQ ID NOs. 1-21 showing the conservation of this cysteine in various animal species) thus results in an increased persistence of HIF-1 α in the cell, leading to higher HIF-1 α activity. Accordingly, inhibitors of HIF-1 α nitrosylation can be employed to reduce HIF-1 activity in cells.

[0136] III.D. Formulations

[0137] The compositions of the presently disclosed subject matter comprise in some embodiments a composition that includes a pharmaceutically acceptable carrier. Any suitable pharmaceutical formulation can be used to prepare the adenovirus vectors for administration to a subject.

[0138] For example, suitable formulations can include aqueous and non-aqueous sterile injection solutions which can contain anti-oxidants, buffers, bacteriostats, bactericidal antibiotics and solutes which render the formulation isotonic with the bodily fluids of the intended recipient; and aqueous and non-aqueous sterile suspensions which can include suspending agents and thickening agents. The formulations can be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and can be stored in a

frozen or freeze-dried (lyophilized) condition requiring only the addition of sterile liquid carrier, for example water for injections, immediately prior to use. Some exemplary ingredients are SDS, in one example in the range of 0.1 to 10 mg/ml, in another example about 2.0 mg/ml; and/or mannitol or another sugar, for example in the range of 10 to 100 mg/ml, in another example about 30 mg/ml; and/or phosphate-buffered saline (PBS).

[0139] It should be understood that in addition to the ingredients particularly mentioned above the formulations of this presently disclosed subject matter can include other agents conventional in the art having regard to the type of formulation in question. For example, sterile pyrogen-free aqueous and non-aqueous solutions can be used.

[0140] The therapeutic regimens and compositions of the presently disclosed subject matter can be used with additional adjuvants or biological response modifiers including, but not limited to, the cytokines IFN- α , IFN- γ , IL2, IL4, IL6, TNF, or other cytokine affecting immune cells. In accordance with this aspect of the presently disclosed subject matter, the disclosed nucleic acid molecules can be administered in combination therapy with one or more of these cytokines.

[0141] III.E. Administration

[0142] Administration of the compositions of the presently disclosed subject matter can be by any method known to one of ordinary skill in the art, including, but not limited to intravenous administration, intrasynovial administration, transdermal administration, intramuscular administration, subcutaneous administration, topical administration, rectal administration, intravaginal administration, intratumoral administration, oral administration, buccal administration, nasal administration, parenteral administration, inhalation, and insufflation. In some embodiments, suitable methods for administration of a composition of the presently disclosed subject matter include, but are not limited to intravenous or intratumoral injection. Alternatively, a composition can be deposited at a site in need of treatment in any other manner, for example by spraying a composition comprising a composition within the pulmonary pathways. The particular mode of administering a composition of the presently disclosed subject matter depends on various factors, including the distribution and abundance of cells to be treated, whether a vector is employed, additional tissue- or cell-targeting features of the vector and/or composition, and mechanisms for metabolism or removal of the composition from its site of administration. For example, relatively superficial tumors can be injected intratumorally. By contrast, internal tumors can be treated by intravenous injection.

[0143] In some embodiments, the method for administration encompasses features for regionalized delivery or accumulation at the site in need of treatment. In some embodiments, a composition is delivered intratumorally. In some embodiments, selective delivery of a composition to a tumor is accomplished by intravenous injection of the composition.

[0144] Alternatively or in addition, a composition of the presently disclosed subject matter can be provided at a pre-determined site using an implantable device containing the composition, whereby longer term delivery of the composition to a target tissue can be accomplished. Representative implantable devices are known in the art. For

example, absorbable thermoplastic elastomers have been developed to address the need in medical device development for an elastic material (e.g., U.S. Pat. Nos. 5,468,253 and 5,713,920). In addition, absorbable polymeric liquids and pastes have been developed to increase the range of physical properties exhibited by the aliphatic polyesters based on glycolide, lactide, p-dioxanone, 5,5-dimethyl-1,3-dioxan-2-one, trimethylene carbonate, and ϵ -caprolactone (e.g., U.S. Pat. Nos. 5,411,554; 5,599,852; 5,631,015; 5,653,992; 5,688,900; 5,728,752; and 5,824,333).

[0145] U.S. Pat. Nos. 5,573,934 and 5,858,746 (both to Hubbell et al.) disclosed the use of photocurable polymers to encapsulate biological materials including drugs, proteins, and cells in a hydrogel. The hydrogel was formed from a water soluble biocompatible macromer containing at least two free radical polymerizable substituents and either a thermal or light activated free radical initiator. An example of such a photoreactive system is an acrylate ester end-capped poly(ethylene glycol) containing ethyl eosin and a tertiary amine. After a series of light activated reactions between ethyl eosin and the amine, the acrylate endgroups polymerize into short segments that result in a crosslinked polymeric network composed of poly(ethylene glycol) chains radiating outward from the acrylate oligomers. The physical and mechanical properties of the resulting hydrogel are dependent on the reproducibility of the free radical oligomerization reaction.

[0146] U.S. Pat. No. 5,410,016 in the form of photocurable, segmented block copolymers composed not only of water soluble segments, such as poly(ethylene glycol), but also of segments with hydrolyzable groups, in particular, with short segments of aliphatic polyesters. In this way, the resulting hydrogel breaks down into soluble units in vitro and in vivo in a controlled fashion. The photochemistry is the same and based on the free radical polymerization of acrylate and methacrylate endgroups.

[0147] Other implantable devices are described in U.S. Pat. Nos. 7,009,034; 7,011,842; and 7,012,126.

[0148] For delivery of compositions to pulmonary pathways, the presently disclosed subject matter can be formulated as an aerosol or coarse spray. Methods for preparation and administration of aerosol or spray formulations can be found, for example, in Cipolla et al., 2000, and in U.S. Pat. Nos. 5,858,784; 6,013,638; 6,022,737; and 6,136,295.

[0149] III.F. Dosage

[0150] An effective dose of a composition of the presently disclosed subject matter is administered to a subject in need thereof. A "therapeutically effective amount" is an amount of the composition sufficient to produce a measurable response (e.g., a cytolytic response in a subject being treated). In some embodiments, an activity that inhibits tumor growth is measured. Actual dosage levels of active ingredients in the compositions of the presently disclosed subject matter can be varied so as to administer an amount of the active compound(s) that is effective to achieve the desired therapeutic response for a particular subject. The selected dosage level can depend upon the activity of the therapeutic composition, the route of administration, combination with other drugs or treatments, the severity of the condition being treated, and the condition and prior medical history of the subject being treated. However, it is within the skill of the

art to start doses of the compositions at levels lower than required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

[0151] The potency of a composition can vary, and therefore a "therapeutically effective" amount can vary. However, one skilled in the art can readily assess the potency and efficacy of a candidate composition of the presently disclosed subject matter and adjust the therapeutic regimen accordingly.

[0152] After review of the disclosure of the presently disclosed subject matter presented herein, one of ordinary skill in the art can tailor the dosages to an individual patient, taking into account the particular formulation, method for administration to be used with the composition, and tumor size. Further calculations of dose can consider patient height and weight, severity and stage of symptoms, and the presence of additional deleterious physical conditions. Such adjustments or variations, as well as evaluation of when and how to make such adjustments or variations, are well known to those of ordinary skill in the art of medicine.

[0153] For example, for local administration of viral expression vectors, previous clinical studies have demonstrated that up to 10^{13} plaque-forming units (pfu) of virus can be injected with minimal toxicity. In human patients, 1×10^9 – 1×10^{13} pfu are routinely used (see Habib et al., 1999). To determine an appropriate dose within this range, preliminary treatments can begin with 1×10^9 pfu, and the dose level can be escalated in the absence of dose-limiting toxicity. Toxicity can be assessed using criteria set forth by the National Cancer Institute and is reasonably defined as any grade 4 toxicity or any grade 3 toxicity persisting more than 1 week. Dose is also modified to maximize anti-tumor or anti-angiogenic activity. Representative criteria and methods for assessing anti-tumor and/or anti-angiogenic activity are described herein below. With replicative virus vectors, a dosage of about 1×10^7 to 1×10^8 pfu can be used in some instances.

IV. Applications

[0154] The presently disclosed subject matter provides methods for inhibiting nitric oxide synthase(s) activity in a cell in a subject. In some embodiments, the methods comprise administering to the cell in the subject a composition comprising (a) an agent selected from the group consisting of (i) an inhibitor of nitric oxide synthase; (ii) a nitric oxide scavenger; (iii) an inhibitor of HIF-1 nitrosylation; and (iv) combinations thereof, whereby nitric oxide synthase activity in the cell is inhibited. This general strategy can be employed in several areas, as disclosed in more detail hereinbelow.

[0155] IV.A. Methods for Inhibiting HIF-1 Activity

[0156] In some embodiments, the presently disclosed subject matter provides methods and compositions for inhibiting HIF-1 activity in a cell. In some embodiments, the cell is a tumor cell, and in some embodiments, the tumor cell is present within a subject including, but not limited to a mammals such as a human.

[0157] As disclosed herein, the presently disclosed subject matter provides compositions for inhibiting HIF-1 activity in a cell by inhibiting nitrosylation of HIF-1 α , which in turn

results in enhanced degradation of HIF-1 α mediated by the ubiquitin-proteasome pathway. As also disclosed herein, a highly conserved cysteine residue has been found to be a site for nitrosylation. This highly conserved cysteine corresponds to the positions listed in Table 1.

TABLE 1

Conserved Cysteines in HIF-1 α from Various Species	
Species	Cys Position
<i>Spalax judaei</i>	520 of SEQ ID NO: 1
<i>Eospalax baileyi</i>	521 of SEQ ID NO: 2
<i>Mus musculus</i>	533 of SEQ ID NO: 3
<i>Rattus norvegicus</i>	520 of SEQ ID NO: 4
<i>Microtus oeconomus</i>	520 of SEQ ID NO: 5
<i>Homo sapiens</i>	520 of SEQ ID NO: 6
<i>Pongo pygmaeus</i>	521 of SEQ ID NO: 7
<i>Macaca fascicularis</i>	520 of SEQ ID NO: 8
<i>Spermophilus tridecemlineatus</i>	520 of SEQ ID NO: 9
<i>Bos taurus</i>	520 of SEQ ID NO: 10
<i>Pantholops hodgsonii</i>	520 of SEQ ID NO: 11
<i>Canis familiaris</i>	520 of SEQ ID NO: 12
<i>Oryctolagus cuniculus</i>	520 of SEQ ID NO: 13
<i>Gallus gallus</i>	518 of SEQ ID NO: 14
<i>Xenopus laevis</i>	516 of SEQ ID NO: 16

Accordingly, the methods and compositions disclosed herein inhibit HIF-1 activity in some embodiments by inhibiting nitrosylation of the listed cysteine residues.

[0158] IV.B. Methods for Treating Tumor Cells and/or Cancer Cells

[0159] The presently disclosed methods and compositions can also be employed for treatment of tumor cells and/or cancer cells. In some embodiments, the methods comprise contacting the tumor cell and/or the cancer cell with the presently disclosed compositions (e.g., by administering the compositions to a subject that has the tumor cell and/or the cancer cell). In some embodiments, the compositions are selected from the group consisting of inhibitors of nitric oxide synthase, nitric oxide scavengers, inhibitors of HIF-1 nitrosylation, and combinations thereof. In some embodiments, the methods and compositions disclosed herein treat the tumor cell and/or the cancer cell by inhibiting HIF-1 activity in the tumor cell and/or the cancer cell.

[0160] As such, the methods and compositions can act directly on the tumor cell and/or the cancer cell to modulate its growth and/or proliferation. However, the methods and compositions can also act indirectly on the tumor cell and/or the cancer cell to modulate its growth and/or proliferation by inhibiting HIF-1 activity in other cells that influence the growth and/or proliferation of the tumor cell and/or the cancer cell. For example, the methods and compositions can inhibit HIF-1 activity in tumor blood vessels (i.e., those blood vessels and other endothelial cells that provide nutrients and remove waste products from the tumor and/or cancer cells) and/or can inhibit tumor angiogenesis modulated by HIF-1 activity. While applicants do not wish to be bound by any particular theory of operation, the methods and compositions disclosed herein can be employed to interfere with the function and/or generation of tumor vasculature, thereby modulating tumor cell and/or cancer cell growth and/or proliferation.

[0161] Additionally, the methods and compositions disclosed herein can be employed for increasing the sensitivity

of a tumor cell and/or a cancer cell to a treatment, such as surgical resection, radiotherapy, and/or chemotherapy, as discussed in more detail hereinbelow. As used herein, the phrase "increasing the sensitivity of a tumor cell and/or a cancer cell to a treatment" refers to an enhancement of the effect that a combination treatment including use of the presently disclosed methods and compositions has on tumor cell and/or cancer cell growth and/or proliferation as compared to the effect that a treatment would have had under the same conditions absent use of the presently disclosed methods and compositions. In some embodiments, the combination treatment employs the methods and/or compositions disclosed herein in conjunction with surgical resection, radiotherapy, and/or chemotherapy, and in some embodiments, the inclusion of a treatment comprising the methods and/or compositions disclosed herein results in a synergistic (i.e., more than additive) effect. Given the current limitations of surgical resection, radiotherapy, and/or chemotherapy, the presently disclosed subject matter provides an additional therapy that can be used to increase the efficacy of medical treatments directed towards modulating tumor cell and/or cancer cell growth and proliferation.

[0162] IV.C. Methods for Inhibiting an Inflammatory Response

[0163] The methods and compositions disclosed herein can also be employed for inhibiting inflammatory responses of cells (i.e., a cell in a subject). Also disclosed herein is the discovery that treatment of macrophages with inflammation-causing agents can result in the stabilization and activation of HIF-1 α . As HIF-1 α has been shown to be important in mediating inflammatory response, methods and compositions that inhibit the nitrosylation and stabilization of HIF-1 α can also be used as anti-inflammatory agents.

[0164] IV.D. Methods and Compositions for Identifying New Nitrosylation Inhibitors

[0165] The presently disclosed subject matter also provides screening methods and compositions that can be employed for identifying potential inhibitors of nitrosylation of an HIF-1 polypeptide. In some embodiments, the methods comprise (a) providing a cell comprising a nucleic acid a nucleotide sequence comprising any of SEQ ID NOs: 18-21; (b) contacting the cell with a compound comprising a potential inhibitor of nitrosylation of an HIF-1 polypeptide; and (c) assaying nitrosylation of a cysteine residue present in the nucleic acid, whereby an inhibitor of nitrosylation of an HIF-1 polypeptide is identified. In some embodiments, the methods further comprise comparing a level of nitrosylation of the cysteine residue present in the nucleic acid to a level of nitrosylation of the cysteine residue present in the nucleic acid prior to the contacting step.

[0166] In some embodiments, the nucleic acid comprises an expression vector in which the nucleic acid is operably linked to a promoter that is active in the cell. In some embodiments, the expression vector is a transgene and the animal is a transgenic animal that expresses the nucleic acid. In some embodiments, the compound is administered to the transgenic animal via a route that results in the compound contacting the cell. In some embodiments, the cell is an in vitro cultured cell and the contacting is performed in vitro.

[0167] In some embodiments, the cell (e.g., an in vitro cultured cell or a cell in a transgenic animal) comprises an

expression construct comprising one or more of SEQ ID NOs: 18-21 operably linked to a promoter. In some embodiments, the cell (e.g., an in vitro cultured cell or a cell in a transgenic animal) comprises an expression construct comprising one or more of SEQ ID NOs: 18-21 operably linked to a promoter, with the proviso that all cysteine residues present within SEQ ID NOs: 18-21 have been replaced with a non-nitrosylatable amino acid. An exemplary non-nitrosylatable amino acid is serine.

[0168] The cells comprising the expression construct comprising one or more of SEQ ID NOs: 18-21 operably linked to a promoter (e.g., an in vitro cultured cell or a cell in a transgenic animal) can be employed for screening candidate compounds for an ability to modulate nitrosylation of HIF-1. As used herein, the terms “candidate substance” and “candidate compound” are used interchangeably and refer to a substance that is believed to be capable of modulating nitrosylation of the conserved cysteine present in any of SEQ ID NOs: 18-21. Exemplary candidate compounds that can be investigated using the methods and compositions disclosed herein include, but are not restricted to, agonists and antagonists of enzymes disclosed herein to influence HIF-1 nitrosylation (e.g., small molecule agonists and antagonists of a NOS and/or a PDH), peptides, small molecules, and antibodies and derivatives thereof, and combinations thereof.

[0169] Assays that can be employed for screening a candidate compound for an ability to modulate HIF-1 nitrosylation are known in the art, and include, but are not limited to the “biotin switch” experiment described in Jaffrey & Snyder, 2001, and in EXAMPLE 6.

V. Combination Therapy

[0170] The presently disclosed subject matter can be employed as a part of a combination therapy. As used herein, the phrase “combination therapy” refers to any treatment wherein the methods and compositions disclosed herein are used in combination with another therapy including, but not limited to radiation therapy (radiotherapy), chemotherapy, surgical therapy (e.g., resection), and combinations thereof.

[0171] V.A. Radiation Treatment

[0172] In some embodiments, the methods and compositions disclosed herein are employed in a combination therapy with radiation treatment. For such treatment of a tumor, the tumor is irradiated concurrent with, or subsequent to, administration of a composition as disclosed herein. In some embodiments, the tumor is irradiated daily for 2 weeks to 7 weeks (for a total of 10 treatments to 35 treatments). Alternatively, tumors can be irradiated with brachytherapy utilizing high dose rate or low dose rate brachytherapy internal emitters.

[0173] The duration for administration of a composition as disclosed herein comprises in some embodiments a period of several months coincident with radiotherapy, but in some embodiments can extend to a period of 1 year to 3 years as needed to effect tumor control. A composition as disclosed herein can be administered about one hour before each fraction of radiation. Alternatively, a composition can be administered prior to an initial radiation treatment and then at desired intervals during the course of radiation treatment (e.g., weekly, monthly, or as required). An initial administration of a composition (e.g., a sustained release drug

carrier) can comprise administering the composition to a tumor during placement of a brachytherapy after-loading device.

[0174] Subtherapeutic or therapeutic doses of radiation can be used for treatment of a radiosensitized tumor as disclosed herein. In some embodiments, a subtherapeutic or minimally therapeutic dose (when administered alone) of ionizing radiation is used. For example, the dose of radiation can comprise in some embodiments at least about 2 Gy ionizing radiation, in some embodiments about 2 Gy to about 6 Gy ionizing radiation, and in some embodiments about 2 Gy to about 3 Gy ionizing radiation. When radiosurgery is used, representative doses of radiation include about 10 Gy to about 20 Gy administered as a single dose during radiosurgery or about 7 Gy administered daily for 3 days (about 21 Gy total). When high dose rate brachytherapy is used, a representative radiation dose comprises about 7 Gy daily for 3 days (about 21 Gy total). For low dose rate brachytherapy, radiation doses typically comprise about 12 Gy administered twice over the course of 1 month. ¹²⁵I seeds can be implanted into a tumor can be used to deliver very high doses of about 110 Gy to about 140 Gy in a single administration.

[0175] Radiation can be localized to a tumor using conformal irradiation, brachytherapy, stereotactic irradiation, or intensity modulated radiation therapy (IMRT). The threshold dose for treatment can thereby be exceeded in the target tissue but avoided in surrounding normal tissues. For treatment of a subject having two or more tumors, local irradiation enables differential drug administration and/or radiotherapy at each of the two or more tumors. Alternatively, whole body irradiation can be used, as permitted by the low doses of radiation required following radiosensitization of the tumor.

[0176] Radiation can also comprise administration of internal emitters, for example ¹³¹I for treatment of thyroid cancer, NETASTRON™ and QUADRAGEN® pharmaceutical compositions (Cytogen Corp., Princeton, N.J., United States of America) for treatment of bone metastases, ³²P for treatment of ovarian cancer. Other internal emitters include ¹²⁵I, iridium, and cesium. Internal emitters can be encapsulated for administration or can be loaded into a brachytherapy device.

[0177] Radiotherapy methods suitable for use in the practice of presently disclosed subject matter can be found in Leibel & Phillips, 1998, among other sources.

[0178] V.B. Chemotherapy Treatment

[0179] In some embodiments, the methods and compositions disclosed herein are employed in a combination therapy with chemotherapy. Particular chemotherapeutic agents are generally chosen based upon the type of tumor to be treated, and such selection is within the skill of the ordinary oncologist.

[0180] Chemotherapeutic agents are generally grouped into several categories including, but not limited to DNA-interactive agents, anti-metabolites, tubulin-interactive agents, hormonal agents, and others such as asparaginase or hydroxyurea. Each of the groups of chemotherapeutic agents can be further divided by type of activity or compound. For a detailed discussion of various chemotherapeutic agents

and their methods for administration, see Dorr et al., 1994, herein incorporated by reference in its entirety.

[0181] In order to reduce the mass of the tumor and/or stop the growth of the cancer cells, a chemotherapeutic agent should prevent the cells from replicating and/or should interfere with the cell's ability to maintain itself. Exemplary agents that accomplish this are primarily the DNA-interactive agents such as Cisplatin, and tubulin interactive agents.

[0182] DNA-interactive agents include, for example, alkylating agents (e.g., Cisplatin, Cyclophosphamide, Altre-tamine); DNA strand-breakage agents (e.g., Bleomycin); intercalating topoisomerase II inhibitors (e.g., Dactinomycin and Doxorubicin); non-intercalating topoisomerase II inhibitors (e.g., Etoposide and Teniposide); and the DNA minor groove binder Plicamycin.

[0183] Generally, alkylating agents form covalent chemical adducts with cellular DNA, RNA, and/or protein molecules, and with smaller amino acids, glutathione, and/or similar biomolecules. These alkylating agents typically react with a nucleophilic atom in a cellular constituent, such as an amino, carboxyl, phosphate, or sulfhydryl group in nucleic acids, proteins, amino acids, or glutathione.

[0184] Anti-metabolites interfere with the production of nucleic acids by either of two major mechanisms. Some of the drugs inhibit production of deoxyribonucleoside triphosphates that are the immediate precursors for DNA synthesis, thus inhibiting DNA replication. Some of the compounds are sufficiently like purines or pyrimidines to be able to substitute for them in the anabolic nucleotide pathways. These analogs can then be substituted into the DNA and RNA instead of their normal counterparts.

[0185] Hydroxyurea appears to act primarily through inhibition of the enzyme ribonucleotide reductase.

[0186] Asparagenase is an enzyme which converts asparagine to nonfunctional aspartic acid and thus blocks protein synthesis in the tumor.

[0187] Tubulin interactive agents act by binding to specific sites on tubulin, a protein that polymerizes to form cellular microtubules. Microtubules are critical cell structure units. When the interactive agents bind on the protein, the cell can not form microtubules. Tubulin interactive agents include Vincristine and Vinblastine, both alkaloids and Paclitaxel.

[0188] Adrenal corticosteroids are derived from natural adrenal cortisol or hydrocortisone. They are used because of their anti-inflammatory benefits as well as the ability of some to inhibit mitotic divisions and to halt DNA synthesis. These compounds include Prednisone, Dexamethasone, Methylprednisolone, and Prednisolone.

[0189] The hormonal agents and leutinizing hormones are not usually used to substantially reduce the tumor mass. However, they can be used in conjunction with the chemotherapeutic agents. Hormonal blocking agents are also useful in the treatment of cancers and tumors. They are used in hormonally susceptible tumors and are usually derived from natural sources. These include, but are not limited to estrogens and conjugated estrogens, progestins, and androgens. Leutinizing hormone releasing hormone agents or gonadotropin-releasing hormone antagonists are used primarily the treatment of prostate cancer. These include leuprolide

acetate and goserelin acetate. They prevent the biosynthesis of steroids in the testes. Other anti-hormonal agents include anti-estrogenic agents, anti-androgen agents, and anti-adrenal agents such as Mitotane and Aminoglutethimide.

[0190] Representative chemotherapeutic agents are presented in Table 2.

TABLE 2

<u>Chemotherapeutic Agents</u>	
Agent Type	Examples
<u>Alkylating Agents</u>	
Nitrogen mustards	Chlorambucil, Cyclophosphamide, Isofamide, Mechlorethamine, Melphalan, Uracil mustard
Aziridines	Thiotepa
Methanesulfonate esters	Busulfan
Nitroso ureas	Carmustine, Lomustine, Streptozocin
Platinum complexes	Cisplatin, Carboplatin
Bioreductive alkylators	Mitomycin, Procarbazine
DNA strand breaking agents	Bleomycin
DNA topoisomerase II inhibitors	Amsacrine, Dactinomycin, Daunorubicin, Doxorubicin, Idarubicin, Mitoxantrone, Etoposide, Teniposide
DNA minor groove binder	Plicamycin
<u>Anti-metabolites</u>	
Folate antagonists	Methotrexate and trimetrexate
Pyrimidine antagonists	Fluorouracil, Fluorodeoxyuridine, CB3717, Azacytidine, Cytarabine, Floxuridine
Purine antagonists	Mercaptopurine, 6-Thioguanine, Fludarabine, Pentostatin
Sugar modified analogs	Cytarabine, Fludarabine
Ribonucleotide reductase inhibitors	Hydroxyurea
Tubulin interactive agents	Vincristine, Vinblastine, Paclitaxel
Adrenal corticosteroids	Prednisone, Dexamethasone, Methylprednisolone, Prednisolone
<u>Hormonal blocking agents</u>	
Estrogens and related	Ethinyl Estradiol, Diethylstilbesterol, Chlorotrianisene, Idenestrol
Progestins	Hydroxyprogesterone caproate, Medroxyprogesterone, Megestrol
Androgens	Testosterone, Testosterone propionate, Fluoxymesterone, Methyltestosterone
Leutinizing hormone releasing hormone agents and/or gonadotropin-releasing hormone antagonists	Leuprolide acetate; Goserelin acetate
Anti-estrogenic agents	Tamoxifen
Anti-androgen agents	Flutamide
Anti-adrenal agents	Mitotane, Aminoglutethimide

[0191] A "potentiator" can be any material that improves or increases the efficacy of a pharmaceutical composition and/or acts on the immune system. Exemplary potentiators are triprolidine and its cis-isomer, which can be used in combination with chemotherapeutic agents. Triprolidine is described in U.S. Pat. No. 5,114,951. Other potentiators are procodazole 1H-Benzimidazole-2-propanoic acid; [β -(2-benzimidazole) propionic acid; 2-(2-carboxyethyl)benzimidazole; propazol) Procodazole is a non-specific active immunoprotective agent against viral and bacterial infec-

tions and can be used with the compositions disclosed herein. Potentiators can improve the efficacy of the disclosed compositions and can be used in a safe and effective amount.

[0192] Antioxidant vitamins such as ascorbic acid, beta-carotene, vitamin A, and vitamin E can also be administered with the compositions disclosed herein.

EXAMPLES

[0193] The following Examples have been included to illustrate modes of the presently disclosed subject matter. In light of the present disclosure and the general level of skill in the art, those of skill will appreciate that the following Examples are intended to be exemplary only and that numerous changes, modifications, and alterations can be employed without departing from the scope of the presently disclosed subject matter.

Materials and Methods used in the Examples

[0194] Cell lines and tissue culture. The 4T1 murine mammary adenocarcinoma cell line and B16F10 murine melanoma cells were obtained from the American Type Culture Collection (ATCC, Manassas, Va., United States of America). The two cell lines were cultured in Dulbeccos's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum.

[0195] Reagents. CoCl₂, S-nitrosoglutathione (GSNO), potassium 2-(4-carboxyphenyl)-4,4,5,5-tetramethylimidazole-1-oxyl-3-oxide (carboxy-PTIO), and carrageenan were purchased from Sigma (St. Louis, Mo., United States of America). The proteasome inhibitor MG-132 was purchased from EMD Biosciences, Inc. (San Diego, Calif., United States of America). Luciferin was obtained from Xenogen (Alameda, Calif., United States of America). N ω -nitro-L-arginine methyl ester (L-NAME) and 1400W were purchased from Cayman Chemical (Ann Arbor, Mich., United States of America). N-(6-[biotinamido]hexyl)-3'-(2'-pyridylidithio)-propionamide (biotin-HPDP) was purchased from Pierce Biotechnology (Rockford, Ill., United States of America).

[0196] Plasmids and cloning procedures. The HIF-1 α bioluminescence reporter (ODD-luc) construct was created by fusing PCR product of ODD domain of HIF-1 α (GENBANK® Accession No. U59496) to the 5' end of firefly luciferase reporter gene. Along with this construct, a luciferase expression vector in which luciferase gene was driven by the CMV promoter was used as a control. AC533S ODD mutation was achieved by in vitro site-directed mutagenesis. ODD fragments with or without the mutation were cloned into pCMV-3Tag-4B Epitope Tagging Mammalian Expression vector (Stratagene, La Jolla, Calif., United States of America). The full length mouse VHL gene (GENBANK® Accession No. S76748) was cloned from mouse tissue and tagged with HA tag by PCR. All constructs were sequence-verified.

[0197] siRNA. siRNA sequences targeted to the VHL gene were designed by using an Internet-based program available at the website of Ambion Inc. (Austin, Tex., United States of America). A retroviral siRNA expression vector (pSilencer 5.1-U6 Retro from Ambion Inc.) was used to stably introduce the following siRNA sequence targeted to VHL gene to 4T1 cells: AACATCACATTGCCAGTGTAT (SEQ ID NO: 17). pSilencer 5.1-U6 Scrambled siRNA (Ambion Inc.) was used as a negative control.

[0198] Imaging luciferase activity. Luciferase expression/activity was detected and quantified as relative light units (RLUs) by using the Xenogen IVIS™ imaging system and associated LIVING IMAGE® software (Xenogen, Alameda, Calif., United States of America). For in vitro observations, cells transfected with luciferase reporter gene constructs were grown in 12-well or 24-well cell culture plates. After cells reached 80% confluence, they were treated with the different chemicals or were cultured in the hypoxic chamber (Sheldon Manufacturing, Inc., Cornelius, Oreg., United States of America). At the times indicated, luciferin (150 μ g/ml) was added and the plates were imaged for luciferase expression. For in vivo experiments, treated tumor-bearing mice received an i.p. injection of luciferin (150 mg/kg) during isoflurane anesthesia. Repeated images of luciferase expression/activity were acquired following manufacturer's specified procedures.

[0199] Animal experiments. Female NIH Swiss nude mice, C57BL/6 mice were purchased from the National Cancer Institute (NCI; Frederick, Md., United States of America). Mice with a genetic disruption of the inducible nitric acid synthase gene (C57BL/6 iNOS^{-/-}) were obtained from the Jackson Laboratory (Bar Harbor, Me., United States of America). Female BALB/c mice were obtained from Charles River Laboratories (Raleigh, N.C., United States of America). Animals were maintained and cared for in accordance with the Duke University Institutional Animal Care and Use Committee guidelines. For tumor implantation, 4T1 tumors were grown in nude mice or in syngeneic BALB/c mice, and B16F10 melanoma in syngeneic C57BL/6 or iNOS^{-/-} mice. About 5 $\times$ 10⁵ wild type- or luciferase reporter gene-transfected tumor cells were injected subcutaneously (s.c.) into mice in 50 μ l of PBS solution in the hind legs of ketamine/xylazine-anesthetized mice. When the tumors reached 6-8 mm in diameter, mice were randomly assigned to experimental groups. To image the luciferase activity, single dose of 6 Gy X-ray was given to the tumor on the right leg and this day was set as day 0. The luciferase activity was imaged everyday or every other day for ten days. For the tumor growth delay assay, radiotherapy was employed in its clinical mode: tumors were treated with 3 fractions of 6 Gy every other day from day 0. Tumor growth was then followed by use of a caliper every 2 days. Tumor volume was calculated using the following formula: volume=(length $\times$ width²)/2. Where indicated, mice received the Pan-NOS inhibitor L-NAME (500 mg/L) or the iNOS-selective inhibitor 1400W (50 mg/L) in the drinking water one day before X-ray treatment. Drinking water was renewed daily until animal sacrifice.

[0200] ELISA and western blot analysis. Tumor homogenates and tissue culture samples were both used for protein analysis. All results were normalized for total sample protein contents, determined by using a Bradford-based assay (BIO-RAD, Hercules, Calif., United States of America). Nuclear extracts were prepared by use of the NUCBUSTER™ Protein Extraction Kit (Novagen, San Diego, Calif., United States of America). HIF-1 binding activities in tumor homogenates were quantified with the TRANSAM™ HIF-1 ELISA Kit (Active Motif, Carlsbad, Calif., United States of America) with the use of an antibody against mouse HIF-1 α (Novus Biologicals, Littleton, Colo., United States of America). VEGF levels were assayed using the mouse VEGF quantikine ELISA Kit (R&D systems, Minneapolis, Minn., United States of America). HIF-1 α levels in tissue

culture samples were determined by Western Blot using a polyclonal rabbit anti-HIF-1 antibody (Novus Biologicals) detected with horseradish peroxidase (HRP)-conjugated donkey anti-rabbit secondary antibody (Santa Cruz Biotechnology, Santa Cruz, Calif., United States of America).

[0201] Macrophage depletion experiments. Macrophage depletion was achieved in mice as described in Muller et al., 2005. Briefly, nude mice received repeated i.p. injections of 2 mg carrageenan at 6, 3, and 1 day before s.c. injection of 4T1 tumor cells, after which mice were injected once per week until the end of the experiment.

[0202] Immunohistochemical stainings. Immunofluorescence stainings were performed on tumors biopsied 5 days after irradiation (6 Gy). Cryoslices were fixed in 4% paraformaldehyde. Endogenous peroxidases were quenched with 3% H₂O₂. Slices were then blocked with 10% normal serum, probed with a primary antibody, and revealed with a secondary antibody coupled to FITC (to reveal CD68 and HIF-1 α) or to TRITC (for iNOS detection; Jackson ImmunoResearch Labs, Inc., West Grove, Pa., United States of America). Primary antibodies were: polyclonal rat anti-CD68 to label macrophages (BD PHARMINGEN™, San Jose, Calif., United States of America), polyclonal goat anti-iNOS (Santa Cruz Biotechnology), and polyclonal rabbit anti-HIF-1 α (Novus Biologicals). Cryoslices were examined with a Zeiss Axioskop microscope equipped for fluorescence. Digitized pictures were overlaid by using the METAMORPH™ software from Molecular Devices Corp. (Sunnyvale, Calif., United States of America).

[0203] Vessel staining was performed on tumor cryoslices by using the VECTASTAIN® ABC and the NOVARED™ kits from Vector Laboratories (Burlingame, Calif., United States of America), according to the manufacturer's protocol. The primary anti-CD31 antibody (Santa Cruz Biotechnology, Santa Cruz, Calif.) was labeled with a secondary biotinylated anti-rabbit antibody (Jackson ImmunoResearch Labs, Inc.). Slices were counterstained using Harris' hematoxylin. Vascular density was determined by counting CD31-positive structures in 5 random fields per tumor.

[0204] In vitro interaction assay for ODD and VHL Protein. 4T1 cells grown in 35 mm dishes were transfected with 3 μ g of pCMV-ODD-3Myc, including c-myc-tagged wild type ODD and its mutant version, or 3 μ g of pCMV-HAVHL encoding HA-tagged full-length VHL by using lipofectamine 2000 (Invitrogen, Carlsbad, Calif., United States of America). 24 hours later, cells were subjected to 1 mM GSNO for 8 hours. Cells were then scraped off the dishes and collected. To each cell pellet 300 μ l lysis buffer (50 mM Tris, 150 mM NaCl, 0.5 μ M ferrous chloride, 0.5% NP-40, 0.5 μ M MG-132, protease inhibitor cocktail, pH 7.5) was added. After centrifugation (15,000 $\times$ g for 30 min), supernatants were transferred to fresh tubes and the input ODD and VHL were detected by Western Blot analysis using antibodies against C-myc tag or HA tag (Novus Biologicals). 0.5 mg of supernatant from ODD-c-myc or C533S-ODD-c-myc-expressing cells were mixed with 0.25 mg of the supernatant from HA-VHL expressing cells and incubated at 4° C. for 2 hours. The co-immunoprecipitation was achieved by the addition of 20 μ l of anti-HA antibody (agarose immobilized; Novus Biologicals). Beads were collected, washed three times with 1 ml washing buffer (20 mM Tris, 100 mM NaCl, 1 mM EDTA, 0.5% NP-40), supple-

mented with 50 μ l 2 $\times$ SDS-PAGE sample buffer, and boiled at 95° C. for 10 minutes. Beads were removed by centrifugation, and supernatants were loaded on 12% SDS-gels. The amount of ODD-c-myc or C533S-ODD-c-myc that had been pulled down with HA-VHL was probed with a primary anti-c-myc antibody (Novus Biologicals), and revealed with an anti-goat secondary antibody (Santa Cruz Biotechnology). After membrane stripping (RESTORE™ Western Blot Stripping Buffer, Pierce Biotechnology), immunoprecipitated HA-VHL was labeled with a primary antibody against HA (Novus Biologicals), and revealed with a secondary anti-goat antibody (Santa Cruz Biotechnology).

[0205] Biotin switch assay. Biotin switch assay was performed as described (Jaffrey & Snyder, 2001). Briefly, 4T1 cells were transfected with pCMV-ODD-3Myc. 24 hours later, cells were treated with 1 mM GSNO for 8 h. Cells were then homogenized by 26G needle in HEN buffer (250 mM HEPES-NaOH pH 7.7, 1 mM EDTA, 0.1 mM Neocuproine), and then centrifuged at 1000 $\times$ g for 10 minutes at 4° C. Supernatant (300 μ g) was added to 4 volumes of blocking buffer (9 volumes of HEN buffer plus 1 vol 25% SDS, 20 mM MMTS) at 50° C. for 20 minutes with frequent vortexing. The MMTS was then removed by desalting three times with the BIO-SPIN® 6 column (Bio-Rad, Hercules, Calif., United States of America) pre-equilibrated in HEN buffer. To the eluate was added biotin-HPDP (final concentration of 2 mM) prepared fresh as a 4 mM stock in DMSO from a 50 mM stock suspension in DMF. Sodium ascorbate was added to a final concentration of 1 mM. After incubation for 1 hour at 25° C., biotinylated proteins were precipitated by streptavidin-agarose beads (Pierce Biotechnology, Rockford, Illinois, United States of America). The streptavidin-agarose beads were then pelleted and washed 5 times with HENS buffer. The biotinylated proteins were eluted by SDS-PAGE sample buffer and subjected to Western blot analysis. The biotinylated ODD was detected by use of an antibody against the c-myc tag.

[0206] Statistics. Student's t test, one-way and two-way ANOVA were used where indicated. In growth delay experiments, the numbers of days for tumors to reach 5 $\times$ their initial volume were used for comparing different treatment groups. P<0.05 was considered to be statistically significant.

Example 1

[0207] A Novel Reporter for Non-Invasive, In Vivo Observation of HIF-1 Activity

[0208] HIF-1 activity is difficult to study in vivo because of the very short half-life of the HIF-1 α subunit (Yu et al., 1998). A strategy was adopted in which the oxygen-dependent degradation (ODD) domain of the protein was fused with the firefly luciferase gene (luc; see FIG. 1A). This approach took advantage of the fact that the stability of the HIF-1 α subunit (and hence the activity of HIF-1) is mainly regulated by the ODD. It has been shown that the modification of key proline residues by proline hydroxylases (PHDs) under normoxic conditions render the HIF-1 α susceptible to binding by VHL and subsequent degradation by the proteasome system.

[0209] Thus, it was reasoned that the fusion protein would recapitulate the regulation of HIF-1 α and serve as noninvasive reporter of HIF-1 α activity. When introduced into several tumor cell lines and evaluated under various treat-

ment conditions, the reporter fulfilled expectations. While background luminescence arising from reporter gene expression was very low, it rose significantly after cellular exposure to hypoxia, CoCl_2 (an established inhibitor of HIF-1 α degradation), or MG132 (a proteasome inhibitor), closely mimicking the known regulation of HIF-1 α (see FIG. 1B).

[0210] The successful recapitulation of HIF-1 α stability regulation was further confirmed by transfecting the reporter cells with a VHL-targeted siRNA. The siRNA effectively reduced the level of the VHL expression (see FIG. 1C), which led to significant increases in ODD-luc level, consistent with the role of VHL as the main mediator of HIF-1 α ubiquitylation and degradation. Western blot analysis showed a parallel increase in endogenous HIF-1 α level after various treatment conditions (FIG. 1D), further validating the ODD-luc reporter as a surrogate marker for HIF-1 α .

Discussion of Example 1

[0211] The ODD-luc reporter had a much better dynamic range for the detection of HIF-1 α levels than previous promoter-based approaches. In previous studies dealing with the radiation-induced HIF-1 α activation (Moeller et al., 2004; Moeller et al., 2005), the *in vivo* data were mostly obtained with a GFP reporter containing an artificial hypoxia-responsive promoter (HRP). As GFP is a very stable protein (with half life exceeding 24 hours) and the artificial HRP promoter is subject to HIF-independent biological influences that lead to high background activity, the sensitivity of the HRP-GFP reporter was very limited, with a dynamic range limited to 1-3 fold over background level. In addition, HIF-GFP-based experiments had to be carried out with invasive tumor models such as the dorsal skinfold window chamber tumors (Huang et al., 1999) to obtain quantitative data or to conduct repeated measurements. With the new reporter, any murine tumor system can be monitored with the dynamic range of the reporter increased from 1 to 100 fold over background, and repeated measurements acquired non-invasively by means of *in vivo* optical imaging such as the Xenogen IVISTM imaging system (Contag et al., 1998; Zhang et al., 2004a).

Example 2

Radiation-induced HIF-1 α Stabilization in Tumors

[0212] In order to observe HIF-1 α regulation after treatment response, 4T1 murine breast tumor cells stably transduced with the ODD-luc reporter gene were implanted subcutaneously into mice. After the tumors reach 6-8 mm in diameter, they were irradiated and followed for ODD-luc expression using the Xenogen IVISTM system. From day 3 after irradiation, the level of HIF-1 α , as determined by ODD-luc, appeared to increase linearly over 3 days. It peaked at around 6 days and fall back to background levels after day 10 (see FIG. 2A). The differences between the irradiated and sham-irradiated groups were highly significant from day 3 ($p < 0.01$).

[0213] Radiation-induced stabilization of ODD-luc was accompanied by increases in HIF-1 promoter binding activities to the corresponding HRE binding element (see FIG. 2B) and upregulation of a downstream target gene, vascular endothelial growth factor (VEGF; see FIG. 2C). Similar

results were obtained with two other tumor models, B16.F10 melanoma model and the CT26 colon cancer model. These results indicated that radiation induced a persistently increasing level of HIF-1 α expression and activity. While radiation has been shown to activate HIF-1 α in previous studies (Moeller et al., 2004), the pattern of *in vivo* induction such as the one disclosed herein had never been observed previously.

Example 3

Role of Nitric Oxide in Mediating Radiation-Induced HIF-1 α Activation

[0214] The cause of radiation-induced HIF-1 α stabilization is not understood. One possibility is that radiation creates a more hypoxic condition in the tumor microenvironment than pre-treatment, which causes the stabilization of HIF-1 α through the prolyl hydroxylase (PHD) pathway. However, this is highly unlikely. Previous studies have indicated no significant changes (Brizel et al., 1999; Brizel et al., 1996) in the level of hypoxia in tumor following radiation. Indeed, it had been shown that tumor oxygen tension actually increases after irradiation due to a reduced cell proliferation and tumor cell death. Measurements of 4T1 tumors after irradiation indicated a similar scenario (Moeller et al., 2004).

[0215] The co-inventors' previous studies had indicated that radiation induced free radicals are at least partially involved in the activation of so-called "stress granules" (Moeller et al., 2004). However, the identity of the free radicals involved in that response is not clear.

[0216] After evaluation with various agents, nitric oxide (NO) was determined to be the main free radical species that was responsible for radiation-induced HIF-1 α activation (see FIG. 3A). The administration of L-NAME, a potent non-specific inhibitor of nitric oxide synthases (NOS), to mice effectively attenuated radiation-induced HIF-1 α stabilization in tumors, as shown by the loss of ODD-luc signal. As NOS are the major source of NO *in vivo*, the presently disclosed results indicated that NO played a pivotal role in radiation-induced HIF-1 α stabilization. Control experiments indicated that NO produced by NOS did not influence the activity of constitutively expressed luciferase activity, confirming the role of NO in regulating ODD (and hence HIF-1 α) stability (see FIG. 3A).

[0217] The role of NO was further confirmed in cell culture assays. Treatment of 4T1-ODD-luc cells with the NO donor S-nitrosoglutathione (GSNO) effectively induced dose-dependent HIF-1 α activation, similar to treatment with ionizing radiation (see FIG. 3B). Western blot analysis of the GSNO treated cells clearly indicated endogenous HIF-1 α induction (FIG. 3C). A NO scavenger, carboxy-PTIO(4-carboxyphenyl-4,4,5,5-tetramethylimidazoline-3-oxide-1-oxyl), effectively suppressed HIF-1 α activation by GSNO (FIG. 3D), demonstrating that NO is directly responsible for the observed ODD-luc accumulation.

Example 4

Inducible Nitric Oxide Synthase as a Major Source of NO in Radiation-Induced HIF-1 α Activation

[0218] The source of the NO that stimulates HIF in irradiated tumors *in vivo* was investigated. Of the three NOS

isoforms, inducible NO synthase (iNOS), is the most likely candidate because, unlike neuronal and endothelial NOS, which are constitutively activated in healthy tissues, it is exclusively expressed and activated in pathological tissues such as tumors, where it can produce high micromolar levels of NO. Moreover, tumors usually contain significant number of macrophages (Colombo & Mantovani, 2005; Lewis & Murdoch, 2005), which express/activate their iNOS as part of their immunoeffector activity and thus provide a ready source of NO upon activation.

[0219] To pinpoint the source of NO, two series of experiments were performed. In the first series of experiments, the iNOS-specific inhibitor, 1400W (Alderton et al., 2001; Thomsen et al., 1997), was used to examine radiation-induced HIF-1 α induction in ODD-luc-transduced 4T1 tumors. The results indicated that 1400W attenuated radiation-induced ODD-luc in 4T1 as potently as the general NOS inhibitor L-NAME (see FIG. 4A). This observation indicated that iNOS is the main mediator of radiation-induced HIF-1 α stabilization.

[0220] In the second series of experiments, C57BL/6 mice with targeted disruption of the iNOS gene (iNOS^{-/-}) were implanted with syngeneic B16F10 melanoma cells stably transduced with ODD-luc gene. The tumors were then irradiated and observed for HIF-1 α activation. A significant attenuation of radiation-induced ODD-luc induction in the tumors grown in iNOS^{-/-} mice compared to wild-type controls was observed. In fact, ODD-luc suppression in iNOS^{-/-} animals and in wild type mice treated with L-NAME were of similar amplitude (see FIG. 4B), indicating that L-NAME-suppressed HIF-1 α activation in the wild type mice was attributable to the inhibition of iNOS.

Example 5

Macrophages are a Major Source of iNOS and NO in Radiation-Induced HIF-1 α Activation

[0221] Previous studies have indicated that macrophages are a rich source of NO and that the tumor microenvironment is abundantly populated with macrophages. In light of this information and together with the results presented hereinabove, it was hypothesized that tumor-associated macrophages might play a significant role in radiation-induced HIF-1 α induction. This would also be consistent with previous findings that tumor-associated macrophages play important roles in regulating tumor angiogenesis, at least partially through NO release (Leek et al., 2000; Leek et al., 2002; Varney et al., 2002).

[0222] In order to investigate the potential involvement of macrophages in HIF-1 α activation, radiation-induced ODD-luc activation in mice that had been chemically depleted of macrophages through the use of carrageenan was measured (Goldmann et al., 2004; Muller et al., 2005; Udonio et al., 1994). The results were very similar to those obtained in iNOS^{-/-} mice (see FIG. 4). A significant reduction in radiation-induced ODD-luc activation and the loss of L-NAME inhibition of the activation in tumors in mice with macrophage depletion was observed (FIG. 5A).

[0223] These results clearly established that iNOS in tumor-associated macrophages was the main source for the NO that was involved in radiation-induced HIF-1 α activation. Immunohistochemistry analysis further confirmed that

irradiation of the tumor increased the number of tumor-associated macrophages and activated the iNOS gene in these macrophages (see FIG. 5B, left panel). The results presented herein also confirmed that activated iNOS gene expression was accompanied by concomitant HIF-1 α activation in tumors (FIG. 5B, right panel).

Example 6

The Molecular Mechanism of HIF-1 α Activation by NO

[0224] The aforementioned experiments provided strong evidence that NO generated by tumor-associated macrophages played critical roles in radiation-induced HIF-1 α activation. However, the exact molecular mechanism of how NO induces stabilization of HIF-1 α remained unclear.

[0225] In theory, there are at least two ways NO can influence HIF-1 α : the inactivation of upstream prolyl hydroxylases and/or the direct modification of the ODD domain. Others have shown that NO can inhibit the activity of the prolyl hydroxylases (PHDs), which can result in the stabilization of HIF-1 α (Metzen et al., 2003). However, inhibition of PHDs did not appear to account for all the NO-induced HIF-1 α activation. It was therefore reasoned that a direct modification of the ODD domain could also participate in HIF-1 activation.

[0226] A previous report suggested that, although all 13 cysteine residues in the purified HIF-1 α protein are susceptible to nitrosylation in test tubes, only 3-4 can be nitrosylated in cells in cultured cells (Sumbayev et al., 2003). However, the biological significance of these nitrosylations on HIF-1 α stability has not been identified.

[0227] Thus, NO might stabilize the HIF-1 α through S-nitrosylation of ODD domain during radiotherapy. To test this hypothesis, a mutant (C533S) involving the only Cys residue in the murine HIF-1 α ODD domain, Cys533, was generated. This residue corresponds to Cys520 (which also is the only Cys the human ODD domain) in human HIF-1 α and is conserved among a wide spectrum of vertebrate species that included human, mouse, rat, frog, etc. (see FIG. 6A). The replacement of the cysteine by a serine was chosen because the only difference between these amino acids is that the thiol (—SH) group of cysteine is replaced by the hydroxyl (—OH) group of serine, thereby preventing S-nitrosylation. The likelihood that the point mutation will alter the 3-D structure of ODD is thus minimal. The C533S ODD domain was then fused with the luciferase reporter gene, transfected into 4T1 tumor cells, and examined for its activation in comparison with wild type ODD-luc in vitro and in vivo.

[0228] In vitro, the background expression level of mutant C533S-ODD-luc was very low, similar to wild type (see FIG. 6B). However, when C533S-ODD-luc was subjected to hypoxia, proteasome inhibition, or CoCl₂ exposure, significant inductions, similar to wild-type, were observed (see FIG. 6B), indicating that the mutation did not cause any gross structural perturbation that would disrupt normal processing by upstream PHDs and downstream VHL and proteasome.

[0229] However, when the mutant C533S-ODD-luc transduced cells were exposed to the NO donor GSNO, induced

ODD-luc expression was almost absent, in sharp contrast to wild type ODD-luc transduced cells, which had significant GSNO induction (see FIG. 6C).

[0230] In vivo, background levels of mutant C533S-ODD-luc transduced tumors were similar to what was observed in wild type ODD-luc transduced tumors (see FIG. 6D). However, the C533S mutation significantly attenuated radiation-induced ODD-luc activation in vivo (see FIG. 6D, days 5, 7, and 10), indicating that S-nitrosylation of the Cys533 residue in the HIF-1 α protein played a critical role in regulating the stabilization of HIF-1 α after radiation therapy.

[0231] The direct proof for S-nitrosylation of HIF-1 α at C533 came from "biotin switch" experiments (Jaffrey & Snyder, 2001) in which direct chemical evidence for the nitrosylation of the ODD domain was sought. In the absence of GSNO treatment, neither wild type ODD nor C533S-ODD was S-nitrosylated (see FIG. 6E). S-nitrosylation was clearly observed in wild type ODD upon GSNO treatment, but completely absent in C533S-ODD after GSNO treatment (see FIG. 6E), demonstrating that C533 was S-nitrosylated in the cellular environment with a sufficient amount of NO.

[0232] The site-directed mutagenesis experiments disclosed herein further suggested that nitrosylation at Cys533 rendered the HIF-1 α protein resistant to degradation by preventing the binding of HIF-1 α by VHL. To examine this possibility, the effects of NO and of C533S on the binding of ODD with VHL in ODD-transfected tumor cells were tested. Co-immunoprecipitation results revealed that the strong binding of wild-type ODD with VHL in the absence of NO was completely abolished in cells exposed to GSNO (see FIG. 6F). Strikingly, this regulation was completely lost in cells expressing C533S-ODD, which is consistent with the continuous degradation of the mutated ODD in the presence of NO (FIG. 6C and 6D). Taken together, the in vivo and in vitro results. (see FIG. 6A-F) provided compelling evidence that NO-mediated stabilization of HIF-1 α is largely mediated by S-nitrosylation of the Cys533 in the ODD domain.

Example 7

The Functional Importance of NO-Mediated HIF-1 α Activation During Cancer Therapy

[0233] As HIF-1 α has been shown to be a key tumor survival factor during cancer therapy, it was postulated that the inhibition of HIF-1 α activation through the prevention of NO production would have anti-tumor efficacy. To examine this hypothesis, tumor growth delay experiments with co-administration of radiotherapy (3{6 Gy) and L-NAME were performed in two aggressive tumor models: 4T1 (murine mammary adenocarcinoma; see FIG. 7A) and B16F10 (murine melanoma; see FIG. 7B). In both models, the inhibition of NO production by L-NAME significantly enhanced the therapeutic efficacy of radiotherapy. In addition, the use of L-NAME in conjunction with radiotherapy significantly reduced tumor vasculature (see FIG. 7C).

[0234] These results suggested that NO-mediated HIF-1 α activation indeed played a critical role in overall tumor response to radiotherapy, consistent with previous reports that the survival of tumor vasculature is key to tumor survival during radiotherapy (Garcia-Barros et al., 2003; Moeller et al., 2004). They further suggested that NOS

inhibitors can be used as therapeutic agents to enhance the efficacy of conventional cancer treatments.

Discussion of Examples 1-7

[0235] Understanding HIF-1 regulation during cancer treatment can provide insights into how tumor responds to therapy. This is because HIF-1 has been shown to be a key tumor survival factor after cancer therapy (Moeller et al., 2004; Zhang et al., 2004b). The presently disclosed discovery of HIF-1 α upregulation through NO generated from tumor-associated macrophages is important for at least two reasons: recognizing the tumor-associated macrophages (TAMs) as a major regulator of HIF-1 and the identification of S-nitrosylation of C533 (human equivalent C520) as a key mechanism for NO-mediated HIF-1 α stabilization. The present disclosure establishes for the first time that TAM is a pivotal mediator of tumor angiogenic activity after radiotherapy while the latter unveils a novel mechanism for HIF-1 α regulation.

[0236] Previous studies have suggested that NO effects on HIF-1 α to be different under hypoxic or normoxic conditions. Under hypoxic conditions, it was shown that the presence of NO can inhibit HIF-1 activity (Sogawa et al., 1998) by inducing the redistribution of intracellular oxygen (Hagen et al., 2003) that increased PHD activity and HIF-1 α degradation.

[0237] Under normoxic conditions, however, NO has been shown to increase the stability and activity of HIF-1 α in at least two different ways. First, NO can directly inhibit the activity of PHDs (Metzen et al., 2003) and thereby inhibiting proteasome-mediated degradation of HIF-1 α . Second, NO can enhance the transcriptional activity of HIF-1 through the nitrosylation of Cys800 (Yasinska & Sumbayev, 2003), which enhances the binding of the HIF-1 α C-terminal transactivation domain (C-TAD) to p300.

[0238] The presently disclosed discovery that NO can regulate HIF-1 α stability through S nitrosylation of Cys533 provides a third avenue for NO-mediated increase in HIF-1 transcriptional activity. It also provides a remarkable example where targeted S-nitrosylation of a single cysteine residue in a protein can significantly influence its interaction with other protein(s), very similar to a recent report (Kim et al., 2005) on nitric oxide regulation of the COX-2 gene activation.

[0239] Of special interest is the fact that HIF-1 has also been known to enhance iNOS gene expression in a variety of cell types (Jung et al., 2000; Matrone et al., 2004; Melillo et al., 1997). Therefore, it is possible that activated iNOS and HIF-1 forms an amplification loop during wound healing or inflammation. Inconsistent with this hypothesis is a recent report that indicate HIF-1 and iNOS do appear to regulate each other positively under normoxic conditions during bacterial infections (Peyssonnaud et al., 2005). This amplification loop might be a key mechanism during inflammatory response. If true, the relationship between NO and HIF-1 α might afford new opportunities of drug development for various inflammatory diseases.

[0240] In terms of cancer therapy, the recognition that NO mediated S-nitrosylation of Cys 533 can upregulate HIF-1 activity during radiation or chemotherapy has important implications as well. This is because quite a few studies have

indicated that HIF-1 plays, critical roles for tumor growth and survival during cancer therapy (Moeller et al., 2004; Yeo et al., 2003). The recognition of the role of NO in the up-regulation of HIF-1 α during cancer therapy suggests a promising strategy to enhance current therapy: the use of NOS inhibitors in conjunction with conventional radiation and chemotherapy modalities. The results presented herein combining NOS inhibitor L-NAME and radiotherapy (see FIG. 7) support for this notion. A similar experiment with 4T1 tumors treated with cyclophosphamide (see FIG. 9) suggests that NOS inhibitors can also augment chemotherapy.

[0241] Although the presented studies were primarily conducted in tumors that were exposed to ionizing radiation, the same NO-mediated HIF-1 activation pathway might operate in other normal cells/tissues. Indeed, the instant co-inventors have observed NO-mediated HIF-1 α activation in macrophages, fibroblasts, and epithelial cells, indicating the general applicability of this pathway.

[0242] In summary, the results presented herein establish the importance of nitric oxide-mediated S-nitrosylation in regulating the stability of HIF-1 α . They indicate that S-nitrosylation of Cys533 (murine equivalent of human Cys520) in HIF-1 α is directly responsible for radiation-induced HIF-1 α stabilization in tumors. The instant disclosure also indicates that modulating HIF-1 α activation through NOS inhibitors is a promising strategy for therapeutic development in a variety of diseases such as cancer and inflammatory diseases where it has been established that both NO and HIF-1 α play prominent roles.

REFERENCES

- [0243] The references listed below, as well as all references cited in the specification, including patents, patent applications, journal articles, and all database entries (e.g., GENBANK® database entries, including any annotations presented in the databases associated with the disclosed sequences), are incorporated herein by reference to the extent that they supplement, explain, provide a background for, or teach methodology, techniques, and/or compositions employed herein.
- [0244] Adelman et al. (1983) *DNA* 2:183-193.
- [0245] Alam & Cook (1990) *Anal Biochem* 188:245-254.
- [0246] Alderton et al. (2001). *Biochem J* 357:593-615.
- [0247] Altschul et al. (1990) *J Mol Biol* 215:403-410.
- [0248] Ausubel et al. (1992) *Current Protocols in Molecular Biology*. Wiley, N.Y., N.Y., United States of America.
- [0249] Ausubel (1995) *Short Protocols in Molecular Biology*, 3rd ed. Wiley, N.Y., N.Y., United States of America.
- [0250] Bass (2001) *Nature* 411:428-429.
- [0251] Brizel et al. (1999). *Radiother Oncol* 53:113-117.
- [0252] Brizel et al. (1996). *Cancer Res* 56:5347-5350.
- [0253] Chan et al. (2002). *J Biol Chem* 277:40112-40117.
- [0254] Chen et al. (2003). *J Biol Chem* 278:13595-13598.
- [0255] Cipolla et al. (2000) *Hum Gene Ther* 11:361-371.
- [0256] Colombo & Mantovani (2005). *Cancer Res* 65:9113-9116.
- [0257] Contag et al. (1998). *Nat Med* 4:245-247.
- [0258] Cubitt et al. (1995) *Trends Biochem Sci* 20:448-455.
- [0259] Cramer et al. (2003). *Cell* 112:645-657.
- [0260] Dorr et al. (1997) *Cancer Chemotherapy Handbook*, 2d edition, Appleton & Lange, Stamford, Conn., United States of America.
- [0261] Elbashir et al. (2001) *Nature* 411:494-498,
- [0262] Fukuda et al. (2002). *J Biol Chem* 277:38205-38211.
- [0263] Fukuda et al. (2003). *Cancer Res* 63:2330-2334.
- [0264] Garcia-Barros et al. (2003). *Science* 300:1155-1159.
- [0265] Giaccia et al. (2003). *Nat Rev Drug Discov* 2:803-811.
- [0266] GENBANK® Accession Nos. AAA62405; AAH69465; AAP43517; AAU14021; AAX89137; AAY27087; ABB17537; BAE01417; BC069465; CAB96628; CAG29396; CAH93355; NM_000625; NP_000616; NP_001521; NP_034561; NP_13_077335; NP_776764; NP_956527; NP_989628; S76748; U17327; U59496; and XP_852278.
- [0267] Glover & Hames (1995) *DNA Cloning: A Practical Approach*, 2nd ed. IRL Press at Oxford University Press, New York, N.Y., United States of America.
- [0268] Goldmann et al. (2004). *Infect Immun* 72:2956-2963.
- [0269] Greenberg et al. (1994) *Mol Endocrinol* 8:230-239.
- [0270] Habib et al. (1999) *Hum Gene Ther* 10:2019-2034.
- [0271] Hagen et al. (2003). *Science* 302:1975-1978.
- [0272] Harris (2002) *Nat Rev Cancer* 2:38-47.
- [0273] Henikoff & Henikoff (1992) *Proc Natl Acad Sci USA* 89:10915-10919.
- [0274] Huang et al. (1999). *Nat Biotechnol* 17:1033-1035.
- [0275] Isaacs et al. (2005). *Cancer Cell* 8:143-153.
- [0276] Ivan et al. (2001). *Science* 292:464-468.
- [0277] Jaakkola et al. (2001). *Science* 292:468-472.
- [0278] Jaffrey & Snyder (2001). *Sci STKE* 2001:PL1.
- [0279] Jeong et al. (2002). *Cell* 111 :709-720.
- [0280] Jiang et al. (1997). *Cancer Res* 57:5328-5335.
- [0281] Jung et al. (2000). *Circ Res* 86:319-325.
- [0282] Karlin & Altschul (1993) *Proc Natl Acad Sci USA* 90:5873-5877.
- [0283] Kim et al. (2005). *Science* 310:1966-1970.
- [0284] Kurihara et al. (2000) *J Clin Invest* 106:763-771.
- [0285] Lando et al. (2002a). *Genes Dev* 16:1466-1471.
- [0286] Lando et al. (2002b). *Science* 295:858-861.

- [0287] Laughner et al. (2001) *Mol Cell Biol* 21:3995-4004.
- [0288] Lee et al. (2000) *Anticancer Res* 20:417-422.
- [0289] Leek et al. (2000). *J Pathol* 190:430-436.
- [0290] Leek et al. (2002). *Cancer Res* 62:1326-1329.
- [0291] Leibel & Phillips (1998) *Textbook of Radiation Oncology*, Saunders, Philadelphia, United States of America.
- [0292] Lewis & Murdoch (2005). *Am J Pathol* 167:627-635.
- [0293] Maltepe et al. (1997). *Nature* 386:403-407.
- [0294] Matrone et al. (2004). *J Neurochem* 90:368-378.
- [0295] Maxwell et al. (2001). *Adv Exp Med Biol* 502:365-376.
- [0296] Maxwell et al. (1999). *Nature* 399:271-275.
- [0297] Melillo (2004). *Cell Cycle* 3:154-155.
- [0298] Melillo et al. (1997). *J Biol Chem* 272:12236-12243.
- [0299] Metzen et al. (2003). *Mol Biol Cell* 14:3470-3481.
- [0300] Moeller et al. (2004). *Cancer Cell* 5:429-441.
- [0301] Moeller et al. (2005). *Cancer Cell* 8:99-110.
- [0302] Muller et al. (2005). *Nat Med* 11:312-319.
- [0303] Needleman & Wunsch (1970) *J Mol Biol* 48:443-453.
- [0304] Ohh et al. (2000). *Nat Cell Biol* 2:423-427.
- [0305] Pause et al. (1999). *Proc Natl Acad Sci USA* 96:9533-9538.
- [0306] PCT International Publication Nos. WO 97/47763; WO 99/07409; WO 99/32619; WO 00/01846; WO 00/44895; WO 00/44914; WO 01/36646; and WO 01/29058.
- [0307] Pearson & Lipman (1988) *Proc Natl Acad Sci USA* 85:2444-2448.
- [0308] Peyssonnaud et al. (2005). *J Clin Invest* 115:1806-1815.
- [0309] Rapisarda et al. (2002). *Cancer Res* 62:4316-4324.
- [0310] Ravi et al. (2000). *Genes Dev* 14:34-44.
- [0311] Rose & Botstein (1983) *Meth Enzymol* 101:167-180.
- [0312] Sanchez-Puig et al. (2005). *Mol Cell* 17:11-21.
- [0313] Sambrook & Russell (2001) *Molecular Cloning: A Laboratory Manual*, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., United States of America.
- [0314] Scharfmann et al. (1991) *Proc Natl Acad Sci USA* 88:4626-4630.
- [0315] Selak et al. (2005). *Cancer Cell* 7:77-85.
- [0316] Semenza (2002). *Trends Mol Med* 8:S62-67.
- [0317] Semenza (2003) *Nat Rev Cancer* 3:721-32.
- [0318] Semenza et al. (2000). *Adv Exp Med Biol* 475:123-130.
- [0319] Silhavy et al. (1984) *Experiments with Gene Fusions*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., United States of America.
- [0320] Smith & Waterman (1981) *Adv Appl Math* 2:482-489.
- [0321] Sogawa et al. (1998). *Proc Natl Acad Sci USA* 95:7368-7373.
- [0322] Sumbayev et al. (2003). *FEBS Lett* 535:106-112.
- [0323] Sutphin et al. (2004). *Cell Cycle* 3:160-163.
- [0324] Thomsen et al. (1997). *Cancer Res* 57:3300-3304.
- [0325] Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology-Hybridization with Nucleic Acid Probes*. Elsevier, N.Y., United States of America.
- [0326] Udono et al. (1994). *Proc Natl Acad Sci USA* 91:3077-3081.
- [0327] U.S. Pat. Nos. 5,114,951; 5,410,016; 5,411,554; 5,468,253; 5,573,934; 5,599,852; 5,631,015; 5,653,992; 5,688,900; 5,713,920; 5,728,752; 5,824,333; 5,858,746; 5,858,784; 6,013,638; 6,022,737; 6,136,295; 7,009,034; 7,011,842; and 7,012,126.
- [0328] Varney et al. (2002). *In Vivo* 16:471-477.
- [0329] Wachsberger et al. (2003). *Clin Cancer Res* 9:1957-1971.
- [0330] Wang & Semenza (1993a). *J Biol Chem* 268:21513-21518.
- [0331] Wang & Semenza (1993b). *Proc Natl Acad Sci USA* 90:4304-4308.
- [0332] Wang & Semenza (1995). *J Biol Chem* 270:1230-1237.
- [0333] Williams et al. (1993) *J Clin Invest* 92:503-508.
- [0334] Yasinska & Sumbayev (2003). *FEBS Lett* 549:105-109.
- [0335] Yeo et al. (2003). *J Natl Cancer Inst* 95:516-525.
- [0336] Yu et al. (1998). *Am J Physiol* 275:L818-826.
- [0337] Yu et al. (1999) *Cancer Res* 59:4200-4203.
- [0338] Zhang et al. (2004a). *Blood* 103:617-626.
- [0339] Zhang et al. (2004b). *Cancer Res* 64:8139-8142.
- [0340] Zhong et al. (2000). *Cancer Res* 60:1541-1545.
- [0341] Zundel et al. (2000). *Genes Dev* 14:391-396.
- [0342] It will be understood that various details of the presently disclosed subject matter can be changed without departing from the scope of the presently disclosed subject matter. Furthermore, the foregoing description is for the purpose of illustration only, and not for the purpose of limitation.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 21

<210> SEQ ID NO 1

<211> LENGTH: 824

<212> TYPE: PRT

<213> ORGANISM: Spalax judaei

<400> SEQUENCE: 1

Met Glu Gly Ala Gly Gly Glu Asn Glu Lys Lys Lys Met Ser Ser Glu
1 5 10 15
Arg Arg Lys Glu Lys Ser Arg Asp Ala Ala Arg Ser Arg Arg Ser Lys
20 25 30
Glu Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro His
35 40 45
Asn Val Ser Ser His Leu Asp Lys Ala Ser Val Met Arg Leu Thr Ile
50 55 60
Ser Tyr Leu Arg Val Arg Lys Leu Leu Asp Ala Gly Asp Leu Asp Ile
65 70 75 80
Glu Asp Glu Met Lys Ala Gln Met Asn Cys Phe Tyr Leu Lys Ala Leu
85 90 95
Asp Gly Phe Val Met Val Leu Thr Asp Asp Gly Asp Met Ile Tyr Ile
100 105 110
Ser Asp Asn Val Asn Lys Tyr Met Gly Leu Thr Gln Phe Glu Leu Thr
115 120 125
Gly His Ser Val Phe Asp Phe Thr His Pro Cys Asp His Glu Glu Met
130 135 140
Arg Glu Met Leu Thr His Arg Asn Gly Pro Val Lys Glu Gly Lys Glu
145 150 155 160
Gln Asn Thr Gln Arg Ser Phe Phe Leu Arg Met Lys Cys Thr Leu Thr
165 170 175
Ser Arg Gly Arg Thr Met Asn Ile Lys Ser Ala Thr Trp Lys Val Leu
180 185 190
His Cys Thr Gly Arg Ile His Val Tyr Asp Thr Asn Ser Asn Gln Pro
195 200 205
Gln Cys Gly Tyr Lys Lys Pro Pro Met Thr Cys Leu Val Leu Ile Cys
210 215 220
Glu Pro Ile Pro His Pro Ser Asn Ile Glu Ile Pro Leu Asp Ser Lys
225 230 235 240
Thr Phe Leu Ser Arg His Ser Leu Asp Met Lys Phe Ser Tyr Cys Asp
245 250 255
Glu Arg Ile Thr Glu Leu Met Gly Tyr Glu Pro Glu Glu Leu Leu Gly
260 265 270
Arg Ser Ile Tyr Glu Tyr Tyr His Ala Leu Asp Ser Asp His Leu Thr
275 280 285
Lys Thr His His Asp Met Phe Thr Lys Gly Gln Val Thr Thr Gly Gln
290 295 300
Tyr Arg Met Leu Ala Lys Arg Gly Gly Tyr Val Trp Val Glu Thr Gln
305 310 315 320
Ala Thr Val Ile Tyr Asn Thr Lys Asn Ser Gln Pro Gln Cys Ile Val
325 330 335
Cys Val Asn Tyr Val Val Ser Gly Ile Ile Gln His Asp Leu Ile Phe

-continued

340							345					350				
Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	Asp	
355							360					365				
Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr	Ser	
370							375					380				
Cys	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Met	Leu	
385							390					395				
Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	Asp	
							410					415				
Asp	Thr	Glu	Thr	Glu	Asp	Gln	Gln	Leu	Glu	Asp	Val	Pro	Leu	Tyr	Asn	
							425					430				
Asp	Val	Met	Phe	Pro	Ser	Ser	Asp	Asp	Lys	Leu	Thr	Ser	Ile	Asn	Leu	
							440					445				
Ala	Met	Ser	Pro	Leu	Pro	Ala	Pro	Glu	Thr	Pro	Lys	Pro	Leu	Arg	Ser	
							455					460				
Asn	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Pro	
465							470					475				
Asn	Ala	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp	
							485					490				
Gln	Pro	Ala	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu	
							500					505				
Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Asp	Met	Val	
							515					520				
Asn	Val	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr	
							530					535				
Glu	Ala	Lys	Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu	Glu	
545							550					555				
Met	Leu	Ala	Pro	Tyr	Ile	Pro	Met	Asp	Asp	Asp	Phe	Gln	Leu	Arg	Leu	
							565					570				
Phe	Asp	Gln	Leu	Ser	Pro	Leu	Glu	Ser	Ser	Ser	Pro	Ser	Pro	Pro	Gly	
							580					585				
Val	Asn	Thr	Ala	Thr	Ala	Phe	Gln	Gln	Thr	Gln	Leu	Gln	Glu	Pro	Thr	
							595					600				
Ile	Ser	Thr	Thr	Thr	Thr	Thr	Thr	Thr	Thr	Asn	Glu	Leu	Lys	Thr	Val	
610							615					620				
Thr	Lys	Asp	Asn	Ile	Glu	Asp	Ile	Lys	Ile	Leu	Ile	Ala	Ser	Pro	Ser	
625							630					635				
Ser	Thr	His	Thr	Pro	Lys	Glu	Thr	Thr	Thr	Ala	Thr	Thr	Ser	Ser	Pro	
							645					650				
Tyr	Ser	Gly	Thr	Gln	Ser	Arg	Thr	Ala	Ser	Pro	Asn	Arg	Ala	Gly	Gln	
							660					665				
Gly	Val	Ile	Glu	Gln	Thr	Glu	Lys	Ser	His	Pro	Arg	Ser	Pro	Asn	Leu	
							675					680				
Leu	Ser	Val	Ser	Leu	Ser	Gln	Arg	Asn	Thr	Val	Pro	Glu	Glu	Glu	Leu	
690							695					700				
Asn	Pro	Lys	Ile	Ile	Ala	Leu	Gln	Asn	Ala	Gln	Arg	Lys	Arg	Lys	Met	
705							710					715				
Glu	His	Asp	Gly	Ser	Leu	Phe	Gln	Ala	Ala	Gly	Ile	Gly	Thr	Leu	Leu	
							725					730				
Gln	Gln	Pro	Asp	Asp	Arg	Ala	Pro	Ala	Thr	Ser	Leu	Ser	Trp	Lys	Arg	
							740					745				

-continued

Val Lys Gly Cys Lys Ser Ser Gly Gln Asn Gly Met Glu Gln Lys Thr
755 760 765

Ile Ile Leu Ile Pro Ser Asp Leu Ala Cys Arg Leu Leu Gly Gln Ser
770 775 780

Met Asp Gly Ser Gly Leu Pro Gln Leu Thr Ser Tyr Asp Cys Glu Val
785 790 795 800

Asn Ala Pro Ile Gln Gly Ser Arg Asn Leu Leu Gln Gly Glu Glu Leu
805 810 815

Leu Arg Ala Leu Asp Gln Val Asn
820

<210> SEQ ID NO 2
<211> LENGTH: 819
<212> TYPE: PRT
<213> ORGANISM: Eospalax baileyi

<400> SEQUENCE: 2

Met Glu Gly Ala Ala Gly Gly Glu Glu Lys Lys Asn Arg Met Ser Ser
1 5 10 15

Glu Arg Arg Lys Glu Lys Ser Arg Asp Ala Ala Arg Ser Arg Arg Ser
20 25 30

Lys Glu Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro
35 40 45

His Asn Val Ser Ser His Leu Asp Lys Ala Ser Val Met Arg Leu Thr
50 55 60

Ile Ser Tyr Leu Arg Val Arg Lys Leu Leu Asp Ala Gly Asp Leu Asp
65 70 75 80

Ile Glu Asp Asp Met Lys Ala Gln Met Asn Cys Phe Tyr Leu Lys Ala
85 90 95

Leu Asp Gly Phe Val Met Val Leu Thr Asp Asp Gly Asp Met Ile Tyr
100 105 110

Ile Ser Asp Asn Val Asn Lys Tyr Met Gly Leu Thr Gln Phe Glu Leu
115 120 125

Thr Gly His Ser Val Phe Asp Phe Thr His Pro Cys Asp His Glu Glu
130 135 140

Met Arg Glu Met Leu Thr His Arg Asn Gly Pro Ile Lys Lys Gly Lys
145 150 155 160

Glu Gln Asn Thr Gln Arg Ser Phe Phe Leu Arg Met Lys Cys Thr Leu
165 170 175

Thr Ser Arg Gly Arg Thr Met Asn Ile Lys Ser Ala Thr Trp Lys Val
180 185 190

Leu His Cys Thr Gly His Ile His Val Tyr Asp Thr Asn Ser Asn Gln
195 200 205

Pro Gln Cys Gly Tyr Lys Lys Pro Pro Met Thr Cys Leu Val Leu Ile
210 215 220

Cys Glu Pro Ile Pro His Pro Ser Asn Ile Glu Ile Pro Leu Asp Ser
225 230 235 240

Lys Thr Phe Leu Ser Arg His Ser Leu Asp Met Lys Phe Ser Tyr Cys
245 250 255

Asp Glu Arg Ile Thr Glu Leu Met Gly Tyr Glu Pro Glu Glu Leu Leu
260 265 270

Gly Arg Ser Ile Tyr Glu Tyr Tyr His Ala Leu Asp Ser Asp His Leu

-continued

275					280					285					
Thr	Lys	Thr	His	His	Asp	Met	Phe	Thr	Lys	Gly	Gln	Val	Thr	Thr	Gly
290						295					300				
Gln	Tyr	Arg	Met	Leu	Ala	Lys	Arg	Gly	Gly	Tyr	Val	Trp	Val	Glu	Thr
305					310					315				320	
Gln	Ala	Thr	Val	Ile	Tyr	Asn	Thr	Lys	Asn	Ser	Gln	Pro	Gln	Cys	Ile
				325					330					335	
Val	Cys	Val	Asn	Tyr	Val	Val	Ser	Gly	Ile	Ile	Gln	His	Asp	Leu	Ile
				340				345					350		
Phe	Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser
		355					360					365			
Asp	Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr
	370					375					380				
Ser	Cys	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu
385					390					395					400
Leu	Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser
				405					410					415	
Asp	Asp	Thr	Glu	Thr	Glu	Asp	Gln	Gln	Leu	Glu	Asp	Val	Pro	Leu	Tyr
			420					425					430		
Asn	Asp	Val	Met	Phe	Pro	Ser	Ser	Asp	Asp	Lys	Leu	Thr	Ser	Ile	Asn
		435					440					445			
Leu	Ala	Met	Ser	Pro	Leu	Pro	Ala	Ser	Glu	Thr	Pro	Lys	Pro	Leu	Arg
	450					455					460				
Ser	Asn	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu
465					470					475					480
Pro	Asn	Ala	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln
				485					490					495	
Asp	Gln	Pro	Ala	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro
			500					505					510		
Glu	Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Asp	Met
		515					520					525			
Val	Asn	Val	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp
	530					535					540				
Thr	Glu	Ala	Lys	Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu
545					550					555					560
Glu	Met	Leu	Ala	Pro	Tyr	Ile	Pro	Met	Asp	Asp	Asp	Phe	Gln	Leu	Arg
				565					570					575	
Ser	Phe	Asp	Gln	Leu	Ser	Pro	Leu	Glu	Ser	Ser	Ser	Pro	Asn	Pro	Pro
			580					585					590		
Ser	Val	Ser	Thr	Ala	Phe	Gln	Gln	Thr	Gln	Leu	Gln	Glu	Pro	Thr	Ile
		595					600					605			
Thr	Thr	Thr	Thr	Thr	Glu	Glu	Leu	Lys	Thr	Val	Thr	Lys	Asp	Ser	Thr
	610					615						620			
Glu	Asp	Ile	Lys	Ile	Leu	Ile	Thr	Ser	Pro	Ser	Ser	Thr	His	Thr	Pro
625					630					635					640
Lys	Glu	Thr	Thr	Thr	Ala	Thr	Thr	Ser	Ser	Pro	Tyr	Ser	Gly	Thr	Gln
				645					650					655	
Ser	Arg	Thr	Ala	Ser	Pro	Asn	Arg	Ala	Gly	Gln	Gly	Val	Ile	Glu	Gln
			660					665					670		
Thr	Glu	Lys	Ser	His	Pro	Arg	Ser	Pro	Asn	Val	Leu	Ser	Val	Thr	Leu
	675						680					685			

-continued

Ser Gln Arg Asn Thr Val Pro Glu Glu Glu Leu Asn Pro Lys Ile Ile
 690 695 700
 Ala Leu Gln Asn Ala Gln Arg Lys Arg Lys Met Glu His Asp Gly Ser
 705 710 715 720
 Leu Phe Gln Ala Ala Gly Ile Gly Thr Leu Leu Gln Gln Pro Asp Asp
 725 730 735
 Arg Ala Pro Ala Thr Ser Leu Ser Trp Lys Arg Val Lys Gly Cys Lys
 740 745 750
 Ser Ser Gly Gln Asn Gly Met Glu Gln Lys Thr Ile Ile Leu Ile Pro
 755 760 765
 Ser Asp Leu Ala Cys Arg Leu Leu Gly Gln Ser Met Asp Gly Ser Gly
 770 775 780
 Leu Pro Gln Leu Thr Ser Tyr Asp Cys Glu Val Asn Ala Pro Ile Gln
 785 790 795 800
 Gly Ser Arg Asn Leu Leu Gln Gly Glu Glu Leu Leu Arg Ala Leu Asp
 805 810 815
 Gln Val Asn

<210> SEQ ID NO 3
 <211> LENGTH: 836
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 3

Met Glu Gly Ala Gly Gly Glu Asn Glu Lys Lys Lys Met Ser Ser Glu
 1 5 10 15
 Arg Arg Lys Glu Lys Ser Arg Asp Ala Ala Arg Ser Arg Arg Ser Lys
 20 25 30
 Glu Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro His
 35 40 45
 Asn Val Ser Ser His Leu Asp Lys Ala Ser Val Met Arg Leu Thr Ile
 50 55 60
 Ser Tyr Leu Arg Val Arg Lys Leu Leu Asp Ala Gly Gly Leu Asp Ser
 65 70 75 80
 Glu Asp Glu Met Lys Ala Gln Met Asp Cys Phe Tyr Leu Lys Ala Leu
 85 90 95
 Asp Gly Phe Val Met Val Leu Thr Asp Asp Gly Asp Met Val Tyr Ile
 100 105 110
 Ser Asp Asn Val Asn Lys Tyr Met Gly Leu Thr Gln Phe Glu Leu Thr
 115 120 125
 Gly His Ser Val Phe Asp Phe Thr His Pro Cys Asp His Glu Glu Met
 130 135 140
 Arg Glu Met Leu Thr His Arg Asn Gly Pro Val Arg Lys Gly Lys Glu
 145 150 155 160
 Leu Asn Thr Gln Arg Ser Phe Phe Leu Arg Met Lys Cys Thr Leu Thr
 165 170 175
 Ser Arg Gly Arg Thr Met Asn Ile Lys Ser Ala Thr Trp Lys Val Leu
 180 185 190
 His Cys Thr Gly His Ile His Val Tyr Asp Thr Asn Ser Asn Gln Pro
 195 200 205
 Gln Cys Gly Tyr Lys Lys Pro Pro Met Thr Cys Leu Val Leu Ile Cys
 210 215 220

-continued

Glu	Pro	Ile	Pro	His	Pro	Ser	Asn	Ile	Glu	Ile	Pro	Leu	Asp	Ser	Lys	225	230	235	240
Thr	Phe	Leu	Ser	Arg	His	Ser	Leu	Asp	Met	Lys	Phe	Ser	Tyr	Cys	Asp	245	250	255	
Glu	Arg	Ile	Thr	Glu	Leu	Met	Gly	Tyr	Glu	Pro	Glu	Glu	Leu	Leu	Gly	260	265	270	
Arg	Ser	Ile	Tyr	Glu	Tyr	Tyr	His	Ala	Leu	Asp	Ser	Asp	His	Leu	Thr	275	280	285	
Lys	Thr	His	His	Asp	Met	Phe	Thr	Lys	Gly	Gln	Val	Thr	Thr	Gly	Gln	290	295	300	
Tyr	Arg	Met	Leu	Ala	Lys	Arg	Gly	Gly	Tyr	Val	Trp	Val	Glu	Thr	Gln	305	310	315	320
Ala	Thr	Val	Ile	Tyr	Asn	Thr	Lys	Asn	Ser	Gln	Pro	Gln	Cys	Ile	Val	325	330	335	
Cys	Val	Asn	Tyr	Val	Val	Ser	Gly	Ile	Ile	Gln	His	Asp	Leu	Ile	Phe	340	345	350	
Ser	Leu	Gln	Gln	Thr	Glu	Ser	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	Asp	355	360	365	
Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr	Ser	370	375	380	
Cys	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu	Leu	385	390	395	400
Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	Asp	405	410	415	
Asp	Thr	Glu	Thr	Glu	Asp	Gln	Gln	Leu	Glu	Asp	Val	Pro	Leu	Tyr	Asn	420	425	430	
Asp	Val	Met	Phe	Pro	Ser	Ser	Asn	Glu	Lys	Leu	Asn	Ile	Asn	Leu	Ala	435	440	445	
Met	Ser	Pro	Leu	Pro	Ser	Ser	Glu	Thr	Pro	Lys	Pro	Leu	Arg	Ser	Ser	450	455	460	
Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Ser	Ser	465	470	475	480
Pro	Glu	Ser	Leu	Gly	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp	Gln	485	490	495	
Pro	Ala	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu	Arg	500	505	510	
Leu	Leu	Gln	Glu	Asn	Val	Asn	Thr	Pro	Asn	Phe	Ser	Gln	Pro	Asn	Ser	515	520	525	
Pro	Ser	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Asp	Met	Val	Asn	Val	Phe	530	535	540	
Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr	Glu	Ala	Lys	545	550	555	560
Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu	Glu	Met	Leu	Ala	565	570	575	
Pro	Tyr	Ile	Pro	Met	Asp	Asp	Asp	Phe	Gln	Leu	Arg	Ser	Phe	Asp	Gln	580	585	590	
Leu	Ser	Pro	Leu	Glu	Ser	Asn	Ser	Pro	Ser	Pro	Pro	Ser	Met	Ser	Thr	595	600	605	
Val	Thr	Gly	Phe	Gln	Gln	Thr	Gln	Leu	Gln	Lys	Pro	Thr	Ile	Thr	Ala	610	615	620	

-continued

Thr Ala Thr Thr Thr Ala Thr Thr Asp Glu Ser Lys Thr Glu Thr Lys
 625 630 635 640
 Asp Asn Lys Glu Asp Ile Lys Ile Leu Ile Ala Ser Pro Ser Ser Thr
 645 650 655
 Gln Val Pro Gln Glu Thr Thr Thr Ala Lys Ala Ser Ala Tyr Ser Gly
 660 665 670
 Thr His Ser Arg Thr Ala Ser Pro Asp Arg Ala Gly Lys Arg Val Ile
 675 680 685
 Glu Gln Thr Asp Lys Ala His Pro Arg Ser Leu Lys Leu Ser Ala Thr
 690 695 700
 Leu Asn Gln Arg Asn Thr Val Pro Glu Glu Glu Leu Asn Pro Lys Thr
 705 710 715 720
 Ile Ala Ser Gln Asn Ala Gln Arg Lys Arg Lys Met Glu His Asp Gly
 725 730 735
 Ser Leu Phe Gln Ala Ala Gly Ile Gly Thr Leu Leu Gln Gln Pro Gly
 740 745 750
 Asp Cys Ala Pro Thr Met Ser Leu Ser Trp Lys Arg Val Lys Gly Phe
 755 760 765
 Ile Ser Ser Glu Gln Asn Gly Thr Glu Gln Lys Thr Ile Ile Leu Ile
 770 775 780
 Pro Ser Asp Leu Ala Cys Arg Leu Leu Gly Gln Ser Met Asp Glu Ser
 785 790 795 800
 Gly Leu Pro Gln Leu Thr Ser Tyr Asp Cys Glu Val Asn Ala Pro Ile
 805 810 815
 Gln Gly Ser Arg Asn Leu Leu Gln Gly Glu Glu Leu Leu Arg Ala Leu
 820 825 830
 Asp Gln Val Asn
 835

<210> SEQ ID NO 4

<211> LENGTH: 823

<212> TYPE: PRT

<213> ORGANISM: Rattus norvegicus

<400> SEQUENCE: 4

Met Glu Gly Ala Gly Gly Glu Asn Glu Lys Lys Asn Arg Met Ser Ser
 1 5 10 15
 Glu Arg Arg Lys Glu Lys Ser Arg Asp Ala Ala Arg Ser Arg Arg Ser
 20 25 30
 Lys Glu Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro
 35 40 45
 His Asn Val Ser Ser His Leu Asp Lys Ala Ser Val Met Arg Leu Thr
 50 55 60
 Ile Ser Tyr Leu Arg Val Arg Lys Leu Leu Gly Ala Gly Asp Leu Asp
 65 70 75 80
 Ile Glu Asp Glu Met Lys Ala Gln Met Asn Cys Phe Tyr Leu Lys Ala
 85 90 95
 Leu Asp Gly Phe Val Met Val Leu Thr Asp Asp Gly Asp Met Ile Tyr
 100 105 110
 Ile Ser Asp Asn Val Asn Lys Tyr Met Gly Leu Thr Gln Phe Glu Leu
 115 120 125
 Thr Gly His Ser Val Phe Asp Phe Thr His Pro Cys Asp His Glu Glu
 130 135 140

-continued

Met	Arg	Glu	Met	Leu	Thr	His	Arg	Asn	Gly	Pro	Val	Arg	Lys	Gly	Lys	145	150	155	160
Glu	Gln	Asn	Thr	Gln	Arg	Ser	Phe	Phe	Leu	Arg	Met	Lys	Cys	Thr	Leu	165	170	175	
Thr	Ser	Arg	Gly	Arg	Thr	Met	Asn	Ile	Lys	Ser	Ala	Thr	Trp	Lys	Val	180	185	190	
Leu	His	Cys	Thr	Gly	His	Ile	His	Val	Tyr	Asp	Thr	Ser	Ser	Asn	Gln	195	200	205	
Pro	Gln	Cys	Gly	Tyr	Lys	Lys	Pro	Pro	Met	Thr	Cys	Leu	Val	Leu	Ile	210	215	220	
Cys	Glu	Pro	Ile	Pro	His	Pro	Ser	Asn	Ile	Glu	Ile	Pro	Leu	Asp	Ser	225	230	235	240
Lys	Thr	Phe	Leu	Ser	Arg	His	Ser	Leu	Asp	Met	Lys	Phe	Ser	Tyr	Cys	245	250	255	
Asp	Glu	Arg	Ile	Thr	Glu	Leu	Met	Gly	Tyr	Glu	Pro	Glu	Glu	Leu	Leu	260	265	270	
Gly	Arg	Ser	Ile	Tyr	Glu	Tyr	Tyr	His	Ala	Leu	Asp	Ser	Asp	His	Leu	275	280	285	
Thr	Lys	Thr	His	His	Asp	Met	Phe	Thr	Lys	Gly	Gln	Val	Thr	Thr	Gly	290	295	300	
Gln	Tyr	Arg	Met	Leu	Ala	Lys	Arg	Gly	Gly	Tyr	Val	Trp	Val	Glu	Thr	305	310	315	320
Gln	Ala	Thr	Val	Ile	Tyr	Asn	Thr	Lys	Asn	Ser	Gln	Pro	Gln	Cys	Ile	325	330	335	
Val	Cys	Val	Asn	Tyr	Val	Val	Ser	Gly	Ile	Ile	Gln	His	Asp	Leu	Ile	340	345	350	
Phe	Ser	Leu	Gln	Gln	Thr	Glu	Ser	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	355	360	365	
Asp	Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr	370	375	380	
Ser	Cys	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu	385	390	395	400
Leu	Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	405	410	415	
Asp	Asp	Thr	Glu	Thr	Glu	Asp	Gln	Gln	Leu	Glu	Asp	Val	Pro	Leu	Tyr	420	425	430	
Asn	Asp	Val	Met	Phe	Pro	Ser	Ser	Asn	Glu	Lys	Leu	Asn	Ile	Asn	Leu	435	440	445	
Ala	Met	Ser	Pro	Leu	Pro	Ala	Ser	Glu	Thr	Pro	Lys	Pro	Leu	Arg	Ser	450	455	460	
Ser	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Ser	465	470	475	480
Ser	Pro	Glu	Ser	Leu	Gly	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp	485	490	495	
Gln	Pro	Ala	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu	500	505	510	
Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Asp	Met	Val	515	520	525	
Asn	Val	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr	530	535	540	

-continued

Glu Ala Lys Asn Pro Phe Ser Ala Gln Asp Thr Asp Leu Asp Leu Glu
 545 550 555 560
 Met Leu Ala Pro Tyr Ile Pro Met Asp Asp Asp Phe Gln Leu Arg Ser
 565 570 575
 Phe Asp Gln Leu Ser Pro Leu Glu Ser Asn Ser Pro Ser Pro Pro Ser
 580 585 590
 Val Ser Thr Val Thr Gly Phe Gln Gln Thr Gln Leu Gln Lys Pro Thr
 595 600 605
 Ile Thr Val Thr Ala Thr Ala Thr Ala Thr Thr Asp Glu Ser Lys Ala
 610 615 620
 Val Thr Lys Asp Asn Ile Glu Asp Ile Lys Ile Leu Ile Ala Ser Pro
 625 630 635 640
 Pro Ser Thr Gln Val Pro Gln Glu Met Thr Thr Ala Lys Ala Ser Ala
 645 650 655
 Tyr Ser Gly Thr His Ser Arg Thr Ala Ser Pro Asp Arg Ala Gly Lys
 660 665 670
 Arg Val Ile Glu Lys Thr Asp Lys Ala His Pro Arg Ser Leu Asn Leu
 675 680 685
 Ser Val Thr Leu Asn Gln Arg Asn Thr Val Pro Glu Glu Glu Leu Asn
 690 695 700
 Pro Lys Thr Ile Ala Leu Gln Asn Ala Gln Arg Lys Arg Lys Met Glu
 705 710 715 720
 His Asp Gly Ser Leu Phe Gln Ala Ala Gly Ile Gly Thr Leu Leu Gln
 725 730 735
 Gln Pro Gly Asp Arg Ala Pro Thr Met Ser Leu Ser Trp Lys Arg Val
 740 745 750
 Lys Gly Tyr Ile Ser Ser Glu Gln Asp Gly Met Glu Gln Lys Thr Ile
 755 760 765
 Phe Leu Ile Pro Ser Asp Leu Ala Cys Arg Leu Leu Gly Gln Ser Met
 770 775 780
 Asp Glu Ser Gly Leu Pro Gln Leu Thr Ser Tyr Asp Cys Glu Val Asn
 785 790 795 800
 Ala Pro Ile Gln Gly Ser Arg Asn Leu Leu Gln Gly Glu Glu Leu Leu
 805 810 815
 Arg Ala Leu Asp Gln Val Asn
 820

<210> SEQ ID NO 5

<211> LENGTH: 828

<212> TYPE: PRT

<213> ORGANISM: Microtus oeconomus

<400> SEQUENCE: 5

Met Glu Gly Ala Gly Gly Glu Asn Glu Lys Lys Lys Met Ser Ser Glu
 1 5 10 15
 Arg Arg Lys Glu Lys Ser Arg Asp Ala Ala Arg Ser Arg Arg Ser Lys
 20 25 30
 Glu Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro His
 35 40 45
 Asn Val Ser Ser His Leu Asp Lys Ala Ser Val Met Arg Leu Thr Ile
 50 55 60
 Ser Tyr Leu Arg Val Arg Lys Leu Leu Asp Ala Gly Asp Leu Asp Ile
 65 70 75 80

-continued

Glu	Asp	Glu	Met	Lys	Ala	Gln	Met	Asn	Cys	Phe	Tyr	Leu	Lys	Ala	Leu	85	90	95
Asp	Gly	Phe	Val	Met	Val	Leu	Thr	Asp	Asp	Gly	Asp	Met	Ile	Tyr	Ile	100	105	110
Ser	Asp	Asn	Val	Asn	Lys	Tyr	Met	Gly	Leu	Thr	Gln	Phe	Glu	Leu	Thr	115	120	125
Gly	His	Ser	Val	Phe	Asp	Phe	Thr	His	Pro	Cys	Asp	His	Glu	Glu	Met	130	135	140
Arg	Glu	Met	Leu	Thr	His	Arg	Asn	Gly	Pro	Val	Lys	Lys	Gly	Lys	Glu	145	150	155
Gln	Asn	Thr	Gln	Arg	Ser	Phe	Phe	Leu	Arg	Met	Lys	Cys	Thr	Leu	Thr	165	170	175
Ser	Arg	Gly	Arg	Thr	Met	Asn	Ile	Lys	Ser	Ala	Thr	Trp	Lys	Val	Leu	180	185	190
His	Cys	Thr	Gly	His	Ile	His	Val	Tyr	Asp	Thr	Asn	Ser	Asn	Gln	Ser	195	200	205
Gln	Cys	Gly	Tyr	Lys	Lys	Pro	Pro	Met	Thr	Cys	Leu	Val	Leu	Ile	Cys	210	215	220
Glu	Pro	Ile	Pro	His	Pro	Ser	Asn	Ile	Glu	Ile	Pro	Leu	Asp	Ser	Lys	225	230	235
Thr	Phe	Leu	Ser	Arg	His	Ser	Leu	Asp	Met	Lys	Phe	Ser	Tyr	Cys	Asp	245	250	255
Glu	Arg	Ile	Thr	Glu	Leu	Met	Gly	Tyr	Glu	Pro	Glu	Glu	Leu	Leu	Gly	260	265	270
Arg	Ser	Ile	Tyr	Glu	Tyr	Tyr	His	Ala	Leu	Asp	Ser	Asp	His	Leu	Thr	275	280	285
Lys	Thr	His	His	Asp	Met	Phe	Thr	Lys	Gly	Gln	Val	Thr	Thr	Gly	Gln	290	295	300
Tyr	Arg	Met	Leu	Ala	Lys	Arg	Gly	Gly	Tyr	Val	Trp	Val	Glu	Thr	Gln	305	310	315
Ala	Thr	Val	Ile	Tyr	Asn	Thr	Lys	Asn	Ser	Gln	Pro	Gln	Cys	Ile	Val	325	330	335
Cys	Val	Asn	Tyr	Val	Val	Ser	Gly	Ile	Ile	Gln	His	Asp	Leu	Ile	Phe	340	345	350
Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	Asp	355	360	365
Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr	Ser	370	375	380
Cys	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu	Leu	385	390	395
Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	Asp	405	410	415
Asp	Thr	Glu	Thr	Glu	Asp	Gln	Gln	Leu	Glu	Asp	Val	Pro	Leu	Tyr	Asn	420	425	430
Asp	Val	Met	Phe	Pro	Ser	Ser	Asn	Glu	Lys	Leu	Thr	Asn	Ile	Asn	Leu	435	440	445
Ala	Leu	Ser	Pro	Leu	Pro	Ala	Ser	Glu	Thr	Pro	Lys	Pro	Leu	Arg	Ser	450	455	460
Ser	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Pro	465	470	475

Ser	Pro	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp	
				485					490					495		
Gln	Pro	Ala	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu	
			500				505				510					
Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Asp	Met	Val	
			515				520				525					
Asn	Val	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr	
			530				535				540					
Glu	Ala	Lys	Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu	Glu	
			545				550				555					
Met	Leu	Ala	Pro	Tyr	Ile	Pro	Met	Asp	Asp	Asp	Phe	Gln	Leu	Arg	Ser	
			565				570				575					
Phe	Asp	Gln	Leu	Ser	Pro	Leu	Glu	Ser	Ser	Ser	Ser	Ser	Pro	Pro	Ser	
			580				585				590					
Val	Ser	Thr	Val	Thr	Gly	Phe	Gln	Gln	Thr	Pro	Leu	Gln	Lys	Pro	Thr	
			595				600				605					
Ile	Thr	Ala	Ala	Ala	Ala	Thr	Thr	Ala	Thr	Thr	Thr	Ala	Thr	Thr	Thr	
			610				615				620					
Asp	Glu	Leu	Lys	Thr	Val	Thr	Lys	Asp	Asn	Ile	Glu	Asp	Ile	Lys	Ile	
			625				630				635					
Leu	Ile	Ala	Ser	Pro	Ser	Ser	Val	His	Val	Pro	Gln	Glu	Thr	Thr	Ala	
			645				650				655					
Ala	Thr	Ala	Ser	Ala	Tyr	Ser	Ala	Pro	His	Ser	Arg	Thr	Ala	Ser	Pro	
			660				665				670					
Asp	Arg	Ala	Gly	Lys	Arg	Val	Arg	Glu	Gln	Ala	Glu	Lys	Ala	His	Pro	
			675				680				685					
Arg	Ser	Pro	Asn	Met	Ser	Val	Thr	Leu	Ser	Gln	Arg	Asn	Thr	Val	Thr	
			690				695				700					
Glu	Glu	Asp	Leu	Asn	Pro	Lys	Ile	Ile	Ala	Leu	Gln	Asn	Ala	Gln	Arg	
			705				710				715					
Lys	Arg	Lys	Met	Glu	His	Asp	Gly	Ser	Leu	Phe	Gln	Ala	Ser	Gly	Ile	
			725				730				735					
Gly	Thr	Leu	Leu	Gln	Gln	Pro	Gly	Asp	Arg	Ala	Pro	Thr	Thr	Ser	Leu	
			740				745				750					
Ser	Trp	Lys	Arg	Val	Lys	Gly	Cys	Lys	Ser	Asn	Glu	Gln	Asn	Gly	Met	
			755				760				765					
Glu	Gln	Lys	Thr	Ile	Ile	Leu	Ile	Pro	Ser	Asp	Leu	Ala	Cys	Arg	Leu	
			770				775				780					
Leu	Gly	Gln	Ser	Met	Asp	Glu	Ser	Gly	Leu	Pro	Gln	Leu	Thr	Ser	Tyr	
			785				790				795					
Asp	Cys	Glu	Val	Asn	Ala	Pro	Val	Gln	Gly	Ser	Arg	Asn	Leu	Leu	Gln	
			805				810				815					
Gly	Glu	Asp	Leu	Leu	Arg	Ala	Leu	Asp	Gln	Val	Asn					
			820				825									

```
<210> SEQ ID NO 6
<211> LENGTH: 826
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

<400> SEQUENCE: 6

Met Glu Gly Ala Gly Gly Ala Asn Asp Lys Lys Lys Ile Ser Ser Glu
1 5 10 15

-continued

Arg	Arg	Lys	Glu	Lys	Ser	Arg	Asp	Ala	Ala	Arg	Ser	Arg	Arg	Ser	Lys	20	25	30
Glu	Ser	Glu	Val	Phe	Tyr	Glu	Leu	Ala	His	Gln	Leu	Pro	Leu	Pro	His	35	40	45
Asn	Val	Ser	Ser	His	Leu	Asp	Lys	Ala	Ser	Val	Met	Arg	Leu	Thr	Ile	50	55	60
Ser	Tyr	Leu	Arg	Val	Arg	Lys	Leu	Leu	Asp	Ala	Gly	Asp	Leu	Asp	Ile	65	70	75
Glu	Asp	Asp	Met	Lys	Ala	Gln	Met	Asn	Cys	Phe	Tyr	Leu	Lys	Ala	Leu	85	90	95
Asp	Gly	Phe	Val	Met	Val	Leu	Thr	Asp	Asp	Gly	Asp	Met	Ile	Tyr	Ile	100	105	110
Ser	Asp	Asn	Val	Asn	Lys	Tyr	Met	Gly	Leu	Thr	Gln	Phe	Glu	Leu	Thr	115	120	125
Gly	His	Ser	Val	Phe	Asp	Phe	Thr	His	Pro	Cys	Asp	His	Glu	Glu	Met	130	135	140
Arg	Glu	Met	Leu	Thr	His	Arg	Asn	Gly	Leu	Val	Lys	Lys	Gly	Lys	Glu	145	150	155
Gln	Asn	Thr	Gln	Arg	Ser	Phe	Phe	Leu	Arg	Met	Lys	Cys	Thr	Leu	Thr	165	170	175
Ser	Arg	Gly	Arg	Thr	Met	Asn	Ile	Lys	Ser	Ala	Thr	Trp	Lys	Val	Leu	180	185	190
His	Cys	Thr	Gly	His	Ile	His	Val	Tyr	Asp	Thr	Asn	Ser	Asn	Gln	Pro	195	200	205
Gln	Cys	Gly	Tyr	Lys	Lys	Pro	Pro	Met	Thr	Cys	Leu	Val	Leu	Ile	Cys	210	215	220
Glu	Pro	Ile	Pro	His	Pro	Ser	Asn	Ile	Glu	Ile	Pro	Leu	Asp	Ser	Lys	225	230	235
Thr	Phe	Leu	Ser	Arg	His	Ser	Leu	Asp	Met	Lys	Phe	Ser	Tyr	Cys	Asp	245	250	255
Glu	Arg	Ile	Thr	Glu	Leu	Met	Gly	Tyr	Glu	Pro	Glu	Glu	Leu	Leu	Gly	260	265	270
Arg	Ser	Ile	Tyr	Glu	Tyr	Tyr	His	Ala	Leu	Asp	Ser	Asp	His	Leu	Thr	275	280	285
Lys	Thr	His	His	Asp	Met	Phe	Thr	Lys	Gly	Gln	Val	Thr	Thr	Gly	Gln	290	295	300
Tyr	Arg	Met	Leu	Ala	Lys	Arg	Gly	Gly	Tyr	Val	Trp	Val	Glu	Thr	Gln	305	310	315
Ala	Thr	Val	Ile	Tyr	Asn	Thr	Lys	Asn	Ser	Gln	Pro	Gln	Cys	Ile	Val	325	330	335
Cys	Val	Asn	Tyr	Val	Val	Ser	Gly	Ile	Ile	Gln	His	Asp	Leu	Ile	Phe	340	345	350
Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	Asp	355	360	365
Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr	Ser	370	375	380
Ser	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu	Leu	385	390	395
Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	Asn	405	410	415

-continued

Asp	Thr	Glu	Thr	Asp	Asp	Gln	Gln	Leu	Glu	Glu	Val	Pro	Leu	Tyr	Asn	
			420					425						430		
Asp	Val	Met	Leu	Pro	Ser	Pro	Asn	Glu	Lys	Leu	Gln	Asn	Ile	Asn	Leu	
		435					440					445				
Ala	Met	Ser	Pro	Leu	Pro	Thr	Ala	Glu	Thr	Pro	Lys	Pro	Leu	Arg	Ser	
	450					455					460					
Ser	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Pro	
465					470					475					480	
Asn	Pro	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp	
				485					490						495	
Gln	Thr	Pro	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu	
			500					505						510		
Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Tyr	Val	Asp	Ser	Asp	Met	Val	
		515					520					525				
Asn	Glu	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr	
	530					535					540					
Glu	Ala	Lys	Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu	Glu	
545					550					555					560	
Met	Leu	Ala	Pro	Tyr	Ile	Pro	Met	Asp	Asp	Asp	Phe	Gln	Leu	Arg	Ser	
			565						570					575		
Phe	Asp	Gln	Leu	Ser	Pro	Leu	Glu	Ser	Ser	Ser	Ala	Ser	Pro	Glu	Ser	
		580						585					590			
Ala	Ser	Pro	Gln	Ser	Thr	Val	Thr	Val	Phe	Gln	Gln	Thr	Gln	Ile	Gln	
		595					600					605				
Glu	Pro	Thr	Ala	Asn	Ala	Thr	Thr	Thr	Thr	Ala	Thr	Thr	Asp	Glu	Leu	
	610					615					620					
Lys	Thr	Val	Thr	Lys	Asp	Arg	Met	Glu	Asp	Ile	Lys	Ile	Leu	Ile	Ala	
625				630						635					640	
Ser	Pro	Ser	Pro	Thr	His	Ile	His	Lys	Glu	Thr	Thr	Ser	Ala	Thr	Ser	
			645						650					655		
Ser	Pro	Tyr	Arg	Asp	Thr	Gln	Ser	Arg	Thr	Ala	Ser	Pro	Asn	Arg	Ala	
		660						665					670			
Gly	Lys	Gly	Val	Ile	Glu	Gln	Thr	Glu	Lys	Ser	His	Pro	Arg	Ser	Pro	
	675					680						685				
Asn	Val	Leu	Ser	Val	Ala	Leu	Ser	Gln	Arg	Thr	Thr	Val	Pro	Glu	Glu	
	690					695					700					
Glu	Leu	Asn	Pro	Lys	Ile	Leu	Ala	Leu	Gln	Asn	Ala	Gln	Arg	Lys	Arg	
705				710						715					720	
Lys	Met	Glu	His	Asp	Gly	Ser	Leu	Phe	Gln	Ala	Val	Gly	Ile	Gly	Thr	
			725					730					735			
Leu	Leu	Gln	Gln	Pro	Asp	Asp	His	Ala	Ala	Thr	Thr	Ser	Leu	Ser	Trp	
		740					745						750			
Lys	Arg	Val	Lys	Gly	Cys	Lys	Ser	Ser	Glu	Gln	Asn	Gly	Met	Glu	Gln	
	755					760						765				
Lys	Thr	Ile	Ile	Leu	Ile	Pro	Ser	Asp	Leu	Ala	Cys	Arg	Leu	Leu	Gly	
	770				775						780					
Gln	Ser	Met	Asp	Glu	Ser	Gly	Leu	Pro	Gln	Leu	Thr	Ser	Tyr	Asp	Cys	
785					790					795					800	
Glu	Val	Asn	Ala	Pro	Ile	Gln	Gly	Ser	Arg	Asn	Leu	Leu	Gln	Gly	Glu	
			805					810						815		
Glu	Leu	Leu	Arg	Ala	Leu	Asp	Gln	Val	Asn							

```

      820                               825

<210> SEQ ID NO 7
<211> LENGTH: 827
<212> TYPE: PRT
<213> ORGANISM: Pongo pygmaeus

<400> SEQUENCE: 7

Met Glu Gly Ala Gly Gly Ala Asn Asp Lys Lys Asn Arg Ile Ser Ser
 1          5          10          15

Glu Arg Arg Lys Glu Lys Ser Arg Asp Ala Ala Arg Ser Arg Arg Ser
          20          25          30

Lys Glu Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro
          35          40          45

His Asn Val Ser Ser His Leu Asp Lys Ala Ser Val Met Arg Leu Thr
          50          55          60

Ile Ser Tyr Leu Arg Val Arg Lys Leu Leu Asp Ala Gly Asp Leu Asp
65          70          75          80

Ile Glu Asp Glu Met Lys Ala Gln Met Asn Cys Phe Tyr Leu Lys Ala
          85          90          95

Leu Asp Gly Phe Val Met Val Leu Thr Asp Asp Gly Asp Met Ile Tyr
          100          105          110

Ile Ser Asp Asn Val Asn Lys Tyr Met Gly Leu Thr Gln Phe Glu Leu
          115          120          125

Thr Gly His Ser Val Phe Asp Phe Thr His Pro Cys Asp His Glu Glu
          130          135          140

Met Arg Glu Met Leu Thr His Arg Asn Gly Pro Val Lys Lys Gly Lys
145          150          155          160

Glu Gln Asn Thr Gln Arg Ser Phe Phe Leu Arg Met Lys Cys Thr Leu
          165          170          175

Thr Ser Arg Gly Arg Thr Met Asn Ile Lys Ser Ala Thr Trp Lys Val
          180          185          190

Leu His Cys Thr Gly His Ile His Val Tyr Asp Thr Asn Ser Asn Gln
          195          200          205

Pro Gln Cys Gly Tyr Lys Lys Pro Pro Met Thr Cys Leu Val Leu Ile
          210          215          220

Cys Glu Pro Ile Pro His Pro Ser Asn Ile Glu Ile Pro Leu Asp Ser
225          230          235          240

Lys Thr Phe Leu Ser Arg His Ser Leu Asp Met Lys Phe Ser Tyr Cys
          245          250          255

Asp Glu Arg Ile Thr Glu Leu Met Gly Tyr Glu Pro Glu Glu Leu Leu
          260          265          270

Gly Arg Ser Ile Tyr Glu Tyr Tyr His Ala Leu Asp Ser Asp His Leu
          275          280          285

Thr Lys Thr His His Asp Met Phe Thr Lys Gly Gln Val Thr Thr Gly
          290          295          300

Gln Tyr Arg Met Leu Ala Lys Arg Gly Gly Tyr Val Trp Val Glu Thr
305          310          315          320

Gln Ala Thr Val Ile Tyr Asn Thr Lys Asn Ser Gln Pro Gln Cys Ile
          325          330          335

Val Cys Val Asn Tyr Val Val Ser Gly Ile Ile Gln His Asp Leu Ile
          340          345          350

```

-continued

Phe	Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser
		355					360					365			
Asp	Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr
		370			375						380				
Ser	Ser	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu
385				390						395			400		
Leu	Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser
				405					410					415	
Asn	Asp	Thr	Glu	Thr	Asp	Asp	Gln	Gln	Leu	Glu	Glu	Val	Pro	Leu	Tyr
		420						425					430		
Asn	Asp	Val	Met	Leu	Pro	Ser	Pro	Asn	Glu	Lys	Leu	Gln	Asn	Ile	Asn
		435					440					445			
Leu	Ala	Met	Ser	Pro	Leu	Pro	Thr	Ala	Glu	Thr	Pro	Lys	Pro	Leu	Arg
450						455					460				
Ser	Ser	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu
465				470						475			480		
Pro	Asn	Pro	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln
				485					490					495	
Asp	Gln	Pro	Pro	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro
		500						505					510		
Glu	Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Tyr	Val	Asp	Ser	Asp	Met
		515					520					525			
Val	Asn	Glu	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp
530						535					540				
Thr	Glu	Ala	Lys	Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu
545				550						555			560		
Glu	Met	Leu	Ala	Pro	Tyr	Ile	Pro	Met	Asp	Asp	Asp	Phe	Gln	Leu	Arg
				565					570					575	
Ser	Phe	Asp	Gln	Leu	Ser	Pro	Leu	Glu	Ser	Ser	Ser	Ala	Ser	Pro	Glu
		580						585					590		
Ser	Ala	Ser	Pro	Gln	Ser	Thr	Val	Thr	Val	Phe	Gln	Gln	Thr	Gln	Ile
595						600						605			
Gln	Glu	Pro	Thr	Ala	Asn	Ala	Thr	Thr	Thr	Thr	Ala	Thr	Thr	Asp	Glu
610						615					620				
Leu	Lys	Thr	Val	Thr	Lys	Asp	Arg	Met	Glu	Asp	Ile	Lys	Ile	Leu	Ile
625				630						635			640		
Ala	Ser	Pro	Ser	Pro	Thr	His	Ile	His	Lys	Glu	Thr	Thr	Ser	Ala	Thr
				645					650					655	
Ser	Ser	Pro	Tyr	Arg	Asp	Thr	Gln	Ser	Arg	Thr	Ala	Ser	Pro	Asn	Arg
		660						665					670		
Ala	Gly	Lys	Gly	Val	Ile	Glu	Gln	Thr	Glu	Lys	Ser	His	Pro	Arg	Ser
675						680						685			
Pro	Asn	Val	Leu	Ser	Val	Ala	Leu	Ser	Gln	Arg	Thr	Thr	Val	Pro	Glu
690						695					700				
Glu	Glu	Leu	Asn	Pro	Lys	Ile	Leu	Ala	Leu	Gln	Asn	Ala	Gln	Arg	Lys
705				710						715			720		
Arg	Lys	Met	Glu	His	Asp	Gly	Ser	Leu	P						

-continued

755	760	765
Gln Lys Thr Ile Ile Leu Ile	Pro Ser Asp Leu Ala Cys Arg Leu Leu	
770	775	780
Gly Gln Ser Met Asp Glu Ser Gly Leu Pro Gln Leu Thr Ser Tyr Asp		
785	790	800
Cys Glu Val Asn Ala Pro Ile Gln Gly Ser Arg Asn Leu Leu Gln Gly		
805	810	815
Glu Glu Leu Leu Arg Ala Leu Asp Gln Val Asn		
820	825	
<210> SEQ ID NO 8		
<211> LENGTH: 826		
<212> TYPE: PRT		
<213> ORGANISM: Macaca fascicularis		
<400> SEQUENCE: 8		
Met Glu Gly Ala Gly Gly Ala Asn Asp Lys Lys Lys Ile Ser Ser Glu		
1	5	10
Arg Arg Lys Glu Lys Ser Arg Asp Ala Ala Arg Ser Arg Arg Ser Lys		
20	25	30
Glu Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro His		
35	40	45
Asn Val Ser Ser His Leu Asp Lys Ala Ser Val Met Arg Leu Thr Ile		
50	55	60
Ser Tyr Leu Arg Val Arg Lys Leu Leu Asp Ala Gly Asp Leu Asp Ile		
65	70	75
Glu Asp Glu Met Lys Ala Gln Met Asn Cys Phe Tyr Leu Lys Ala Leu		
85	90	95
Asp Gly Phe Val Met Val Leu Thr Asp Asp Gly Asp Met Ile Tyr Ile		
100	105	110
Ser Asp Asn Val Asn Lys Tyr Met Gly Leu Thr Gln Phe Glu Leu Thr		
115	120	125
Gly His Ser Val Phe Asp Phe Thr His Pro Cys Gly His Glu Glu Met		
130	135	140
Arg Glu Met Leu Thr His Arg Asn Gly Pro Val Lys Lys Gly Lys Glu		
145	150	155
Gln Asn Thr Gln Arg Ser Phe Phe Leu Arg Met Lys Cys Thr Leu Thr		
165	170	175
Ser Arg Gly Arg Thr Met Asn Ile Lys Ser Ala Thr Trp Lys Val Leu		
180	185	190
His Cys Thr Gly His Ile His Val Tyr Asp Thr Asn Ser Asn Gln Pro		
195	200	205
Gln Cys Gly Tyr Lys Lys Pro Pro Met Thr Cys Leu Val Leu Ile Cys		
210	215	220
Glu Pro Ile Pro His Pro Ser Asn Ile Glu Ile Pro Leu Asp Ser Lys		
225	230	235
Thr Phe Leu Ser Arg His Ser Leu Asp Met Lys Phe Ser Tyr Cys Asp		
245	250	255
Glu Arg Ile Thr Glu Leu Met Gly Tyr Glu Pro Glu Glu Leu Leu Gly		
260	265	270
Arg Ser Ile Tyr Glu Tyr Tyr His Ala Leu Asp Ser Asp His Leu Thr		
275	280	285

-continued

Lys	Thr	His	His	Gly	Met	Phe	Thr	Lys	Gly	Gln	Val	Thr	Thr	Gly	Gln
290						295					300				
Tyr	Arg	Met	Leu	Ala	Lys	Arg	Gly	Gly	Tyr	Val	Trp	Val	Glu	Thr	Gln
305					310					315					320
Ala	Thr	Val	Ile	Tyr	Asn	Thr	Lys	Asn	Ser	Gln	Pro	Gln	Cys	Ile	Val
				325					330					335	
Cys	Val	Asn	Tyr	Val	Val	Ser	Gly	Ile	Ile	Gln	His	Asp	Leu	Ile	Phe
			340					345					350		
Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	Asp
		355					360					365			
Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr	Ser
	370					375					380				
Ser	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu	Leu
385					390					395					400
Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	Asn
				405					410					415	
Asp	Thr	Glu	Thr	Asp	Asp	Gln	Gln	Leu	Glu	Glu	Val	Pro	Leu	Tyr	Asn
			420					425					430		
Asp	Val	Met	Leu	Pro	Ser	Ser	Asn	Glu	Lys	Leu	Gln	Asn	Ile	Asn	Leu
		435					440					445			
Ala	Met	Ser	Pro	Leu	Pro	Thr	Ser	Glu	Thr	Pro	Lys	Pro	Leu	Arg	Ser
	450					455					460				
Ser	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Pro
465					470					475					480
Asn	Pro	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp
			485					490						495	
Gln	Pro	Pro	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu
			500					505					510		
Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Tyr	Val	Asp	Ser	Asp	Met	Val
		515					520					525			
Asn	Glu	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr
	530					535					540				
Glu	Ala	Lys	Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu	Glu
545					550					555					560
Met	Leu	Ala	Pro	Tyr	Ile	Pro	Met	Asp	Asp	Asp	Phe	Gln	Leu	Arg	Ser
			565					570						575	
Phe	Asp	Gln	Leu	Ser	Pro	Leu	Glu	Ser	Ser	Ser	Ala	Ser	Pro	Glu	Ser
			580					585					590		
Ala	Ser	Pro	Gln	Ser	Thr	Val	Thr	Val	Phe	Gln	Gln	Thr	Gln	Ile	Gln
		595					600					605			
Glu	Pro	Thr	Ala	Asn	Ala	Thr	Thr	Thr	Thr	Ala	Thr	Thr	Asp	Glu	Leu
	610					615					620				
Lys	Thr	Val	Thr	Lys	Asp	Arg	Met	Glu	Asp	Ile	Lys	Ile	Leu	Ile	Ala
625					630					635					640
Ser	Pro	Ser	Ser	Thr	His	Ile	His	Lys	Glu	Thr	Thr	Ser	Ala	Thr	Ser
			645						650					655	
Ser	Pro	Tyr	Arg	Asp	Thr	Gln	Ser	Arg	Thr	Ala	Ser	Pro	Asn	Arg	Ala
			660					665					670		
Gly	Lys	Gly	Val	Ile	Glu	Gln	Thr	Glu	Lys	Ser	His	Pro	Arg	Ser	Pro
		675					680					685			
Asn	Val	Leu	Ser	Val	Thr	Leu	Ser	Gln	Arg	Thr	Thr	Val	Pro	Glu	Glu

-continued

690	695	700
Glu Leu Asn Pro Lys	Ile Leu Ala Leu Gln Asn Ala Gln Arg Lys Arg	
705	710	715 720
Lys Met Glu His Asp Gly Ser Leu Phe Gln Ala Val Gly Ile Gly Thr		
	725	730 735
Leu Leu Gln Gln Pro Asp Asp His Ala Ala Thr Thr Ser Leu Ser Trp		
	740	745 750
Lys Arg Val Lys Gly Cys Lys Ser Ser Glu Gln Asn Gly Met Glu Gln		
	755	760 765
Lys Thr Ile Ile Leu Ile Pro Ser Asp Leu Ala Cys Arg Leu Leu Gly		
	770	775 780
Gln Ser Met Asp Glu Ser Gly Leu Pro Gln Leu Thr Ser Tyr Asp Cys		
	785	790 795 800
Glu Val Asn Ala Pro Ile Gln Gly Ser Arg Asn Leu Leu Gln Gly Glu		
	805	810 815
Glu Leu Leu Arg Ala Leu Asp Gln Val Asn		
	820	825

<210> SEQ ID NO 9
 <211> LENGTH: 821
 <212> TYPE: PRT
 <213> ORGANISM: *Spermophilus tridecemlineatus*
 <220> FEATURE:
 <221> NAME/KEY: misc feature
 <222> LOCATION: (49)..(49)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 9

Met Glu Gly Ala Gly Gly Thr Asn Asp Lys Lys Lys Ile Ser Ser Glu		
1	5	10 15
Arg Arg Lys Glu Lys Ser Arg Asp Ala Ala Arg Ser Arg Arg Ser Lys		
	20	25 30
Glu Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro His		
	35	40 45
Xaa Val Ser Ser His Leu Asp Lys Ala Ser Val Met Arg Leu Thr Ile		
	50	55 60
Ser Tyr Leu Leu Val Arg Lys Leu Leu Asp Ala Gly Asp Leu Asp Ile		
	65	70 75 80
Glu Asp Glu Met Lys Ala Gln Met Asn Cys Phe Tyr Leu Lys Ala Leu		
	85	90 95
Asp Gly Phe Val Met Val Leu Thr Asp Asp Gly Asp Met Ile Tyr Ile		
	100	105 110
Ser Asp Asn Val Asn Lys Tyr Met Gly Leu Thr Gln Phe Glu Leu Thr		
	115	120 125
Gly His Ser Val Phe Asp Phe Thr His Pro Cys Asp His Glu Glu Met		
	130	135 140
Arg Glu Met Leu Thr His Arg Asn Gly Pro Val Lys Lys Gly Lys Glu		
	145	150 155 160
Gln Asn Thr Gln Arg Ser Phe Phe Leu Arg Met Lys Cys Thr Leu Thr		
	165	170 175
Ser Arg Gly Arg Thr Met Asn Ile Lys Ser Ala Thr Trp Lys Val Leu		
	180	185 190
His Cys Thr Gly His Ile His Val Tyr Asp Thr Asn Ser Asn Gln Ser		
	195	200 205

-continued

Gln	Cys	Gly	Tyr	Lys	Lys	Pro	Pro	Met	Thr	Cys	Leu	Val	Leu	Ile	Cys
210						215					220				
Glu	Pro	Ile	Pro	His	Pro	Ser	Asn	Ile	Glu	Ile	Pro	Leu	Asp	Ser	Lys
225					230				235						240
Thr	Phe	Leu	Ser	Arg	His	Ser	Leu	Asp	Met	Lys	Phe	Ser	Tyr	Cys	Asp
				245					250					255	
Glu	Arg	Ile	Thr	Glu	Leu	Met	Gly	Tyr	Glu	Pro	Glu	Glu	Leu	Leu	Gly
			260					265					270		
Arg	Ser	Ile	Tyr	Glu	Tyr	Tyr	His	Ala	Leu	Asp	Ser	Asp	His	Leu	Thr
		275					280					285			
Lys	Thr	His	His	Asp	Met	Phe	Thr	Lys	Gly	Gln	Val	Thr	Thr	Gly	Gln
	290					295					300				
Tyr	Arg	Met	Leu	Ala	Lys	Arg	Gly	Gly	Tyr	Val	Trp	Val	Glu	Thr	Gln
305					310					315					320
Ala	Thr	Val	Ile	Tyr	Asn	Thr	Lys	Asn	Ser	Gln	Pro	Gln	Cys	Ile	Val
				325					330					335	
Cys	Val	Asn	Tyr	Val	Val	Ser	Gly	Ile	Ile	Gln	His	Gly	Leu	Ile	Phe
		340						345					350		
Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	Asp
		355					360					365			
Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr	Ser
	370					375					380				
Ser	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu	Leu
385					390				395						400
Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	Asn
				405					410					415	
Asp	Thr	Glu	Thr	Glu	Asp	Gln	Gln	Leu	Glu	Glu	Val	Pro	Leu	Tyr	Asn
		420						425					430		
Asp	Val	Met	Phe	Pro	Ser	Ser	Ser	Glu	Lys	Leu	Gln	Asn	Ile	Asn	Leu
		435					440					445			
Ala	Met	Ser	Pro	Leu	Pro	Ala	Ser	Glu	Thr	Pro	Lys	Pro	Leu	Arg	Ser
	450					455					460				
Ser	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Pro
465					470				475						480
Asn	Pro	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp
			485						490					495	
Gln	Pro	Ala	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu
		500						505					510		
Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Asp	Met	Val
	515						520					525			
Asn	Val	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr
	530					535					540				
Glu	Ala	Lys	Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu	Glu
545					550				555						560
Met	Leu	Ala	Pro	Tyr	Ile	Pro	Met	Asp	Asp	Asp	Phe	Gln	Leu	Arg	Ser
			565						570					575	
Phe	Asp	Gln	Leu	Ser	Pro	Leu	Glu	Ser	Ser	Ser	Ser	Ser	Pro	Gln	Ser
		580						585					590		
Val	Ser	Thr	Ile	Ser	Val	Phe	Gln	Gln	Thr	Gln	Ile	Gln	Glu	Pro	Thr
	595						600					605			

```

<210> SEQ ID NO 10
<211> LENGTH: 823
<212> TYPE: PRT
<213> ORGANISM: Bos taurus

<400> SEQUENCE: 10

Met  Glu  Gly  Ala  Gly  Gly  Ala  Asn  Asp  Lys  Lys  Lys  Ile  Ser  Ser  Glu
1          5          10          15
Arg  Arg  Lys  Glu  Lys  Ser  Arg  Asp  Ala  Ala  Arg  Ser  Arg  Arg  Ser  Lys
          20          25          30
Glu  Ser  Glu  Val  Phe  Tyr  Glu  Leu  Ala  His  Gln  Leu  Pro  Leu  Pro  His
          35          40          45
Asn  Val  Ser  Ser  His  Leu  Asp  Lys  Ala  Ser  Val  Met  Arg  Leu  Thr  Ile
          50          55          60
Ser  Tyr  Leu  Arg  Val  Arg  Lys  Leu  Leu  Asp  Ala  Gly  Asp  Leu  Asp  Ile
65          70          75          80
Glu  Asp  Glu  Met  Lys  Ala  Gln  Met  Asn  Cys  Phe  Tyr  Leu  Lys  Ala  Leu
          85          90          95
Asp  Gly  Phe  Val  Met  Val  Leu  Thr  Asp  Asp  Gly  Asp  Met  Ile  Tyr  Ile
          100          105          110
Ser  Asp  Asn  Val  Asn  Lys  Tyr  Met  Gly  Leu  Thr  Gln  Phe  Glu  Leu  Thr
          115          120          125
Gly  His  Ser  Val  Phe  Asp  Phe  Thr  His  Pro  Cys  Asp  His  Glu  Glu  Met
          130          135          140

```

-continued

Arg	Glu	Met	Leu	Thr	His	Arg	Asn	Gly	Leu	Val	Lys	Lys	Gly	Lys	Glu
145					150					155					160
Gln	Asn	Thr	Gln	Arg	Ser	Phe	Phe	Leu	Arg	Met	Lys	Cys	Thr	Leu	Thr
			165						170					175	
Ser	Arg	Gly	Arg	Thr	Met	Asn	Ile	Lys	Ser	Ala	Thr	Trp	Lys	Val	Leu
		180						185					190		
His	Cys	Thr	Gly	His	Ile	His	Val	Tyr	Asp	Thr	Asn	Ser	Asn	Gln	Ser
		195					200					205			
Gln	Cys	Gly	Tyr	Lys	Lys	Pro	Pro	Met	Thr	Cys	Leu	Val	Leu	Ile	Cys
	210					215					220				
Glu	Pro	Ile	Pro	His	Pro	Ser	Asn	Ile	Glu	Ile	Pro	Leu	Asp	Ser	Lys
225					230					235					240
Thr	Phe	Leu	Ser	Arg	His	Ser	Leu	Asp	Met	Lys	Phe	Ser	Tyr	Cys	Asp
				245					250					255	
Glu	Arg	Ile	Thr	Glu	Leu	Met	Gly	Tyr	Glu	Pro	Glu	Glu	Leu	Leu	Gly
			260					265					270		
Arg	Ser	Ile	Tyr	Glu	Tyr	Tyr	His	Ala	Leu	Asp	Ser	Asp	His	Leu	Thr
		275					280					285			
Lys	Thr	His	His	Asp	Met	Phe	Thr	Lys	Gly	Gln	Val	Thr	Thr	Gly	Gln
	290					295					300				
Tyr	Arg	Met	Leu	Ala	Lys	Arg	Gly	Gly	Tyr	Val	Trp	Ile	Glu	Thr	Gln
305					310					315					320
Ala	Thr	Val	Ile	Tyr	Asn	Thr	Lys	Asn	Ser	Gln	Pro	Gln	Cys	Ile	Val
				325					330					335	
Cys	Val	Asn	Tyr	Val	Val	Ser	Gly	Ile	Ile	Gln	His	Asp	Leu	Ile	Phe
		340						345					350		
Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	Asp
		355					360					365			
Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr	Ser
	370					375					380				
Ser	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu	Leu
385					390				395						400
Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	Asn
			405						410					415	
Asp	Thr	Glu	Thr	Asp	Asp	Gln	Gln	Leu	Glu	Glu	Val	Pro	Leu	Tyr	Asn
			420					425					430		
Asp	Val	Met	Leu	Pro	Ser	Ser	Asn	Glu	Lys	Leu	Gln	Asn	Ile	Asn	Leu
		435					440					445			
Ala	Met	Ser	Pro	Leu	Pro	Ala	Ser	Glu	Thr	Pro	Lys	Pro	Leu	Arg	Ser
	450					455					460				
Ser	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Pro
465					470				475						480
Asn	Pro	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp
			485					490						495	
Gln	Pro	Ala	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu
		500						505					510		
Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Asp	Met	Val
	515						520					525			
Asn	Glu	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr
	530					535					540				

-continued

Glu Ala Lys Asn Pro Phe Ser Thr Gln Asp Thr Asp Leu Asp Leu Glu
 545 550 555 560
 Met Leu Ala Pro Tyr Ile Pro Met Asp Asp Asp Phe Gln Leu Arg Ser
 565 570 575
 Phe Asp Gln Leu Ser Pro Leu Glu Asn Ser Ser Thr Ser Pro Gln Ser
 580 585 590
 Ala Ser Thr Asn Thr Val Phe Gln Pro Thr Gln Met Gln Lys Pro Pro
 595 600 605
 Ile Ala Thr Val Thr Thr Thr Ala Thr Ser Asp Glu Leu Lys Thr Val
 610 615 620
 Thr Lys Asp Gly Met Glu Asp Ile Lys Ile Leu Ile Ala Phe Pro Ser
 625 630 635 640
 Pro Pro His Val Pro Lys Glu Pro Pro Cys Ala Thr Thr Ser Pro Tyr
 645 650 655
 Ser Asp Thr Gly Ser Arg Thr Ala Ser Pro Asn Arg Ala Gly Lys Gly
 660 665 670
 Val Ile Glu Gln Thr Glu Lys Ser His Pro Arg Ser Pro Asn Val Leu
 675 680 685
 Ser Val Ala Leu Ser Gln Arg Thr Thr Ala Pro Glu Glu Glu Leu Asn
 690 695 700
 Pro Lys Ile Leu Ala Leu Gln Asn Ala Gln Arg Lys Arg Lys Ile Glu
 705 710 715 720
 His Asp Gly Ser Leu Phe Gln Ala Val Gly Ile Gly Thr Leu Leu Gln
 725 730 735
 Gln Pro Asp Asp Arg Ala Thr Thr Thr Ser Leu Ser Trp Lys Arg Val
 740 745 750
 Lys Gly Cys Lys Ser Ser Glu Gln Asn Gly Met Glu Gln Lys Thr Ile
 755 760 765
 Ile Leu Ile Pro Ser Asp Leu Ala Cys Arg Leu Leu Gly Gln Ser Met
 770 775 780
 Asp Glu Ser Gly Leu Pro Gln Leu Thr Ser Tyr Asp Cys Glu Val Asn
 785 790 795 800
 Ala Pro Ile Gln Gly Ser Arg Asn Leu Leu Gln Gly Glu Glu Leu Leu
 805 810 815
 Arg Ala Leu Asp Gln Val Asn
 820

<210> SEQ ID NO 11

<211> LENGTH: 823

<212> TYPE: PRT

<213> ORGANISM: *Pantholops hodgsonii*

<400> SEQUENCE: 11

Met Glu Gly Ala Gly Gly Ala Asn Asp Lys Lys Lys Ile Ser Ser Glu
 1 5 10 15
 Arg Arg Lys Glu Lys Ser Arg Asp Ala Ala Arg Ser Arg Arg Ser Lys
 20 25 30
 Glu Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro His
 35 40 45
 Asn Val Ser Ser His Leu Asp Lys Ala Ser Val Met Arg Leu Thr Ile
 50 55 60
 Ser Tyr Leu Arg Val Arg Lys Leu Leu Asp Ala Gly Asp Leu Asp Ile
 65 70 75 80

-continued

Glu	Asp	Glu	Met	Lys	Ala	Gln	Met	Asn	Cys	Phe	Tyr	Leu	Lys	Ala	Leu
				85					90					95	
Asp	Gly	Phe	Val	Met	Val	Leu	Thr	Asp	Asp	Gly	Asp	Met	Ile	Tyr	Ile
			100					105					110		
Ser	Asp	Asn	Val	Asn	Lys	Tyr	Met	Gly	Leu	Thr	Gln	Phe	Glu	Leu	Thr
		115					120					125			
Gly	His	Ser	Val	Phe	Asp	Phe	Thr	His	Pro	Cys	Asp	His	Glu	Glu	Met
	130					135					140				
Arg	Glu	Met	Leu	Thr	His	Arg	Asn	Gly	Leu	Val	Lys	Lys	Gly	Lys	Glu
145					150					155					160
Gln	Asn	Thr	Gln	Arg	Ser	Phe	Phe	Leu	Arg	Met	Lys	Cys	Thr	Leu	Thr
				165					170						175
Ser	Arg	Gly	Arg	Thr	Met	Asn	Ile	Lys	Ser	Ala	Thr	Trp	Lys	Val	Leu
			180				185						190		
His	Cys	Thr	Gly	His	Ile	His	Val	Tyr	Asp	Thr	Asn	Ser	Asn	Gln	Ser
	195						200					205			
Gln	Cys	Gly	Tyr	Lys	Lys	Pro	Pro	Met	Thr	Cys	Leu	Val	Leu	Ile	Cys
	210					215					220				
Glu	Pro	Ile	Pro	His	Pro	Ser	Asn	Ile	Glu	Ile	Pro	Leu	Asp	Ser	Lys
225					230					235					240
Thr	Phe	Leu	Ser	Arg	His	Ser	Leu	Asp	Met	Lys	Phe	Ser	Tyr	Cys	Asp
				245					250					255	
Glu	Arg	Ile	Thr	Glu	Leu	Met	Gly	Tyr	Glu	Pro	Glu	Glu	Leu	Leu	Gly
			260					265					270		
Arg	Ser	Ile	Tyr	Glu	Tyr	Tyr	His	Ala	Leu	Asp	Ser	Asp	His	Leu	Thr
	275						280					285			
Lys	Thr	His	His	Asp	Met	Phe	Thr	Lys	Gly	Gln	Val	Thr	Thr	Gly	Gln
	290					295					300				
Tyr	Arg	Met	Leu	Ala	Lys	Arg	Gly	Gly	Tyr	Val	Trp	Ile	Glu	Thr	Gln
305					310					315					320
Ala	Thr	Val	Ile	Tyr	Asn	Thr	Lys	Asn	Ser	Gln	Pro	Gln	Cys	Ile	Val
				325					330					335	
Cys	Val	Asn	Tyr	Val	Val	Ser	Gly	Ile	Ile	Gln	His	Asp	Leu	Ile	Phe
			340					345					350		
Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	Asp
		355					360					365			
Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr	Ser
	370					375					380				
Ser	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu	Leu
385					390					395					400
Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	Asn
				405					410					415	
Asp	Ala	Glu	Thr	Asp	Asp	Gln	Gln	Leu	Glu	Glu	Val	Pro	Leu	Tyr	Asn
			420					425					430		
Asp	Val	Met	Leu	Pro	Ser	Ser	Asn	Glu	Lys	Leu	Gln	Asn	Ile	Asn	Leu
		435					440					445			
Ala	Met	Ser	Pro	Leu	Pro	Ala	Ser	Glu	Thr	Pro	Lys	Pro	Leu	Arg	Ser
	450					455					460				
Ser	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Pro
465					470					475					480

-continued

Asn	Pro	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp	485	490	495
Gln	Pro	Ala	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu	500	505	510
Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Asp	Met	Val	515	520	525
Asn	Glu	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr	530	535	540
Glu	Ala	Lys	Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu	Glu	545	550	555
Met	Leu	Ala	Pro	Tyr	Ile	Pro	Met	Asp	Asp	Asp	Phe	Gln	Leu	Arg	Ser	565	570	575
Phe	Asp	Gln	Leu	Ser	Pro	Leu	Glu	Asn	Ser	Ser	Thr	Ser	Pro	Gln	Ser	580	585	590
Ala	Ser	Thr	Asp	Thr	Val	Phe	Gln	Pro	Thr	Gln	Met	Gln	Glu	Pro	Pro	595	600	605
Ile	Ala	Thr	Ala	Thr	Thr	Thr	Ala	Thr	Asn	Asp	Glu	Leu	Lys	Thr	Val	610	615	620
Thr	Lys	Asp	Gly	Met	Glu	Asp	Ile	Lys	Ile	Leu	Ile	Ala	Phe	Pro	Ser	625	630	635
Pro	Pro	His	Val	Pro	Lys	Glu	Pro	Pro	Cys	Ala	Thr	Thr	Ser	Pro	Tyr	645	650	655
Ser	Asp	Thr	Gly	Ser	Arg	Thr	Ala	Ser	Pro	Asn	Arg	Ala	Gly	Lys	Gly	660	665	670
Val	Ile	Glu	Gln	Thr	Glu	Lys	Ser	His	Pro	Arg	Ser	Pro	Asn	Val	Leu	675	680	685
Ser	Val	Ala	Leu	Ser	Gln	Arg	Thr	Thr	Ala	Pro	Glu	Glu	Glu	Leu	Asn	690	695	700
Pro	Lys	Ile	Leu	Ala	Leu	Gln	Asn	Ala	Gln	Arg	Lys	Arg	Lys	Ile	Glu	705	710	715
His	Asp	Gly	Ser	Leu	Phe	Gln	Ala	Val	Gly	Ile	Gly	Thr	Leu	Leu	Gln	725	730	735
Gln	Pro	Asp	Asp	Arg	Ala	Thr	Thr	Thr	Ser	Leu	Ser	Trp	Lys	Arg	Val	740	745	750
Lys	Gly	Cys	Lys	Ser	Ser	Glu	Gln	Asn	Gly	Met	Glu	Gln	Lys	Thr	Ile	755	760	765
Ile	Leu	Ile	Pro	Ser	Asp	Leu	Ala	Cys	Arg	Leu	Leu	Gly	Gln	Ser	Met	770	775	780
Asp	Glu	Ser	Gly	Leu	Pro	Gln	Leu	Thr	Ser	Tyr	Asp	Cys	Glu	Val	Asn	785	790	795
Ala	Pro	Ile	Gln	Gly	Ser	Arg	Asn	Leu	Leu	Gln	Gly	Glu	Glu	Leu	Leu	805	810	815
Arg	Ala	Leu	Asp	Gln	Val	Asn										820		

<210> SEQ ID NO 12

<211> LENGTH: 823

<212> TYPE: PRT

<213> ORGANISM: Canis familiaris

<400> SEQUENCE: 12

Met	Glu	Gly	Ala	Gly	Gly	Ala	Asn	Asp	Lys	Lys	Lys	Ile	Ser	Ser	Glu	1	5	10	15
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	---	---	----	----

-continued

Arg	Arg	Lys	Glu	Lys	Ser	Arg	Asp	Ala	Ala	Arg	Ser	Arg	Arg	Ser	Lys	20	25	30	
Glu	Ser	Glu	Val	Phe	Tyr	Glu	Leu	Ala	His	Gln	Leu	Pro	Leu	Pro	His	35	40	45	
Asn	Val	Ser	Ser	His	Leu	Asp	Lys	Ala	Ser	Val	Met	Arg	Leu	Thr	Ile	50	55	60	
Ser	Tyr	Leu	Arg	Val	Arg	Lys	Leu	Leu	Asp	Ala	Gly	Asp	Leu	Asp	Ile	65	70	75	80
Glu	Asp	Glu	Met	Lys	Ala	Gln	Met	Asn	Cys	Phe	Tyr	Leu	Lys	Ala	Leu	85	90	95	
Asp	Gly	Phe	Val	Met	Val	Leu	Thr	Asp	Asp	Gly	Asp	Met	Ile	Tyr	Ile	100	105	110	
Ser	Asp	Asn	Val	Asn	Lys	Tyr	Met	Gly	Leu	Thr	Gln	Phe	Glu	Leu	Thr	115	120	125	
Gly	His	Ser	Val	Phe	Asp	Phe	Thr	His	Pro	Cys	Asp	His	Glu	Glu	Met	130	135	140	
Arg	Glu	Met	Leu	Thr	His	Arg	Asn	Gly	Leu	Val	Lys	Lys	Gly	Lys	Glu	145	150	155	160
Gln	Asn	Thr	Gln	Arg	Ser	Phe	Phe	Leu	Arg	Met	Lys	Cys	Thr	Leu	Thr	165	170	175	
Ser	Arg	Gly	Arg	Thr	Met	Asn	Ile	Lys	Ser	Ala	Thr	Trp	Lys	Val	Leu	180	185	190	
His	Cys	Thr	Gly	His	Ile	His	Val	Tyr	Asp	Thr	Asn	Ser	Asn	Gln	Ser	195	200	205	
Gln	Cys	Gly	Tyr	Lys	Lys	Pro	Pro	Met	Thr	Cys	Leu	Val	Leu	Ile	Cys	210	215	220	
Glu	Pro	Ile	Pro	His	Pro	Ser	Asn	Ile	Glu	Ile	Pro	Leu	Asp	Ser	Lys	225	230	235	240
Thr	Phe	Leu	Ser	Arg	His	Ser	Leu	Asp	Met	Lys	Phe	Ser	Tyr	Cys	Asp	245	250	255	
Glu	Arg	Ile	Thr	Glu	Leu	Met	Gly	Tyr	Glu	Pro	Glu	Glu	Leu	Leu	Gly	260	265	270	
Arg	Ser	Ile	Tyr	Glu	Tyr	Tyr	His	Ala	Leu	Asp	Ser	Asp	His	Leu	Thr	275	280	285	
Lys	Thr	His	His	Asp	Met	Phe	Thr	Lys	Gly	Gln	Val	Thr	Thr	Gly	Gln	290	295	300	
Tyr	Arg	Met	Leu	Ala	Lys	Arg	Gly	Gly	Tyr	Val	Trp	Val	Glu	Thr	Gln	305	310	315	320
Ala	Thr	Val	Ile	Tyr	Asn	Thr	Lys	Asn	Ser	Gln	Pro	Gln	Cys	Ile	Val	325	330	335	
Cys	Val	Asn	Tyr	Val	Val	Ser	Gly	Ile	Ile	Gln	His	Asp	Leu	Ile	Phe	340	345	350	
Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	Asp	355	360	365	
Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr	Ser	370	375	380	
Ser	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu	Leu	385	390	395	400
Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	Asn	405	410	415	

Asp	Thr	Glu	Thr	Asp	Asp	Gln	Gln	Leu	Glu	Glu	Val	Pro	Leu	Tyr	Asn	
			420					425					430			
Asp	Val	Met	Leu	Pro	Ser	Ser	Asn	Glu	Lys	Leu	Gln	Asn	Ile	Asn	Leu	
		435					440					445				
Ala	Met	Ser	Pro	Leu	Pro	Ala	Ser	Glu	Thr	Pro	Lys	Pro	Leu	Arg	Ser	
	450					455					460					
Ser	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Pro	
465					470					475					480	
Asn	Pro	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp	
				485					490					495		
Gln	Pro	Ala	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu	
			500					505					510			
Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Asp	Met	Val	
		515					520					525				
Asn	Glu	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr	
	530					535					540					
Glu	Ala	Lys	Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu	Glu	
545					550					555					560	
Met	Leu	Ala	Pro	Tyr	Ile	Pro	Met	Asp	Asp	Asp	Phe	Gln	Leu	Arg	Ser	
				565					570					575		
Phe	Asp	Gln	Leu	Ser	Pro	Leu	Glu	Ser	Asn	Ser	Thr	Ser	Pro	Gln	Ser	
			580					585					590			
Ala	Ser	Thr	Ile	Thr	Val	Phe	Gln	Pro	Thr	Pro	Met	Gln	Glu	Pro	Pro	
		595					600					605				
Leu	Thr	Thr	Thr	Ser	Thr	Thr	Ala	Thr	Thr	Asp	Glu	Leu	Lys	Thr	Val	
	610					615					620					
Thr	Lys	Asp	Gly	Ile	Glu	Asp	Ile	Lys	Ile	Leu	Ile	Ala	Ala	Pro	Ser	
625					630					635					640	
Pro	Thr	His	Val	Pro	Lys	Val	Thr	Thr	Ser	Ala	Thr	Thr	Ser	Pro	Tyr	
				645					650					655		
Ser	Asp	Thr	Gly	Ser	Arg	Thr	Ala	Ser	Pro	Asn	Arg	Ala	Gly	Lys	Gly	
			660					665					670			
Val	Ile	Glu	Gln	Thr	Glu	Lys	Ser	His	Pro	Arg	Ser	Pro	Asn	Val	Leu	
		675					680					685				
Ser	Val	Thr	Leu	Ser	Gln	Arg	Thr	Thr	Ile	Pro	Glu	Glu	Glu	Leu	Asn	
	690					695					700					
Pro	Lys	Ile	Leu	Ala	Leu	Gln	Asn	Ala	Gln	Arg	Lys	Arg	Lys	Ile	Glu	
705					710					715					720	
His	Asp	Gly	Ser	Leu	Phe	Gln	Ala	Val	Gly	Ile	Gly	Thr	Leu	Leu	Gln	
				725					730					735		
Gln	Pro	Asp	Asp	Arg	Ala	Thr	Thr	Thr	Ser	Leu	Ser	Trp	Lys	Arg	Val	
			740					745					750			
Lys	Gly	Cys	Lys	Ser	Ser											

-continued

820

<210> SEQ ID NO 13

<211> LENGTH: 819

<212> TYPE: PRT

<213> ORGANISM: Oryctolagus cuniculus

<400> SEQUENCE: 13

```

Met Glu Gly Ala Gly Gly Ala Asn Asp Lys Lys Lys Ile Ser Ser Glu
1      5      10      15

Arg Arg Lys Glu Lys Ser Arg Asp Ala Arg Ser Arg Arg Ser Lys
20     25     30

Glu Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro His
35     40     45

Asn Val Ser Ser His Leu Asp Lys Ala Ser Val Met Arg Leu Thr Ile
50     55     60

Ser Tyr Leu Arg Val Arg Lys Leu Leu Asp Ala Gly Asp Leu Asp Ile
65     70     75     80

Glu Asp Glu Met Lys Ala Gln Met Asn Cys Phe Tyr Leu Lys Ala Leu
85     90     95

Asp Gly Phe Val Met Val Leu Thr Asp Asp Gly Asp Met Ile Tyr Ile
100    105    110

Ser Asp Asn Val Asn Lys Tyr Met Gly Leu Thr Gln Phe Glu Leu Thr
115    120    125

Gly His Ser Val Phe Asp Phe Thr His Pro Cys Asp His Glu Glu Met
130    135    140

Arg Glu Met Leu Thr His Arg Asn Gly Pro Val Lys Lys Gly Lys Glu
145    150    155    160

Gln Asn Thr Gln Arg Ser Phe Phe Leu Arg Met Lys Cys Thr Leu Thr
165    170    175

Ser Arg Gly Arg Thr Met Asn Ile Lys Ser Ala Thr Trp Lys Val Leu
180    185    190

His Cys Thr Gly His Ile His Val Tyr Asp Thr Asn Ser Asn Gln Ser
195    200    205

Gln Cys Gly Tyr Lys Lys Pro Pro Met Thr Cys Leu Val Leu Ile Cys
210    215    220

Glu Pro Ile Pro His Pro Ser Asn Ile Glu Ile Pro Leu Asp Ser Lys
225    230    235    240

Thr Phe Leu Ser Arg His Ser Leu Asp Met Lys Phe Ser Tyr Cys Asp
245    250    255

Glu Arg Ile Thr Glu Leu Met Gly Tyr Glu Pro Glu Glu Leu Leu Gly
260    265    270

Arg Ser Ile Tyr Glu Tyr Tyr His Ala Leu Asp Ser Asp His Leu Thr
275    280    285

Lys Thr His His Asp Met Phe Thr Lys Gly Gln Val Thr Thr Gly Gln
290    295    300

Tyr Arg Met Leu Ala Lys Arg Gly Gly Tyr Val Trp Val Glu Thr Gln
305    310    315    320

Ala Thr Val Ile Tyr Asn Thr Lys Asn Ser Gln Pro Gln Cys Ile Val
325    330    335

Cys Val Asn Tyr Val Val Ser Gly Ile Ile Gln His Asp Leu Ile Phe
340    345    350

```

-continued

Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	Asp
	355						360					365			
Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Ala	Asp	Thr	Ser
	370					375					380				
Ser	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu	Leu
385					390					395					400
Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	Asn
				405					410					415	
Asp	Thr	Glu	Thr	Asp	Asp	Gln	Gln	Leu	Glu	Glu	Val	Pro	Leu	Tyr	Asn
				420				425					430		
Asp	Val	Met	Phe	Pro	Ser	Ser	Asn	Glu	Lys	Leu	Gln	Asp	Ile	Asn	Leu
		435					440					445			
Ala	Met	Ser	Pro	Leu	Pro	Ala	Ser	Glu	Ser	Pro	Lys	Pro	Leu	Arg	Ser
	450					455					460				
Ser	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Pro
465					470					475					480
Asn	Pro	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp
				485					490					495	
Gln	Pro	Ala	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu
			500					505					510		
Pro	Thr	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Asp	Met	Val
		515					520					525			
Asn	Val	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr
	530					535					540				
Glu	Ala	Lys	Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu	Glu
545					550					555					560
Met	Leu	Ala	Pro	Tyr	Ile	Pro	Met	Asp	Asp	Asp	Phe	Gln	Leu	Arg	Ser
			565						570					575	
Phe	Asp	Gln	Leu	Ser	Pro	Leu	Glu	Ser	Ser	Ser	Ala	Gly	Ala	Val	Thr
			580					585					590		
Val	Phe	Pro	Gln	Thr	Gln	Gly	Gln	Glu	Pro	Ala	Ala	Asn	Thr	Ala	Pro
		595					600					605			
Cys	Ala	Pro	Pro	Pro	Leu	Thr	Ile	Lys	Thr	Ser	Ala	Lys	Asp	Ser	Val
	610					615					620				
Glu	Asp	Met	Lys	Val	Leu	Ile	Ala	Ser	Pro	Ser	Pro	Asn	His	Ile	Pro
625					630					635					640
Lys	Glu	Thr	Ala	Ser	Ala	Thr	Thr	Ser	Pro	Pro	Tyr	Arg	Asp	Thr	Pro
			645						650					655	
Ser	Arg	Thr	Ala	Ser	Pro	Asn	Thr	Ala	Gly	Lys	Gly	Val	Ile	Glu	Gln
			660					665					670		
Thr	Glu	Lys	Ser	His	Pro	Arg	Ser	Pro	Asn	Val	Leu	Ala	Val	Thr	Leu
		675					680					685			
Ser	Gln	Arg	Thr	Thr	Ala	Pro	Glu	Glu	Glu	Leu	Asn	Pro	Lys	Ile	Leu
	690					695					700				
Ala	Leu	Gln	Asn	Ala	Gln	Arg	Lys	Arg	Lys	Met	Glu	Ser	Asp	Gly	Ser
705					710					715					720
Leu	Phe	Gln	Ala	Val	Gly	Ile	Gly	Thr	Leu	Leu	Gln	Gln	Pro	Asp	Asp
			725					730						735	
Arg	Thr	Thr	Thr	Ala	Ser	Leu	Ser	Trp	Lys	Arg	Val	Lys	Gly	Cys	Lys
			740					745					750		
Ser	Ser	Glu	Gln	Asn	Gly	Met	Glu	Gln	Lys	Thr	Ile	Ile	Leu	Ile	Pro

-continued

755					760					765					
Ser	Asp	Leu	Ala	Cys	Arg	Leu	Leu	Gly	Gln	Ser	Met	Asp	Glu	Ser	Gly
770						775					780				
Leu	Pro	Gln	Leu	Thr	Ser	Tyr	Asp	Cys	Glu	Val	Asn	Ala	Pro	Ile	Gln
785					790					795					800
Gly	Ser	Arg	Asn	Leu	Leu	Gln	Gly	Glu	Glu	Leu	Leu	Arg	Ala	Leu	Asp
			805					810						815	
Gln Val Asn															
<210> SEQ ID NO 14															
<211> LENGTH: 811															
<212> TYPE: PRT															
<213> ORGANISM: Gallus gallus															
<400> SEQUENCE: 14															
Met	Asp	Ser	Pro	Gly	Gly	Val	Thr	Asp	Lys	Lys	Arg	Ile	Ser	Ser	Glu
1				5					10				15		
Arg	Arg	Lys	Glu	Lys	Ser	Arg	Asp	Ala	Ala	Arg	Cys	Arg	Arg	Ser	Lys
		20						25					30		
Glu	Ser	Glu	Val	Phe	Tyr	Glu	Leu	Ala	His	Gln	Leu	Pro	Leu	Pro	His
		35				40						45			
Thr	Val	Ser	Ala	His	Leu	Asp	Lys	Ala	Ser	Ile	Met	Arg	Leu	Thr	Ile
	50				55					60					
Ser	Tyr	Leu	Arg	Met	Arg	Lys	Leu	Leu	Asp	Ala	Gly	Glu	Leu	Glu	Thr
65				70					75					80	
Glu	Ala	Asn	Met	Glu	Lys	Glu	Leu	Asn	Cys	Phe	Tyr	Leu	Lys	Ala	Leu
			85					90						95	
Asp	Gly	Phe	Val	Met	Val	Leu	Ser	Glu	Asp	Gly	Asp	Met	Ile	Tyr	Met
		100						105				110			
Ser	Glu	Asn	Val	Asn	Lys	Cys	Met	Gly	Leu	Thr	Gln	Phe	Asp	Leu	Thr
	115				120						125				
Gly	His	Ser	Val	Phe	Asp	Phe	Thr	His	Pro	Cys	Asp	His	Glu	Glu	Leu
	130				135						140				
Arg	Glu	Met	Leu	Thr	His	Arg	Asn	Gly	Pro	Val	Lys	Lys	Gly	Lys	Glu
145				150					155						160
Gln	Asn	Thr	Glu	Arg	Ser	Phe	Phe	Leu	Arg	Met	Lys	Cys	Thr	Leu	Thr
		165						170						175	
Ser	Arg	Gly	Arg	Thr	Val	Asn	Ile	Lys	Ser	Ala	Thr	Trp	Lys	Val	Leu
	180							185					190		
His	Cys	Thr	Gly	His	Ile	Arg	Val	Tyr	Asp	Thr	Cys	Asn	Asn	Gln	Thr
	195				200						205				
His	Cys	Gly	Tyr	Lys	Lys	Pro	Pro	Met	Thr	Cys	Leu	Val	Leu	Ile	Cys
	210				215					220					
Glu	Pro	Ile	Pro	His	Pro	Ser	Asn	Ile	Glu	Val	Pro	Leu	Asp	Ser	Lys
225				230					235					240	
Thr	Phe	Leu	Ser	Arg	His	Ser	Leu	Asp	Met	Lys	Phe	Ser	Tyr	Cys	Asp
		245						250					255		
Glu	Arg	Ile	Thr	Glu	Leu	Met	Gly	Tyr	Glu	Pro	Glu	Glu	Leu	Leu	Gly
	260						265						270		
Arg	Ser	Ile	Tyr	Glu	Tyr	Tyr	His	Ala	Leu	Asp	Ser	Asp	His	Leu	Thr
	275						280				285				
Lys Thr His His Asp Met Phe Thr Lys Gly Gln Val Thr Thr Gly Gln															

-continued

290					295					300					
Tyr 305	Arg	Met	Leu	Ala	Lys 310	Gln	Gly	Gly	Tyr	Val 315	Trp	Val	Glu	Thr	Gln 320
Ala	Thr	Val	Ile	Tyr 325	Asn	Thr	Lys	Asn	Ser 330	Gln	Pro	Gln	Cys	Ile	Val 335
Cys	Val	Asn	Tyr 340	Val	Leu	Ser	Gly	Ile	Val 345	Gln	Lys	Asp	Leu	Ile	Phe 350
Ser	Leu	Gly 355	Gln	Thr	Glu	Cys	Met 360	Leu	Lys	Pro	Val	Glu 365	Ser	Pro	Glu
Met 370	Lys	Met	Thr	Lys	Ile 375	Phe	Ser	Lys	Asp	Asp 380	Trp	Asp	Asp	Thr	Asn
Ser 385	Leu	Phe	Glu	Lys 390	Leu	Lys	Gln	Glu	Pro	Asp 395	Ala	Leu	Thr	Val	Leu 400
Ala	Pro	Ala	Ala	Gly 405	Asp	Thr	Ile	Ile	Ser 410	Leu	Asp	Phe	Ser	Ser	Asn 415
Glu	Ser	Asp	Glu 420	Gln	Gln	Cys	Asp 425	Glu	Val	Pro	Leu	Tyr	Asn 430	Asp	Val
Met	Leu	Pro 435	Ser	Ser	Ser	Glu	Lys 440	Leu	Gln	Asn	Ile	Asn 445	Ile	Ala	Met
Ser	Pro	Leu 450	Pro	Ala	Ser	Glu 455	Thr	Thr	Lys	Pro	Leu 460	Arg	Ser	Asn	Ala
Asp 465	Pro	Ala	Leu	Asn 470	Arg	Glu	Val	Val	Ser	Lys 475	Leu	Glu	Pro	Asn	Thr 480
Glu	Thr	Leu	Glu 485	Leu	Ser	Phe	Thr	Met	Pro 490	Gln	Val	Gln	Glu 495	Gln	Pro
Thr	Ser	Pro 500	Ser	Asp	Ala	Ser	Thr	Ser	Gln 505	Ser	Ser	Pro	Glu 510	Pro	Ser
Ser	Pro	Asn 515	Asp	Tyr	Cys	Phe	Asp 520	Val	Asp	Asn	Asp	Met 525	Ala	Asn	Glu
Phe 530	Lys	Leu	Glu	Leu	Val	Glu 535	Lys	Leu	Phe	Ala	Ile 540	Asp	Thr	Glu	Ala
Lys 545	Asn	Pro	Phe	Ser 550	Thr	Gln	Glu	Thr	Asp	Leu 555	Asp	Leu	Glu	Met	Leu 560
Ala	Pro	Tyr	Ile 565	Pro	Met	Asp	Asp	Asp	Phe 570	Gln	Leu	Arg	Ser	Phe	Asp 575
Gln	Leu	Ser	Pro 580	Leu	Glu	Ser	Ser	Ser	Ser 585	Gly	Ser	Gln	Asn 590	Ala	Ala
Thr	Ile 595	Thr	Ile	Leu	Gln	Gln	Thr 600	Gln	Thr	Pro	Ser	Thr 605	Ala	Ala	Asp
Glu 610	Ile	Lys	Pro	Val	Ala	Glu 615	Arg	Val	Asp	Asp	Val 620	Lys	Ala	Leu	Ile
Val 625	Pro	Ser	Ser	Pro 630	Val	His	Val	Ile	Asn	Asp 635	Thr	Ser	Ser	Ala	Pro 640
Ala	Ser	Pro	Tyr 645	Ser	Gly	Asn	Arg	Ser	Arg 650	Thr	Ala	Ser	Pro	Ile	Arg 655
Ala	Gly	Lys	Gly 660	Thr	Leu	Glu	Gln	Thr	Glu 665	Lys	Ser	Cys	Pro 670	Gly	Ala
Pro	Ser	Leu 675	Ile	Thr	Val	Thr 680	Leu	Asn	Lys	Arg	Ser	Thr 685	Ala	Met	Asp
Glu 690	Glu	Leu	Asn	Pro	Lys 695	Met	Leu	Ala	Leu	His 700	Asn	Ala	Gln	Arg	Lys

-continued

Arg Lys Met Glu His Asp Gly Ser Leu Phe Gln Ala Val Gly Ile Gly
 705 710 715 720
 Ser Leu Phe Gln Gln Thr Gly Asp Arg Gly Gly Asn Ala Ser Leu Ala
 725 730 735
 Trp Lys Arg Val Lys Ala Cys Lys Thr Asn Gly His Asn Gly Val Glu
 740 745 750
 Gln Lys Thr Ile Ile Leu Leu Ser Thr Asp Ile Ala Ser Lys Leu Leu
 755 760 765
 Gly Gln Ser Met Asp Glu Ser Gly Leu Pro Gln Leu Thr Ser Tyr Asp
 770 775 780
 Cys Glu Val Asn Ala Pro Ile Gln Gly Asn Arg Asn Leu Leu Gln Gly
 785 790 795 800
 Glu Glu Leu Leu Arg Ala Leu Asp Gln Val Asn
 805 810

<210> SEQ ID NO 15

<211> LENGTH: 533

<212> TYPE: PRT

<213> ORGANISM: Danio rerio

<400> SEQUENCE: 15

Met Asp Thr Gly Val Val Thr Glu Lys Lys Arg Val Ser Ser Glu Arg
 1 5 10 15
 Arg Lys Gly Lys Ser Arg Asp Ala Ala Arg Ser Arg Arg Gly Lys Glu
 20 25 30
 Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro His Asn
 35 40 45
 Val Thr Ser His Leu Asp Lys Ala Ser Ile Met Arg Leu Thr Ile Ser
 50 55 60
 Tyr Leu Arg Met Arg Lys Leu Leu Asn Ser Asp Glu Lys Glu Glu Lys
 65 70 75 80
 Glu Glu Asn Glu Leu Glu Ser Gln Leu Asn Gly Phe Tyr Leu Lys Ala
 85 90 95
 Leu Glu Gly Phe Leu Met Val Leu Ser Glu Asp Gly Asp Met Val Tyr
 100 105 110
 Leu Ser Glu Asn Val Ser Lys Ser Met Gly Leu Thr Gln Phe Asp Leu
 115 120 125
 Thr Gly His Ser Ile Phe Glu Phe Ser His Pro Cys Asp His Glu Glu
 130 135 140
 Leu Arg Glu Met Leu Val His Arg Thr Gly Ser Lys Lys Thr Lys Glu
 145 150 155 160
 Gln Asn Thr Glu Arg Ser Phe Phe Leu Arg Met Lys Cys Thr Leu Thr
 165 170 175
 Ser Arg Gly Arg Thr Val Asn Ile Lys Ser Ala Thr Trp Lys Val Leu
 180 185 190
 His Cys Ala Gly His Val Arg Val His Glu Gly Ser Glu Ala Ser Glu
 195 200 205
 Asp Ser Gly Phe Lys Glu Pro Pro Val Thr Tyr Leu Val Leu Ile Cys
 210 215 220
 Glu Pro Ile Pro His Pro Ser Asn Ile Glu Val Pro Leu Asp Ser Lys
 225 230 235 240
 Thr Phe Leu Ser Arg His Thr Leu Asp Met Lys Phe Ser Tyr Cys Asp

```

<210> SEQ ID NO 16
<211> LENGTH: 805
<212> TYPE: PRT
<213> ORGANISM: Xenopus laevis

<400> SEQUENCE: 16

Met Glu Gly Ser Val Val Val Ser Glu Lys Lys Arg Ile Ser Ser Glu
1          5          10          15
Arg Arg Lys Glu Lys Ser Arg Asp Ala Ala Arg Cys Arg Arg Ser Asn
          20          25          30
Glu Ser Glu Val Phe Tyr Glu Leu Ser His Glu Leu Pro Leu Pro His
          35          40          45
Asn Val Ser Ser His Leu Asp Lys Ala Ser Ile Met Arg Leu Asp His
          50          55          60

```

-continued

Gln	Leu	Pro	Ala	Val	Glu	Lys	Val	Ala	Asp	Ala	Gly	Asp	Leu	Asp	Gly	65	70	75	80
Glu	Thr	Glu	Leu	Asp	Lys	Gln	Leu	Asn	Cys	Phe	Tyr	Leu	Lys	Ala	Leu	85	90	95	
Glu	Gly	Phe	Val	Leu	Val	Leu	Thr	Glu	Glu	Gly	Asp	Met	Ile	Tyr	Leu	100	105	110	
Ser	Glu	Asn	Val	Asn	Lys	Cys	Met	Gly	Leu	Thr	Gln	Phe	Glu	Leu	Thr	115	120	125	
Gly	His	Ser	Val	Phe	Asp	Phe	Thr	His	Pro	Cys	Asp	His	Glu	Glu	Leu	130	135	140	
Arg	Glu	Met	Leu	Thr	Phe	Arg	Asn	Gly	Pro	Ala	Lys	Lys	Arg	Lys	Arg	145	150	155	160
Thr	Asn	His	Arg	Glu	Lys	Phe	Leu	Pro	Ser	Tyr	Glu	Met	Tyr	Ile	Asn	165	170	175	
Gln	Ser	Trp	Lys	Asn	Arg	Glu	Tyr	Lys	Val	Ser	His	Met	Glu	Gly	Pro	180	185	190	
Ser	Leu	Tyr	Arg	Thr	His	Ala	Cys	Ile	Tyr	Asp	Asn	Ala	Asn	Asn	Gln	195	200	205	
Asn	His	Cys	Gly	Tyr	Lys	Lys	Pro	Pro	Met	Thr	Cys	Met	Val	Val	Ile	210	215	220	
Cys	Glu	Pro	Ile	Pro	His	Pro	Ser	Asn	Ile	Glu	Phe	Pro	Leu	Asp	Ser	225	230	235	240
Lys	Thr	Phe	Leu	Ser	Arg	His	Ser	Leu	Asp	Met	Lys	Phe	Ser	Tyr	Cys	245	250	255	
Asp	Glu	Arg	Val	Thr	Glu	Leu	Val	Gly	Tyr	Glu	Pro	Asp	Glu	Leu	Leu	260	265	270	
Gly	Arg	Ser	Val	Tyr	Glu	Tyr	Tyr	His	Ala	Leu	Asp	Ser	Asp	His	Leu	275	280	285	
Thr	Lys	Pro	Asn	Tyr	Asn	Met	Phe	Thr	Lys	Gly	Gln	Val	Thr	Thr	Gly	290	295	300	
Gln	Tyr	Arg	Met	Leu	Ala	Lys	Lys	Gly	Gly	Tyr	Val	Trp	Val	Glu	Thr	305	310	315	320
Gln	Ala	Thr	Val	Ile	Tyr	Asn	Ser	Lys	Asn	Ser	Gln	Pro	Gln	Cys	Ile	325	330	335	
Val	Cys	Val	Asn	Tyr	Val	Leu	Ser	Glu	Val	Val	Glu	Lys	Asp	Leu	Ile	340	345	350	
Leu	Ser	Leu	Gly	Gln	Thr	Ala	Ser	Val	Leu	Ile	Pro	Val	Glu	Ser	Gln	355	360	365	
Glu	Ile	Lys	Met	Pro	Glu	Ile	Phe	Thr	Glu	Leu	Asn	Glu	Glu	Asn	Asn	370	375	380	
Ser	Glu	Cys	Leu	Phe	Asp	Lys	Leu	Lys	Gln	Glu	Pro	Glu	Ser	Leu	Thr	385	390	395	400
Val	Leu	Ala	Pro	Asp	Ala	Gly	Asp	Glu	Ile	Ile	Pro	Leu	Asp	Phe	Ser	405	410	415	
Ser	Gly	Asp	Ser	Asp	Lys	Pro	Tyr	Glu	Asp	Val	Pro	Leu	Tyr	Asn	Asp	420	425	430	
Val	Met	Leu	His	Ser	Thr	Ser	Asn	Lys	Leu	Glu	Ser	Thr	Pro	Ile	Thr	435	440	445	
Pro	Leu	Pro	Ala	Pro	Glu	Met	Pro	Lys	Pro	Leu	Arg	Ser	Asn	Val	Asp	450	455	460	
Pro	Ala	Leu	Asn	Arg	Glu	Val	Val	Ile	Lys	Met	Glu	Ser	Asn	Pro	Arg				

-continued

465	470	475	480
Thr Thr Cys Ala Ser	Ile His His Ser	Thr Ala Ile Gln Ala Arg	Gln
	485	490	495
Pro Phe Arg Tyr Gln Phe Gln Ser	Glu Pro Ser Thr	Glu Pro Asn Thr	
	500	505	510
Pro Glu Tyr Cys Phe Asp Val Asp	Ser Glu Met Ala Ser	Glu Phe Lys	
	515	520	525
Leu Asp Leu Val Glu Lys Leu Phe Ala Ile Asp	Thr Glu Ala Lys Ala		
	530	535	540
Pro Phe Tyr Tyr Pro Gly Asn Asp Leu Asp Leu Glu Met Leu Ala Pro			
545	550	555	560
Tyr Ile Pro Met Asp Asp Asp Phe Gln Leu Arg Thr Phe Asp Gln Leu			
	565	570	575
Ser Ser Leu Glu Cys Asp Ser Ser Ile Pro Gln Thr Leu Gly Ser Met			
	580	585	590
Thr Thr Leu Phe His Gln Ser Leu Ser Pro Ser Thr Ser Asp Phe Lys			
	595	600	605
Pro Glu Asp Ala Met Ser Asp Leu Lys Thr Ile Ile Gln Ser Pro Val			
610	615	620	
His Met Met Lys Glu Ser Thr Ser Ala Pro Val Ser Pro Tyr Asn Gly			
625	630	635	640
Asn Arg Ser Arg Thr Ser Ser Pro Val Arg Pro Ala Lys Ala Val Val			
	645	650	655
Asp Lys Thr Glu Lys Ser Arg Pro Gly Thr Pro Asn Leu Pro Val Pro			
	660	665	670
Leu Asn Lys Arg Cys Thr Ile Leu Asp Glu Glu Leu Asn Pro Lys Met			
	675	680	685
Ile Cys Phe Thr Gln Cys Thr Ala Glu Lys Arg Lys Met Glu Ser Asp			
690	695	700	
Gly Pro Leu Phe Gln Ala Ile Gly Ile Gly Thr Leu Phe Gln Thr Asn			
705	710	715	720
Val Asp Pro Gly Pro Asn Ser Ser Leu Gln Trp Lys Arg Val Lys Gly			
	725	730	735
Ser Asp Ser Glu Arg Leu Ser Ser Ala Glu Gln Arg Thr Ile Leu Leu			
	740	745	750
Leu Ser Thr Asp Met Ala Ser Gln Leu Leu Gly Gln Ser Phe Asp Gly			
	755	760	765
Thr Val Leu Pro Gln Leu Thr Gly Tyr Asp Cys Glu Val Asn Ala Pro			
770	775	780	
Val His Gly Thr Arg Asn Leu Leu Gln Gly Glu Glu Leu Leu Arg Ala			
785	790	795	800
Leu Asp Gln Ala Asn			
	805		

<210> SEQ ID NO 17

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: siRNA directed against mouse VHL factor, including 3' AT

<400> SEQUENCE: 17

-continued

aacatcacat tgccagtgtat

21

<210> SEQ ID NO 18

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Tyr	Val	Asp	Ser	Asp	Met
1				5					10				15	

<210> SEQ ID NO 19

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 19

Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Asp	Met
1				5					10				15	

<210> SEQ ID NO 20

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Carassius carassius

<400> SEQUENCE: 20

Pro	Asn	Ser	Pro	Met	Glu	Tyr	Cys	Phe	Gln	Val	Asp	Ser	Asp	Ile
1				5					10				15	

<210> SEQ ID NO 21

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Xenopus laevis

<400> SEQUENCE: 21

Glu	Pro	Asn	Thr	Pro	Glu	Tyr	Cys	Phe	Asp	Val	Asp	Ser	Glu	Met
1				5					10				15	

What is claimed is:

1. A method for increasing a sensitivity of a tumor in a subject to a treatment, the method comprising administering to the tumor a composition comprising an effective amount of an inhibitor of nitric oxide synthase, a nitric oxide scavenger, an inhibitor of HIF-1 nitrosylation, or a combination thereof, wherein:

- (i) the tumor is resistant to radiation therapy, chemotherapy, or both radiation therapy and chemotherapy; and
- (ii) the administering increases the sensitivity of the tumor to the radiotherapy, the chemotherapy, or both the radiotherapy and the chemotherapy.

2. The method of claim 1, wherein the inhibitor of nitric oxide synthase is selected from the group consisting of L-N(6)-(1-iminoethyl)lysine tetrazole-amide (SC-51); aminoguanidine (AG); guanidinoethyldisulfide; L-NG-nitroarginine methyl ester; mercaptoethylguanidine (MEG); N^ω-nitro-L-arginine methyl ester (L-NAME); N-(3-(aminomethyl)benzyl)acetamidine (1400W); N^G-monomethyl-L-arginine (L-NMMA); 7-nitroindazole (7-NI).

3. The method of claim 1, wherein the nitric oxide scavenger is selected from the group consisting of hydroxo-

cobalamin; 2-(4-carboxyphenyl)-4,4,5,5-tetramethylimidazole-1-oxyl-3-oxide (carboxy-PTIO); diethyldithiocarbamate; AMD6221 (ruthenium [hydrogen(diethylenetrinitrilo) pentaacetato]chloride); and N-dithiocarboxy-sarcosine (DTCS).

4. The method of claim 1, wherein the administering comprises administering a minimally therapeutic dose of an inhibitor of inducible nitric oxide synthase (iNOS).

5. The method of claim 1, wherein the subject is a mammal.

6. The method of claim 5, wherein the mammal is a human.

7. The method of claim 6, wherein the composition inhibits nitrosylation of Cys520 of SEQ ID NO: 6.

8. A method for delaying tumor growth in a subject, the method comprising:

- (a) administering to the subject a composition comprising an effective amount of an inhibitor of nitric oxide synthase, a nitric oxide scavenger, an inhibitor of HIF-1 nitrosylation, or a combination thereof; and
- (b) treating a tumor that is resistant to radiation therapy, chemotherapy, or both radiation therapy and chemo-

therapy, with radiation therapy, chemotherapy, or both radiation therapy and chemotherapy,

whereby tumor growth in the subject is delayed.

9. The method of claim 8, wherein the subject is a mammal.

10. The method of claim 9, wherein the mammal is a human.

11. The method of claim 10, wherein the composition inhibits nitrosylation of Cys520 of SEQ ID NO: 6.

12. The method of claim 8, wherein the treating comprises treating the tumor with a subtherapeutic dose of ionizing radiation.

13. The method of claim 8, wherein the treating comprises administering to the subject a therapeutically effective amount of cyclophosphamide.

14. The method of claim 8, further comprising promoting tumor regression.

15. A method for inhibiting tumor blood vessel growth in a subject, the method comprising:

(a) administering to the subject a composition comprising an effective amount of an inhibitor of nitric oxide synthase, a nitric oxide scavenger, an inhibitor of HIF-1 nitrosylation, or a combination thereof; and

(b) treating a tumor that is resistant to radiation therapy, chemotherapy, or both radiation therapy and chemotherapy, with radiation therapy, chemotherapy, or both radiation therapy and chemotherapy,

whereby tumor blood vessel growth is inhibited.

16. The method of claim 15, wherein the subject is a mammal.

17. The method of claim 16, wherein the mammal is a human.

18. The method of claim 17, wherein the composition inhibits nitrosylation of Cys520 of SEQ ID NO: 6.

19. The method of claim 15, further comprising delaying tumor growth in the subject.

20. The method of claim 15, further comprising promoting tumor regression in the subject.

21. A method for inhibiting HIF-1 activity in a cell, the method comprising contacting the cell with a composition comprising an effective amount of an inhibitor of nitric oxide synthase, a nitric oxide scavenger, an inhibitor of HIF-1 nitrosylation, or a combination thereof, whereby HIF-1 activity in the cell is inhibited.

22. The method of claim 21, wherein the cell is a tumor cell.

23. The method of claim 21, wherein the tumor cell is present in a subject.

24. The method of claim 21, wherein the subject is a mammal.

25. The method of claim 21, wherein the mammal is a human.

26. The method of claim 21, wherein the composition inhibits nitrosylation of Cys520 of SEQ ID NO: 6.

27. The method of claim 21, further comprising exposing the tumor cell to a treatment selected from the group consisting of radiation therapy, chemotherapy, and combinations thereof.

28. A method for inhibiting an inflammatory response in a cell, the method comprising contacting the cell with a composition comprising an effective amount of an inhibitor of nitric oxide synthase, a nitric oxide scavenger, an inhibi-

tor of HIF-1 nitrosylation, or a combination thereof, whereby an inflammatory response in the cell is inhibited.

29. The method of claim 28, wherein the cell is present in a subject.

30. The method of claim 28, wherein the subject is a mammal.

31. The method of claim 30, wherein the mammal is a human.

32. The method of claim 30, wherein the agent inhibits nitrosylation of Cys520 of SEQ ID NO: 6.

33. The method of any of claims 1, 8, 15, 22, and 28, wherein the composition is provided to the subject in an implantable device.

34. A method for identifying an inhibitor of nitrosylation of an HIF-1 polypeptide, the method comprising:

(a) providing a cell comprising a nucleic acid a nucleotide sequence comprising any of SEQ ID NOs: 18-21;

(b) contacting the cell with a compound comprising a potential inhibitor of nitrosylation of an HIF-1 polypeptide; and

(c) assaying nitrosylation of a cysteine residue present in the nucleic acid,

whereby an inhibitor of nitrosylation of an HIF-1 polypeptide is identified.

35. The method of claim 34, wherein the cell is present in an animal.

36. The method of claim 35, wherein the animal is a mammal.

37. The method of claim 36, wherein the mammal is a human.

38. The method of claim 34, wherein the nucleic acid comprises an expression vector in which the nucleic acid is operably linked to a promoter that is active in the cell.

39. The method of claim 38, wherein the expression vector is a transgene and the animal is a transgenic animal that expresses the nucleic acid.

40. The method of claim 39, wherein the compound is administered to the transgenic animal via a route that results in the compound contacting the cell.

41. The method of claim 35, further comprising comparing a level of nitrosylation of the cysteine residue present in the nucleic acid to a level of nitrosylation of the cysteine residue present in the nucleic acid prior to the contacting step.

42. The method of claim 35, wherein the cell is an in vitro cultured cell and the contacting is performed in vitro.

43. An expression construct comprising one or more of SEQ ID NOs: 18-21 operably linked to a promoter.

44. An expression construct comprising one or more of SEQ ID NOs: 18-21 operably linked to a promoter, with the proviso that all cysteine residues present within SEQ ID NOs: 18-21 have been replaced with a non-nitrosylatable amino acid.

45. The expression construct of claim 44, wherein the non-nitrosylatable amino acid is serine.

46. A host cell comprising the expression construct of claim 43.

47. A host cell comprising the expression construct of claim 44.

48. A transgenic, non-human animal comprising the expression construct of claim 43.

49. A transgenic, non-human animal comprising the expression construct of claim 44.

50. The use of inhibitors of nitric oxide synthases to prevent activation of HIF-1 activity in tumors by cancer therapy that include radiation and chemotherapy.

51. The use of nitric oxide scavengers to prevent activation of HIF-1 activity in tumors by cancer therapy that include radiation and chemotherapy.

52. The use of agents that can reduce the production of NO to sensitize tumors to radiotherapy and/or chemotherapy.

53. The use of nitric oxide synthase or nitric oxide scavengers to inhibit inflammatory reaction through the inhibition of HIF-1 activation.

54. The use of agents that can block the nitrosylation of HIF-1 α cysteine 520 for the purpose of enhancing cancer therapy.

55. The use of agents that can block the nitrosylation of HIF-1 α cysteine 520 for the purpose of inhibiting/attenuating inflammatory response.

* * * * *