

Arctic Sea Ice in CMIP6

SIMIP Community*

(Detailed listing according to AGU guidelines for community papers is in the appendix. Repeated here for reference: Dirk Notz, Jakob Dörr, David A Bailey, Ed Blockley, Mitchell Bushuk, Jens Boldingh Debernard, Evelien Dekker, Patricia DeRepentigny, David Docquier, Neven S. Fučkar, John C. Fyfe, Alexandra Jahn, Marika Holland, Elizabeth Hunke, Doroteaciro Iovino, Narges Khosravi, François Massonnet, Gurvan Madec, Siobhan O'Farrell, Alek Petty, Arun Rana, Lettie Roach, Erica Rosenblum, Clement Rousset, Tido Semmler, Julianne Stroeve, Bruno Tremblay, Takahiro Toyoda, Hiroyuki Tsujino, Martin Vancoppenolle)

Key Points:

- CMIP6 model simulations of Arctic sea-ice area capture the observational record in the multi-model ensemble spread
- The sensitivity of Arctic sea ice to changes in the forcing is better captured by CMIP6 models than by CMIP5 and CMIP3 models
- The majority of available CMIP6 simulations lose most September sea ice for the first time before 2050 in all scenarios

*See appendix for detailed listing

Corresponding author: Dirk Notz, dirk.notz@uni-hamburg.de

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1029/2019GL086749

Abstract

We examine CMIP6 simulations of Arctic sea-ice area and volume. We find that CMIP6 models produce a wide spread of mean Arctic sea-ice area, capturing the observational estimate within the multi-model ensemble spread. The CMIP6 multi-model ensemble mean provides a more realistic estimate of the sensitivity of September Arctic sea-ice area to a given amount of anthropogenic CO₂ emissions and to a given amount of global warming, compared with earlier CMIP experiments. Still, most CMIP6 models fail to simulate at the same time a plausible evolution of sea-ice area and of global mean surface temperature. In the vast majority of the available CMIP6 simulations, the Arctic Ocean becomes practically sea-ice free (sea-ice area < 1 million km²) in September for the first time before the year 2050 in each of the four emission scenarios SSP1-1.9, SSP1-2.6, SSP2-4.5 and SSP5-8.5 examined here.

Plain Language Summary

We examine simulations of Arctic sea ice from the latest generation of global climate models. We find that the observed evolution of Arctic sea-ice area lies within the spread of model simulations. In particular, the latest generation of models performs better than models from previous generations at simulating the sea-ice loss for a given amount of CO₂ emissions and for a given amount of global warming. In most simulations, the Arctic Ocean becomes practically sea-ice free (sea-ice area < 1 million km²) in September for the first time before the year 2050.

1 Introduction

In recent decades, Arctic sea-ice area has decreased rapidly, and the signal of a forced sea-ice retreat has clearly emerged from the background noise of year-to-year variability. Because of this, the ability of climate models to plausibly simulate the observed changes in Arctic sea-ice coverage has become a central measure of model performance in Arctic-focused climate model intercomparisons (e.g., Koenig et al., 2014; Massonnet et al., 2012; Melia et al., 2015; Olonscheck & Notz, 2017; Shu et al., 2015; Stroeve et al., 2007, 2012, 2014). In this contribution, we extend these earlier studies that examined model performance in the third and fifth phases of the Coupled Model Intercomparison Project (CMIP3 and CMIP5) by examining model simulations from the sixth phase of the Coupled Model Intercomparison Project (CMIP6, Eyring et al., 2015). For CMIP6, the Sea-Ice Model

Intercomparison Project (SIMIP, Notz et al., 2016) designed a specific set of diagnostics that allow for detailed analyses of sea-ice related processes and thus a process-based evaluation of sea-ice simulations of the participating models. To lay the foundation for such analyses, we here provide an initial overview of CMIP6 model performance by examining some large-scale, pan-Arctic metrics of model performance and future sea-ice evolution, including a comparison to CMIP5 and CMIP3 simulations. A similar analysis for Antarctic sea ice is given by (Roach et al., under review).

2 Analysis Method

In this contribution, we examine two large-scale integrated quantities that describe the time evolution of Arctic sea ice. These are the Northern Hemisphere total sea-ice area (SIA) and total sea-ice volume (SIV), which can be calculated readily from SIMIP variables as follows.

To obtain sea-ice area for CMIP6 model simulations, we use the SIMIP variable of Northern Hemisphere sea-ice area `siarean` when provided. If `siarean` is not provided, we calculate the sea-ice area by multiplying sea-ice concentration on the ocean grid (`siconc`, preferred) or on the atmospheric grid (`siconca`) with individual grid-cell area and then sum over the Northern Hemisphere. Note that we use sea-ice area as our primary variable to describe sea-ice coverage instead of sea-ice extent, which is usually calculated as the total area of all grid cells with at least 15% sea-ice concentration. Our choice to focus on sea-ice area derives primarily from the fact that sea-ice extent is a strongly grid-dependent, non-linear quantity, making it difficult to meaningfully compare between model output and satellite observations (compare Notz, 2014). In addition, the observational spread across different satellite products is smaller for trends in sea-ice area than it is for trends in sea-ice extent (Comiso et al., 2017).

To calculate sea-ice volume for CMIP6 models, we (1) directly use the SIMIP variable of Northern Hemisphere sea-ice volume `sivoln` when provided, or (2) multiply the sea-ice volume per grid-cell area `sivol` by individual grid-cell area and sum over the Northern Hemisphere, or (3) multiply sea-ice-concentration `siconc`, sea-ice thickness `sithick` and individual grid-cell area and then sum over the Northern Hemisphere. For CMIP5, only the sea-ice volume per grid-cell area (also called “equivalent sea-ice thickness”, `sit`) is available, so we use method (2) for all CMIP5 models. We were unable to obtain sea-

ice volume data for CMIP3 models, so volume comparisons in the following are limited to CMIP5 and CMIP6 model simulations.

To meaningfully estimate model performance relative to the real evolution of the sea-ice cover in the Arctic, we must take internal variability into account (see, for example, England et al., 2019; Kay et al., 2011; Notz, 2015; Olonscheck & Notz, 2017; Swart et al., 2015). Internal variability describes the spread in plausible climate trajectories in response to a given forcing scenario, owing to the chaotic nature of our climate system. The observational record is just one such plausible trajectory, and no single model simulation can ever be expected to perfectly agree with it because of its chaotic nature. Therefore, most CMIP6 models have been run several times with slightly different initial conditions to estimate the range of trajectories that are compatible with a given model's physics. In the following, we take two different approaches to examine whether a given model provides a plausible simulation of the observational record in light of internal variability.

First, for CMIP6 models, we estimate a best-guess CMIP6-average internal variability σ_{cmip6} by averaging across the individual ensemble spread of those models that provide three or more ensemble members (see Table S3 for details). In calculating the standard deviation, we correct for small sample size n by using Bessel's correction and then dividing the resulting standard deviation by the scale mean of the chi distribution with $n-1$ degrees of freedom. We then define all simulations that lie within the range of $2\sigma = \pm 2\sqrt{\sigma_{cmip6}^2 + \sigma_{obs}^2}$ around the observational estimate as plausible simulations (compare Olonscheck & Notz, 2017). Here, σ_{obs}^2 refers to the observational uncertainty explained below. This approach allows us to also examine the plausibility of those models that only provide a single ensemble member. In addition to considering internal variability explicitly, we reduce its impact by examining model performance relative to a time average over several years. We take the first twenty years of the satellite record (1979–1998) for comparing mean values, as those twenty years provide a compromise between using as many years as possible and using a period with no strong trend in Arctic sea-ice area and volume. However, even on multi-decadal time scales internal variability affects the Arctic sea-ice cover, so averaging over 20 years is not long enough an averaging period to remove the impact of internal variability entirely. To compare trends, we examine the overlap period 1979–2014 of the satellite record, which begins in 1979, and the historical period of CMIP6, which ends in 2014.

Second, in order to select a subset of models for estimating a best guess of the future evolution of the Arctic sea-ice cover, we take the more strict approach to define a model as plausible if its ensemble spread includes the observational record, considering observational uncertainty. These models are referred to as “selected models” hereafter.

To obtain an observational estimate of sea-ice area, we use observational records of sea-ice concentration from the OSI SAF (Lavergne et al., 2019), NASA-Team (Cavalieri et al., 1997) and Bootstrap (Comiso et al., 1997) algorithms. Sea-ice area is then calculated by multiplying the sea-ice concentration with individual grid-cell area and summing over the Northern Hemisphere. For the NASA-Team and Bootstrap algorithms, we filled the observational pole hole with the average sea-ice concentration around its edge (Olason & Notz, 2014). For OSI SAF, we used the filled pole hole of the product itself. We take the spread of the three algorithms obtained this way as the observational uncertainty σ_{obs} .

For sea-ice volume, we do not compare models with an observational estimate due to substantial uncertainties for reanalysed and observed estimates of Arctic sea-ice thickness and thus volume (e.g. Bunzel et al., 2018; Chevallier et al., 2017; Zygmuntowska et al., 2014).

For global-mean surface temperature (GMST), we use the average of NOAA GlobalTemp v5.0.0 (Vose et al., 2012), GISTemp v4 (GISTEMP Team, 2019; Lenssen et al., 2019), HadCRUT4.6.0.0 (Morice et al., 2012) and Berkeley (Rohde et al., 2013) time-series as an estimate for the mean evolution, and the spread across these four records as an estimate for observational uncertainty. We calculate anomalies relative to the period 1850–1900, except for the shorter record of NOAA GlobalTemp where we calculate anomalies relative to 1880–1900. Because the 20-year running-mean temperature fluctuations during these periods are less than 0.1 °C, our results are largely insensitive to this choice of baseline period (Figure S2). We take the spread of the four products as the observational uncertainty σ_{obs} .

Historical anthropogenic CO₂ emissions are taken from the historical budget of (Global Carbon Project, 2019). Future anthropogenic CO₂ emissions for CMIP6 simulations are taken from the respective SSP scenarios described by (Riahi et al., 2017).

3 CMIP6 Model Performance

3.1 Mean Quantities

We start with an analysis of the mean sea-ice fields simulated by individual CMIP3, CMIP5 and CMIP6 models (Figure 1a, b, e, f) over the period 1979–1998. To allow for a fair comparison across the three CMIP phases, in this section we analyze only the first ensemble member of each model. Given the large number of participating models, this results in a fair comparison: for models with several ensemble members, the first ensemble member is as likely to be above a model’s ensemble mean as below.

For sea-ice area, we find a large spread across CMIP6 simulations both in March and in September (Figure 1a, b), which usually are the months of maximum and minimum sea-ice coverage in the Arctic, respectively. In March, the 1979–1998 mean sea-ice area simulated by CMIP6 models ranges from around 12 million km² to more than 20 million km² and thus includes the observational estimate of 14.4 million km² (Figure 1a, Table S3). Out of the 40 CMIP6 models, 21 are within the $2\sigma = \pm 1.29$ million km² plausibility range around the observational estimate given by the CMIP6-average internal variability and observational uncertainty as introduced in section 2 (Figure 1a, Table S3). CMIP3 and CMIP5 simulations also show a large spread in mean March sea-ice area, and include the observational estimate within their multi-model ensemble spread (Figure 1a, Tables S1 and S2). However, in CMIP3 and CMIP5, the multi-model ensemble spread is more evenly distributed around the observational estimate than in CMIP6, where most models lie above it.

For the mean September sea-ice area over the period 1979–1998, the CMIP6 ensemble also shows a large spread of individual simulations, ranging from around 3 million km² to around 10 million km² (Figure 1b, Table S3). The observed value of around 6 million km² lies well within the range, and 25 out of 40 CMIP6 models are within the plausible range of $2\sigma = \pm 1.49$ million km² around this value (Table S3). The CMIP6 multi-model ensemble mean is very close to the observational estimate and well within the plausible range. The same holds for CMIP3 and CMIP5, with their individual models also spanning a wide range around the observational estimate (Figure 1b, Tables S1 and S2).

For sea-ice volume, we lack data for CMIP3 models and thus can only compare CMIP6 results to CMIP5 results (see tables S2 and S3 for a detailed overview). For both phases of CMIP, the models produce a similar spread of simulated Arctic sea-ice volume from less than 20,000 km³ to more than 40,000 km³ in March (Figure 1e), and from less than 5,000 km³ to more than 30,000 km³ in September (Figure 1f). Given a simulated average spread from internal variability of around 2,000 km³, the large spread in sea ice volume from CMIP6 models can not be explained by internal variability alone. Instead, it is caused by the models' large spread in simulated sea-ice area and thickness.

Based on this analysis of mean Arctic sea-ice quantities, we find that there is little difference in overall model performance between CMIP3, CMIP5 and CMIP6. The multi-model spread of the mean quantities remains large, the observational record lies within the multi-model ensemble spread, and many models simulate plausible values of mean sea-ice area when considering the impact of internal variability and observational uncertainty. The multi-model ensemble means of the past three phases of CMIP are relatively similar to each other and largely consistent with the observational record.

3.2 Sensitivity

In addition to their plausible simulation of mean quantities, the models' adequacy for simulating reality hinges critically on their ability to realistically simulate the response of a given climate metric to changes in external forcing. Internal variability causes a large spread of plausible climate trajectories in response to a given change in the forcing and must carefully be taken into account when interpreting a possible mismatch between a simulation and a given observational sea-ice record (Jahn et al., 2016; Kay et al., 2011; Notz, 2015; Olonscheck & Notz, 2017; Swart et al., 2015). We find this to remain valid for CMIP6 simulations.

For our analysis of the simulated sensitivity of Arctic sea ice to changes in external forcing, we calculate two distinct quantities: first, the change in sea-ice area for a given change in cumulative anthropogenic CO₂ emissions over the period 1979–2014 (Figure 1c); second, the change in sea-ice area for a given change in global mean surface temperature (GMST) over the period 1979–2014 (Figure 1d). Both quantities can be calculated from the previously demonstrated linear relationships of sea-ice area to cumulative CO₂ emissions (Herrington & Zickfeld, 2014; Notz & Stroeve, 2016; Zickfeld et al.,

205 2012) and to GMST (e.g., Gregory et al., 2002; Mahlstein & Knutti, 2012; Rosenblum
 206 & Eisenman, 2016; Stroeve & Notz, 2015; Winton, 2011). Together, these two quanti-
 207 ties allow us to estimate whether CMIP6 models simulate changes in sea ice with the cor-
 208 rect sensitivity to changes in external forcing, and whether they potentially do so for the
 209 right reason. This is because the relationship between sea-ice area and cumulative an-
 210 thropogenic CO₂ emissions is an almost linear proxy for the long-term time evolution
 211 of Arctic sea-ice area, as cumulative emissions map monotonously to time. In contrast,
 212 the sensitivity of sea-ice area to GMST changes is a proxy for the sensitivity of the sea-
 213 ice cover to one particular response of the climate system to changes in external forc-
 214 ing.

215 Our analysis reveals that over the historical period 1979–2014, 28 out of 40 CMIP6
 216 models simulate a sensitivity of the Arctic sea-ice area to cumulative anthropogenic CO₂
 217 emissions that is within the plausible range of 2.73 ± 1.37 m² of sea-ice loss per ton of CO₂
 218 emissions (Figure 1c, Table S3). In addition to the larger spread of the CMIP6 multi-
 219 model ensemble, a major difference between CMIP5 and CMIP6 models is that, in their
 220 first ensemble member analyzed here, only 3 out of 40 CMIP5 models simulate a larger
 221 loss of sea-ice area per ton of CO₂ emissions than observed. This number increases to
 222 10 out of 40 models for CMIP6. This results in the CMIP6 multi-model ensemble mean
 223 being closer to the observational estimate than the CMIP5 and the CMIP3 multi-model
 224 ensemble means. It is however unclear whether this reflects an improvement of model
 225 physics or primarily arises from the change in historical forcing in CMIP6 relative to CMIP5
 226 (compare Rosenblum & Eisenman, 2016). For example, in CMIP6 the historical ozone
 227 radiative forcing is about 80 % higher than it was in CMIP5 (Checa-Garcia et al., 2018).
 228 In contrast, black carbon emissions in the CMIP6 historical forcing are substantially higher
 229 over the past years than prescribed in the CMIP5 RCP8.5 scenario (Gidden et al., 2019).
 230 The impact of these changes in non-CO₂ climate drivers is confounded into the sensi-
 231 tivity of sea-ice area to CO₂ emissions (again, compare Rosenblum & Eisenman, 2016).
 232 Emissions of CO₂ itself, and of methane, are largely unchanged over the historical pe-
 233 riod for CMIP5 and CMIP6. However, for the future simulations the CMIP6 SSP5-8.5
 234 scenario assumes higher CO₂ emissions and lower methane emissions than the CMIP5
 235 RCP8.5 scenario (Gidden et al., 2019).

236 Examining the sea-ice loss per degree of global warming, we find that only 11 out
 237 of 40 CMIP6 models are within the plausible range of 4.01 ± 1.28 million m² of sea-ice

loss per degree of warming (Figure 1d, Table S3). This is comparable to CMIP5, where 9 out of 40 models were within this plausible range (Figure 1d, Table S2). In CMIP3, not a single model provided a plausible sensitivity (Figure 1d). Also, the CMIP6 multi-model ensemble mean of Arctic sea-ice loss for a given amount of global warming is closer to (but still outside) the plausible range than the multi-model ensemble mean of both CMIP5 and CMIP3. This might indicate an improvement of CMIP6 models over previous CMIP phases on a process level, given that the main physical link of sea-ice loss to any change in external forcing is given by a change in temperature. However, as before, this might also be a reflection of a more realistic historical forcing of CMIP6 compared to CMIP5 and CMIP3.

While the more realistic simulation of these two sensitivities might indicate progress in CMIP6 models' capability to simulate the ongoing loss of Arctic sea ice, as in CMIP5 (Rosenblum & Eisenman, 2017) few CMIP6 models are able to simulate a plausible amount of sea-ice loss and simultaneously a plausible change in global mean temperature over time (or cumulative anthropogenic CO₂ emissions). Of the CMIP6 models analyzed here, these are ACCESS-CM2, BCC-CSM2-MR, CNRM-CM6-1-HR, FGOALS-f3-L, FIO-ESM-2-0, GFDL-ESM4, GISS-E2-1-G, GISS-E2-1-G-CC, MPI-ESM-1-2-HAM, MPI-ESM1-2-HR, MPI-ESM1-2-LR, MRI-ESM2-0 and NorESM2-MM. For the other CMIP6 models, those models that have a reasonable sea-ice loss tend to have too much global warming, while those models that simulate reasonable global warming simulate too little sea-ice loss (Figure 1g, Table S3). In particular, the models with a high sensitivity of Arctic sea-ice area to anthropogenic CO₂ emissions also display a high sensitivity of global mean temperature to CO₂ emissions. Hence, understanding this high climate sensitivity is most likely key to understanding why some CMIP6 models display such rapid loss of Arctic sea ice. A recent study suggested this high sensitivity to be caused by stronger cloud feedbacks (Zelinka et al., 2020).

If we plot the two sensitivity metrics against each other, it is generally impossible to distinguish a given CMIP6 model from the cloud given by CMIP5 models, with the exception of the highly sensitive CMIP6 simulations that clearly fall outside the cloud of previous CMIP phases (Figure 1g). The lack of both such high-sensitive simulations and of very low-sensitive simulations in CMIP5 might be one reason for why the correlation between the two metrics is lower for CMIP5 than for CMIP3 and CMIP6.

In summary, we find that over the period 1979–2014, CMIP6 models on average simulate a sensitivity of Arctic sea ice that is closer to the observed value than CMIP5 and CMIP3 models, both relative to a given CO₂ emission (as a proxy for time) and to a given warming. However, only few models are able to simulate a plausible sea-ice loss sensitivity to cumulative CO₂ emissions and simultaneously a plausible rise in global mean surface temperature.

4 Projections of Future Arctic Sea Ice

The identified spread of CMIP models in simulating the past mean state and sensitivity to warming and CO₂ emissions introduces significant model uncertainty into future projections of the evolution of the Arctic sea-ice cover. This model uncertainty remains large in CMIP6.

To address this issue when analyzing projections of when Arctic sea-ice area might drop below 1 million km², a commonly used threshold for an ice-free Arctic, we take the following approach. First, we examine the full range of CMIP6 model simulations, noting that the model spread provides a wide spectrum of the possible future evolution of Arctic sea-ice area. Second, we narrow the range by considering only those models that have the observations within their ensemble spread simultaneously for two key metrics (compare Massonnet et al., 2012): (a) the 2005–2014 September mean sea-ice area and (b) the observed sensitivity of sea-ice area to cumulative CO₂ emissions over the period 1979–2014. We choose these metrics because they correlate with the first sea-ice free year at a correlation of $R > 0.5$ for all scenarios over the entire CMIP6 multi-model ensemble. Note, however, that care must be taken when interpreting the range of selected models, as the relationship between past and future evolution of a climate model is not always clear (Jahn et al., 2016; Stroeve & Notz, 2015). On the other hand, it becomes more important that a model plausibly captures the observed mean state of Arctic sea-ice area the lower that mean state becomes, because initial conditions become more important as the observed sea-ice state approaches ice-free conditions and the simulations start entering the realm of decadal predictions. We hence trust that the range of uncertainty given by the selected models gives a more realistic estimate of the true model uncertainty than that given by the full CMIP6 multi-model ensemble. The selected models are printed in bold in table S4.

In analyzing the future relationship between sea-ice loss and changes in the forcing, we find that the simulated correlation between winter Arctic sea-ice area and cumulative CO₂ emissions remains high well into the future (Figure 2a). For summer, the linear relationship eventually decreases as more and more years of zero Arctic sea-ice coverage are averaged into the multi-model mean (Figure 2d). In interpreting these results quantitatively, it is of course important to note that CO₂, while being the most important external driver of observed changes in Arctic sea-ice coverage, is not the only cause of observed and future changes. Its dominant role, however, holds well into the future and/or the additional impacts of other anthropogenic forcings, such as methane and aerosols, remain roughly stable over time. Otherwise the correlation between March Arctic sea-ice area and cumulative CO₂ emissions would not remain as stable over time and would not be as independent of the specific forcing scenario (Figure 2a).

We also find that the simulated correlation of temperature with winter Arctic sea-ice area remains high well into the future (Figure 2b), while again in summer the correlation eventually decreases as more models lose their sea ice completely (Figure 2e).

The high correlation between sea-ice loss and changes in the forcing allows us to estimate the cumulative future CO₂ emissions, warming level and eventually year at which the Arctic Ocean will practically be sea-ice free for the first time, defined as the first year in which the monthly mean September sea-ice area drops below 1 million km².

We find that CMIP6 models simulate a large spread of cumulative future CO₂ emissions at which the Arctic could first become practically sea-ice free in September (Figure 3a). The simulated future emissions for the first occurrence of a practically sea-ice free Arctic Ocean range from 450 Gt CO₂ below to more than 5000 Gt CO₂ above present cumulative emissions. However, 158 out of 243 simulations become practically sea-ice free before future cumulative CO₂ emissions reach 1000 GtCO₂ above that of 2019 (equivalent to about 3400 GtCO₂ cumulative emissions since 1850). Considering only the models with ensemble members within the plausible range of observed sea-ice evolution, we find a reduced range of 170 Gt below to 2200 Gt above cumulative future anthropogenic CO₂ emissions when Arctic sea-ice area is projected to drop below 1 million km². Of these members from the selected models, the vast majority (101 out of 128) become practically sea-ice free at future cumulative CO₂ emissions less than 1000 Gt. This compares favourably with the range of 800±300 Gt estimated from a direct analysis of the observed

sensitivity (Notz & Stroeve, 2018). In combination, these estimates make it appear likely that the Arctic Ocean will practically lose its sea ice cover in September for the first time at future anthropogenic CO₂ emissions of between 200 and 1100 Gt above that of 2019.

As a function of GMST, ice-free conditions occur across the entire CMIP6 multi-model ensemble at a global warming of between 0.9 and 3.2 °C above pre-industrial conditions of each individual model (Figure 3b). If we select only those models with a reasonable simulation of past Arctic sea-ice conditions, the estimated temperature range decreases slightly to 1.3 to 2.9 °C. The upper end of this range is higher than the range of 1.7±0.4 °C estimated from a direct analysis of the observed sensitivity (Notz & Stroeve, 2018) and higher than estimates from bias-corrected simulations that all project the first ice-free Arctic at temperatures below 2 °C (Jahn, 2018; Niederdrenk & Notz, 2018; Ridley & Blockley, 2018; Screen & Williamson, 2017; Sigmond et al., 2018). This high bias is probably a reflection of the CMIP6 models' weak sensitivity of sea-ice area loss to global warming, resulting in too high estimates of the warming at which the Arctic becomes practically sea-ice free in summer.

In the CMIP6 ensemble, the sea-ice area loss per cumulative CO₂ emissions and degree of global warming does barely depend on the forcing scenario (Figure 3a, b). Scenario dependence is also very small regarding the near-term future evolution of Arctic summer sea ice as a function of time until about 2040 (Figures 2f and 3c). This is related to the fact that until 2040, the scenarios evolve quite similarly (O'Neill et al., 2016). Furthermore, given that the current sea-ice area is much smaller than it used to be, the importance of internal variability increases relative to the forced change necessary to lose the remaining sea-ice cover in September. As a consequence, for some models the sea ice disappears earlier for the low-emissions scenarios than for the high-emissions scenarios in the ensemble members provided to the CMIP6 archive (Table S4). For all scenarios, the first year of practically sea-ice-free conditions ranges from some years before present to the end of this century (Table S4), with a clear majority of models reaching ice-free conditions before 2050. This finding remains valid for the selected models. From the middle of the century onward, scenario dependence becomes more and more evident. For example, the loss of sea-ice area in March occurs much faster from 2050 onward in scenario SSP5-8.5 than in other scenarios (Figure 2c).

5 Conclusion

Based on the analyzed evolution of Arctic sea-ice area and volume in CMIP6 models, in this contribution we have found the following:

- CMIP6 model performance in simulating Arctic sea ice is similar to CMIP3 and CMIP5 model performance in many aspects. This includes models simulating a wide spread of mean sea-ice area and volume in March and September; the multi-model ensemble spread capturing the observed mean sea-ice area in March and September; the models' general underestimation of the sensitivity of September sea-ice area to a given amount of global warming; as well as most models' failure to simulate at the same time a plausible evolution of sea-ice area and of global mean surface temperature.
- CMIP6 model performance differs from CMIP3 and CMIP5 in some aspects. These include a larger fraction of CMIP6 models capturing the observed sensitivity of Arctic sea ice to anthropogenic CO₂ emissions and the CMIP6 multi-model ensemble mean being closer to the observed sensitivity of Arctic sea ice to global warming. It is unclear to what degree these improvements are caused by a change in the forcing versus improvement of model physics.
- The CMIP6 models simulate a large spread for when Arctic sea-ice area is predicted to drop below 1 million km², such that the Arctic Ocean becomes practically sea-ice free. However, the clear majority of all models, and of those models that best capture the observed evolution, project that the Arctic will become practically sea-ice free in September before the year 2050 at future anthropogenic CO₂ emissions of less than 1000 GtCO₂ above that of 2019 in all scenarios.





Figure 1. Comparison of sea-ice metrics as simulated by the first ensemble members of CMIP3 (blue), CMIP5 (orange) and CMIP6 (green) models. The individual panels show the mean Arctic sea-ice area (SIA) in (a) March and (b) September for 1979–1998; mean Arctic sea-ice volume (SIV) in (e) March and (f) September for 1979–1998; and (c-d) the sensitivity over the period 1979–2014 of September sea-ice area to (c) CO₂ emissions and (d) global annual mean surface temperature (GMST). (g) The sensitivity of Arctic sea-ice area to CO₂ emissions scattered against the sensitivity of GMST to CO₂ emissions. In (a-f), horizontal dashes represent the first ensemble member of each model and crosses represent the multi-model ensemble mean. The thick dashed black lines denote the average of the observational satellite products, where available. The dotted lines denote one standard deviation of observational uncertainty. The green dashed lines denote the 2σ plausible range including internal variability and observational uncertainty as defined in section 2. The gray shadings around the lines denote overlays of estimated internal variability from all CMIP6 models with three or more ensemble members, with each overlay representing the 1-standard-deviation spread of a single model. Hence, the darker the shading, the more models agree on internal variability to cover a certain range.

March



September




Figure 2. Evolution of Arctic sea-ice area over the historical period and following three scenario projections in (a-c) March and (d-f) September as a function of (a,d) cumulative anthropogenic CO₂ emissions, (b,e) global annual mean surface temperature anomaly and (c,f) time for all available CMIP6 models. Thick lines denote the multi-model ensemble mean, where all models are represented by their first ensemble member, and the shading around the lines indicates one standard deviation around the multi-model mean. Faint dots denote the first ensemble member of each model and thick black lines and crosses denote observations. Note that discontinuities in the multi-model ensemble mean arise from a different number of available models for the historical period and the scenario simulations.




Figure 3. CMIP6 projections of (a) future cumulative CO₂ emissions, (b) global annual mean surface temperature anomaly and (c) year when September-mean sea-ice area drops below 1 million km² for the first time in each simulation. The numbers at the top of the panels denote the number of simulations that do not simulate a sea-ice cover below 1 million km² by 2100 (top row) and the total number of simulations (bottom row) for each scenario. Each dot represents a single simulation, with all available CMIP6 simulations shown in the figure.

Appendix A Authors

All authors contributed to discussions and the writing of the paper, as well as implementation or analysis of SIMIP variables in CMIP6 models. Additional contributions are listed below.

Dirk Notz, Center for Earth System Research and Sustainability (CEN), University of Hamburg and Max Planck Institute for Meteorology, Hamburg, Germany; co-chair of SIMIP, lead the development of this paper, contributed to implementing the SIMIP protocol in MPI-ESM

Jakob Dörr, Max Planck Institute for Meteorology, Hamburg, Germany; carried out all data analysis for this paper; compiled all figures and tables

David A Bailey, Climate and Global Dynamics Laboratory, National Center for Atmospheric Research, Boulder, Colorado, United States of America

Ed Blockley, Met Office Hadley Centre, Exeter, United Kingdom ; contributed to the sea ice component of the UKESM and HadGEM3 models

Mitchell Bushuk, Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey, United States of America

Jens Boldingh Debernard, Norwegian Meteorological Institute, Norway; contribution to the sea ice component of NorESM2-LM

Evelien Dekker, Rossby Centre, Swedish Meteorological and Hydrological Institute, Norrköping, Meteorological Department at Stockholm University, Sweden

Patricia DeRepentigny, Department of Atmospheric and Oceanic Sciences and Institute of Arctic and Alpine Research, University of Colorado Boulder, Boulder, Colorado, United States of America

David Docquier, Rossby Centre, Swedish Meteorological and Hydrological Institute, Norrköping, Sweden

Neven S. Fućkar, Environmental Change Institute, University of Oxford, Oxford, UK, and Earth Sciences Department, Barcelona Supercomputing Center, Barcelona, Spain

John C. Fyfe, Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada

Alexandra Jahn, University of Colorado Boulder, Department of Atmospheric and Oceanic Sciences and Institute of Arctic and Alpine Research, Boulder, Colorado, United States of America; co-chair of SIMIP

Marika Holland, Climate and Global Dynamics Laboratory, National Center for Atmospheric Research, Boulder, Colorado, United States of America; SIMIP steering-group member

Elizabeth Hunke, Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico, United States of America; SIMIP steering-group member

Doroteaciro Iovino, Ocean Modeling and Data Assimilation Division, Centro Euro-Mediterraneo sui Cambiamenti Climatici, Italy

Narges Khosravi, Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

Gurvan Madec, Sorbonne Universits, UPMC Paris 6, LOCEAN-IPSL, CNRS/IRD/MNHN, Paris, France

François Massonnet, Georges Lematre Centre for Earth and Climate Research, Earth and Life Institute, Universit catholique de Louvain, Louvain-la-Neuve, Belgium; SIMIP steering-group member

Siobhan O'Farrell, CSIRO Oceans and Atmosphere, Aspendale, Victoria, Australia

Alek Petty, Cryospheric Sciences Laboratory, NASA Goddard Space Flight Center, Greenbelt, Maryland, United States of America, and Earth System Science Interdisciplinary Center, University of Maryland, College Park, Maryland, United States of America

Arun Rana, Georges Lematre Centre for Earth and Climate Research, Earth and Life Institute, Universit catholique de Louvain, Louvain-la-Neuve, Belgium

Lettie Roach, Atmospheric Sciences, University of Washington, Seattle, Washington, United States of America

Erica Rosenblum, Centre for Earth Observation Science, University of Manitoba, Winnipeg, Manitoba, Canada; contributed to the preliminary data analysis

Clement Rousset, Sorbonne Universits, UPMC Paris 6, LOCEAN-IPSL, CNRS/IRD/MNHN,
Paris, France

Tido Semmler, Alfred Wegener Institute, Helmholtz Centre for Polar and Marine
Research, Bremerhaven, Germany

Julienne Stroeve, University College London, London, United Kingdom and Na-
tional Snow and Ice Data Center, Boulder, Colorado, United States of America; SIMIP
steering-group member

Takahiro Toyoda, Meteorological Research Institute, Japan Meteorological Agency,
Japan; contributed to carry out the MRI-ESM2 experiments and to prepare the output
for SIMIP analyses

Bruno Tremblay, Department of Atmospheric and Oceanic Sciences, McGill Uni-
versity, Montreal, Canada; SIMIP steering-group member.

Hiroyuki Tsujino, Meteorological Research Institute, Japan Meteorological Agency,
Japan; contributed to carry out the MRI-ESM2 experiments and to prepare the output
for SIMIP analyses

Martin Vancoppenolle, Sorbonne Universits, UPMC Paris 6, LOCEAN-IPSL, CNRS/
IRD/ MNHN, Paris, France; SIMIP steering-group member; contributed to the sea ice
component of IPSL-CM and EC-Earth

Acknowledgments

We thank two anonymous reviewers for their valuable feedback that helped improving
this manuscript.

We are grateful to all modeling centers for carrying out CMIP6 simulations used
here. The data used for this study are freely available from the Earth System Grid Fed-
eration (ESGF) at esgf-node.llnl.gov/search/cmip6. See supporting information for
a detailed listing of all CMIP6 data sets used in this study, including their dois. The scripts
for analysis and plotting of the data are available from https://github.com/jakobdoerr/SIMIP_2020.

EC-Eart3-Veg simulations that are not published on ESGF yet are locally stored
at the Swedish Meteorological and Hydrological Institute (cmip6-data@ecearth.org). We

thank the EC-Earth consortium that realized the development of EC-Earth. The EC-Earth3-Veg simulations were done as part of the European Union's Horizon 2020 research and innovation programme under grant agreement No 641816 (CRESCENDO) on resources provided by the Swedish National Infrastructure for Computing (SNIC) at PDC and NSC.

Previous and current CESM versions are freely available at www.cesm.ucar.edu/models. The CESM project is supported primarily by the National Science Foundation (NSF). This material is based upon work supported by the National Center for Atmospheric Research, which is a major facility sponsored by the NSF under Cooperative Agreement No. 1852977. Computing and data storage resources, including the Cheyenne supercomputer (doi:10.5065/D6RX99HX), were provided by the Computational and Information Systems Laboratory (CISL) at NCAR. We thank all the scientists, software engineers, and administrators who contributed to the development of CESM2.

The ACCESS-CM2 CMIP6 submission was jointly funded through CSIRO and the Earth Systems and Climate Change Hub of the Australian Government's National Environmental Science Program, with support from the Australian Research Council Centre of Excellence for Climate System Science. The ACCESS-ESM1.5 CMIP6 submission was supported by the CSIRO Climate Science Centre.

Parts of the work described in this paper has received funding from the European Union's Horizon 2020 Research and Innovation programme through grant agreement No. 727862 APPLICATE. The content of the paper is the sole responsibility of the authors and it does not represent the opinion of the European Commission, and the Commission is not responsible for any use that might be made of information contained.

We thank the WCRP-CliC Project for supporting the SIMIP project.

E. Blockley was supported by the Joint UK BEIS/Defra Met Office Hadley Centre Climate Programme (GA01101)

J. B. Debernard is supported by the Research Concile of Norway through INES (270061).

E. Dekker is supported by the Arctic Across Scales Project through the Knut and Alice Wallenberg Foundation (KAW2016.0024)

P. DeRepentigny is supported by the Natural Sciences and Engineering Council of Canada, the Fond de recherche du Qubec – Nature et Technologies and the Canadian Meteorological and Oceanographic Society through PhD scholarships and NSF-OPP award 1847398.

D. Docquier is funded by the EU Horizon 2020 OSeaIce project, under the Marie Skłodowska-Curie grant agreement no. 834493.

J. Dörr is funded by the German Ministry for Education and Research through the project “Meereis bei +1.5 °C”.

N.S. Fućkar acknowledges support of H2020 MSCA IF (Grant ID 846824).

E. Hunke is supported by the Regional and Global Modeling and Analysis program of the Department of Energy’s Biological and Environmental Research division.

A. Jahn’s contribution is supported by NSF-OPP award 1847398.

F. Massonnet is a F.R.S.-FNRS Research Fellow.

D. Notz is funded by the Deutsche Forschungsgemeinschaft under Germanys Excellence Strategy EXC 2037 ‘CLICCS - Climate, Climatic Change, and Society’ Project Number: 390683824, contribution to the Center for Earth System Research and Sustainability (CEN) of Universität Hamburg.

L. Roach was supported by the National Science Foundation grant PLR-1643431 and National Oceanic and Atmospheric Administration grant NA18OAR4310274.

J. Stroeve and E. Rosenblum are supported by the Canada C150 Chair Program

This work is a contribution to NSF-OPP award 1504023 awarded to B. Tremblay.

References

- Bunzel, F., Notz, D., & Pedersen, L. T. (2018). Retrievals of Arctic Sea-Ice Volume and Its Trend Significantly Affected by Interannual Snow Variability. *Geophysical Research Letters*, 45(21), 11,751–11,759. doi: 10.1029/2018GL078867
- Cavalieri, D. J., Parkinson, C. L., Gloersen, P., & Zwally, H. J. (1997). *Arctic and Antarctic Sea Ice Concentrations from Multichannel Passive-microwave Satellite Data Sets: October 1978 to December 1996* (Technical Memorandum No.

- 104647). NASA.
- Checa-Garcia, R., Hegglin, M. I., Kinnison, D., Plummer, D. A., & Shine, K. P. (2018). Historical tropospheric and stratospheric ozone radiative forcing using the CMIP6 database. *Geophysical Research Letters*, 45(7), 3264–3273. doi: 10.1002/2017GL076770
- Chevallier, M., Smith, G. C., Dupont, F., Lemieux, J.-F., Forget, G., Fujii, Y., ... Wang, X. (2017). Intercomparison of the Arctic sea ice cover in global oceansea ice reanalyses from the ORA-IP project. *Clim. Dynam.*, 49(3), 1107–1136. doi: 10.1007/s00382-016-2985-y
- Comiso, J. C., Cavalieri, D. J., Parkinson, C. L., & Gloersen, P. (1997). Passive microwave algorithms for sea ice concentration: A comparison of two techniques. *Remote Sens. Env.*, 60(3), 357–384. doi: 10.1016/S0034-4257(96)00220-9
- Comiso, J. C., Meier, W. N., & Gersten, R. (2017). Variability and trends in the Arctic sea ice cover: Results from different techniques. *Journal of Geophysical Research: Oceans*, 122(8), 6883–6900.
- England, M., Jahn, A., & Polvani, L. (2019). Nonuniform contribution of internal variability to recent Arctic sea ice loss. *J. Climate*, 32(13), 4039–4053. doi: 10.1175/JCLI-D-18-0864.1
- Eyring, V., Bony, S., Meehl, G. A., Senior, C., Stevens, B., Stouffer, R. J., & Taylor, K. E. (2015). Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organisation. *Geosci. Model Dev. Discuss.*, 8(12), 10539–10583. doi: 10.5194/gmdd-8-10539-2015
- Gidden, M., Riahi, K., Smith, S., Fujimori, S., Luderer, G., Kriegler, E., ... others (2019). Global emissions pathways under different socioeconomic scenarios for use in CMIP6: a dataset of harmonized emissions trajectories through the end of the century. *Geoscientific model development*, 12(4), 1443–1475. doi: 10.5194/gmd-12-1443-2019
- GISTEMP Team. (2019). *GISS Surface Temperature Analysis (GISTEMP), version 4*. NASA Goddard Institute for Space Studies [date accessed: XX/XX/XX]. Retrieved from <https://data.giss.nasa.gov/gistemp/>
- Global Carbon Project. (2019). *Supplemental data of global carbon budget 2019 (version 1.0 [data set]*. doi: doi.org/10.18160/gcp-2019
- Gregory, J. M., Stott, P. A., Cresswell, D. J., Rayner, N. A., Gordon, C., & Sex-

- ton, D. M. H. (2002). Recent and future changes in Arctic sea ice simulated by the HadCM3 AOGCM. *Geophys. Res. Lett.*, 29(24), 2175, doi:10.1029/2001GL014575.
- Herrington, T., & Zickfeld, K. (2014). Path independence of climate and carbon cycle response over a broad range of cumulative carbon emissions. *Earth Syst. Dynam.*, 5, 409422. doi: 10.5194/esd-5-409-2014
- Jahn, A. (2018). Reduced probability of ice-free summers for 1.5 °C compared to 2 °C warming. *Nat. Clim. Change*. doi: 10.1038/s41558-018-0127-8
- Jahn, A., Kay, J. E., Holland, M. M., & Hall, D. M. (2016). How predictable is the timing of a summer ice-free Arctic? *Geophys. Res. Lett.*, 2016GL070067. doi: 10.1002/2016GL070067
- Kay, J. E., Holland, M. M., & Jahn, A. (2011). Inter-annual to multi-decadal Arctic sea ice extent trends in a warming world. *Geophys. Res. Lett.*, 38. doi: 10.1029/2011GL048008
- Koenigk, T., Devasthale, A., & Karlsson, K.-G. (2014). Summer Arctic sea ice albedo in CMIP5 models. *Atmos. Chem. Phys.*, 14(4), 1987–1998. doi: 10.5194/acp-14-1987-2014
- Lavergne, T., Srensen, A. M., Kern, S., Tonboe, R., Notz, D., Aaboe, S., ... Pedersen, L. T. (2019). Version 2 of the EUMETSAT OSI SAF and ESA CCI sea-ice concentration climate data records. *The Cryosphere*, 13(1), 49–78. doi: https://doi.org/10.5194/tc-13-49-2019
- Lenssen, N. J., Schmidt, G. A., Hansen, J. E., Menne, M. J., Persin, A., Ruedy, R., & Zyss, D. (2019). Improvements in the GISTEMP uncertainty model. *Journal of Geophysical Research: Atmospheres*, 124(12), 6307–6326. doi: 10.1029/2018JD029522
- Mahlstein, I., & Knutti, R. (2012). September Arctic sea ice predicted to disappear near 2 °C global warming above present. *J. Geophys. Res.*, 117, D06104. doi: 10.1029/2011JD016709
- Massonnet, F., Fichet, T., Goosse, H., Bitz, C. M., Philippon-Berthier, G., Holland, M. M., & Barriat, P.-Y. (2012). Constraining projections of summer Arctic sea ice. *The Cryosphere*, 6(6), 1383–1394. doi: 10.5194/tc-6-1383-2012
- Melia, N., Haines, K., & Hawkins, E. (2015). Improved Arctic sea ice thickness projections using bias-corrected CMIP5 simulations. *The Cryosphere*, 9(6), 2237–

2251. doi: 10.5194/tc-9-2237-2015
- Morice, C. P., Kennedy, J. J., Rayner, N. A., & Jones, P. D. (2012). Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: The HadCRUT4 data set. *Journal of Geophysical Research Atmospheres*, 117(8), 1–22. doi: 10.1029/2011JD017187
- Niederdrenk, A. L., & Notz, D. (2018). Arctic Sea Ice in a 1.5°C Warmer World. *Geophys. Res. Lett.*, 45(4), 1963–1971. doi: 10.1002/2017GL076159
- Notz, D. (2014). Sea-ice extent and its trend provide limited metrics of model performance. *The Cryosphere*, 8(1), 229–243. doi: 10.5194/tc-8-229-2014
- Notz, D. (2015). How well must climate models agree with observations? *Phil. Trans. R. Soc. A*, 373(2052), 20140164. doi: 10.1098/rsta.2014.0164
- Notz, D., Jahn, A., Holland, M., Hunke, E., Massonnet, F., Stroeve, J., ... Van-coppenolle, M. (2016). Sea Ice Model Intercomparison Project (SIMIP): Understanding sea ice through climate-model simulations. *Geoscientific Model Development Discussions*, 1–29. doi: 10.5194/gmd-2016-67
- Notz, D., & Stroeve, J. (2016). Observed Arctic sea-ice loss directly follows anthropogenic CO₂ emission. *Science*, aag2345. doi: 10.1126/science.aag2345
- Notz, D., & Stroeve, J. (2018). The trajectory towards a seasonally ice-free Arctic Ocean. *Current Climate Change Reports*, 4(4), 407–416. doi: 10.1007/s40641-018-0113-2
- Olason, E., & Notz, D. (2014). Drivers of variability in Arctic sea-ice drift speed. *Journal of Geophysical Research: Oceans*, 119(9), 5755–5775.
- Olonscheck, D., & Notz, D. (2017). Consistently Estimating Internal Climate Variability from Climate Model Simulations. *J. Clim.*, 30(23), 9555–9573. doi: 10.1175/JCLI-D-16-0428.1
- O'Neill, B. C., Tebaldi, C., Van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., ... Sanderson, B. M. (2016). The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6. *Geoscientific Model Development*, 9(9), 3461–3482. doi: 10.5194/gmd-9-3461-2016
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., ... Tavoni, M. (2017). The shared socioeconomic pathways and their energy, land use, and greenhouse gas emissions implications: An overview. *Global Environmental Change*, 42, 153 – 168. doi: 10.1016/j.gloenvcha.2016.05.009

- Ridley, J., & Blockley, E. (2018). Solar radiation management not as effective as CO₂ mitigation for Arctic sea ice loss in hitting the 1.5 and 2 °C COP climate targets. *The Cryosphere*(12), 3355–3360. doi: 10.5194/tc-12-3355-2018
- Roach, L. A., Dörr, J., Holmes, C. R., Massonnet, F., Blockley, E. W., Notz, D., ... Bitz, C. M. (2020). Antarctic sea ice in CMIP6. *Geophys. Res. Lett.*, 2019GL086729.
- Rohde, R., Muller, R., Jacobsen, R., Perlmutter, S., & Mosher, S. (2013). Berkeley Earth Temperature Averaging Process. *Geoinformatics & Geostatistics: An Overview*, 01(02). doi: 10.4172/2327-4581.1000103
- Rosenblum, E., & Eisenman, I. (2016). Faster Arctic sea ice retreat in CMIP5 than in CMIP3 due to volcanoes. *J. Climate*, 29(24), 9179–9188. doi: 10.1175/JCLI-D-16-0391.1
- Rosenblum, E., & Eisenman, I. (2017). Sea Ice Trends in Climate Models Only Accurate in Runs with Biased Global Warming. *J. Clim.*, 30(16), 6265–6278. doi: 10.1175/JCLI-D-16-0455.1
- Screen, J. A., & Williamson, D. (2017). Ice-free Arctic at 1.5 °C? *Nat. Clim. Change*, 7(4), 230–231. doi: 10.1038/nclimate3248
- Shu, Q., Song, Z., & Qiao, F. (2015). Assessment of sea ice simulations in the CMIP5 models. *The Cryosphere*, 9(1), 399–409. doi: 10.5194/tc-9-399-2015
- Sigmond, M., Fyfe, J. C., & Swart, N. C. (2018). Ice-free Arctic projections under the Paris Agreement. *Nat. Clim. Change*. doi: 10.1038/s41558-018-0124-y
- Stroeve, J., Barrett, A., Serreze, M., & Schweiger, A. (2014). Using records from submarine, aircraft and satellites to evaluate climate model simulations of Arctic sea ice thickness. *The Cryosphere*, 8(5), 1839–1854. doi: 10.5194/tc-8-1839-2014
- Stroeve, J., Holland, M. M., Meier, W., Scambos, T., & Serreze, M. (2007). Arctic sea ice decline: Faster than forecast. *Geophys. Res. Lett.*, 34, L09501, doi:10.1029/2007GL029703.
- Stroeve, J., Kattsov, V., Barrett, A., Serreze, M., Pavlova, T., Holland, M., & Meier, W. N. (2012). Trends in Arctic sea ice extent from CMIP5, CMIP3 and observations. *Geophys. Res. Lett.*, 39(16), L16502. doi: 10.1029/2012GL052676
- Stroeve, J., & Notz, D. (2015). Insights on past and future sea-ice evolution from combining observations and models. *Glob. Planet Change*, 135, 119–132. doi:

10.1016/j.gloplacha.2015.10.011

Swart, N. C., Fyfe, J. C., Hawkins, E., Kay, J. E., & Jahn, A. (2015). Influence of internal variability on Arctic sea-ice trends. *Nat. Clim. Change*, 5(2), 86–89.

doi: 10.1038/nclimate2483

Vose, R. S., Arndt, D., Banzon, V. F., Easterling, D. R., Gleason, B., Huang, B., ... Wuertz, D. B. (2012). NOAA’s merged land-ocean surface temperature analysis. *Bulletin of the American Meteorological Society*, 93(11), 1677–1685.

doi: 10.1175/BAMS-D-11-00241.1

Winton, M. (2011). Do Climate Models Underestimate the Sensitivity of Northern Hemisphere Sea Ice Cover? *J. Clim.*, 24(15), 3924–3934.

Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., ... Taylor, K. E. (2020). Causes of higher climate sensitivity in cmip6 models. *Geophysical Research Letters*, 47(1), e2019GL085782. doi:

10.1029/2019GL085782

Zickfeld, K., Arora, V. K., & Gillett, N. P. (2012). Is the climate response to CO₂ emissions path dependent? *Geophys. Res. Lett.*, 39, L05703. doi: 10.1029/2011GL050205

Zygmuntowska, M., Rampal, P., Ivanova, N., & Smedsrud, L. H. (2014). Uncertainties in Arctic sea ice thickness and volume: new estimates and implications for trends. *The Cryosphere*, 8(2), 705–720. doi: 10.5194/tc-8-705-2014