

Review article

Current evidence regarding 2D ultrasonography monitoring of intrinsic foot muscle properties: A systematic review

Nicolas Haelewyn^{a,*}, Jean-Louis Peters Dickie^b, Filip Staes^c, Evie Vereecke^d, Kevin Deschamps^{a,e}

^a Department of Rehabilitation Sciences, Musculoskeletal Rehabilitation Research Group, KU Leuven, Spoorwegstraat 12, 8200 Brugge, Belgium

^b Neuromusculoskeletal Lab (NMSK), Secteur des Sciences de la Santé, Institut de Recherche Expérimentale et Clinique, Université Catholique de Louvain, Brussels, Belgium

^c Department of Rehabilitation Sciences, Musculoskeletal Rehabilitation Research Group, KU Leuven, Tervuursevest 101, 3000 Leuven, Belgium

^d Department of Development & Regeneration, Muscles & Movement Group, KU Leuven, Etienne Sabbelaan 53, 8500 Kortrijk, Belgium

^e Haute Ecole Leonard De Vinci, Division of Podiatry, Avenue E. Mounier 84, 1200 Sint-Lambrechts-Woluwe, Belgium

ARTICLE INFO

Keywords:

Intrinsic foot muscles

Reference values

Reliability

Systematic review

Ultrasound imaging

ABSTRACT

Background: Ultrasonography can discriminate between intrinsic and extrinsic foot muscle properties and has therefore gained considerable popularity as an indirect strength evaluation. However, an overview on the use of ultrasound for assessing intrinsic foot musculature (IFM) is currently lacking.

Research question: What is the current evidence regarding (1) 2D ultrasonography protocols and its reliability? (2) Reference values for cross-sectional area and dorso-plantar thickness evaluation in asymptomatic and symptomatic persons?

Methods: The PRISMA guidelines were used to conduct this systematic review. Eight databases (PubMed, Embase, Web of Science, Cochrane Library, Scopus, CINAHL, SPORTDiscus and EuropePMC) were searched up to November 1, 2021. Studies reporting quantitative 2D ultrasound findings of the intrinsic foot muscles with no limitation to sex, BMI, ethnicity or physical activity were included. Studies were assessed for methodological quality using the Downs and Black checklist.

Results: Fifty-three studies were retained. Protocols showed an overall good to great reliability, suggesting limits of agreement between 8 and 30% of relative muscle size with minimal detectable changes varying from 0.10 to 0.29 cm² for cross-sectional area and 0.03–0.23 cm for thickness. Reference values are proposed for both cross-sectional area and thickness measurements of the abductor hallucis, flexor digitorum brevis, flexor hallucis brevis and quadratus plantae in asymptomatic persons. This could not be performed in the symptomatic studies due to a limited number of relevant studies addressing the symptomatic population, therefore a clinical overview is outlined. Clinically, IFM properties have been studied in ten distinct pathological conditions, predominantly pointing towards decreased muscle properties of the abductor hallucis.

Significance: We provide a clear and comprehensive overview of the literature regarding 2D ultrasonography of the IFM, making the available evidence more accessible to decision makers and researchers.

* Corresponding author.

E-mail addresses: nicolas.haelewyn@kuleuven.be (N. Haelewyn), jean-louis.peters@uclouvain.be (J.-L. Peters Dickie), filip.staes@kuleuven.be (F. Staes), evie.vereecke@kuleuven.be (E. Vereecke), kevin.deschamps@kuleuven.be (K. Deschamps).

<https://doi.org/10.1016/j.heliyon.2023.e18252>

Received 21 June 2023; Received in revised form 10 July 2023; Accepted 12 July 2023

Available online 17 July 2023

2405-8440/© 2023 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The intrinsic foot muscles (IFM) play an important role in sports activities as they contribute to the elastic energy absorption and generation of the medial longitudinal arch while working in conjunction with the plantar aponeurosis, ligaments and extrinsic foot muscles [1–4]. According to the ‘foot core’ paradigm, proposed by McKeon and colleagues, the IFM act as principal local stabilizers of the foot arches in a similar way the deep trunk muscles do in stabilizing the vertebral segments [5]. They contribute to balance performance and contract eccentrically during the early stance phase of rearfoot running and shorten during the late stance phase to produce positive work [2]. There is increasing evidence towards an association of IFM weakness and atrophy across the clinical course of a number of musculoskeletal conditions, including plantar fasciitis [6], posteromedial shin pain [7] and chronic ankle instability [7–11], and systemic diseases, such as diabetic mellitus [12]. IFM strengthening has therefore been recommended as an integral part of foot and lower limb rehabilitation programs [9,13,14] as well as sport-specific strength training and prevention exercises [15].

In order to assess IFM strength or their response to training, a valid and reliable strength evaluation is required. However, intrinsic and extrinsic foot muscles share similar functions, which makes a clinical differentiation among both muscle groups very challenging [16]. Currently, clinicians rely on direct and indirect methods. Direct methods include clinical tests such as the paper grip test [17] or dynamometry [18] but these evaluate several muscles at a time, not allowing the clinician to assess a single intrinsic foot muscle. Indirect methods commonly use medical imaging to quantify individual muscle size. As a muscle’s cross-sectional area (CSA) and force production are strongly correlated, the CSA can be used to estimate muscle strength [19,20].

Magnetic resonance imaging (MRI) is considered the method of choice, but MRI is neither cost-effective nor feasible for daily clinical practice. Ultrasonography (US) may provide a more clinical friendly and portable alternative [21]. However, ultrasound imaging is highly dependent on the sonographer’s level of technical skill and professional background. Furthermore, patient position, probe location and gain, depth or frequency settings may influence image capturing [22–24]. Medical and paramedical staff should be informed about this diversity in protocols when choosing the most appropriate one for a given specific objective. Determining reliability and repeatability is therefore highly relevant since it provides the basis for establishing reference values.

This systematic review aims to (1) provide an overview of 2-dimensional morphology-oriented ultrasound evaluation (2D-MOUSE) protocols and their reliability and repeatability; and (2) put forward reference values concerning cross-sectional area and dorsoplantar thickness of the abductor hallucis, flexor digitorum brevis, flexor hallucis brevis and quadratus plantae.

2. Methods

This systematic review was written according to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) checklist [25] and followed the methods of the Cochrane Handbook for Systematic Reviews of Interventions [26] where appropriate. The protocol was prospectively registered in the PROSPERO database (CRD42021286578).

2.1. Search strategy

The literature search was conducted during the first half of November 2021. An experienced biomedical research librarian aided in developing a sensitive search strategy. Eight different electronic databases were used: PubMed, Embase, Web of Science Core Collection, Cochrane Library, Scopus, CINAHL, SPORTdiscus and EuropePMC. Search terms were related to “ultrasonography” and “intrinsic foot muscles” and their synonyms. Search equations are provided in **Appendix 1**. Reference lists of each included study and similar reviews were screened for additional papers.

2.2. Study selection

In order to perform this systematic review, the literature search was performed using the PICO method [27]. The following inclusion criteria were set for approaching the problem:

- Population - Adults, aged 18–65 years [28], symptomatic or asymptomatic persons with no limitation to sex, BMI, ethnicity or physical activity.
- Intervention - Protocols using 2D morphology-oriented ultrasound evaluation (2D-MOUSE) of the intrinsic foot muscles.
- Comparison – None.
- Outcome measures – 2D morphology measures of intrinsic foot muscles: cross-sectional area and dorso-plantar thickness, as determined by B-mode ultrasound.

Papers were excluded for the following reasons: 1) publication type: opinion, report, meeting summary, symposium summary, letters to the editor, commentaries, guidelines, editorials, abstracts 2) exclusive use of following ultrasound modalities: three-dimensional, Doppler, elastography, ultrasound palpation, M-mode or therapeutic ultrasound 3) population: <18 or >65 years and 4) language: other than English.

The search results were imported in EndNote Library, where a deduplication process was performed following Bramer et al. [29]. After this process, the eligible papers were exported to Rayyan [30], an online collaborative tool to review papers. Blinded for each other’s decisions, the eligibility assessment was performed by two reviewers (NH and JLPD) during the month of November 2021.

After each reviewer completed their selection, blinding was turned off. Conflicted papers were resolved through consensus. If not, a third reviewer (KD) made a final decision.

2.3. Methodological quality assessment

The Downs and Black Checklist [31] was used to assess the methodological quality of the non-interventional studies, this was done independently by two reviewers (NH, JLPD) [31]. It addresses completeness of reporting (ten items), external validity (three items), internal validity (thirteen items) and power (one item). For single group studies, items 5, 21 and 22 do not apply. Item 27, regarding the power of the study, was transformed into a dichotomous scale indicating whether or not a sample size calculation was reported. Nine items are not applicable for the assessment of intervention studies (items 4, 8, 13, 14, 17, 19, 23, 24 and 26). To allow comparison between the categories of studies and adjusting for the varying total items scored, the results are expressed as percentage scores. Percentage scores of >75% are considered strong methodological quality, a score of 50–74% is considered moderate, a score between 25 and 49% is considered limited and a score <25% is considered poor [9].

Quality assessment was performed to provide an overview, not to in- or exclude. In case of continued disagreement, a third reviewer was available for arbitration (KD).

2.4. Data extraction

The following data of each study were extracted independently by two reviewers (NH and JLPD) in a self-made data extraction sheet in Excel: participant characteristics, cross-sectional area and thickness of the abductor hallucis (ABH), flexor digitorum brevis (FDB), abductor digiti minimi (ADM), flexor hallucis brevis (FHB), quadratus plantae (QP), adductor hallucis (ADH), lumbricals (LUM), extensor digitorum brevis (EDB) and muscles of the first interstitial space (MIL). In addition, the reliability properties of every 2D MOUSE protocol were included in the data extraction sheet; limits of agreement (LoA), standard error of the mean (SEM), minimal detectable change (MDC) and intra-class correlation coefficients (ICC). We decided to use the following qualitative benchmarks: ICC values less than 0.5, between 0.5 and 0.75, between 0.75 and 0.9, and greater than 0.90 are indicative of poor, moderate, good, and excellent reliability, respectively [32]. Extraction sheet is available in **Appendix 2**.

2.5. Data synthesis

We pooled the morphology measures from asymptomatic persons for the most commonly scanned muscles: Abductor Hallucis




Fig. 1. PRISMA flowchart illustrating the process for the selection of the included articles for the systematic review.

(ABH), Flexor Digitorum Brevis (FDB), Flexor Hallucis Brevis (FHB) and Quadratus Plantae (QP). Descriptive statistics were calculated based on the extraction sheet provided in **Appendix 2** and expressed as the pooled mean $\pm$ two standard deviations to obtain a set of reference values for the CSA and thickness of the ABH, FDB, FHB and QP. We used the means from (i) asymptomatic studies, (ii) control groups from symptomatic studies and (iii) baseline measurements from intervention studies involving asymptomatic persons. If a study did not provide a single mean (e.g. a study where means of both the right and left foot of the same individual were provided), an author-based mean calculation was performed, as indicated by an asterisk (*).

Table 1

Characteristics of studies involving only asymptomatic participants.

Study with Downs & Black score	Participants						2D-MOUSE	
	Activity level		Sex (M/F)	Age (yrs)	BMI ^(*) (kg/m ²)		Muscle	Outcome measure
Abe et al., 2016 [41]	57%	Sports activity min. 2/week: judo, resistance exercise, soccer, running, canoeing (n = 34)	17/17	24 (4)	22.6 (3.4)		ABH, FDB	CSA
Abraham et al., 2020 (a) [34]	71%	UTD (n = 65)	24/41	21–82	UTD		ABH, ADM, EDB	Thickness
Battaglia et al., 2016	64%	UTD (n = 34)	9/17	25.5 (3.8)	28* (7.8)		ABH, FDB, QP	CSA, Thickness
Cameron et al., 2008 [42]	64%	UTD (n = 30)	10/20	28.2 (10.2)	23.5*		ABH	CSA, thickness
Croft et al., 2014 [23]	64%	UTD (n = 10)	4/6	29.1 (7.2)	25.5 (4.8)		ABH, FDB	CSA & Thickness
Frannetovich Smith et al., 2019 [43]	71%	UTD (n = 24)	(–)	31 (9)	24.8 (3.2)		ABH, FIS	CSA & Thickness
Fraser et al., 2018 [44]	57%	Recreationally active, measured with Godin Leisure Time Questionnaire (n = 24)	12/12	21.5 (4.8)	23.5 (2.9)		ABH, FDB, FHB, QP	CSA & Thickness
Hing et al., 2009 [45]	64%	UTD (n = 30)	10/20	28.2 (10.2)	23.5*		ABH	CSA, Thickness
Johnson et al., 2020 [46]	57%	UTD (n = 12)	8/4	23.5 (1.9)	22.7*		ABH, FDB, FHB, QP	CSA: ABH, FDB, QP Thickness: FHB
Koyama et al., 2019 [39]	76%	Elite judo athletes, holding a black belt for >6 years (n = 24) Control, physically active: 2h per session, >2/week (n = 24)	24/0	19.8 (1.3)	25.5 (2.1)		ABH, FDB, FHB, QP	Thickness
Latey et al., 2018 [22]	79%	UTD (n = 21)	6/15	20.2 (1.7)	23.7 (2.6)			
Maeda et al., 2021 [47]	79%	At least 150 min of moderate activity per week for at least 6 months (n = 27)	27	39.5 (10)	23.8 (3.3)		ABH, FHB	CSA
Mickle et al., 2013 [24]	64%	UTD (n = 10)	5/5	21.9 (1.9)	21.7 (2.3)		ABH, FDB, FHB	CSA & Thickness
Mickle et al., 2016 [48]	72%	UTD (n = 41)	19/22	32.1 (10.1)	UTD		ABH, FDB, FHB, ADM, QP	CSA & Thickness
Morikawa et al., 2021 [36]	71%	At least 150 min of moderate activity for at least 6 months (n = 23)	UTD	28.8 (8.2)	26.0 (4.1)		ABH, FDB, FHB, ADM, QP	CSA & Thickness
Seok et al., 2016 [33]	79%	UTD (n = 80)	40/40	23.3 (1.9)	20.7 (2.2)		ABH, FDB, FHB	CSA & Thickness
Tanaka et al., 2019 [40]	59%	Well-trained sprinters, sprint training minimum 5/week (n = 26) Non-sprinters, recreationally active (n = 26)	26/0	39.5 (11)	22.5 (2.3)		ABH, EDB	Thickness
Taş et al., 2019 (a) [49]	64%	UTD (n = 41)	0/41	21.1 (2.1)	21.7*		ABH, FDB, FHB	Thickness
Taş et al., 2020 [50]	79%	Sedentary, no exercise in last 6 months (n = 40)	0/40	21.9 (1.4)	22.2*			
Verhulst et al., 2011 [51]	71%	UTD (n = 60)	(–)	26.9 (3.73)	21.9 (2.9)		ABH, FDB, FHB	CSA & Thickness
Zhang et al., 2017 (a) [52]	57%	Recreational runners, min 10 km/week (n = 28)	18/13	26.2 (5.1)	21.3 (2.2)		ABH, FDB, FHB	CSA & Thickness
				(–)	(–)		ABH, FDB	Thickness
							ABH, FDB, QP	CSA: ABH, FDB Thickness: ABH, FDB, QP

Abbreviations: M = male, F = female, yrs = years, kg = kilograms, m = metres, UTD = unable to determine, IFM = intrinsic foot muscles, ABH = abductor hallucis, FDB = flexor digitorum brevis, FHB = flexor hallucis brevis, QP = quadratus plantae, EDB = extensor digitorum brevis, FIS = first interstitial space, * = author-based calculation of BMI.

Table 2

Characteristics of studies involving symptomatic subjects or subjects with a different foot posture.

Study with Downs & Black score		Participants			BMI (*) (kg/m ²)	Significant differences (p < 0.05)
		Characteristics	Sex (M/F)	Age (yrs)		Mean (±SD)
Abraham et al., 2020 (b) [53]	76%	SMA patients (n = 31)	13/18	32	23.8 (21.9,	FDI: 0.66 (0.52, 0.78) ADM: 0.67 (0.56, 0.82) ABH: 0.99
		ALS patients (n = 91)	51/40	(24–42)	28.9)	(0.81, 1.13) EDB: 0.59 (0.41, 0.70)
		Asymptomatic (n = 59)	16/35	66	24.2 (21.8,	FDI: 0.79 (0.57, 0.97) ADM: 0.84 (0.70, 0.98) ABH: 0.98
				(54–72)	27.3)	(0.87, 1.08) EDB: 0.50 (0.29, 0.66)
Aiyer et al., 2015 [54]	82%	HV 20–44 yrs (n = 21)	7/14	46	23.5 (21.6,	FDI: 1.10 (0.99, 1.26) ADM: 0.97 (0.85, 1.07) ABH: 1.17
		HV 45–64 yrs (n = 17)	1/16	(31–60)	28.2)	(1.03, 1.34) EDB: 0.74 (0.67, 0.86)
		HV 65+ yrs (n = 21)	2/19	26.0	24.5 (3.3)	ABH CSA: 3.53 (6.71) Th: 1.38 (0.21)
				(5.9)	27.0 (5.4)	ABH CSA: 3.16 (6.88) Th: 1.24 (0.21)
Calvo Lobo et al., 2016 [55]	84%	HV (n = 20)	4/16	54.2	28.4 (8.1)	ABH CSA: 2.89 (9.73) Th: 1.19 (0.23)
		Control (n = 20)	2/18	(4.8)		
				76.4		
				(10.1)		
Stewart et al., 2013 [56]	79%	HV grade 1–3 (n = 64)	52/12	46.2	24.7 (3.38)	ABH CSA: 2.22 (0.49) Th: 0.91 (0.23); FHB CSA: 1.57 (0.41)
		Control (n = 64)		(11.3)	22.9 (3.50)	Th: 0.93 (0.14)
				42.1		ABH CSA: 2.74 (0.64) Th: 1.10 (0.26); FHB CSA: 2.13 (0.65)
				(12.2)		Th: 1.09 (0.18)
Taş et al., 2019 (c) [57]	61%	HV grade 1–3 (n = 64)	52/12	60.3	26.3 (5.2)	ABH CSA grade 1: 3.0 (0.46) grade 3: 2.73 (0.62) ABH Th
		Control (n = 64)		(20.5)		grade 2: 1.14 (0.16) grade 3: 1.13 (0.17)
		HV (n = 30)	3/27	36.5	23.5 (3.2)	ABH CSA: 3.39 (0.56) Th: 1.33 (0.2)
		Control (n = 30)		(12.4)	24.0 (3.3)	ABH CSA: 2.1 (0.3) Th: 0.97 (0.21); FHB CSA: 2.3 (0.4) Th:
Angin et al., 2014 [58]	71%	Pes planus (n = 49)	29/20	36.5	23.8 (3.8)	1.4 (0.14); FDB CSA: 2.1 (0.4) Th: 0.97 (0.12)
		Control (n = 49)	29/20	(10.5)	23.6 (3.6)	ABH CSA: 2.4 (0.5) Th: 1.14 (0.22); FHB CSA: 2.7 (0.5) Th:
				24.1		1.65 (0.20); FDB CSA: 1.9 (0.6) Th: 0.89 (0.18)
				(5.5)		
Angin et al., 2018 [59]	71%	Pes planus (n = 43)	25/18	23.4	23.8 (3.8)	ABH CSA: 2.36 ± 0.47 Th: 1.18 ± 0.11; FHB CSA: 2.66 ±
		Control (n = 68)	36/32	(4.2)	23.2 (3.4)	0.48 Th: 1.30 ± 0.18
				23.7	23.3 (3.2)	ABH CSA: 2.75 ± 0.35 Th: 1.27 ± 0.10; FHB CSA: 2.97 ±
				(4.8)		0.47 Th: 1.43 ± 0.21
Okamura et al., 2021 [60]	78%	Pronators (n = 13)	4/9	24.7	20.4 (1.5)	(–)
		Overpronators (n = 13)	4/9	(6.3)	20.4 (1.7)	
				20.5		
				(1.7)		
Sakamoto et al., 2020 [61]	61%	Flat foot (n = 43)	25/18	21.7	21.2*	ABH CSA: 2.36 ± 0.51 Th: 1.28 ± 0.19; ADM CSA: 0.95 ±
		Typical feet (n = 34)	14/17	(3.2)	21.2*	0.25 Th: 0.66 ± 0.11
				20.9		ABH CSA: 2.14 ± 0.64 Th: 1.12 ± 0.19; ADM CSA: 1.15 ±
				(0.4)		0.33 Th: 0.82 ± 0.11
Taş et al., 2018 [62]	78%	Flat foot (n = 40)	20/60	26	23 (20–25)	ABH Th: 1.22 (1.07–1.34)
		Control (n = 40)		(21–33)	22 (21–25)	ABH Th: 0.99 (0.89–1.10)
				27		
				(22–37)		
Zhang et al., 2017 (b) [63]	67%	Asx overpronators (n = 9)	15/11	22.6	21.8 (1.7)	ABH Th: 1.29 (0.08); ADM CSA: 1.89 (0.18) Th: 0.93 (0.06);
		Normal posture (n = 17)		(1.9)	21.8 (2.6)	FDB CSA: 2.55 (0.35)
				25.9		ABH Th: 1.20 (0.12); ADM CSA: 2.11 (0.21) Th: 1.06 (0.08);
				(6.4)		FDB CSA: 2.15 (0.41)
Zhang et al., 2019 [64]	61%	Sx pronators (n = 15)	9/6	23.3	22.6 (1.6)	ABH CSA: 2.28 ± 0.48 Th: 1.10 ± 0.13
		Asx pronators (n = 15)	10/5	(5.1)	22.7 (1.9)	ABH CSA: 2.71 ± 0.35 Th: 1.25 ± 0.07
				25 (5.3)		
				(9.7)		
Henderson et al., 2020 [38]	83%	DM + NP (n = 15)	15/0	61.4	33*	ABH CSA: 1.40 ± 0.76; FDB CSA: 1.42 ± 0.47; QP CSA:
		Control (n = 15)	15/0	(12.4)	29.6*	1.16 ± 0.30; FHB Th: 1.36 ± 0.18
				61.6		ABH CSA: 1.96 ± 0.47; FDB CSA: 2.16 ± 0.35; QP CSA:
				(9.7)		1.72 ± 0.71; FHB Th: 1.57 ± 0.21
Kumar et al., 2015 [65]	65%	DM + PN (n = 30)	13/5	56.2	23.6 (2.7)	EDB CSA: 1.72 ± 0.42; LUM1 Th: 1.29 ± 0.44; INT1 Th:
		Control (n = 30)	17/13	(6.6)	24.6 (1.7)	1.29 ± 0.45
				54.4		EDB CSA: 2.17 ± 0.42; LUM1 Th: 1.57 ± 0.50; INT1 Th:
				(9.8)		1.57 ± 0.50
Severinsen et al., 2007 [66]	72%	DM (n = 26)	16/10	50	23.5*	EDB CSA: 1.16 ± 0.65; EDB Th: 0.64 ± 0.21 MIL Th: 2.96
		Control (n = 26)	16/10	(26–64)	25.2*	± 0.83
				49		EDB CSA: 1.16 ± 0.65; EDB Th: 0.9 ± 0.10 MIL Th: 4.02 ±
				(25–67)		0.36

(continued on next page)

Table 2 (continued)

Study with Downs & Black score	Participants					Significant differences ($p < 0.05$) Mean ($\pm$ SD)
	Characteristics	Sex (M/F)	Age (yrs)	BMI (*)(kg/m^2)		
Wang et al., 2014 [67]	56% T2DM + PN (n = 56) T2DM without PN (n = 50) Control (n = 50)	30/26 26/24 25/25	63 (7) 59 (10) 59 (7)	UTD		EDB CSA - Th $\downarrow$ FIS Th $\downarrow$ in both groups compared to control
Calvo Lobo et al., 2018 (a) [68]	89% Hemiparesis foot Contralateral foot (n = 11)	4/7	62 (5)	19.5 (7.9)	(–)	
Calvo Lobo et al., 2018 (b) [69]	89% Hemiparesis feet (n = 20) Contralateral feet (n = 20) Control feet (n = 20)	10/10 10/10 8/12	61.5 (11.5) 61.5 (11.5)	21.6 (4.1) 21.6 (4.1) 22.6 (3.3)		FHB Th $\uparrow$ in hemiparesis and contralateral feet compared to control feet
Fraser et al., 2021 [70]	57% CAI (n = 20) LAS (n = 17) Copers (n = 21) Control (n = 22)	5/15 8/9 8/13 9/13	20.9 (4.7) 21.8 (4.1) 20.8 (2.9) 19.6 (0.9)	25.1 (4.5) 24.1 (3.7) 23.7 (2.9) 22.5 (3.2)		No between-group differences in resting CSA
Hogan et al., 2020 [71]	78% Plantar heel pain (n = 16) Matched controls (n = 16)	3/13 3/13	25.0 (2.2) 26.06 (1.7)	24.5* 24.1*	(–)	
Jaffri et al., 2019 [72]	76% MTP I arthrodesis Contralateral feet (n = 9)	3/6	57.56 (9.0)	30.6*		ABH CSA - Th $\downarrow$ FHB CSA - Th $\downarrow$ FDB - Th $\downarrow$ FDB Th $\downarrow$
Pompeo et al., 2021 [73]	72% Patellofemoral pain (n = 20) Control (n = 20)	0/20 0/20	30.0 (5.6) 28.5 (5.1)	21.8 (2.4) 21.8 (1.3)		
Romero-Morales et al., 2019 [74]	72% Achilles tendinopathy (n = 71) Control (n = 70)	(–)	45 (13) 35 (18.7)	24.8 (2.1) 23.88 (3.6)		ABH Th $\uparrow$ FDB CSA - Th $\uparrow$ FHB CSA $\uparrow$ FHB Th $\downarrow$

Abbreviations: M = male, F = female, yrs = years, kg = kilograms, m = meters, CSA = cross-sectional area, Th. = thickness, IFM = intrinsic foot muscles, ABH = abductor hallucis, FDB = flexor digitorum brevis, FHB = flexor hallucis brevis, QP = quadratus plantae, EDB = extensor digitorum brevis, FIS = first interstitial space, LUM1 = lumbrical 1, INT1 = interosseus 1, FDI = first dorsal interosseus, ADHO = oblique head of adductor hallucis, SMA = spinal muscular atrophy, ALS = amyotrophic lateral sclerosis, DM = diabetic mellitus, T2DM = type 2 diabetic mellitus, PN = polyneuropathy, CAI = chronic ankle instability, LAS = lateral ankle sprain * = author-based calculation of BMI.


3. Results

3.1. Study selection

The initial search yielded 2075 records. After removing duplicates, this was reduced to 968 records. Articles were then screened by title, leaving 124 records eligible for full-text analysis. After applying the selection criteria, 53 papers were included for qualitative synthesis. No additional authors were contacted. A flow diagram is summarised in Fig. 1.

3.2. Characteristics of included studies

Descriptive data of asymptomatic participants are presented in Table 1, whilst descriptive data of symptomatic participants are included in Table 2. This also includes participants with a different foot posture, not necessarily with symptoms. A total of 21 studies involving asymptomatic participants have been retrieved, with a sample size ranging from 10 [23] to 80 participants [33]. Age of the asymptomatic participants ranges between 20 and 82 years. Abraham et al. did not include an average age but an age range, from 21 to 82 years [34], with six subjects older than 70 years. We included this study as the authors themselves state that only 6 out of 65 participants had an age above 70. BMI of all participants was normal [35], with the lowest BMI being 20.7 kg/m^2 [36] and highest BMI 28 kg/m^2 [37]. One study exceeded this significantly, with an average BMI of 29.6 kg/m^2 [38]. Nevertheless, it was taken for synthesis. Two studies assessed the IFM in elite sport athletes. Koyama et al. [39] assessed muscle thickness of ABH, FDB, FHB and QP in elite judo athletes and found no significant difference with matched physically active subjects. Tanaka et al. [40] found larger foot muscle thicknesses in well-trained male sprinters compared to non-sprinters. However, despite greater foot muscularity, muscle thickness was



(caption on next page)

Fig. 2. Reference values for cross-sectional area of the (a) abductor hallucis (c) flexor digitorum brevis (e) flexor hallucis brevis (g) quadratus plantae and dorsoplantar thickness of the (b) abductor hallucis (d) flexor digitorum brevis (f) flexor hallucis brevis and (h) quadratus plantae. Athletic populations are indicated with a square in front of the author's name.

not positively related to sprint performance. As an athletic population is characterized by muscle properties that may deviate from the general population, we have indicated this population with a square in Fig. 2.

Studies involving symptomatic participants exhibited a variety of foot pathologies and lower limb overuse injuries including neuromuscular diseases [53], hallux valgus [54,56,57,75], pes planus [58,60–62], diabetic neuropathy [38,65–67], hemiparesis [69, 68], ankle sprain/instability [70], plantar heel pain [71], first MTP joint arthrodesis [72], patellofemoral pain [73] and Achilles tendinopathy [74].

3.3. Methodological quality assessment

A total of 53 studies were assessed against the Downs's and Black checklist and attained a score of either moderate or high quality. Of the 53 articles, a total of 17 were of high quality (score >75%), the other 36 articles were of moderate quality (score: 50–74%). The average score was 71% and highest score 89% [69]. The majority of studies clearly reported inclusion/exclusion criteria and procedures for measuring the IFM. External validity was poor, with only a few studies stating the proportion of participating subjects compared to the total invited population [33,56,76,50]. Internal bias was low, with a total of 29 studies where the assessor was blinded to the majority of measures and procedures. Full quality assessment can be retrieved in **Appendix 3**.

3.4. Summary of results

3.4.1. 2D-MOUSE protocols

A plethora of 2D-MOUSE protocols have been used so far and an overview of the reference papers is presented in **Table 3** whereas **Table 4** provides an overview of their reported reliability measurements. **Table 5** represents the intervention studies. We presented the reliability measures as described in the respective study. We were able to identify six studies that have developed and used reliable 2D-MOUSE protocols for measuring the IFM. Other studies were primarily based on these six protocols. The first study reporting reliability outcome measures was by Cameron et al., in 2008 [42]. They evaluated within and between-session reliability of ABH dorso-plantar thickness and CSA. Within-session reliability demonstrated subsequently excellent ICC's of 0.97 (95% CI: 0.99–0.99) for thickness and 0.98 (95% CI: 0.98–0.99) for CSA. Between-session reliability demonstrated excellent ICC's of 0.97 (95% CI: 0.95 to 0.98) for thickness and a good ICC of 0.79 (95% CI 0.65 to 0.88) for CSA.

In 2013, Mickle et al. proposed a protocol to identify thickness and CSA of ABH, FDB, FHB, ABDM and QP. This protocol was found to have good to excellent intra-tester reliability (ICC = 0.89–0.99), with limits of agreement between 8 and 30% of the relative muscle size [24]. However, aforementioned protocols only report intra-operator reliability. Therefore, Crofts et al. proposed a novel protocol and reported inter-rater reliability, with excellent ICC's for muscle thickness (ICC range 0.90–0.97) and CSA (ICC range 0.91–0.98) [23]. They report limits of agreement of 13–17% of relative muscle size. This protocol however only encompasses CSA and thickness measurements of the ABH, FDB and FHB. In 2018, Fraser et al. established test-retest reliability at rest and during active, resisted contraction and during IFM exercises. Reliability of ABH, FDB, QP, and FHB CSA and thickness resting measures were good to excellent

Table 3

Overview of current relaxed, non-weight bearing ultrasound protocols.

Author	Examiner	Device & Probe	Frequency	Number of images	Days between measures
Cameron et al., 2008	Novice researcher with training in US imaging over 3 months	Philips HD11 Ultrasound machine; Linear, 50 mm	5–12 MHz	3	3–7
Mickle et al., 2013	Chief investigator, with 8 years of experience in musculoskeletal ultrasound	Venue 40 portable musculoskeletal ultrasound system (GE Healthcare, United Kingdom); Linear	6–10.7 MHz	3	1–6
Crofts et al., 2014	One experienced (8 years) and one inexperienced (4 weeks) operator	Venue 40 portable musculoskeletal ultrasound system (GE Healthcare, UK); Linear, 12.7 mm × 47.1 mm	5–13 MHz	3	n.a.
Battaglia et al., 2016	2 years and 4 years of experience	GE Logiq E9 (GE Healthcare, Wauwatosa, WI); Linear (ML6-15)	9–12 MHz	/	/
Fraser et al., 2018	Physical therapist with 14 years of practice experience and 2 months of USI experience	Siemens acuson freestyle US system (Siemens, Mountain View, CA); Linear 3.8 cm	8 MHz	A previously measured US image was used for reference in identification of fascial borders and spatial orientation of each muscle	/
Franettovich Smith et al., 2019	1 year and 8 years of experience	LOGIQ S7 Expert (GE Healthcare); Linear	4.2–12 MHz	/	6.8

Abbreviations: MHz = Megahertz, LoA = limits of agreement.

Table 4

Overview of reliability measures of current relaxed, non-weight bearing 2D ultrasound protocols for the intrinsic foot muscles.

Author	Population	Outcome parameter		Reliability measures				Limits of agreement (LoA)	Standard error of the mean (SEM)	Minimal detectable change (MDC)
				Intraclass Correlation Coefficients (ICC's)			Interrater			
				Intrarater						
				Within-session	Between session	Test-retest				
Cameron et al., 2008 [42]	Asymptomatic	ABH	CSA Thickness	0.97 (0.99–0.99) 0.98 (0.98–0.99)	0.79 (0.65–0.88) 0.97 (0.95–0.98)	n/a	n/a	n/a		
Mickle et al., 2013 [24]	Asymptomatic	ABH	CSA	n/a	n/a	0.95	n/a	30%		
		FDB	Thickness			0.95		18%		
		FHB	CSA			0.99		8%		
		ABDM	Thickness			0.95		16%		
		QP	CSA			0.89		28%		
			Thickness			0.97		17%		
			CSA			0.98		16%		
			Thickness			0.97		18%		
			CSA			0.99		9%		
			Thickness			0.97		18%		
Croft et al., 2014 [23]	Asymptomatic	ABH	CSA	n/a	n/a	n/a	0.91	16%		
		FDB	Thickness				0.92	13%		
		FHB	CSA				0.98	17%		
			Thickness				0.96	13.5%		
			CSA				0.95	17%		
Battaglia et al., 2016 [37]	Asymptomatic	ABH	CSA	0.97 (0.94–0.99)	n/a	n/a	0.95 (0.88–0.98)	n/a		
		FDB	Thickness				0.72 (0.12–0.89)			
		QP	CSA	0.95 (0.88–0.96)			0.92 (0.78–0.96)			
			Thickness	0.97 (0.93–0.99)			0.74 (0.29–0.89)			
			CSA	0.85 (0.70–0.93)			0.77 (0.49–0.90)			
			Thickness	0.92 (0.83–0.96)			0.57 (–0.19–0.84)			
			CSA	0.35 (–0.03–0.65)						
		ABH	CSA	n/a	n/a	0.97–0.98	n/a	n/a	0.09 cm ²	0.26 cm ²
		FDB	Thickness			0.88–0.91			0.07 cm	0.18 cm
		FHB	CSA			0.91–0.93			0.13 cm ²	0.36 cm ²
Fraser et al., 2018 [44]	Asymptomatic	QP	Thickness			0.87–0.89			0.06 cm	0.15 cm
			CSA			0.95–0.98			0.14 cm ²	0.40 cm ²
			Thickness			0.76–0.83			0.09 cm	0.24 cm
			CSA			0.97–0.98			0.09 cm ²	0.25 cm ²
			Thickness			0.90–0.92			0.05 cm	0.14 cm
		ABH	CSA	0.99	0.97	n/a	0.98	n/a	0.06–0.13 cm ²	0.10–0.29 cm ²
Franetovich Smith et al., 2019 [43]	Asymptomatic	FIS	Thickness	0.97	0.92		0.85		0.03–0.08 cm	0.10–0.23 cm
			Thickness	0.98	0.81		0.86		0.05–0.17 cm	0.14–0.48 cm

Abbreviations: SEM = standard error of the mean; MDC = minimal detectable change, CSA = cross-sectional area, Th. = thickness, ABH = abductor hallucis, FDB = flexor digitorum brevis, FHB = flexor hallucis brevis, QP = quadratus plantae, n/a = not applicable.

Table 5

Summary of intervention studies using 2D-MOUSE outcome measures.

Study	Participants	Intervention group	Control group	2D-MOUSE outcome measures
Campitelli et al., 2016 [77]	Healthy individuals (n = 41)	Minimalist shoes: Restricted walking (n = 11); Unlimited walking (n = 11); Running (n = 10); 24 weeks	Traditional shod (n = 9)	ABH Thickness
Johnson et al., 2015 [78]	Recreational runners (n = 37)	Transitioning to minimalist shoes during 10 weeks (n = 18)	Traditional running shoes (n = 19)	CSA of ABH and FDB
Ridge et al., 2018 [79]	Recreational runners (n = 57)	Minimalist shoe walking (n = 19)	No intervention (n = 19)	Thickness of FHB and EDB
Reeves et al., 2021 [80]	Healthy participants (n = 41)	Foot strengthening 8 weeks (n = 19)	No intervention (n = 18)	CSA of ABH, FDB and QP
Protopapas et al., 2020 [81]	Pes planus (n = 18)	Prefabricated orthoses, 3 months (n = 23)	No intervention (n = 9)	Thickness of FHB
Kim et al., 2015 [82]	Hallux valgus (n = 24)	Custom-made foot orthotics for 12 weeks (n = 9)	Orthosis only (n = 12)	CSA of ABH and FDB
Jung et al., 2011 [83]	Pes planus (n = 28)	Orthosis + toe-spread-out exercise, 8 weeks, 20min/day, 4 day/week (n = 12)	Orthosis only (n = 14)	CSA of ABH, FDB and QP
Namsawang et al., 2019 [84]	Flexible flatfoot (n = 36)	Orthosis + short foot exercise, 8 weeks, 3 × 5 reps, 2x/day, everyday (n = 14)	Short foot exercise + placebo NMES	Thickness of ABH and FDB
Okamura et al., 2020 [85]	Pes planus (n = 20)	Short foot exercise + NMES, 4 weeks	Control group	Thickness of ABH, FDB and FHB
		Short foot exercise, 8 weeks, 3 day/week		CSA of ABH
				CSA of ABH
				CSA of ABH
				Thickness of FDB and FHB

Abbreviations: ABH = abductor hallucis, FDB = flexor digitorum brevis, FHB = flexor hallucis brevis, EDB = extensor digitorum brevis, AbDM = abductor digiti minimi, CSA = cross-sectional area.

(0.76–0.98). Active and resisted measures and IFM exercises had moderate to excellent reliability (0.66–0.99). Standard error of measurement (SEM) of resting CSA thickness measures ranged from 0.09 to 0.14 cm² and 0.05–0.09 cm. Minimal detectable change (MDC) for CSA resting measures ranged from 0.25 to 0.40 cm² and 0.14–0.24 cm for thickness measures.

As the IFM function primarily during weight-bearing [1], Battaglia et al. proposed both a weight-bearing and non-weight-bearing protocol to measure CSA and thickness of the ABH, FDB and QP muscles [37]. Inter-rater reliability was moderate to good for all CSA measurements (ICC 0.57–0.95) and poor to moderate in thickness measurements (ICC range 0.49–0.74). However, a custom-built platform is required so the participant can support his/her weight while permitting the researcher access to image the medial and plantar foot muscles with an US probe. In response, Franettovich Smith et al. developed a more clinically feasible protocol in weight bearing while the foot is in contact with the ground. They demonstrated excellent within-session reliability (ICC > 0.94) and good between-session reliability (ICC > 0.81), as well as interrater reliability (ICC > 0.82) [43]. It is suggested that changes of 10–18% should be considered to exceed measurement error.

At the time of searching, no study was found reporting the validity of 2D-MOUSE measurements involving the IFM.

3.4.2. Reference values

2D-MOUSE measurements from ABH, FDB, FHB and QP are expressed as the pooled mean $\pm$ 2 standard deviations (SD) and is presented in Fig. 2. Total participants considered for pooled ABH CSA measurement were 781, for pooled FDB CSA measurement 595, for pooled FHB CSA measurement 465 and for pooled QP CSA measurement 124. Pooled samples included 1033 individuals for ABH thickness, for pooled FDB thickness measurement 647, for pooled FHB thickness measurement 495 and for pooled QP thickness measurement 127.

Based on the number of participants and each study mean, pooled means $\pm$ 2 SD for CSA measurements of the IFM were: ABH: 2.43 cm² $\pm$ 0.78, FDB: 2.01 cm² $\pm$ 0.34, FHB: 2.36 cm² $\pm$ 1.2 and QP: 1.75 cm² $\pm$ 0.54. When measuring muscle thickness, reference values of the IFM were: ABH: 1.11 cm $\pm$ 0.28, FDB: 0.87 cm $\pm$ 0.28, FHB: 1.38 cm $\pm$ 0.44 and QP: 0.98 cm $\pm$ 0.12.

3.4.3. Clinical outline in symptomatic populations

Table 2 provides an overview of studies where the morphology of the IFM was investigated in symptomatic individuals as providing reference values was not possible in the symptomatic population due to the limited number of studies. Significant differences are presented in relaxed, unloaded position.

Abraham et al. investigated relaxed and contracted muscle thickness in patients with amyotrophic lateral sclerosis and spinal muscular atrophy and found a smaller thickness in ADM ($p < 0.001$) and first dorsal interosseus space ($p < 0.001$) [53].

Four studies could be retrieved in subjects with hallux valgus [54,56,57,55]. Taş et al. investigated morphological features of ABH, FDB, FHB and found significantly smaller thickness and CSA of ABH ($p = 0.002$, $p = 0.003$) and FHB ($p < 0.001$, $p = 0.001$) in the hallux valgus population [57]. On the contrary, FDB thickness and CSA were larger (respectively, $p = 0.027$ and $p = 0.006$) [57]. Similar results were found by Calvo Lobo et al. where smaller thickness and CSA of ABH ($p = 0.02$, $p < 0.01$) and FHB ($p < 0.01$, $p < 0.01$) are reported. However, no significant difference in FDB thickness and CSA ($p = 0.19$, $p = 0.14$) was reported [55]. Aiyer et al. compared morphology of the ABH in hallux valgus between a 20–44 years old age group, a 45–64 years old age group and a 65+ age group. Increasing age was significantly associated with a reduction in thickness ($r = -0.27$, $p = 0.008$) and CSA ($r = -0.24$, $p = 0.019$) [54]. Stewart et al. imaged the ABH muscle in relationship to hallux valgus severity. Significant differences in thickness ($p = 0.001$)

and CSA ($p < 0.001$) between grade 0 and grade 2 and in thickness ($p < 0.001$) and CSA ($p < 0.001$) between grade 0 and 3 were noticed [56].

A total of seven studies could be retrieved where morphology of the IFM was assessed in participants with a different foot posture: pes planus, flat foot or pronation [58,60–62,59,63,64]. Angin et al. [58] reported significantly lower values in the pes planus group, ranging from -12.8% for CSA of the ABH muscle and -8.9% for CSA of the FHB. For thickness, percentual difference for ABH was -6.8% and -7.6% for FHB. No difference in CSA or thickness measurements of the FDB muscle could be observed. Sakamoto et al. [61] found significantly larger thickness of the ABH muscle in the flat foot group ($p < 0.01$). Thickness and CSA of ABDM ($p < 0.01$, $p < 0.01$) and thickness of the oblique head of adductor hallucis were significantly smaller ($p < 0.01$) [61]. Taş et al. found larger ABH thickness ($p < 0.001$) in the flat foot group, whereas both groups were similar in terms of FHB ($p = 0.627$) and FDB thickness ($p = 0.212$) [62]. Zhang and colleagues compared recreational runners who over-pronated with runners without over-pronation and showed that runners with over-pronated feet have 7.5% larger ABH thickness and 18.7% FDB CSA, whilst also having a -10.3% smaller CSA and a -12.3% thickness of the ABDM [63]. Two years later, they compared IFM morphology between recreational runners with a pronated foot posture and overuse injuries (symptomatic) and those without overuse injuries (asymptomatic). Symptomatic pronators demonstrated 19% smaller CSA and 14% smaller thickness of the ABH than their asymptomatic counterparts ($p < 0.05$) [64]. Okamura et al. compared muscle thicknesses of FDB and FHB and found no between-group differences between a pronated group and an overpronated group [60].

Four studies could be retrieved in diabetic (poly)neuropathy [38,65–67]. Dimensions of the EDB muscle were consistently significantly smaller in all studies compared to healthy controls except for the study of Kumar et al. [65], who reported no significant differences in EDB thickness. Wang et al. [67] and Kumar et al. [65] reported a combined measurement of the first metatarsal interspace while Severinsen et al. [66] reported discrete measurements of the first interspace (first lumbrical, first dorsal interosseus and the adductor hallucis). Combined thickness of the first metatarsal space was significantly smaller in diabetic patients compared to healthy controls, and even more small in diabetic patients with neuropathy [65–67]. Recently, Henderson et al. [38] also included evaluation of the thickness of the FDB and CSA of the ABH, FDB and QP muscle and found smaller IFM in early stage diabetic neuropathy.

Two studies compared CSA and thickness in poststroke patients [69,68]. Calvo Lobo et al. (a) [68] compared CSA of ABH, FDB and FHB of the hemiparesis foot to the contralateral foot. No between-group differences in CSA were found ($p > 0.167$). Later, they compared hemiparesis and contralateral feet to control feet, and found larger FHB thickness in the hemiparesis and contralateral feet ($p < 0.01$) [69].

Fraser et al. [70] assessed IFM dimensions in ankle injuries. No difference was found in CSA of the ABH, FDB, FHB and QP between a chronic ankle instability group, a lateral ankle sprain group, copers and a control group using ultrasound measurements in a relaxed, unloaded position.

Hogan et al. [71] compared ABH morphology in individuals with and without plantar heel pain. No significant group differences were found in CSA and thickness.

Jaffri et al. [72] determined differences in the size of the ABH, FDB and FHB after arthrodesis of the 1st metatarsophalangeal joint by comparing the surgical with the contralateral asymptomatic foot during sitting and standing positions. The CSA and thickness of the ABH and FHB of the surgical foot were significantly smaller than that of the contralateral foot, and this in both positions ($p < 0.05$). For the FDB, only a significant difference was found during standing, with the surgical foot having a smaller thickness compared to the contralateral foot ($p < 0.05$).

Pompeo et al. [73] compared morphology of the ABH, FDB and FHB muscles in women with and without patellofemoral pain. Women with patellofemoral pain had smaller FDB thickness ($p < 0.01$).

Romero-Morales et al. [74] compared thickness and CSA of the ABH, FDB and FHB in persons with and without chronic midportion Achilles tendinopathy. ABH ($p < 0.001$) and FDB ($p < 0.001$) thickness, as well as FDB ($p = 0.005$) and FHB CSA ($p = 0.048$), were larger in the tendinopathy group. On the contrary, FHB muscle thickness ($p < 0.001$) was larger in the control group.

4. Discussion

We identified 53 studies that used 2D-MOUSE to assess the IFM and pooled these results to obtain reference values in asymptomatic adults. We provide a visualization of the CSA and thickness of the IFM reported in current literature. For each muscle, means and SD of each study are visualised, along with indication of the weighted mean. This provides a precise illustration of the variation in IFM dimensions across the population and a valuable resource for future research and more clinically oriented studies. Pooled sample sizes were quite variable, ranging from 127 participants for QP thickness to 1033 for ABH thickness. It should be noted that there is a high variability in dimensions of the IFM, which can mainly be explained by the heterogeneity of the different populations and the lack of normalisation for other variables: sex, height, body size, foot posture and physical activity. Another factor could be anatomical variability. This is especially true when evaluating the flexor hallucis brevis muscle, which delineation is more difficult to identify on US due to its obliquely oriented muscle fascicles.

Various measuring protocols have been put forward, with good to excellent established reliability in both an unloaded [23,24,37,42] and a loaded position [37,86]. We reported good reliability between and within operators for several US protocols. Cameron et al. [42], Mickle et al. [24], Crofts et al. [23] and Battaglia et al. [37] assess the ABH muscle in a supine position and other plantar muscles in a prone position. Frannetovich Smith et al. [43] position the participant in a sitting rather than lying position, with both feet placed on the floor, which may be more comfortable for those suffering from e.g. back problems. Weightbearing did not reduce intra- or interexaminer reliability and all IFM exhibited a significant increase in CSA [37]. In summary, we notice a general trend of good to

excellent reliability in every protocol. This despite the many variables which can influence the protocol as summarised in Table 3: the difference in education and experience of the sonographer and the particular details of the protocol itself; device and type of probe, frequency, number of images taken and days between measurements. However, we would still advocate structured training and education in operator skills.

At the time of searching, we could not retrieve a study determining the validity of US measurements for the IFM. However, recently, a study was published by Swanson et al. [87], demonstrating strong agreement between US imaging and MRI for the ABH, FDB, QP and ADM muscles, though MRI retained higher precision and reliability. This underlines that dedicated training in US imaging remains important as this will undoubtedly lead to better image capturing and more reliable measurements of the IFM.

Differences in IFM morphology have not been studied extensively in symptomatic populations. Only single studies exist investigating the morphology in ankle instability, plantar heel pain, MTP I arthrodesis, patellofemoral pain, and Achilles tendinopathy. Therefore it's yet impossible to detect a clear trend and further research in these populations is warranted. In populations with hallux valgus, pes planus, diabetic neuropathy and stroke patients more studies could be retrieved. Thickness and CSA of the ABH and FHB muscle are significantly smaller in every retrieved hallux valgus study. Calvo Lobo et al. [75] stated no significant differences in FDB muscle size, while Taş et al. [57] showed greater CSA and thickness of the FDB in individuals with hallux valgus. A higher CSA of the FDB muscle may suggest a compensation against mechanical changes in the foot, thus increasing the loading on the FDB muscle, causing hypertrophy.

4.1. Limitations

This study has several limitations. First, the last item of the Downs and Black checklist examining the power of results was simplified to a score of 0 (no sample size calculation) or 1 (sample size calculation reported). This dichotomous approach is an oversimplification of study power and is a clear limitation in the quality assessment process.

Second, a single reference value as presented here, only paints a limited picture of reality as several factors such as age, sex, BMI, athletic status and foot phenotype affect intrinsic foot muscle size [70,88]. More studies including a larger population are needed to further investigate the relation between these factors and IFM dimensions and would also allow us to determine reference values for each subpopulation. However, the number of studies investigating IFM morphology is too limited to investigate the impact of these different variables via normalisation. This would be an excellent goal for future studies, but the current manuscript aims to provide reference values for the dimensions of the intrinsic foot muscles in the populations studied up to now. Nevertheless, we are the first to provide an example of reference values for the intrinsic foot muscles and highly recommend to take these variables into close consideration when comparing to our reference values.

Thirdly, values reported here are for an asymptomatic population of recreationally active adults with cut off age at 65 and cannot be used as reference values for an elderly population. Several studies have underlined the importance of intrinsic foot muscle strength in balance and falls in older adults [89]. For this reason we placed an upper limit of 65 years and restricted our target population as we believe this might be an interesting venue for future research.

Finally, we summarised the clinical applicability of 2D ultrasound for the IFM by comparing it to a symptomatic population. These comparative studies however are cross-sectional in nature and therefore causality cannot be determined. Ultimately, evaluating the intervention studies could be a topic for another review to work out in depth. Therefore, we recommend to be careful when interpreting these data. We should also note that there are more studies using 2D ultrasound to evaluate intrinsic foot muscles than the 9 studies we have presented here. However, the outcome measures of these studies are different from what we have examined. For example, Fraser 2019 and Lee 2019 use 2D ultrasound to evaluate intrinsic foot muscles but their outcome measures are ultrasound activation ratios (contracted/relaxed) [13,14].

4.2. Clinical implications

Scientific interest into IFM function has increased significantly in the last decade since there is growing evidence that adequate IFM strength is a cornerstone of efficient sport and daily life performance. Therefore there is great merit in investigating the value of 2D-MOUSE in assessing the IFM as its low cost, non-invasiveness, accessibility and reliability are perfectly suitable in clinical practice.

5. Conclusion

Current protocols show an overall good to great reliability. We have formulated reference values for musculoskeletal ultrasonography of the ABH, FDB, FHB and QP muscles in an asymptomatic adult population. We encourage these data to be used by medical and paramedical practitioners to guide detailed assessment of the intrinsic foot musculature. In addition, we characterised clinical differences in IFM morphology in symptomatic populations. Based on current literature there is evidence of different IFM size in several foot pathologies however, limited evidence still exists in common overuse injuries such as plantar heel pain, patellofemoral pain, and Achilles tendinopathy.

Author contributions

Nicolas Haelewyn: Conceptualization, Database search, Quality evaluation, Data extraction Writing – original draft, Jean-Louis Peters Dickie: Quality evaluation, Data extraction, Filip Staes: Conceptualization, Writing – review & editing, Evie Vereecke:

Writing – reviewing & editing, Visualization Kevin Deschamps: Conceptualization, Writing – review & editing Supervision, Project Administration. All authors have made substantial contributions to all of the following: (1) the conception and design of the study, or acquisition of data, or analysis and interpretation of data, (2) drafting the article or revising it critically for important intellectual content, (3) final approval of the version to be submitted.

Data availability statement

Data included in article/supp. material/referenced in article.

Additional information

Supplementary content related to this article has been published online at [URL].

Funding

This research did not receive any specific grant from any funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors would like to thank the librarian Krizia Tuand for assisting in the design and implementation of the electronic literature search.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.heliyon.2023.e18252>.

References

- [1] L.A. Kelly, A.G. Cresswell, S. Racinais, R. Whiteley, G. Lichtwark, Intrinsic foot muscles have the capacity to control deformation of the longitudinal arch, in eng, *J. R. Soc. Interface* 11 (93) (Apr 6 2014), 20131188, <https://doi.org/10.1098/rsif.2013.1188>.
- [2] L.A. Kelly, G. Lichtwark, A.G. Cresswell, Active regulation of longitudinal arch compression and recoil during walking and running, in eng, *J. R. Soc. Interface* 12 (102) (Jan 6 2015), 20141076, <https://doi.org/10.1098/rsif.2014.1076>.
- [3] K.E. Zelik, V. La Scaleia, Y.P. Ivanenko, F. Lacquaniti, Coordination of intrinsic and extrinsic foot muscles during walking, in eng, *Eur. J. Appl. Physiol.* 115 (4) (Apr 2015) 691–701, <https://doi.org/10.1007/s00421-014-3056-x>.
- [4] P. Fiolkowski, D. Brunt, M. Bishop, R. Woo, M. Horodyski, Intrinsic pedal musculature support of the medial longitudinal arch: an electromyography study, in eng, *J. Foot Ankle Surg.* 42 (6) (Nov-Dec 2003) 327–333, <https://doi.org/10.1053/j.jfas.2003.10.003>.
- [5] P.O. McKeon, J. Hertel, D. Bramble, I. Davis, The foot core system: a new paradigm for understanding intrinsic foot muscle function, *Br. J. Sports Med.* 49 (5) (Mar 2015) 290, <https://doi.org/10.1136/bjsports-2013-092690>.
- [6] R.T. Cheung, L.K. Sze, N.W. Mok, G.Y. Ng, Intrinsic foot muscle volume in experienced runners with and without chronic plantar fasciitis, in eng, *J. Sci. Med. Sport* 19 (9) (Sep 2016) 713–715, <https://doi.org/10.1016/j.jsams.2015.11.004>.
- [7] W.P. Garth Jr., S.T. Miller, Evaluation of claw toe deformity, weakness of the foot intrinsics, and posteromedial shin pain, in eng, *Am. J. Sports Med.* 17 (6) (Nov-Dec 1989) 821–827, <https://doi.org/10.1177/036354658901700617>.
- [8] R.T.H. Cheung, L.K.Y. Sze, N.W. Mok, G.Y.F. Ng, Intrinsic foot muscle volume in experienced runners with and without chronic plantar fasciitis, in english, *J. Sci. Med. Sport* 19 (9) (2016) 713–715, <https://doi.org/10.1016/j.jsams.2015.11.004>.
- [9] D. Huffer, W. Hing, R. Newton, M. Clair, Strength training for plantar fasciitis and the intrinsic foot musculature: a systematic review, in eng, *Phys. Ther. Sport* 24 (Mar 2017) 44–52, <https://doi.org/10.1016/j.ptsp.2016.08.008>.
- [10] M.A. Feger, et al., Diminished foot and ankle muscle volumes in young adults with chronic ankle instability, in eng, *Orthop J Sports Med* 4 (6) (Jun 2016), 2325967116653719, <https://doi.org/10.1177/2325967116653719>.
- [11] R. Chang, J.A. Kent-Braun, J. Hamill, Use of MRI for volume estimation of tibialis posterior and plantar intrinsic foot muscles in healthy and chronic plantar fasciitis limbs, in eng, *Clin. Biomech.* 27 (5) (Jun 2012) 500–505, <https://doi.org/10.1016/j.clinbiomech.2011.11.007>.
- [12] K. Severinsen, H. Andersen, Evaluation of atrophy of foot muscles in diabetic neuropathy – a comparative study of nerve conduction studies and ultrasonography, *Clin. Neurophysiol.* 118 (10) (Oct 2007) 2172–2175, <https://doi.org/10.1016/j.clinph.2007.06.019>.
- [13] J.J. Fraser, J. Hertel, Effects of a 4-week intrinsic foot muscle exercise program on motor function: a preliminary randomized control trial, in eng, *J. Sport Rehabil.* 28 (4) (May 1 2019) 339–349, <https://doi.org/10.1123/jsr.2017-0150>.
- [14] D.R. Lee, Y.E. Choi, Effects of a 6-week intrinsic foot muscle exercise program on the functions of intrinsic foot muscle and dynamic balance in patients with chronic ankle instability, in eng, *J Exerc Rehabil* 15 (5) (Oct 2019) 709–714, <https://doi.org/10.12965/jer.1938488.244>.
- [15] A.B. Matias, et al., Effects of foot-core training on foot-ankle kinematics and running kinetics in runners: secondary outcomes from a randomized controlled trial, in eng, *Front. Bioeng. Biotechnol.* 10 (2022), 890428, <https://doi.org/10.3389/fbioe.2022.890428>.
- [16] A. Soysa, C. Hiller, K. Refshauge, J. Burns, Importance and challenges of measuring intrinsic foot muscle strength, *J. Foot Ankle Res.* 5 (1) (2012) 29, <https://doi.org/10.1186/1757-1146-5-29>.

- [17] M.M. de Win, W.J. Theuvsen, P.W. Roche, R.A. de Bie, H. van Mameren, The paper grip test for screening on intrinsic muscle paralysis in the foot of leprosy patients, in eng, *Int. J. Lepr. Other Mycobact. Dis.* 70 (1) (Mar 2002) 16–24.
- [18] M.J. Spink, M.R. Fotoohabadi, H.B. Menz, Foot and ankle strength assessment using hand-held dynamometry: reliability and age-related differences, in eng, *Gerontology* 56 (6) (2010) 525–532, <https://doi.org/10.1159/000264655>.
- [19] B. Mohseny, T.H. Nijhuis, C.A. Hundedoel, W.G. Janssen, R.W. Selles, J.H. Coert, Ultrasonographic quantification of intrinsic hand muscle cross-sectional area; reliability and validity for predicting muscle strength, in eng, *Arch. Phys. Med. Rehabil.* 96 (5) (May 2015) 845–853, <https://doi.org/10.1016/j.apmr.2014.11.014>.
- [20] T. Kurihara, J. Yamauchi, M. Otsuka, N. Tottori, T. Hashimoto, T. Isaka, Maximum toe flexor muscle strength and quantitative analysis of human plantar intrinsic and extrinsic muscles by a magnetic resonance imaging technique, in eng, *J. Foot Ankle Res.* 7 (2014) 26, <https://doi.org/10.1186/1757-1146-7-26>.
- [21] G.P. Kuo, J.A. Carrino, Skeletal muscle imaging and inflammatory myopathies, in eng, *Curr. Opin. Rheumatol.* 19 (6) (Nov 2007) 530–535, <https://doi.org/10.1097/BOR.0b013e3282efdc66>.
- [22] P.J. Latey, J. Burns, E.J. Nightingale, J.L. Clarke, C.E. Hiller, Reliability and correlates of cross-sectional area of abductor hallucis and the medial belly of the flexor hallucis brevis measured by ultrasound, *J. Foot Ankle Res.* 11 (2018) 28, <https://doi.org/10.1186/s13047-018-0259-0>.
- [23] G. Crofts, S. Angin, K.J. Mickle, S. Hill, C.J. Nester, Reliability of ultrasound for measurement of selected foot structures, *Gait Posture* 39 (1) (Jan 2014) 35–39, <https://doi.org/10.1016/j.gaitpost.2013.05.022>.
- [24] K.J. Mickle, C.J. Nester, G. Crofts, J.R. Steele, Reliability of ultrasound to measure morphology of the toe flexor muscles, *J. Foot Ankle Res.* 6 (1) (Apr 4 2013) 12, <https://doi.org/10.1186/1757-1146-6-12>.
- [25] M.J. Page, et al., The PRISMA 2020 statement: an updated guideline for reporting systematic reviews, in eng, *Bmj* 372 (Mar 29 2021) n71, <https://doi.org/10.1136/bmj.n71>.
- [26] J.P. Higgins, et al., *Cochrane Handbook for Systematic Reviews of Interventions*, John Wiley & Sons, 2019.
- [27] S. Aslam, P. Emmanuel, Formulating a researchable question: a critical step for facilitating good clinical research, in eng, *Indian J. Sex. Transm. Dis.* 31 (1) (Jan 2010) 47–50, <https://doi.org/10.4103/0253-7184.69003>.
- [28] L. Willemse, E.J.M. Wouters, M.F. Pisters, B. Vanwanseele, Intra-assessor reliability and measurement error of ultrasound measures for foot muscle morphology in older adults using a tablet-based ultrasound machine, in eng, *J. Foot Ankle Res.* 15 (1) (Jan 25 2022) 6, <https://doi.org/10.1186/s13047-022-00510-1>.
- [29] W.M. Bramer, D. Giustini, G.B. de Jonge, L. Holland, T. Bekhuis, De-duplication of database search results for systematic reviews in EndNote, in eng, *J. Med. Libr. Assoc.: JMLA* 104 (3) (2016) 240–243, <https://doi.org/10.3163/1536-5050.104.3.014>.
- [30] M. Ouzzani, H. Hammady, Z. Fedorowicz, A. Elmagarmid, Rayyan—a web and mobile app for systematic reviews, *Syst. Rev.* 5 (2016), <https://doi.org/10.1186/s13643-016-0384-4>.
- [31] S.H. Downs, N. Black, The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions, in eng, *J. Epidemiol. Community Health* 52 (6) (Jun 1998) 377–384, <https://doi.org/10.1136/jech.52.6.377>.
- [32] T.K. Koo, M.Y. Li, A guideline of selecting and reporting intraclass correlation coefficients for reliability research, in eng, *J. Chiropr Med* 15 (2) (Jun 2016) 155–163, <https://doi.org/10.1016/j.jcm.2016.02.012>.
- [33] J.I. Seok, F.O. Walker, S.G. Kwak, Evaluation of extensor digitorum brevis thickness in healthy subjects: a comparative analysis of nerve conduction studies and ultrasound scans, in eng, *Clin. Neurophysiol.* 127 (2) (Feb 2016) 1664–1668, <https://doi.org/10.1016/j.clinph.2015.07.025>.
- [34] A. Abraham, V.E. Drory, Y. Fainmesser, L.E. Lovblom, V. Bril, Quantitative sonographic evaluation of muscle thickness and fasciculation prevalence in healthy subjects, in eng, *Muscle Nerve* 61 (2) (Feb 2020) 234–238, <https://doi.org/10.1002/mus.26758>.
- [35] *Obesity: preventing and managing the global epidemic. Report of a WHO consultation*, in eng, *World Health Organ. Tech. Rep. Ser.* 894 (2000) 1–253.
- [36] M. Morikawa, et al., Contribution of plantar fascia and intrinsic foot muscles in a single-leg drop landing and repetitive rebound jumps: an ultrasound-based study, in eng, *Int. J. Environ. Res. Publ. Health* 18 (9) (Apr 23 2021), <https://doi.org/10.3390/ijerph18094511>.
- [37] P.J. Battaglia, R. Mattox, B. Winchester, N.W. Kettner, Non-weight-bearing and weight-bearing ultrasonography of select foot muscles in young, asymptomatic participants: a descriptive and reliability study, in eng, *J. Manip. Physiol. Ther.* 39 (9) (Nov-Dec 2016) 655–661, <https://doi.org/10.1016/j.jmpt.2016.10.001>.
- [38] A.D. Henderson, et al., Early-stage diabetic neuropathy reduces foot strength and intrinsic but not extrinsic foot muscle size, in eng, *J. Diabetes Res.* 2020 (2020), 9536362, <https://doi.org/10.1155/2020/9536362>.
- [39] K. Koyama, M. Hirokawa, Y. Yoshitaka, J. Yamauchi, Toe flexor muscle strength and morphological characteristics of the foot in judo athletes, in eng, *Int. J. Sports Med.* 40 (4) (Apr 2019) 263–268, <https://doi.org/10.1055/a-0796-6679>.
- [40] T. Tanaka, et al., Characteristics of lower leg and foot muscle thicknesses in sprinters: does greater foot muscles contribute to sprint performance?, in eng, *Eur. J. Sport Sci.* 19 (4) (May 2019) 442–450, <https://doi.org/10.1080/17461391.2018.1534991>.
- [41] T. Abe, K. Tayashiki, M. Nakatani, H. Watanabe, Relationships of ultrasound measures of intrinsic foot muscle cross-sectional area and muscle volume with maximum toe flexor muscle strength and physical performance in young adults, in eng, *J. Phys. Ther. Sci.* 28 (1) (Jan 2016) 14–19, <https://doi.org/10.1589/jpts.28.14>.
- [42] A.F. Cameron, K. Rome, W.A. Hing, Ultrasound evaluation of the abductor hallucis muscle: reliability study, *J. Foot Ankle Res.* 1 (1) (Sep 25 2008) 12, <https://doi.org/10.1186/1757-1146-1-12>.
- [43] M.M. Frantovich Smith, J.A. Hides, P.W. Hodges, N.J. Collins, Intrinsic foot muscle size can be measured reliably in weight bearing using ultrasound imaging, *Gait Posture* 68 (Feb 2019) 369–374, <https://doi.org/10.1016/j.gaitpost.2018.12.012>.
- [44] J.J. Fraser, L.C. Mangum, J. Hertel, Test-retest reliability of ultrasound measures of intrinsic foot motor function, *Phys. Ther. Sport* 30 (Mar 2018) 39–47, <https://doi.org/10.1016/j.ptsp.2017.11.032>.
- [45] W.A. Hing, K. Rome, A.F. Cameron, Reliability of measuring abductor hallucis muscle parameters using two different diagnostic ultrasound machines, *J. Foot Ankle Res.* 2 (2009) 33, <https://doi.org/10.1186/1757-1146-2-33>.
- [46] A.W. Johnson, et al., Use of cine loops and structural landmarks in ultrasound image processing improves reliability and reduces error in the assessment of foot and leg muscles, in eng, *J. Ultrasound Med.* 39 (6) (Jun 2020) 1107–1116, <https://doi.org/10.1002/jum.15192>.
- [47] N. Maeda, et al., Intrinsic foot muscle hardness is related to dynamic postural stability after landing in healthy young men, in eng, *Gait Posture* 86 (May 2021) 192–198, <https://doi.org/10.1016/j.gaitpost.2021.03.005>.
- [48] K.J. Mickle, C.J. Nester, Size of the toe flexor muscles in older people with toe deformities, in english, *Foot Ankle Surg., Conference Abstract* 22 (2) (2016) 78, <https://doi.org/10.1016/j.fas.2016.05.194>.
- [49] S. Taş, A. Çetin, An investigation of the relationship between plantar pressure distribution and the morphologic and mechanic properties of the intrinsic foot muscles and plantar fascia, in eng, *Gait Posture* 72 (Jul 2019) 217–221, <https://doi.org/10.1016/j.gaitpost.2019.06.021>.
- [50] S. Taş, N. Ünlüer, A. Çetin, Thickness, cross-sectional area, and stiffness of intrinsic foot muscles affect performance in single-leg stance balance tests in healthy sedentary young females, in eng, *J. Biomech.* 99 (Jan 23 2020), 109530, <https://doi.org/10.1016/j.jbiomech.2019.109530>.
- [51] F.V. Verhulst, A.E.E.P.M. Leeuwesteijn, J.W.K. Louwerens, A.C.H. Geurts, N. Van Alfen, S. Pillen, Quantitative ultrasound of lower leg and foot muscles: feasibility and reference values, *Foot Ankle Surg.* 17 (3) (2011/09/01/2011) 145–149, <https://doi.org/10.1016/j.fas.2010.04.002>.
- [52] X. Zhang, K.H. Schütte, B. Vanwanseele, Foot muscle morphology is related to center of pressure sway and control mechanisms during single-leg standing, in eng, *Gait Posture* 57 (Sep 2017) 52–56, <https://doi.org/10.1016/j.gaitpost.2017.05.027>.
- [53] A. Abraham, Y. Fainmesser, L.E. Lovblom, V. Bril, V.E. Drory, Superiority of sonographic evaluation of contracted versus relaxed muscle thickness in motor neuron diseases, in eng, *Clin. Neurophysiol.* 131 (7) (Jul 2020) 1480–1486, <https://doi.org/10.1016/j.clinph.2020.04.003>.
- [54] A. Aiyer, S. Stewart, K. Rome, The effect of age on muscle characteristics of the abductor hallucis in people with hallux valgus: a cross-sectional observational study, in eng, *J. Foot Ankle Res.* 8 (2015) 19, <https://doi.org/10.1186/s13047-015-0078-5>.
- [55] C.C. Lobo, et al., Ultrasound evaluation of intrinsic plantar muscles and fascia in hallux valgus: a case-control study, *Medicine (Baltim.)* 95 (45) (Nov 2016) e5243, <https://doi.org/10.1097/MD.0000000000005243>.

- [56] S. Stewart, R. Ellis, M. Heath, K. Rome, Ultrasonic evaluation of the abductor hallucis muscle in hallux valgus: a cross-sectional observational study, *BMC Musculoskel. Disord.* 14 (45) (Jan 28 2013), <https://doi.org/10.1186/1471-2474-14-45>.
- [57] S. Taş, A. Çetin, Mechanical properties and morphologic features of intrinsic foot muscles and plantar fascia in individuals with hallux valgus, in eng, *Acta Orthop. Traumatol. Turcica* 53 (4) (Jul 2019) 282–286, <https://doi.org/10.1016/j.aott.2019.03.009>.
- [58] S. Angin, G. Crofts, K.J. Mickle, C.J. Nester, Ultrasound evaluation of foot muscles and plantar fascia in pes planus, *Gait Posture* 40 (1) (2014) 48–52, <https://doi.org/10.1016/j.gaitpost.2014.02.008>.
- [59] S. Angin, K.J. Mickle, C.J. Nester, Contributions of foot muscles and plantar fascia morphology to foot posture, in eng, *Gait Posture* 61 (Mar 2018) 238–242, <https://doi.org/10.1016/j.gaitpost.2018.01.022>.
- [60] K. Okamura, K. Egawa, T. Ikeda, K. Fukuda, S. Kanai, Relationship between foot muscle morphology and severity of pronated foot deformity and foot kinematics during gait: a preliminary study, in eng, *Gait Posture* 86 (May 2021) 273–277, <https://doi.org/10.1016/j.gaitpost.2021.03.034>.
- [61] K. Sakamoto, S. Kudo, Morphological characteristics of intrinsic foot muscles among flat foot and normal foot using ultrasonography, *Acta Bioeng. Biomech.* 22 (4) (2020) 161–166, <https://doi.org/10.37190/abb-01713-2020-02>.
- [62] S. Taş, N. Ünlüer, F. Korkusuz, Morphological and mechanical properties of plantar fascia and intrinsic foot muscles in individuals with and without flat foot, in eng, *J. Orthop. Surg.* 26 (3) (May-Aug 2018), 2309499018802482, <https://doi.org/10.1177/2309499018802482>.
- [63] X.Y. Zhang, J. Aeles, B. Vanwanseele, Comparison of foot muscle morphology and foot kinematics between recreational runners with normal feet and with asymptomatic over-pronated feet, *Gait Posture* 54 (May 2017) 290–294, <https://doi.org/10.1016/j.gaitpost.2017.03.030>.
- [64] X. Zhang, R. Puael, K. Deschamps, I. Jonkers, B. Vanwanseele, Differences in foot muscle morphology and foot kinematics between symptomatic and asymptomatic pronated feet, *Scand. J. Med. Sci. Sports* 29 (11) (2019) 1766–1773, <https://doi.org/10.1111/sms.13512>.
- [65] C.G. Kumar, K.V. Rajagopal, H.M. Hande, A.G. Maiya, S.S. Mayya, Intrinsic foot muscle and plantar tissue changes in type 2 diabetes mellitus, in eng, *J. Diabetes* 7 (6) (Nov 2015) 850–857, <https://doi.org/10.1111/1753-0407.12254>.
- [66] K. Severinsen, A. Obel, J. Jakobsen, H. Andersen, Atrophy of foot muscles in diabetic patients can be detected with ultrasonography, *Diabetes Care* 30 (12) (Dec 2007) 3053–3057, <https://doi.org/10.2337/dc07-0108>.
- [67] X. Wang, L. Chen, W. Liu, B. Su, Y. Zhang, Early detection of atrophy of foot muscles in Chinese patients of type 2 diabetes mellitus by high-frequency ultrasonography, *J. Diabetes Res.* 2014 (2014) 1–6, <https://doi.org/10.1155/2014/927069>.
- [68] C. Calvo-Lobo, et al., Quantitative ultrasound imaging pixel analysis of the intrinsic plantar muscle tissue between hemiparesis and contralateral feet in post-stroke patients, in eng, *Int. J. Environ. Res. Publ. Health* 15 (11) (Nov 11 2018), <https://doi.org/10.3390/ijerph15112519>.
- [69] C. Calvo-Lobo, et al., Rehabilitative ultrasound imaging of the bilateral intrinsic plantar muscles and fascia in post-stroke survivors with hemiparesis: a case-control study, in eng, *Int. J. Med. Sci.* 15 (9) (2018) 907–914, <https://doi.org/10.7150/ijms.25836>.
- [70] J.J. Fraser, R. Koldenhoven, J. Hertel, Ultrasound measures of intrinsic foot muscle size and activation following lateral ankle sprain and chronic ankle instability, *J. Sport Rehabil.* (Apr 9 2021) 1–11, <https://doi.org/10.1123/jsr.2020-0372>.
- [71] K.K. Hogan, J.A. Prince, M.C. Hoch, The evaluation of the foot core system in individuals with plantar heel pain, in eng, *Phys. Ther. Sport* 42 (Mar 2020) 75–81, <https://doi.org/10.1016/j.ptsp.2019.11.011>.
- [72] A.H. Jaffri, J. Hertel, S. Saliba, Ultrasound examination of intrinsic foot muscles in patients with 1st metatarsophalangeal joint arthrodesis, *Foot* 41 (Dec 2019) 79–84, <https://doi.org/10.1016/j.foot.2019.08.009>.
- [73] K.D. Pompeo, et al., Proximal, local, and distal muscle morphology in women with patellofemoral pain, *J. Diagn. Med. Sonogr.* 37 (2) (Mar 2021) 120–131, <https://doi.org/10.1177/8756479320972097>.
- [74] C. Romero-Morales, et al., Intrinsic foot muscles morphological modifications in patients with Achilles tendinopathy: a novel case-control research study, in eng, *Phys. Ther. Sport* 40 (Nov 2019) 208–212, <https://doi.org/10.1016/j.ptsp.2019.09.014>.
- [75] C.C. Lobo, et al., Ultrasound evaluation of intrinsic plantar muscles and fascia in hallux valgus A case-control study, *Medicine* 95 (45) (Nov 2016), e5243, <https://doi.org/10.1097/md.0000000000005243>.
- [76] K. Yahata, et al., Effects of a high-volume static stretching programme on plantar-flexor muscle strength and architecture, in eng, *Eur. J. Appl. Physiol.* 121 (4) (Apr 2021) 1159–1166, <https://doi.org/10.1007/s00421-021-04608-5>.
- [77] N.A. Campitelli, S.A. Spencer, K. Bernhard, K. Heard, A. Kidon, Effect of vibram FiveFingers minimalist shoes on the abductor hallucis muscle, in eng, *J. Am. Podiatr. Med. Assoc.* 106 (5) (Sep 2 2016) 344–351, <https://doi.org/10.7547/14-084>.
- [78] A.W. Johnson, J.W. Myrer, U.H. Mitchell, I. Hunter, S.T. Ridge, The effects of a transition to minimalist shoe running on intrinsic foot muscle size, *Int. J. Sports Med.* 37 (2) (2015) 154–158, <https://doi.org/10.1055/s-0035-1559685>.
- [79] S.T. Ridge, et al., Walking in minimalist shoes is effective for strengthening foot muscles, in eng, *Med. Sci. Sports Exerc.* 51 (1) (Jan 2019) 104–113, <https://doi.org/10.1249/mss.0000000000001751>.
- [80] J. Reeves, R. Jones, A. Liu, L. Bent, A. Martinez-Santos, C. Nester, No change in foot soft tissue morphology and skin sensitivity after three months of using foot orthoses that alter plantar pressure, *Footwear Sci.* 13 (3) (2021) 233–244, <https://doi.org/10.1080/19424280.2021.1961880>.
- [81] K. Protopoulos, S.D. Perry, The effect of a 12-week custom foot orthotic intervention on muscle size and muscle activity of the intrinsic foot muscle of young adults during gait termination, in eng, *Clin. Biomech.* 78 (Aug 2020), 105063, <https://doi.org/10.1016/j.clinbiomech.2020.105063>.
- [82] M.H. Kim, C.H. Yi, J.H. Weon, H.S. Cynn, D.Y. Jung, O.Y. Kwon, Effect of toe-spread-out exercise on hallux valgus angle and cross-sectional area of abductor hallucis muscle in subjects with hallux valgus, in eng, *J. Phys. Ther. Sci.* 27 (4) (Apr 2015) 1019–1022, <https://doi.org/10.1589/jpts.27.1019>.
- [83] D.Y. Jung, E.K. Koh, O.Y. Kwon, Effect of foot orthoses and short-foot exercise on the cross-sectional area of the abductor hallucis muscle in subjects with pes planus: a randomized controlled trial, *J. Back Musculoskeletal Rehabilitation, Journal Article; Randomized Controlled Trial* 24 (4) (2011) 225–231, <https://doi.org/10.3233/BMR-2011-0299>.
- [84] J. Namsawang, W. Eungpinichpong, R. Vichiansiri, S. Rattanathongkom, Effects of the short foot exercise with neuromuscular electrical stimulation on navicular height in flexible flatfoot in Thailand: a randomized controlled trial, *J. Preventive Med. Public Health = Yebang Uihakhoe chi, Journal Article; Randomized Controlled Trial* 52 (4) (2019) 250–257, <https://doi.org/10.3961/jpmph.19.072>.
- [85] K. Okamura, K. Fukuda, S. Oki, T. Ono, S. Tanaka, S. Kanai, Effects of plantar intrinsic foot muscle strengthening exercise on static and dynamic foot kinematics: a pilot randomized controlled single-blind trial in individuals with pes planus, in eng, *Gait Posture* 75 (Jan 2020) 40–45, <https://doi.org/10.1016/j.gaitpost.2019.09.030>.
- [86] A. Smith, J.A. Hides, P.W. Hodges, N.J. Collins, Intrinsic foot muscle size can be measured reliably in weight bearing using ultrasound imaging, *Gait Posture* 68 (Feb 2019) 369–374, <https://doi.org/10.1016/j.gaitpost.2018.12.012>.
- [87] D.C. Swanson, et al., Validity of ultrasound imaging for intrinsic foot muscle cross-sectional area measurements demonstrated by strong agreement with MRI, in eng, *BMC Musculoskel. Disord.* 23 (1) (Feb 14 2022) 146, <https://doi.org/10.1186/s12891-022-05090-6>.
- [88] K.J. Mickle, S. Angin, G. Crofts, C.J. Nester, Effects of age on strength and morphology of toe flexor muscles, in eng, *J. Orthop. Sports Phys. Ther.* 46 (12) (Dec 2016) 1065–1070, <https://doi.org/10.2519/jospt.2016.6597>.
- [89] L. Willemse, E.J.M. Wouters, H.M. Bronts, M.F. Pisters, B. Vanwanseele, The effect of interventions anticipated to improve plantar intrinsic foot muscle strength on fall-related dynamic function in adults: a systematic review, in eng, *J. Foot Ankle Res.* 15 (1) (Jan 20 2022) 3, <https://doi.org/10.1186/s13047-021-00509-0>.