
No Decrease of the β -Cell Mass in Type 2 Diabetic Patients

Yves Guiot, Christine Sempoux, Pierre Moulin, and Jacques Rahier

BACKGROUND AND AIMS

Both insulin resistance and a β -cell defect contribute to type 2 diabetes. The nature of this β -cell defect is unknown, and the eventual role of a β -cell mass decrease remains controversial, probably due to differences in methodology, sampling, and ways of expressing the results between the studies. Only total β -cell mass per pancreas has been thoroughly evaluated. The oldest studies did not distinguish between type 1 and type 2 diabetes, and in many cases, they only considered small groups of patients. The present study aimed to quantify total β -cell mass in a large series of diabetic patients in relation to obesity and treatment.

MATERIALS AND METHODS

Total β -cell mass, proinsulin, and relative insulin content were evaluated in pancreata obtained at autopsy from 39 type 2 diabetic and 28 normoglycemic patients on consecutive sections stained by hematoxylin-eosin, proinsulin, and insulin antibodies. Total β -cell mass was calculated from the weight of the pancreas and from the β -cell volume density measured by point counting, and the relative insulin content

was evaluated by insulin immunodensitometry. Diabetic patients were subdivided into four groups according to their BMI ($>$ or <25 kg/m²) and their eventual insulin requirement.

RESULTS

Pancreatic weight was significantly lower in the group of lean insulin-requiring type 2 diabetic patients than in others. β -Cell volume density and insulin content were similar in all groups. Mean total β -cell mass did not significantly differ in normoglycemic and type 2 diabetic patients when not subdivided according to the treatment. However, it was significantly lowered in obese and lean type 2 diabetic patients who required insulin treatment. Whether this is the cause or the consequence of the insulin treatment is uncertain. Proinsulin synthesis was still active in islets of type 2 diabetic patients.

CONCLUSIONS

Because β -cell mass was preserved in 82% of type 2 diabetic patients, and because β -cell insular stock and cellular synthesis capacity were maintained, it is likely that the pathogenesis of type 2 diabetes results from an abnormal coupling of the insulin secretion to glycemia rather than to a significant β -cell loss.

From the Department of Pathology, Université Catholique de Louvain, Brussels, Belgium.

Address correspondence and reprint requests to Dr. Jacques Rahier, Department of Pathology, Université Catholique de Louvain, 1200 Brussels, Belgium. E-mail: rahier@anps.ucl.ac.be.

Received for publication 3 July 2000 and accepted 3 August 2000.

This article is based on a symposium. The symposium and the publication of this article were made possible by an unrestricted educational grant from Les Laboratoires Servier.