

Chapter III: Is the NF-κB pathway activated in peritoneal endometriosis in women?

1. Article 1: Nuclear factor-kappaB (NF-κB) is constitutively activated in peritoneal endometriosis

Nuclear factor-kappa B is constitutively activated in peritoneal endometriosis

Reinaldo González-Ramos¹, Jacques Donnez^{1,2}, Sylvie Defrère¹, Isabelle Leclercq², Jean Squifflet¹, Jean-Christophe Lousse¹ and Anne Van Langendonck¹

¹Department of Gynecology, Université Catholique de Louvain, Cliniques Universitaires St Luc, Avenue Hippocrate 10, 1200 Brussels, Belgium and ²Department of Internal Medicine, Gastroenterology Unit, Université Catholique de Louvain, Brussels, Belgium

³Correspondence address. Tel: +32-2-764-95-06; Fax: +32-2-764-95-07; E-mail: donnez@gyne.ucl.ac.be

Mol Hum Reprod (2007), *in press, advance access on line.*

Running title: NF-κB and ICAM-1 in peritoneal endometriosis

Keywords: endometriosis / ICAM-1 / IkappaB / inflammation / NF-kappaB

Chapter IV: Involvement of NF-κB in the development of endometriotic lesions

1. Article 2: Agents blocking the nuclear factor-kappaB (NF-κB) pathway are effective inhibitors of endometriosis in an *in vivo* experimental model

Reinaldo González-Ramos¹, Anne Van Langendonck¹, Sylvie Defrère¹, Jean-Christophe Lousse¹, Marcel Mettlen², Alain Guillet³ and Jacques Donner¹. *Gynecol Obstet Invest*, *in press*.

¹ Department of Gynecology, Université Catholique de Louvain, Brussels, Belgium

² CELL Unit, Université Catholique de Louvain and Christian de Duve Institute of Cellular Pathology, Brussels, Belgium

³ Institute of Statistics, Université Catholique de Louvain, Louvain-La-Neuve, Belgium

Short title: Nuclear factor-κB inhibition reduces endometriosis *in vivo*

Keywords: apoptosis / BAY 11-7085 / cell proliferation / endometriosis / inflammatory response / ICAM-1 / NF-kappaB inhibition / nude mouse model / SN-50