

Secondary hyperalgesia is mediated by heat-insensitive A-fibre nociceptors

Emanuel N. van den Broeke*, Cédric Lenoir* and André Mouraux

Institute of Neuroscience, Université catholique de Louvain, B-1200, Brussels, Belgium

Key points

- It is believed that secondary hyperalgesia (the increased sensitivity to mechanical nociceptive stimuli that develops after cutaneous tissue injury in the surrounding uninjured skin) is mediated by a subclass of nociceptors: the slowly adapting A-fibre mechano-heat nociceptors (AMH-type I). Here we tested this hypothesis.
- By using intense long-lasting heat stimuli, which are known to activate these slowly adapting AMH-type I nociceptors, we show that the perceived intensity elicited by these stimuli is *not* increased in the area of secondary hyperalgesia.
- Moreover, we show that during an A-fibre nerve conduction block the perception elicited by the long-lasting heat stimuli is significantly reduced in a time window that matches the response profile of the AMH-type I nociceptors.
- AMH-type I nociceptors contribute to the perception of sustained heat, but they do not mediate secondary hyperalgesia. Therefore, we propose that secondary hyperalgesia is mediated by high threshold mechanoreceptors.

Abstract Secondary hyperalgesia refers to the increase in sensitivity to mechanical nociceptive stimuli delivered outside the area of tissue injury. Previous studies have suggested that secondary hyperalgesia is mediated by a specific class of myelinated nociceptors: slowly adapting A-fibre mechano- and heat-sensitive (AMH) type I nociceptors. Here, we tested this hypothesis by examining whether long-lasting heat stimuli, which are known to activate AMH-type I nociceptors, elicit enhanced responses when delivered to the area of secondary hyperalgesia induced by high frequency electrical stimulation of the skin (HFS). Before and 20 min after HFS, sustained 30 s radiant heat stimuli were delivered to the area of increased mechanical pinprick sensitivity while participants continuously rated intensity of perception using an online visual analog scale (0–100 mm). After HFS, no significant enhancement of heat perception was observed in the area of increased pinprick sensitivity. To establish that myelinated nociceptors actually contribute to the perception of sustained heat, we conducted a second experiment in which sustained heat stimuli were presented before and during an A-fibre nerve conduction block, achieved by applying a rubber band with weights which compresses the superficial radial nerve against the radius. During the block, heat perception was significantly reduced 17–33 s after the onset of the heat stimulus (before: mean = 53 mm, during: mean = 31 mm; $P = 0.03$), matching the response profile of AMH-type I nociceptors. These results support the notion that AMH-type I nociceptors contribute to the perception of sustained heat, but also show that these afferents do not mediate secondary hyperalgesia.

*These authors contributed equally.

(Received 11 April 2016; accepted after revision 1 July 2016; first published online 5 July 2016)

Corresponding author E. N. van den Broeke: Institute of Neuroscience (IoNS), Faculty of Medicine, Université catholique de Louvain, B-1200, Brussels, Belgium. E-mail: Emanuel.vandenbroeke@uclouvain.be

Abbreviations AMH, A fibre mechano-heat nociceptor; HFS, high frequency stimulation; HTM, high threshold mechanoreceptor; NRS, numerical rating scale; RT, reaction time; TRPV1, transient receptor potential, vanilloid subfamily, member 1; VAS, visual analog scale.

Introduction

Cutaneous tissue injury produces inflammation and hyperalgesia. This hyperalgesia is thought to result from a combination of peripheral and central sensitization of the nociceptive system. Peripheral sensitization refers to the enhanced responsiveness of peripheral nociceptors innervating the injured or inflamed tissues. Central sensitization refers to an increased responsiveness of nociceptive neurons in the CNS (Loeser & Treede, 2008).

A key feature of central sensitization is the development of an increased sensitivity to mechanical pinprick stimuli that spreads beyond the site of injury or inflammation, a phenomenon also referred to as secondary hyperalgesia. In humans, secondary hyperalgesia can be induced experimentally by activating nociceptors in a sustained and intense fashion, for example using an intradermal injection of capsaicin to selectively activate the transient receptor potential, vanilloid subfamily, member 1 (TRPV1) receptor (LaMotte *et al.* 1991), or using high frequency electrical stimulation of the skin (Klein *et al.* 2004). With these methods, studies have demonstrated that, contrasting with the marked enhanced sensitivity to mechanical pinprick stimuli, the sensitivity to short-lasting heat stimuli delivered to the area of secondary hyperalgesia is not enhanced (Ali *et al.* 1996). Studies have also shown that selective blockade of myelinated fibres suppresses secondary hyperalgesia (Ziegler *et al.* 1999). Furthermore, increased pinprick sensitivity also developed in skin pre-treated with capsaicin to induce a denervation of fibres expressing TRPV1 (Magerl *et al.* 2001). Based on these results, it is generally accepted that secondary hyperalgesia is mediated by myelinated nociceptors that respond to mechanical pinprick stimuli but do not respond to short-duration heat stimuli and capsaicin (Ali *et al.* 1996; Ziegler *et al.* 1999; Magerl *et al.* 2001).

Based on their response to sustained heat (30 s, 53°C), teased-fibre recordings in monkeys have shown that myelinated nociceptors can be categorized into quickly adapting nociceptors [A-fibre mechano- and heat-sensitive (AMH)-type II], slowly adapting nociceptors (AMH-type I) and heat-insensitive nociceptors (Treede *et al.* 1995, 1998). AMH-type II nociceptors respond immediately at the onset of the sustained heat stimulus (mean heat threshold around 47°C), but their rate of discharge fades out very quickly over time. These

nociceptors probably mediate the 'first' pain elicited by a short-lasting heat stimulus (Treede *et al.* 1995, 1998). In contrast, AMH-type I nociceptors do not respond immediately after the onset of the sustained heat stimulus, and they do not respond to short-lasting 1 s heat stimuli, even at 53°C. When exposed to sustained heat, they start responding after a few seconds, with an average peak latency around 16 or 21 s, depending on the studies (Treede *et al.* 1995, 1998). Finally, heat-insensitive nociceptors do not respond at all to heat. Because they do respond to mechanical nociceptive stimuli, they are also called high-threshold mechanoreceptors (HTMs).

Treede *et al.* (1998) showed that, when long-duration heat stimuli are used (e.g. 30 s) AMH-type I and AMH-type II nociceptors have similar heat thresholds.

As most AMH-type I nociceptors respond to mechanical stimuli but not to short-lasting heat stimuli (Treede *et al.* 1998) or capsaicin (Ringkamp *et al.* 2001) it is believed that secondary pinprick hyperalgesia is mediated by this class of nociceptor (Iannetti *et al.* 2013). However, an alternative possibility could be that secondary pinprick hyperalgesia is mediated by HTMs (Magerl *et al.* 2001).

The aim of the present study was to test whether secondary hyperalgesia is mediated by AMH-type I nociceptors by examining whether long-duration heat stimuli (30 s) delivered at an intensity above the threshold of AMH-type II elicit increased heat pain in the heterotopic area of increased pinprick sensitivity with a similar time course as the response profile of the AMH-type I nociceptors.

In a first experiment, we show that, in the area of increased pinprick sensitivity induced by HFS, the perception of long-lasting heat stimuli is not significantly enhanced. In a second study, we show that, during an A-fibre block, the perception of long-lasting heat is significantly reduced within a time window corresponding to the expected response profile of AMH-type I. Taken together, the results of these two experiments suggest strongly that the AMH-type I nociceptors do not mediate mechanical secondary hyperalgesia.

Methods

Ethical approval

The experiments were conducted according to the *Declaration of Helsinki*. Approval for the two experiments

was obtained from the local Ethical Committee (Hospital and Departmental Ethics Committee, Saint-Luc - Université catholique de Louvain). All participants signed an informed consent form.

Participants

Fourteen healthy volunteers took part in experiment 1 [8 men and 6 women; aged 21–33 years; 25.1 ± 3.6 years (mean $\pm$ SD)]. Six healthy volunteers took part in experiment 2 (3 men and 3 women; aged 24–38 years; 27.8 ± 5.1 years). The participants in experiment 1 received financial compensation for their participation.

Experiment 1

The design of the experiment is summarized in Fig. 1. During the experiment, participants were comfortably seated in a chair with their arms resting on a table in front of them.

Induction of increased pinprick sensitivity. Transcutaneous high frequency electrical stimulation (HFS) of the left or right volar forearm was used to induce secondary hyperalgesia. Several studies have shown that this technique induces a robust and long-lasting increase in mechanical pinprick sensitivity, extending beyond the area of stimulation (Klein *et al.* 2004, 2008; van den Broeke & Mouraux, 2014; Henrich *et al.* 2015). Similar to other commonly used methods to induce secondary hyperalgesia (e.g. intradermal injection of capsaicin), HFS-induced secondary hyperalgesia depends on the activation of capsaicin-sensitive fibres (Magerl *et al.* 2001; Henrich *et al.* 2015). HFS stimulation was applied to the left or right volar forearm, 10 cm distal to the cubital fossa, and consisted of five trains of 100 Hz (pulse width: 2 ms), each lasting 1 s. The time interval between each train was 10 s. The intensity of stimulation was individually adjusted to 20 times the detection threshold to a single

pulse [0.25 ± 0.10 mA (mean $\pm$ SD)]. The electrical pulses were triggered by a programmable pulse generator (Master-8; AMPI Israel, Jerusalem, Israel), generated by a constant current electrical stimulator (Digitimer DS7A, Digitimer UK, Welwyn Garden City, UK), and delivered to the skin using a specifically designed electrode built at the Centre for Sensory-Motor Interaction (Aalborg University, Denmark). The electrode consists of 16 blunt stainless steel pins (diameter: 0.2 mm) protruding 1 mm from the base. The pins are placed in a 10 mm diameter circle and serve as cathode. A stainless steel anode electrode is concentrically located around the steel pins (inner diameter: 22 mm; outer diameter: 40 mm). To avoid any confounding effect of handedness, the arm onto which HFS was applied (dominant *vs.* non dominant) was counterbalanced across participants. Handedness was assessed using the Flinders Handedness Survey (Nicholls *et al.* 2013).

Mapping the area of increased pinprick sensitivity. To map the area of increased mechanical pinprick sensitivity after HFS, a calibrated pinprick stimulator exerting a normal force of 128 mN (MRC Systems, Heidelberg, Germany) was applied perpendicular to the skin, starting at a distance from the conditioning stimulation area and repeated every 1 cm along a proximal–distal and lateral–medial direction towards the conditioning site. During this procedure, participants did not view the arm. Participants were instructed to indicate the point at which the perception was changed. This point was then indicated on the skin with a marker.

Quantifying changes in mechanical pinprick sensitivity. To assess changes in pinprick sensitivity, the same pinprick stimuli used to map the area of increased pinprick sensitivity were applied three times onto the skin. To avoid sensitization of the stimulated skin, the target of each pinprick stimulus was displaced after each stimulus. Participants were asked to report the intensity




Figure 1. Time-line of the experiment

Before (T0) and after (T1) applying HFS on the left or right forearm, the sensitivity to mechanical pinprick stimuli and long-lasting heat stimuli was assessed at both arms (control *vs.* HFS). The area of increased mechanical pinprick sensitivity was mapped immediately before T1.

of perception elicited by the pinprick stimulation on a numerical rating scale (NRS) ranging from 0 (no perception) to 100 (maximal pain), with 50 representing the transition from non-painful to painful domains of sensation. Stimuli were applied to the HFS-conditioned arm and the contralateral control arm, before applying HFS (T0) and 20 min after applying HFS (T1).

Quantifying changes in sensitivity to long-duration heat stimuli. To assess changes in heat sensitivity, long-duration (30 s) radiant heat stimuli were delivered onto the skin of the HFS-conditioned arm and the contralateral control arm, before applying HFS (T0) and 20 min after applying HFS (T1). The heat stimuli were generated by an infrared CO₂ laser stimulator (wavelength 10.6 μ m) whose power output is regulated using a closed-loop control of pulse-width modulation based on an online measurement of skin temperature performed using a radiometer whose field of view is collinear with the laser beam (Laser Stimulation Device; SIFEC, Ferrières, Belgium). The stimuli consisted of a 10 ms heating ramp during which the skin was brought to the desired target temperature, followed by a plateau during which skin temperature was maintained at target temperature for 30 s. The diameter of the laser beam at the target site was 6 mm. At each stimulation site and time point, a total of five stimuli were delivered. To avoid skin overheating and minimize stimulus-induced sensitization or habituation, the target of the laser beam was displaced manually by the experimenter after each stimulus. In order to rate continuously the perception elicited by the long-duration heat stimulus, a custom made visual analog scale (VAS) box was used, consisting of a 100 mm vertical slider connected to a computer. The lower extremity of the slider was labelled 'no perception' and the upper extremity of the slider was labelled 'maximal pain'. An anchor at the middle of the scale marked the transition between non-painful and painful domains of sensation. Participants were instructed to rate continuously the intensity of perception, as soon as they perceived a sensation, by moving the slider. The recording lasted 40 s from the onset of the stimulus. During heat stimulation, participants were instructed to keep their gaze on the VAS, and view of the arm was prevented. Before starting the heat stimulation during the post-stimulation measurement, subjects were explicitly told that although there was increased sensitivity to mechanical pinprick stimulation (as experienced during the mapping procedure), this was not necessarily going to be the case for thermal stimulation.

At the beginning of the experiment, for each participant, the target temperature for the long-duration heat stimuli was defined as follows. The individual thermal detection threshold of quickly adapting AMH-type II nociceptors was estimated for each arm using an adaptive staircase procedure. Participants were requested to press a button

held in the contralateral hand as soon as they detected a short-lasting heat stimulus (10 ms heating ramp followed by a 90 ms plateau). Reaction times (RTs) were used as a criterion to distinguish between detections mediated by myelinated AMH-type II (RT < 650 ms) vs. unmyelinated C fibres (RT $\geq$ 650 ms) (Churyukanov *et al.* 2012). The staircase algorithm started at 46°C. If the stimulus was detected with an RT < 650 ms, the temperature of the subsequent stimulus was decreased by 1°C. Conversely, if it was detected with an RT $\geq$ 650 ms or if it was undetected, the temperature of the subsequent stimulus was increased by 1°C. The staircase procedure ended after four reversals (when a previous stimulus was undetected and the subsequent one was detected or the reverse). The threshold was defined as the stimulus temperature of the fourth reversal. The target of the laser beam was displaced manually by the experimenter after each stimulus. During the threshold estimation procedure, the view of the arm was prevented. Considering that previous studies have suggested that, when long-duration heat stimuli are used, the thermal activation thresholds of AMH-type I and AMH-type II are similar (Treede *et al.* 1998), the actual target temperature used for the long-duration heat stimuli was set to 1°C above the estimated AMH-type II threshold (with a maximum of 50°C for safety reasons).

Data analysis. The effect of HFS on the intensity of the percept elicited by the mechanical pinprick stimuli was assessed using a repeated measures ANOVA with *time* (T0 and T1) and *side* (control vs. HFS arm) as within-subject factors. The interaction between the two factors was taken as a measure of the effect of HFS. The assumption of sphericity was tested using Mauchly's test. If the data violated the assumption of sphericity, F-values were corrected using the Greenhouse–Geisser procedure. *Post hoc* tests were performed using paired-sample *t*-tests in which *P*-values were Bonferroni corrected for the number of tests. The level of significance was set at $P < 0.05$ (two-sided). The statistical analyses were conducted using SPSS 18 (SPSS Inc., Chicago, IL, USA).

Analysis of the effect of HFS on the continuous ratings of the percept elicited by the long-lasting heat stimuli was conducted using Letswave6 (<http://www.nocions.org/letswave6/>). First, we computed the average of the five rating waveforms for each participant and condition. Second, we computed for each participant *difference waveforms* assessing the change in heat perception at T1 vs. T0 at the HFS arm (HFS_{T1} – HFS_{T0}) and at the control arm (control_{T1} – control_{T0}). Finally, we performed a temporal non-parametric cluster-based permutation test (Maris & Oostenveld, 2007; Groppe *et al.* 2011) using the *difference waveforms* of the two arms, thereby testing the interaction between *time* (T1 vs. T0) and *side* (HFS vs. control arm). The permutation test consisted of the following

steps. First, the *difference* waveforms were compared by means of a point-by-point paired-sample *t*-test. Second, samples above the critical *t*-value for a parametric two-sided test that were adjacent in time were identified and clustered. An estimate of the magnitude of each cluster was then obtained by computing the sum of the absolute *t*-values constituting each cluster (cluster-level statistic). Random permutation testing (2000 times) of the subject-specific *difference* waveforms of the two arms (performed independently for every subject) was then used to obtain a reference distribution of *maximum* cluster magnitude. Finally, the proportion of random partitions that resulted in a larger cluster-level statistic than the one observed (i.e. *p*-value) was calculated. Clusters in the observed data were regarded as significant if they had a magnitude exceeding the threshold of the 95.0th percentile of the permutation distribution (corresponding to a critical alpha-level of 0.05).

Experiment 2

To assess the relative contribution of myelinated fibres to the perception of sustained heat, we delivered long-duration heat stimuli in the innervation territory of the left superficial branch of the radial nerve, before and during an A-fibre conduction block of that nerve.

A-fibre nerve conduction block. To block selectively the conduction of myelinated fibres, pressure was applied to the left superficial branch of the radial nerve, at the level of the wrist (Magerl *et al.* 1999; Nahra & Plaghki, 2003; Mouraux *et al.* 2010). Participants were comfortably seated in a chair with the left forearm immobilized on an armrest in a neutral pronation/supination position with a handle bar in their left hand preventing any pronation or supination of the hand and forearm. The pressure was delivered by means of two weights of 700 g each hanging on a 1.5 cm rubber band applied on the forearm proximally to the wrist.

Development of the nerve conduction blockade was monitored by assessing, every 10 min, the ability of the participants to detect tactile stimuli (calibrated nylon filament exerting a target force of 5.9 mN; Semmens-Weinstein Von Frey Aesthesiometers, evaluator size 3.84, Stoelting Co., Wood Dale, IL, USA) and short-lasting cold stimuli (10°C stimuli applied during 100 ms using a 125 mm² surface; custom-built contact probe based on micro Peltier elements; Prof. André Dufour, Université de Strasbourg, France). As soon as the tactile and cold perception disappeared, we then assessed RTs to short-lasting heat stimuli (55°C stimuli lasting 100 ms, delivered using the temperature-controlled infrared CO₂ laser stimulator, laser beam diameter 6 mm) and mechanical pinprick stimuli (custom-made calibrated

pinprick stimulator exerting a normal force of 90 mN with a 0.35 mm probe diameter). Such as in Experiment 1, RTs to these stimuli were used to distinguish detections triggered by sensory input conveyed by myelinated A-fibres (RT < 650 ms) and unmyelinated C-fibres (RT ≥ 650 ms). The area of hypoesthesia was then defined as the area where (1) tactile stimuli and cold stimuli were not detected and (2) both the short-lasting heat stimuli and the pinprick stimuli were either undetected or detected with RTs ≥ 650 ms. During the assessments the view of the hand was prevented.

Quantifying changes in sensitivity to long-duration heat stimuli. The assessment of heat perception elicited by the long-duration heat stimuli before and during the nerve conduction blockade was performed in the same way as in Experiment 1.

Data analysis. To test the effect of the A-fibre nerve conduction block on the percept elicited by the long-duration heat stimuli, we used the cluster-based permutation testing also used in Experiment 1 (Maris & Oostenveld, 2007; Groppe *et al.* 2011) to compare the average intensity rating waveforms obtained before *vs.* during the block. Such as in Experiment 1, clusters were regarded as significant if their magnitude exceeded the threshold of the 95.0th percentile of the permutation distribution (corresponding to a critical alpha-level of 0.05).

Results

Experiment 1

Thermal detection thresholds to short-lasting heat stimuli. The AMH-type II detection threshold estimated at the beginning of experiment was on average (± SD) 47.6 ± 1.0°C (range 45–49°C) at the control arm and 47.6 ± 1.2°C (46–49°C) at the HFS-treated arm.

HFS-induced change in mechanical pinprick sensitivity. HFS induced a clear increase in the intensity of the percept elicited by mechanical pinprick stimuli delivered to the HFS-treated arm (Fig. 2). The repeated-measures ANOVA showed a significant *time* × *side* interaction ($F_{1,13} = 38.208$, $P < 0.001$, $\eta^2 = 0.746$). *Post hoc* tests showed that the perceived intensity after HFS at the treated arm was significantly increased (paired *t*-test; $t_{13} = 7.177$, $P < 0.001$). In contrast, no significant differences were observed for the control arm ($P = 0.468$).

HFS-induced change in the sensitivity to long-lasting heat stimuli. None of the participants showed any visible sign of a burn lesion after the long-duration heat stimuli. The

group-level average waveforms of intensity of perception are shown in Fig. 3. After HFS, the percept elicited by long-lasting heat stimuli delivered to the HFS-treated arm was similar to the percept elicited by stimulation of the control arm, except during the first 5 s which tended to show a slight enhancement of perception at the HFS arm. Nevertheless, the permutation test did not reveal any cluster having a *P*-value smaller than the critical alpha level of 0.05. This means that, after HFS, the heat perception elicited throughout the long-duration heat stimuli delivered to the HFS-treated arm was not significantly different as compared to baseline and control side.

Experiment 2

Thermal detection thresholds to short-lasting heat stimuli. The AMH-type II detection threshold estimated at the beginning of the experiment was on average ($\pm$ SD) $47.0 \pm 1.5^\circ\text{C}$ (range 45–49).

Time required to establish the A-fibre nerve conduction block. In five of six participants, the block was established 110 min after the start of the procedure. In one participant the block was established 130 min after the procedure.

Effects of the A-fibre nerve conduction block on the perception of brief tactile, heat, pinprick and cold stimuli. Before the start of the block, tactile, pinprick, cold and heat stimuli were all detected by every participant. Once the block was established, four out of six subjects spontaneously reported a mild burning sensation in their hand. Five out of six subjects did not detect any of the tactile stimuli and one subject detected 1/5 stimuli. Five out of six subjects were no longer able to detect the cold stimuli, and one subject detected 2/5 cold stimuli (Fig. 4C). Heat and pinprick stimuli were either not detected or detected with increased RTs (heat stimuli: 325.5 ± 65.0 vs. 958.5 ± 256.8 ms; pinprick stimuli: 261.9 ± 77.2 vs. 1065.0

± 454.4 ms) (Fig. 4A and B). All participants recovered completely after the release of the nerve conduction block.

Effect of the A-fibre nerve conduction block on the perception of long-lasting heat stimuli. None of the participants showed any visible sign of a burn after the long-duration heat stimuli. On average, ratings were decreased during the block, especially during the second half of the stimulus (Fig. 5). The permutation test comparing the rating waveforms before *versus* during the block revealed one cluster having a *P*-value smaller than the critical alpha level of 0.05 extending between 17 and 33 s after stimulus onset ($P = 0.03$). During this interval, heat perception was significantly reduced during the block.

Discussion

Although there was a clear increase in the perception of mechanical pinprick stimuli after HFS, no significant change in the perception of long-lasting heat stimuli delivered to the area of secondary hyperalgesia was observed. This indicates that, unlike previously proposed, increased pinprick sensitivity is not mediated by slowly adapting AMH-type I nociceptors.

The results of our second experiment support the view that myelinated fibres do contribute to sustained heat perception. Indeed, heat perception *during the second half* of the sustained heat stimulus was significantly decreased during the A-fibre nerve conduction block. Interestingly, the perception of long-lasting heat was not completely abolished, suggesting that sustained heat perception also involves *unmyelinated* afferents.

If AMH-type I nociceptors are not the primary mediators of secondary hyperalgesia, then the most likely candidate are HTMs (Treede & Magerl, 2000; Magerl *et al.* 2001; Meyer & Treede, 2004). Indeed, HTMs are mechano-sensitive but do not respond to short- or long-lasting heat stimuli and capsaicin (Treede *et al.* 1998; Ringkamp *et al.* 2001).




Figure 2. Effect of HFS on mechanical pinprick sensitivity

A, individual (thin black lines) and group-level average (thick black line) area of increased pinprick sensitivity at the forearm, estimated 20 min after applying HFS. B, group-level mean (and SD) intensity of perception elicited by pinprick stimulation of the control arm (grey line) and the HFS arm (black line), before (T0) and 20 min after (T1) applying HFS. At the HFS arm, pinprick sensitivity was significantly increased at T1 vs. T0 (** $P < 0.001$).




Figure 3. Effect of HFS on the perception of long-lasting heat stimuli

A, example of the skin heating profiles generated by the temperature-controlled laser stimulator and measured by the radiometer collinear with the laser beam. The horizontal dashed line indicates baseline skin temperature. The vertical dashed line indicates the end of the heat stimulation. B, group-level mean rating waveforms obtained during the long-duration heat stimulation for all conditions. The horizontal dashed line indicates the transition from non-painful to painful domains of sensation. The vertical dashed line indicates the end of the heat stimulation. C, group-level average difference rating waveforms (T1 minus T0) for both arms (control and HFS). The vertical lines represent error bars expressing standard deviation.




Figure 4. Effect of the A-fibre nerve conduction block on the reaction times to short-lasting heat stimuli, the reaction times to mechanical pinprick stimuli, and the detection of tactile and cold stimuli

A, stacked reaction time distribution for short-lasting (100 ms) 55°C heat stimuli applied on the dorsum of the left hand before and during the A-fibre nerve block. The vertical dashed line represents the cut-off used to distinguish reaction times compatible with the conduction velocities of myelinated vs. unmyelinated fibres. B, stacked reaction time distribution for mechanical pinprick stimulation. Note that during the A-fibre block the reaction times to both short-lasting heat stimuli and mechanical pinprick stimuli are shifted towards the right. C, during the A-fibre block, tactile and cold stimuli were, in almost all cases, not detected.

Peripheral vs. central sensitization

In rats, Reeh *et al.* (1987) showed that mechanical injury of the tail skin leads to peripheral sensitization of HTMs, both when the injury is applied within their receptive fields, or when applied closely outside the border of their receptive fields. Most of the sensitized HTMs developed spontaneous activity, expanded their receptive fields and exhibited lower thresholds to punctate mechanical stimuli (Reeh *et al.* 1987). These results suggest that peripheral sensitization of HTMs could contribute to the increased mechanical sensitivity surrounding the site of tissue injury. However, it remains unclear whether this peripheral phenomenon can explain the increased mechanical sensitivity commonly observed at a greater

distance from the injured site (i.e. the area of secondary hyperalgesia). Moreover, it remains unknown whether peripheral sensitization of HTMs can be induced in the absence of tissue injury.

On the other hand, there is ample evidence that secondary mechanical hyperalgesia results, at least partly, from a facilitation of nociceptive transmission at the level of the spinal cord, i.e. *central sensitization* (Kenshalo *et al.* 1982; McMahon & Wall, 1984; Woolf & King, 1990; Baumann *et al.* 1991; LaMotte *et al.* 1991; Simone *et al.* 1991). For example, McMahon & Wall (1984) showed in rats that the receptive field of dorsal horn lamina I neurons increases after a burn injury applied outside the receptive field of these neurons. Moreover, activation thresholds to mechanical stimuli delivered within their receptive fields were lower after the burn injury. These lamina I neurons also responded more strongly to transcutaneous electrical stimuli delivered to their cutaneous receptive fields. Because such stimuli bypass peripheral transduction processes, these effects can only be explained by a central mechanism.

The present study shows that, unlike mechanical pinprick perception, long-lasting heat perception is not significantly increased in the area of secondary hyperalgesia. This suggests that the central mechanism underlying increased pinprick sensitivity involves the facilitation of a modality-specific pathway dedicated to noxious mechanical input. Is there any evidence for such a pathway? In cats (Christensen & Perl, 1970; Light & Perl, 1979) and monkeys (Light & Perl, 1979), it was shown that HTMs project to the superficial layer (lamina I) of the dorsal horn. Moreover, the presence of modality-specific nociceptive neurons has been demonstrated in lamina I. For example, Han *et al.* (1998) showed, in cats, that 18 out of 38 lamina I neurons were nociceptive-specific. Eight of these neurons responded *only* to mechanical nociceptive stimulation (pinch) (Han *et al.* 1998). In a follow-up study, Andrew & Craig (2002) found that four out of 20 nociceptive-specific lamina I neurons responded only to noxious mechanical stimulation (Andrew & Craig, 2002). Taken together, these results demonstrate, at least in some animals, the existence of neurons specifically relaying mechanical nociceptive input at the level of the dorsal horn.

At first glance, the lack of increase in perception to long-lasting heat observed in the present study seems at odds with our previous results showing that HFS can enhance the percept elicited by short-lasting heat stimuli delivered to the skin closely surrounding the area onto which HFS is applied (van den Broeke & Mouraux, 2014). In the present study, the heat stimuli were often applied at a greater distance from the site of HFS, corresponding to the area of increased pinprick sensitivity. Possibly, the increase in heat sensitivity reported in our previous study involved a small area not covering the entire area of secondary hyperalgesia.




Figure 5. Effect of the A-fibre nerve conduction block on the percept elicited by long duration heat stimuli

A, group-level average rating waveforms obtained before and during the A-fibre nerve conduction block. During the A-fibre nerve conduction block heat perception was significantly decreased 17–33 s after the onset of the 30 s heat stimuli (grey area, $P < 0.05$). The horizontal dashed line indicates the transition from non-painful to painful domains of sensation. The vertical dashed line indicates the end of stimulation. Inset: note that during the block the delay between the increase in skin temperature and the increase in rating was delayed by approximately 1 s, compatible with the blockade of myelinated afferents. **B**, individual ratings representing the mean calculated within the time window 17–33 s, before and during the A-fibre block.

To conclude, we provide evidence that the increased sensitivity to mechanical pinprick stimulation in the area of secondary hyperalgesia is not conveyed by AMH-type I nociceptors. For this reason, we propose that HTMs are the primary mediators of mechanical secondary hyperalgesia. The lack of increase in perception to heat in the area of secondary hyperalgesia further suggests that the central mechanisms underlying secondary hyperalgesia involve the facilitation of CNS neurons specifically involved in the transmission of mechanical nociceptive input conveyed by HTMs.

References

- Ali Z, Meyer RA & Campbell JN. (1996). Secondary hyperalgesia to mechanical but not heat stimuli following a capsaicin injection in hairy skin. *Pain* **68**, 401–411.
- Andrew D & Craig AD. (2002). Quantitative responses of spinothalamic lamina I neurones to graded mechanical stimulation in the cat. *J Physiol* **545**, 913–931.
- Baumann TK, Simone DA, Shain CN & LaMotte RH. (1991). Neurogenic hyperalgesia: the search for the primary cutaneous afferent fibers that contribute to capsaicin-induced pain and hyperalgesia. *J Neurophysiol* **66**, 212–227.
- Christensen BN & Perl ER. (1970). Spinal neurons specifically excited by noxious or thermal stimuli: marginal zone of the dorsal horn. *J Neurophysiol* **33**, 293–307.
- Churyukanov M, Plaghki L, Legrain V & Mouraux A. (2012). Thermal detection thresholds of A δ - and C-fibre afferents activated by brief CO₂ laser pulses applied onto the human hairy skin. *PLoS One* **7**, e35817.
- Groppe DM, Urbach TP & Kutas M. (2011). Mass univariate analysis of event-related brain potentials/fields I: a critical tutorial review. *Psychophysiology* **48**, 1711–1725.
- Han ZS, Zhang ET & Craig AD. (1998). Nociceptive and thermoreceptive lamina I neurons are anatomically distinct. *Nat Neurosci* **1**, 218–225.
- Henrich F, Magerl W, Klein T, Greffrath W & Treede RD. (2015). Capsaicin-sensitive C- and A-fibre nociceptors control long-term potentiation-like pain amplification in humans. *Brain* **138**, 2505–2520.
- Iannetti GD, Baumgartner U, Tracey I, Treede RD & Magerl W. (2013). Pinprick-evoked brain potentials: a novel tool to assess central sensitization of nociceptive pathways in humans. *J Neurophysiol* **110**, 1107–1116.
- Kenshalo DR, Jr, Leonard RB, Chung JM & Willis WD. (1982). Facilitation of the response of primate spinothalamic cells to cold and to tactile stimuli by noxious heating of the skin. *Pain* **12**, 141–152.
- Klein T, Magerl W, Hopf HC, Sandkuhler J & Treede RD. (2004). Perceptual correlates of nociceptive long-term potentiation and long-term depression in humans. *J Neurosci* **24**, 964–971.
- Klein T, Stahn S, Magerl W & Treede RD. (2008). The role of heterosynaptic facilitation in long-term potentiation (LTP) of human pain sensation. *Pain* **139**, 507–519.
- LaMotte RH, Shain CN, Simone DA & Tsai EF. (1991). Neurogenic hyperalgesia: psychophysical studies of underlying mechanisms. *J Neurophysiol* **66**, 190–211.
- Light AR & Perl ER. (1979). Spinal termination of functionally identified primary afferent neurons with slowly conducting myelinated fibers. *J Comp Neurol* **186**, 133–150.
- Loeser JD & Treede RD. (2008). The Kyoto protocol of IASP Basic Pain Terminology. *Pain* **137**, 473–477.
- Magerl W, Ali Z, Ellrich J, Meyer RA & Treede RD. (1999). C- and A δ -fiber components of heat-evoked cerebral potentials in healthy human subjects. *Pain* **82**, 127–137.
- Magerl W, Fuchs PN, Meyer RA & Treede RD. (2001). Roles of capsaicin-insensitive nociceptors in cutaneous pain and secondary hyperalgesia. *Brain* **124**, 1754–1764.
- Maris E & Oostenveld R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *J Neurosci Methods* **164**, 177–190.
- McMahon SB & Wall PD. (1984). Receptive fields of rat lamina I projection cells move to incorporate a nearby region of injury. *Pain* **19**, 235–247.
- Meyer RA & Treede RD. (2004). Mechanisms of secondary hyperalgesia: a role for myelinated nociceptors in punctate hyperalgesia. In *Hyperalgesia: Molecular Mechanisms and Clinical Implications (Progress in Pain Research and Management)*, ed. Brune K & Handwerker HO, pp. 143–155. IASP Press, Seattle.
- Mouraux A, Iannetti GD & Plaghki L. (2010). Low intensity intra-epidermal electrical stimulation can activate A δ -nociceptors selectively. *Pain* **150**, 199–207.
- Nahra H & Plaghki L. (2003). The effects of A-fiber pressure block on perception and neurophysiological correlates of brief non-painful and painful CO₂ laser stimuli in humans. *Eur J Pain* **7**, 189–199.
- Nicholls ME, Thomas NA, Loetscher T & Grimshaw GM. (2013). The Flinders Handedness survey (FLANDERS): a brief measure of skilled hand preference. *Cortex* **49**, 2914–2926.
- Reeh PW, Bayer J, Kocher L & Handwerker HO. (1987). Sensitization of nociceptive cutaneous nerve fibers from the rat's tail by noxious mechanical stimulation. *Exp Brain Res* **65**, 505–512.
- Ringkamp M, Peng YB, Wu G, Hartke TV, Campbell JN & Meyer RA. (2001). Capsaicin responses in heat-sensitive and heat-insensitive A-fiber nociceptors. *J Neurosci* **21**, 4460–4468.
- Simone DA, Sorkin LS, Oh U, Chung JM, Owens C, LaMotte RH & Willis WD. (1991). Neurogenic hyperalgesia: central neural correlates in responses of spinothalamic tract neurons. *J Neurophysiol* **66**, 228–246.
- Treede RD & Magerl W. (2000). Multiple mechanisms of secondary hyperalgesia. *Prog Brain Res* **129**, 331–341.
- Treede RD, Meyer RA & Campbell JN. (1998). Myelinated mechanically insensitive afferents from monkey hairy skin: heat-response properties. *J Neurophysiol* **80**, 1082–1093.
- Treede RD, Meyer RA, Raja SN & Campbell JN. (1995). Evidence for two different heat transduction mechanisms in nociceptive primary afferents innervating monkey skin. *J Physiol* **483**, 747–758.

- van den Broeke EN & Mouraux A. (2014). High-frequency electrical stimulation of the human skin induces heterotopical mechanical hyperalgesia, heat hyperalgesia, and enhanced responses to nonnociceptive vibrotactile input. *J Neurophysiol* **111**, 1564–1573.
- Woolf CJ & King AE. (1990). Dynamic alterations in the cutaneous mechanoreceptive fields of dorsal horn neurons in the rat spinal cord. *J Neurosci* **10**, 2717–2726.
- Ziegler EA, Magerl W, Meyer RA & Treede RD. (1999). Secondary hyperalgesia to punctate mechanical stimuli. Central sensitization to A-fibre nociceptor input. *Brain* **122**, 2245–2257.

Additional information

Competing interests

All authors declare that they have no competing interests

Author contributions

ENvdB, CL and AM were involved in the conception and design of the study. ENvdB and CL collected and analysed the data. ENvdB, CL and AM interpreted the data and drafted the manuscript. All authors approved the final version of the manuscript, and agreed to be accountable for all aspects of the work. All authors meet the criteria for authorship.

Funding

ENvdB, CL and AM are supported by the ERC ‘Starting Grant’ (PROBING PAIN 336130).