

# Tight Convergence Rates in Gradient Mapping for the Difference-of-Convex Algorithm

Teodor Rotaru<sup>1,2</sup>[0000–0003–2039–6228], Panagiotis  
Patrinos<sup>1</sup>[0000–0003–4824–7697], and François Glineur<sup>2</sup>[0000–0002–5890–1093]

<sup>1</sup> Department of Electrical Engineering (ESAT-STADIUS), KU Leuven, Kasteelpark  
Arenberg 10, Leuven, 3001, Belgium

{teodor.rotaru, panos.patrinis}@esat.kuleuven.be

<sup>2</sup> Department of Mathematical Engineering (ICTEAM-INMA), UCLouvain, Av.  
Georges Lemaître 4, Louvain-la-Neuve, 1348, Belgium

{teodor.rotaru, francois.glineur}@uclouvain.be

**Abstract.** We establish new theoretical convergence guarantees for the difference-of-convex algorithm (DCA), where the second function is allowed to be weakly-convex, measuring progress via composite gradient mapping. Based on a tight analysis of two iterations of DCA, we identify six parameter regimes leading to sublinear convergence rates toward critical points and establish those rates by proving adapted descent lemmas. We recover existing rates for the standard difference-of-convex decompositions of nonconvex-nonconcave functions, while for all other curvature settings our results are new, complementing recently obtained rates on the gradient residual. Three of our sublinear rates are tight for any number of DCA iterations, while for the other three regimes we conjecture exact rates, using insights from the tight analysis of gradient descent and numerical validation using the performance estimation methodology. Finally, we show how the equivalence between proximal gradient descent (PGD) and DCA allows the derivation of exact PGD rates for any constant stepsize.

**Keywords:** Difference-of-Convex Algorithm · Performance Estimation.

## 1 Introduction

We investigate the difference-of-convex (DC) formulation

$$\underset{x \in \mathbb{R}^d}{\text{minimize}} F(x) := f_1(x) - f_2(x),$$

where  $f_i \in \mathcal{F}_{\mu_i, L_i}$ ,  $i = \{1, 2\}$ , denoting proper, lower semicontinuous (l.s.c.) functions  $f_i : \mathbb{R}^d \rightarrow \mathbb{R}$  having curvature in the interval  $[\mu_i, L_i]$ , i.e.,  $\frac{L_i \|\cdot\|^2}{2} - f_i$  and  $f_i - \frac{\mu_i \|\cdot\|^2}{2}$  are convex. If  $f_i \in \mathcal{C}^2$ , then  $\mu_i$  and  $L_i$  correspond to the minimum and maximum eigenvalues of the Hessian. When  $L_i < \infty$ , then  $f_i$  is smooth. Moreover,  $f_i$  is  $\mu_i$ -convex, including weakly-convexity under  $\mu_i < 0$ . We assume  $\mu_1 \geq 0$  and allow  $\mu_2 < 0$ , which extends the standard DC case

**Algorithm 1:** Difference-of-convex algorithm (DCA)

---

**Data:**  $f_1 \in \mathcal{F}_{\mu_1, L_1}$ ,  $f_2 \in \mathcal{F}_{\mu_2, L_2}$ , with  $\mu_1 \geq 0$ ,  $\mu_1 + \mu_2 > 0$  or  $\mu_1 = \mu_2 = 0$ ;  
 $N \geq 1$  iterations starting from  $x^0 \in \text{dom } \partial f_1$

**1** for  $k = 0, \dots, N$  **do**  
**2**   Select  $g_2^k \in \partial f_2(x^k)$  ;  
**3**   Select  $x^{k+1} \in \text{argmin}_{w \in \mathbb{R}^d} \{f_1(w) - \langle g_2^k, w \rangle\}$

**Result:** Best iterate  $x^k$  with  $k = \text{argmin}_{0 \leq k \leq N} \{\|x^k - x^{k+1}\|^2\}$

---

where both functions are convex. The difference-of-convex algorithm (DCA), stated in [Algorithm 1](#), is a standard method to solve the DC problem. Various work provide extensive analysis of DCA [[6,20,10,13](#)], which is also referred as the convex-concave procedure (CCCP) [[27,11,14](#)]. In this work we focus on improving the theoretical guarantees. For the interested reader, a comprehensive list of DCA applications is found in the review paper [[13](#)].

If the entire sequence of iterates  $\{x^k\}$  is bounded, then each accumulation point of it is a critical point of  $F$  [[6](#), Theorem 3(iv)]. Inspired by this result, we aim to find the best iterate  $x^k$  providing the lowest value of  $\{\|x^k - x^{k+1}\|\}$  over  $N$  iterations. This metric is referred as the *best (composite) gradient mapping* [[15,16](#)] or the *prox-gradient mapping* [[5](#)], and is complementary to the *best residual gradient norm*  $\{\|g_1^k - g_2^k\|\}$  over  $N$  iterations, where  $g_i \in \partial f_i(x^k)$ ,  $i = \{1, 2\}$ , are any subgradients of  $f_i$  evaluated at  $x^k$ . On this metric, [[3](#)] provides the first tight rates for any convex functions  $f_i$  when  $F$  is nonconvex and nonconcave, using the performance estimation problem (PEP) methodology [[7,21,22](#)]. The analysis is later refined and extended in [[19](#)] to accommodate weakly-convex functions  $f_2$ . For the standard DC setup, [[8](#)] provides convergence rates in terms of Bregman distance, viewing DCA as a instance of the Bregman proximal point algorithm, while [[12,26](#)] provides linear rates under the Polyak-Łojasiewicz (PL) inequality. The Frank-Wolfe algorithm [[28](#)] and the proximal gradient descent (PGD) [[13](#)] are also equivalent to DCA.

*Contributions.* In [Theorem 1](#) we establish new theoretical convergence guarantees for DCA and PGD, measuring progress via composite gradient mapping. When both  $f_1$  and  $f_2$  are convex and  $F$  is nonconvex-nonconcave, we refine the analysis from [[3](#)], providing standard descent lemmas in [Section 2.1](#). Our results extend existing work by introducing new rates for all other curvature settings, complementing the best gradient residual rates from [[3,19](#)]. Building on insights from gradient descent (GD), we conjecture in [Conjecture 1](#) the exact behavior of DCA across all curvature choices, with some regimes rigorously proven ([Theorems 2 and 3](#)) and others remaining open. As shown in [Section 4](#), the equivalence between PGD and DCA enables us to formulate precise PGD rates for any constant stepsize, including the ones exceeding the inverse Lipschitz constant. Finally, we propose a general adaptive curvature shifting procedure for DCA, leveraging PGD stepsize schedules aligned with established GD strategies.

Our derivations are aided by the PEP technique, allowing to establish the tightness of the regimes and validate our conjecture. The setup of PEP for DCA is described in [3,19] and can be solved using dedicated software packages [23,9].

*Theoretical setup and running assumptions.* Let  $f$  be a proper, convex, l.s.c. function. Its subdifferential at  $x \in \mathbb{R}^d$  is  $\partial f(x) := \{g \in \mathbb{R}^d \mid f(y) \geq f(x) + \langle g, y - x \rangle, \forall y \in \mathbb{R}^d\}$ . If  $f$  is weakly-convex with  $\mu < 0$ , let  $\tilde{f}(x) := f(x) - \mu \frac{\|x\|^2}{2}$  be a convex function. Since its subdifferential  $\partial \tilde{f}(x)$  is well-defined, we take  $\partial f(x) := \partial \tilde{f}(x) + \mu x$  [4, Proposition 6.3], which is well-defined. For  $f$  differentiable at  $x$  we have  $\partial f(x) = \{\nabla f(x)\}$ . The domain and range of  $f$  are  $\text{dom } f := \{x \in \mathbb{R}^d : f(x) < \infty\}$  and  $\text{range } f := \{y \in \mathbb{R} : \exists x \in \text{dom } f \text{ with } y = f(x)\}$ . The domain and range of the subdifferential are  $\text{dom } \partial f = \{x \in \mathbb{R}^d : \partial f(x) \neq \emptyset\}$  and  $\text{range } \partial f = \cup \{\partial f(x) : x \in \text{dom } \partial f\}$ . The convex conjugate of  $f$  l.s.c. function is defined as  $f^*(y) := \sup_{x \in \text{dom } f} \{\langle y, x \rangle - f(x)\}$ , where  $f^*$  is closed and convex.

**Assumption 1** *The objective function  $F = f_1 - f_2$  is lower bounded by  $F_{l_0} := \inf_x F$ , where  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$  with  $0 \leq \mu_1 < L_1 \leq \infty$ ,  $\mu_2 \in (-\infty, \infty)$  and  $L_2 \in (\mu_2, \infty]$ . Additionally, assume  $\emptyset \neq \text{dom } \partial f_1 \subseteq \text{dom } \partial f_2$  and  $\text{range } \partial f_2 \subseteq \text{range } \partial f_1$ .*

Note that  $F \in \mathcal{F}_{\mu_1 - L_2, L_1 - \mu_2}$ . The second part of [Assumption 1](#) implies that the DCA iterations are well-defined, meaning there exists a sequence  $\{x^k\}$ , starting from  $x^0 \in \text{dom } \partial f_1$ , generated by [Algorithm 1](#), such that  $x^{k+1} \in \partial f_1^*(\partial f_2(x^k))$ . By definition, this condition implies the existence of  $g_1^{k+1} \in \partial f_1(x^{k+1})$  such that  $g_1^{k+1} = g_2^k$ , where  $g_2^k \in \partial f_2(x^k)$ . This is the only characterization of DCA iterations employed in our derivations.

DCA finds critical points  $x^*$  satisfying  $\partial f_2(x^*) \cap \partial f_1(x^*) \neq \emptyset$ . These are stationary only if  $f_2$  is smooth [17, Exercise 10.10]. If only  $f_1$  is smooth, then only the inclusion  $\partial(-f_2)(x^*) \subseteq -\partial f_2(x^*)$  is guaranteed [17, Corollary 9.21], hence  $\partial F(x^*) \subseteq \nabla f_1(x^*) - \partial f_2(x^*)$  and critical points are not necessary stationary.

We assume that at each iteration a subgradient from  $\partial f_1^*$  is available and only focus on the progress of the iterations  $\min_{0 \leq k \leq N} \{\|x^k - x^{k+1}\|^2\}$  subject to the initial condition  $F(x^0) - F(x^{N+1}) \leq \Delta$ , with  $\Delta > 0$ . To remove the dependence of the initial condition on the last iterate, in all our rates the initial objective gap can be further upper bounded by  $F(x^0) - F_{l_0}$ .

## 2 Sublinear convergence rates

**Theorem 1.** *Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$  satisfying [Assumption 1](#) and  $\mu_1 + \mu_2 > 0$ . Then after  $N + 1 \geq 2$  iterations of DCA starting from  $x^0$  it holds*

$$\frac{1}{2} \min_{0 \leq k \leq N} \{\|x^k - x^{k+1}\|^2\} \leq \frac{F(x^0) - F(x^{N+1})}{(\mu_1 + \mu_2) + p_i(\mu_1, L_1, \mu_2, L_2)N}, \quad (1)$$

where the expressions of the regimes  $p_{i=\{1, \dots, 6\}}$  are given in [Table 1](#).

The proof is deferred to [Section 2.2](#). All regimes are exact for  $N = \{0, 1\}$  and their domains span the entire range of parameters satisfying the condition  $\mu_1 + \mu_2 > 0$ , which is sufficient for decreasing the objective (see [Proposition 1](#)).

**Table 1.** Coefficients  $p_i$  from [Theorem 1](#). Their domains cover all parameter space under the condition  $\mu_1 + \mu_2 > 0$ . Regimes  $p_{\{1,2,3\}}$  are tight for any number of iterations. (s.c.: strongly convex; n.c.: nonconvex; ccv.: concave; n.ccv.: nonconcave)

Regime	Domain		Description
$p_1 = \mu_1 + \mu_2 + \frac{(\mu_1 - \mu_2)^2}{L_2 - \mu_2}$	$\mu_1 \geq \mu_2 \geq 0$	$L_2 > \mu_1$	$f_1, f_2$ convex $F$ n.c.-n.ccv.
$p_2 = \mu_1 + \mu_2 + \frac{(\mu_1 - \mu_2)^2}{L_1 - \mu_1}$	$\mu_2 \geq \mu_1 \geq 0$	$L_1 > \mu_2$	$f_1, f_2$ convex $F$ n.c.-n.ccv.
$p_3 = \frac{(\mu_1 + \mu_2)(L_2 + \mu_1)}{L_2 + \mu_2}$	$\mu_2 \in [\frac{-L_2\mu_1}{L_2 + \mu_1}, 0)$	$L_2 > \mu_1 > 0$	$f_1$ s.c., $f_2$ n.c. $F$ n.c.-n.ccv.
$p_4 = \frac{\mu_1^2(L_2 + \mu_1)}{L_2^2}$	$\mu_2 \in [\frac{-L_2\mu_1}{L_2 + \mu_1}, L_2)$	$L_2 \in [0, \mu_1]$	$f_1$ s.c., $f_2$ n.c. $F$ convex
$p_5 = \frac{\mu_1^2(\mu_1 + \mu_2)}{\mu_2^2}$	$\mu_2 < 0, \mu_2 \in (-\mu_1, \frac{-L_2\mu_1}{L_2 + \mu_1}]$		$f_1$ s.c., $f_2$ n.c. $F$ n.ccv.
$p_6 = \frac{\mu_2^2(L_1 + \mu_2)}{L_1^2}$	$\mu_2 \geq \mu_1 \geq 0$	$L_1 \in (0, \mu_2]$	$f_1$ convex, $f_2$ s.c. $F$ ccv.

Based on extensive numerical experiments using PEP, we determine that regimes  $p_{\{1,2,3\}}$  are tight for any number of iterations. Regimes  $p_1$  and  $p_2$  are derived in [[3](#), Proposition 3.1], by applying Toland duality [[25](#)] on the rates measuring the best gradient residual criterion. [Theorem 1](#) gives more explicit expressions and includes in the proofs a simpler derivation based on [Lemmas 2](#) and [3](#). These regimes assume  $F$  is nonconvex-nonconcave ( $L_1 > \mu_2$  and  $L_2 > \mu_1$ ), with the active one indexed by  $\operatorname{argmax}_i\{\mu_1, \mu_2\}$ . All other regimes are new.

When  $F$  is nonconvex-nonconcave, regime  $p_3$  is active when  $f_2$  is weakly-convex with  $\mu_2 \in [\frac{-L_2\mu_1}{L_2 + \mu_1}, 0)$ . When  $\mu_2$  is even lower, then regime  $p_5$  is active. This is however not tight for  $N \geq 2$ , where a more complicated analysis is required. The exact rates expressions in this case are given in [Conjecture 1](#).

When  $F$  is (strongly) convex ( $\mu_1 \geq L_2$ ), the sublinear rates of regimes  $p_4$  and  $p_5$  are tight only for  $N = 1$ . These regimes are separated in this case by the threshold  $\mu_2 = \frac{-L_2\mu_1}{L_2 + \mu_1}$ . When  $N \geq 2$ , we prove a linear rate for regime  $p_4$  in [Theorem 2](#), which is confirmed to be exact using PEP. Within the particular range of  $p_5$  satisfying the condition  $L_2 + \mu_2 \leq 0$ , we demonstrate a linear rate in [Theorem 3](#). For the rest domain of  $p_5$ , we conjecture the tight rates in [Conjecture 1](#). Finally, regime  $p_6$  characterizes the iterates over *concave* objectives  $F$  and is included for a complete covering of the parameter space.

## 2.1 Lemmas to prove sublinear rates

For compactness of the proofs, at each iteration  $k$  we denote  $\Delta x^k := x^k - x^{k+1}$ ,  $f_1^k = f_1(x^k)$ ,  $f_2^k = f_2(x^k)$ ,  $\Delta F(x^k) := F(x^k) - F(x^{k+1})$ ,  $g_1^k \in \partial f_1(x^k)$ ,  $g_2^k \in \partial f_2(x^k)$ ,  $G^k := g_1^k - g_2^k$ . Recall that  $g_1^{k+1} = g_2^k$  for all DCA iterations.

**Lemma 1.** *Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$ , with  $L_2 > \mu_2$  and  $L_1 > \mu_1 \geq 0$ . Consider one DCA iteration connecting  $x^k$  and  $x^{k+1}$ . Let  $g_1^j \in \partial f_1(x^j)$ ,  $g_2^j \in$*

$\partial f_2(x^j)$  and  $G^j = g_1^j - g_2^j$ , where  $j = \{k, k+1\}$ . The following inequalities hold:

$$\Delta F(x^k) \geq \frac{\mu_1 + \mu_2}{2} \|\Delta x^k\|^2 + \frac{\|G^{k+1} - \mu_2 \Delta x^k\|^2}{2(L_2 - \mu_2)} + \frac{\|G^k - \mu_1 \Delta x^k\|^2}{2(L_1 - \mu_1)}; \quad (B^k)$$

$$\langle G^k, \Delta x^k \rangle \geq \mu_1 \|\Delta x^k\|^2 + \frac{\|G^k - \mu_1 \Delta x^k\|^2}{L_1 - \mu_1}; \quad (C_{f_1}^k)$$

$$\langle G^{k+1}, \Delta x^k \rangle \geq \mu_2 \|\Delta x^k\|^2 + \frac{\|G^{k+1} - \mu_2 \Delta x^k\|^2}{L_2 - \mu_2}. \quad (C_{f_2}^k)$$

*Proof.* Given  $\mu < L$  and  $f \in \mathcal{F}_{\mu, L}$ , from [21, Theorem 4], [22, Theorem 3.10], [18, Theorem 3.2] it holds  $\forall x, y \in \mathbb{R}^d$ , with  $g^x \in \partial f(x)$ ,  $g^y \in \partial f(y)$ , that

$$f(x) - f(y) - \langle g^y, x - y \rangle \geq \frac{\mu}{2} \|x - y\|^2 + \frac{1}{2(L - \mu)} \|g^x - g^y - \mu(x - y)\|^2. \quad (2)$$

We exploit the property  $g_1^{k+1} = g_2^k$ , implying  $G^k = g_1^k - g_2^k = g_1^k - g_1^{k+1}$  and  $G^{k+1} = g_1^{k+1} - g_2^{k+1} = g_2^k - g_2^{k+1}$ . By writing inequality (2) for:  $f_1$  with the iterates  $(x^k, x^{k+1})$  and  $(x^{k+1}, x^k)$  we get (3) and (4), respectively;  $f_2$  with the iterates  $(x^k, x^{k+1})$  and  $(x^{k+1}, x^k)$  we obtain (5) and (6), respectively:

$$f_1^k - f_1^{k+1} - \langle g_1^{k+1}, \Delta x^k \rangle \geq \frac{\mu_1}{2} \|\Delta x^k\|^2 + \frac{\|G^k - \mu_1 \Delta x^k\|^2}{2(L_1 - \mu_1)}; \quad (3)$$

$$f_1^{k+1} - f_1^k + \langle g_1^k, \Delta x^k \rangle \geq \frac{\mu_1}{2} \|\Delta x^k\|^2 + \frac{\|G^k - \mu_1 \Delta x^k\|^2}{2(L_1 - \mu_1)}; \quad (4)$$

$$f_2^k - f_2^{k+1} - \langle g_2^{k+1}, \Delta x^k \rangle \geq \frac{\mu_2}{2} \|\Delta x^k\|^2 + \frac{\|G^{k+1} - \mu_2 \Delta x^k\|^2}{2(L_2 - \mu_2)}; \quad (5)$$

$$f_2^{k+1} - f_2^k + \langle g_2^k, \Delta x^k \rangle \geq \frac{\mu_2}{2} \|\Delta x^k\|^2 + \frac{\|G^{k+1} - \mu_2 \Delta x^k\|^2}{2(L_2 - \mu_2)}. \quad (6)$$

Then summing (3) and (6) gives  $(B^k)$ ; inequality  $(C_{f_1}^k)$  results by summing (3) and (4); inequality  $(C_{f_2}^k)$  is obtained by summing (5) and (6).  $\square$

**Proposition 1 (Decrease after one iteration).** *Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$  satisfying Assumption 1. Consider one DCA iteration connecting  $x^k$  and  $x^{k+1}$ . Then it holds*

$$F(x^k) - F(x^{k+1}) \geq \frac{\mu_1 + \mu_2}{2} \|x^k - x^{k+1}\|^2. \quad (7)$$

Consequently, the objective function  $F$  decreases after each DCA iteration if  $\mu_1 + \mu_2 \geq 0$  and the decrease is strict if  $\mu_1 + \mu_2 > 0$ , unless  $x^{k+1} = x^k$ .

*Proof.* Inequality (7) results by ignoring the mixed squares in  $(B^k)$ .  $\square$

Proposition 1 was first proved in [6, Theorem 3, Proposition 2].

*Remark 1 (Proofs structure).* The proofs are based on multiplying the inequalities from Lemma 1 with nonnegative weights. For regimes  $p_{1,3,4,5}$  we sum:

$$B^k + \beta_1 \times C_{f_1}^{k+1} + \beta_2 \times C_{f_2}^k, \quad (8)$$

while for the proofs of  $p_{2,6}$  we sum:  $B^{k+1} + \beta_1 \times C_{f_1}^k + \beta_2 \times C_{f_2}^{k+1}$ , where  $B^{\{k, k+1\}}$ ,  $C_{f_1}^{\{k, k+1\}}$ ,  $C_{f_2}^{\{k, k+1\}}$  are instances of inequalities  $(B^k)$ ,  $(C_{f_1}^k)$  and  $(C_{f_2}^k)$ , respectively. The weights  $\beta_1$  and  $\beta_2$  are functions of the curvature parameters.

The demonstrations rewrite the inequalities from [Remark 1](#) by building specific mixed squared norms of  $G$  and  $\Delta x$  at specific indices  $k$ , using [Property 1](#). For space reasons, we only give the final inequalities containing all the squares.

*Property 1.* The following identities hold, where  $i, j \in \{1, 2\}$  and  $G$  and  $\Delta x$  are specialized when building the squares within the proofs:

$$-\langle G, \Delta x \rangle = \frac{1}{2\mu_j} [\|G - \mu_j \Delta x\|^2 - \|G\|^2 - \|\mu_j \Delta x\|^2] \text{ and}$$

$$\|G - \mu_i \Delta x\|^2 = \frac{\mu_i}{\mu_j} \|G - \mu_j \Delta x\|^2 - \frac{\mu_i}{\mu_j} (1 - \frac{\mu_i}{\mu_j}) \|\mu_j \Delta x\|^2 + (1 - \frac{\mu_i}{\mu_j}) \|G\|^2.$$

**Lemma 2 (Regime  $p_1$ ).** *Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$ , with  $L_2 > \mu_1 \geq \mu_2 \geq 0$ . Then after two DCA iterations connecting  $x^k$ ,  $x^{k+1}$  and  $x^{k+2}$  it holds  $\Delta F(x^k) \geq (\mu_1 + \mu_2 \frac{L_2 - \mu_1}{L_2 - \mu_2}) \frac{1}{2} \|\Delta x^k\|^2 + \mu_1 \frac{\mu_1 - \mu_2}{L_2 - \mu_2} \frac{1}{2} \|\Delta x^{k+1}\|^2$ .*

*Proof.* Let  $\beta_2 = 0$  and  $\beta_1 := \frac{\mu_1 - \mu_2}{L_2 - \mu_2}$ , which is nonnegative since  $\mu_1 \geq \mu_2$ . After replacing in [\(8\)](#), building squares and performing simplifications we get:

$$\begin{aligned} \Delta F(x^k) &\geq (\mu_1 + \mu_2 \frac{L_2 - \mu_1}{L_2 - \mu_2}) \frac{1}{2} \|\Delta x^k\|^2 + \mu_1 \frac{\mu_1 - \mu_2}{L_2 - \mu_2} \frac{1}{2} \|\Delta x^{k+1}\|^2 + \\ &\quad \frac{\|G^k - \mu_1 \Delta x^k\|^2}{2(L_1 - \mu_1)} + \frac{\mu_2 \|G^{k+1} - \mu_1 \Delta x^k\|^2}{2\mu_1(L_2 - \mu_2)} + \frac{(L_1 + \mu_1)(\mu_1 - \mu_2) \|G^{k+1} - \mu_1 \Delta x^{k+1}\|^2}{2\mu_1(L_1 - \mu_1)(L_2 - \mu_2)}. \end{aligned}$$

Since  $\mu_1 \geq \mu_2 \geq 0$ , all coefficients of the squares are positive and the conclusion follows by dropping the mixed terms.  $\square$

**Lemma 3 (Regime  $p_2$ ).** *Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$ , with  $L_1 > \mu_2 \geq \mu_1 \geq 0$ . Then after two DCA iterations connecting  $x^k$ ,  $x^{k+1}$  and  $x^{k+2}$  it holds  $\Delta F(x^{k+1}) \geq \mu_2 \frac{\mu_2 - \mu_1}{L_1 - \mu_1} \frac{1}{2} \|\Delta x^k\|^2 + (\mu_2 + \mu_1 \frac{L_1 - \mu_2}{L_1 - \mu_1}) \frac{1}{2} \|\Delta x^{k+1}\|^2$ .*

*Proof.* Let  $\beta_1 = 0$  and  $\beta_2 := \frac{\mu_2 - \mu_1}{L_1 - \mu_1}$ , nonnegative since  $\mu_2 \geq \mu_1$ , be the specific multipliers from [Remark 1](#). Building the mixed squares we obtain:

$$\begin{aligned} \Delta F(x^{k+1}) &\geq (\mu_2 + \mu_1 \frac{L_1 - \mu_2}{L_1 - \mu_1}) \frac{1}{2} \|\Delta x^{k+1}\|^2 + \mu_2 \frac{\mu_2 - \mu_1}{2(L_1 - \mu_1)} \|\Delta x^k\|^2 + \\ &\quad \frac{\|G^{k+2} - \mu_2 \Delta x^{k+1}\|^2}{2(L_2 - \mu_2)} + \frac{\mu_1 \|G^{k+1} - \mu_2 \Delta x^{k+1}\|^2}{2\mu_2(L_1 - \mu_1)} + \frac{(L_2 + \mu_2)(\mu_2 - \mu_1) \|G^{k+1} - \mu_2 \Delta x^k\|^2}{2\mu_2(L_2 - \mu_2)(L_1 - \mu_1)}. \end{aligned}$$

Since  $\mu_2 \geq \mu_1 \geq 0$ , all coefficients of the squares are positive and the conclusion follows by dropping the mixed terms.  $\square$

*Property 2 ( $T_1$ ).* When  $\mu_2 < 0$ , we define  $T_1 := \mu_1 (\frac{L_2 + \mu_1}{L_2} - \frac{\mu_1 + \mu_2}{\mu_2^2})$ . One can check that  $T_1 \geq 0$  if  $\mu_2 \in (-\mu_1, \frac{-L_2 \mu_1}{L_2 + \mu_1}]$  and  $T_1 \leq 0$  if  $\mu_2 \in [\frac{-L_2 \mu_1}{L_2 + \mu_1}, L_2)$ .

**Lemma 4 (Regime  $p_3$ ).** *Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$ , with  $L_2 > \mu_1 > 0$ ,  $\mu_2 \in [\frac{-L_2 \mu_1}{L_2 + \mu_1}, 0)$  and  $\mu_1 + \mu_2 > 0$ . Consider two DCA iterations connecting  $x^k$ ,  $x^{k+1}$  and  $x^{k+2}$ . Then  $\Delta F(x^k) \geq (\mu_1 + \frac{L_2 \mu_2}{L_2 + \mu_2}) \frac{1}{2} \|\Delta x^k\|^2 + \frac{\mu_1^2}{2(L_2 + \mu_2)} \|\Delta x^{k+1}\|^2$ .*

*Proof.* Let  $\beta_1 := \frac{\mu_1}{L_2 + \mu_2}$  and  $\beta_2 := \frac{-\mu_2}{L_2 + \mu_2}$ , which are nonnegative since  $\mu_1 > -\mu_2 > 0$  and  $L_2 + \mu_2 \geq \frac{-\mu_2 L_2}{\mu_1} > 0$ . After replacing in (8), building squares and performing simplifications we obtain:

$$\begin{aligned} \Delta F(x^k) \geq & \left(\mu_1 + \frac{L_2 \mu_2}{L_2 + \mu_2}\right) \frac{1}{2} \|\Delta x^k\|^2 + \frac{\mu_1^2}{2(L_2 + \mu_2)} \|\Delta x^{k+1}\|^2 + \\ & \frac{\|G^k - \mu_1 \Delta x^k\|^2}{2(L_1 - \mu_1)} + \frac{(L_1 + \mu_1) \|G^{k+1} - \mu_1 \Delta x^{k+1}\|^2}{2(L_1 - \mu_1)(L_2 + \mu_2)}. \end{aligned}$$

Since  $L_2 > -\mu_2$ , the coefficients of the mixed terms are positive and the conclusion follows by dropping them. The coefficient of  $\|\Delta x^k\|^2$  is positive if  $T_1 \leq 0$  as it can be factorized as  $\frac{L_2^2 \mu_2^2}{\mu_1(L_2^2 - \mu_2^2)}(-T_1)$ . The inequality holds even for  $T_1 > 0$ .  $\square$

**Lemma 5 (Regime  $p_4$ ).** *Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$ , with  $L_2 \in [0, \mu_1]$ ,  $\mu_2 \in [\frac{-L_2 \mu_1}{L_2 + \mu_1}, L_2)$  and  $\mu_1 + \mu_2 > 0$ . Consider two DCA iterations connecting  $x^k$ ,  $x^{k+1}$  and  $x^{k+2}$ . Then it holds  $\Delta F(x^k) \geq \frac{\mu_1^2(L_2 + \mu_1)}{L_2^2} \frac{1}{2} \|\Delta x^{k+1}\|^2$ .*

*Proof.* Let  $\beta_1 := \frac{\mu_1(\mu_1 + L_2)}{L_2^2}$  and  $\beta_2 := \frac{\mu_1}{L_2}$ , both positive. After replacing in (8), building the mixed squares and performing simplifications we get:

$$\begin{aligned} \Delta F(x^k) \geq & \frac{\mu_1^2(L_2 + \mu_1)}{L_2^2} \frac{1}{2} \|\Delta x^{k+1}\|^2 + \frac{\|G^k - \mu_1 \Delta x^k\|^2}{2(L_1 - \mu_1)} + \\ & \frac{(L_1 + \mu_1)(L_2 + \mu_1) \|G^{k+1} - \mu_1 \Delta x^{k+1}\|^2}{L_2^2(L_1 - \mu_1)} + \frac{\mu_2^2(-T_1) \|G^{k+1} - L_2 \Delta x^k\|^2}{\mu_1(L_2 - \mu_2)^2}. \end{aligned}$$

Since  $T_1 \leq 0$  (Property 2), the conclusion follows.  $\square$

**Lemma 6 (Regime  $p_5$ ).** *Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$ , with  $\mu_1 > -\mu_2 > 0$  such that  $\mu_2 \in (-\mu_1, \frac{-L_2 \mu_1}{L_2 + \mu_1}]$ . Consider two DCA iterations connecting  $x^k$ ,  $x^{k+1}$  and  $x^{k+2}$ . Then it holds  $\Delta F(x^k) \geq \frac{\mu_1^2(\mu_1 + \mu_2)}{\mu_2^2} \frac{1}{2} \|\Delta x^{k+1}\|^2$ .*

*Proof.* Let  $\beta_1 := \frac{\mu_1(\mu_1 + \mu_2)}{\mu_2^2}$  and  $\beta_2 := \frac{\mu_1 + \mu_2}{-\mu_2}$ , both positive. After replacing in (8), building the mixed squares and performing simplifications we get:

$$\begin{aligned} \Delta F(x^k) \geq & \frac{\mu_1^2(\mu_1 + \mu_2)}{2\mu_2^2} \|\Delta x^{k+1}\|^2 + \frac{\|G^k - \mu_1 \Delta x^k\|^2}{2(L_1 - \mu_1)} + \\ & \frac{(L_1 + \mu_1)(\mu_1 + \mu_2) \|G^{k+1} - \mu_1 \Delta x^{k+1}\|^2}{\mu_2^2(L_1 - \mu_1)} + \frac{L_2^2 T_1 \|G^{k+1} - \mu_2 \Delta x^k\|^2}{\mu_1(L_2 - \mu_2)^2}. \end{aligned}$$

Since  $T_1 \geq 0$  (Property 2), the conclusion follows.  $\square$

**Lemma 7 (Regime  $p_6$ ).** *Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$ , with  $\mu_2 \geq \mu_1 \geq 0$  and  $L_1 \in (0, \mu_2]$ . Then after two DCA iterations connecting  $x^k$ ,  $x^{k+1}$  and  $x^{k+2}$  we have  $\Delta F(x^{k+1}) \geq \frac{\mu_2^2(L_1 + \mu_2)}{L_1^2} \frac{1}{2} \|\Delta x^k\|^2$ .*

*Proof.* Let  $\beta_2 := \frac{\mu_2(\mu_2 + L_1)}{L_1^2}$  and  $\beta_1 := \frac{\mu_2}{L_1}$ , both positive. After replacing in (8), building the mixed squares and performing simplifications we get:

$$\begin{aligned} \Delta F(x^{k+1}) \geq & \frac{\mu_2^2(L_1 + \mu_2)}{L_1^2} \frac{1}{2} \|\Delta x^k\|^2 + \frac{\|G^{k+2} - \mu_2 \Delta x^{k+1}\|^2}{2(L_2 - \mu_2)} + \\ & \frac{(L_1 + \mu_2)(L_2 + \mu_2) \|G^{k+1} - \mu_2 \Delta x^k\|^2}{L_1^2(L_2 - \mu_2)} + \frac{\mu_1 \mu_2 + L_1(\mu_1 + \mu_2) \|G^{k+1} - L_1 \Delta x^{k+1}\|^2}{L_1^2(L_1 - \mu_1)}. \end{aligned}$$

The conclusion follows by dropping all mixed squares terms.  $\square$

## 2.2 Proof of Theorem 1

*Proof (of Theorem 1).* For regimes  $p_1$ ,  $p_3$ ,  $p_4$  and  $p_5$ , by telescoping inequalities from Lemma 2, Lemma 4, Lemma 5 and Lemma 6, respectively, for  $k = \{0, \dots, N-1\}$  and taking the minimum over iterations differences norms:

$$F(x^0) - F(x^N) \geq p_i(\mu_1, L_1, \mu_2, L_2)N \frac{1}{2} \min_{0 \leq k \leq N} \{\|x^k - x^{k+1}\|^2\}, \quad i = \{1, 3, 4, 5\},$$

where  $p_i$  denotes the sum of the two coefficients multiplying  $\frac{1}{2}\|x^k - x^{k+1}\|^2$  and  $\frac{1}{2}\|x^{k+1} - x^{k+2}\|^2$  in each inequality. Writing (7) for the iterates  $x^N$  and  $x^{N+1}$ , we have  $F(x^N) - F(x^{N+1}) \geq \frac{\mu_1 + \mu_2}{2}\|x^N - x^{N+1}\|^2$ . Then the rate results by summing it to the previous inequality and taking the minimum in the right-hand side. For regimes  $p_2$  and  $p_6$ , by telescoping inequalities from Lemmas 3 and 7, respectively, for  $k = \{1, \dots, N\}$  and taking the minimum over iterations:

$$F(x^1) - F(x^{N+1}) \geq p_i(\mu_1, L_1, \mu_2, L_2)N \frac{1}{2} \min_{0 \leq k \leq N} \{\|x^k - x^{k+1}\|^2\}, \quad i = \{2, 6\}.$$

Writing (7) for the iterates  $x^0$  and  $x^1$ , summing it up and taking the minimum in the right-hand side leads to the rate.  $\square$

## 3 Tight convergence rates for any number of iterations

**Notation.** We define  $E_k : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ ,  $E_k(z) := \sum_{j=1}^{2k} z^{-j}$ , which is  $\frac{-1+z^{-2k}}{1-z}$ , if  $z \neq 1$ , and  $2k$  is  $z = 1$ . Note that  $E_0(z) = 0$ . We denote  $[z]_+ := \max\{0, z\}$ .

### 3.1 Subdomains with proved rates

**Theorem 2.** Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$ , with  $L_2 \in [0, \mu_1]$ ,  $\mu_1 + \mu_2 > 0$  and  $\mu_2 \in [\frac{-L_2\mu_1}{L_2+\mu_1}, L_2)$ . Then after  $N+1 \geq 2$  iterations of DCA starting from  $x^0$  it holds:  $\frac{1}{2}\|x^N - x^{N+1}\|^2 \leq \frac{F(x^0) - F(x^{N+1})}{\mu_1 + \mu_2 + \mu_1 E_N(\frac{L_2}{\mu_1})}$ .

*Proof.* We take  $\beta_1 = E_{k+1}(\frac{L_2}{\mu_1})$  and  $\beta_2 = -1 + \frac{L_2}{\mu_1} E_{k+1}(\frac{L_2}{\mu_1})$  in (8), where  $k = 0, \dots, N-1$ . After building the squares and simplifications we get:

$$\begin{aligned} \Delta F(x^k) &\geq -E_k(\frac{L_2}{\mu_1})\frac{\mu_1}{2}\|\Delta x^k\|^2 + E_{k+1}(\frac{L_2}{\mu_1})\frac{\mu_1}{2}\|\Delta x^{k+1}\|^2 + \frac{\|G^k - \mu_1 \Delta x^k\|^2}{2(L_1 - \mu_1)} + \\ &\quad \frac{(L_1 + \mu_1)E_{k+1}(\frac{L_2}{\mu_1})\|G^{k+1} - \mu_1 \Delta x^{k+1}\|^2}{2\mu_1(L_1 - \mu_1)} + \frac{[-1 + \frac{L_2 + \mu_2}{\mu_1} E_{k+1}(\frac{L_2}{\mu_1})]\|G^{k+1} - L_2 \Delta x^k\|^2}{2(L_2 - \mu_2)}. \end{aligned}$$

We have  $-1 + \frac{L_2 + \mu_2}{\mu_1} E_{k+1}(\frac{L_2}{\mu_1}) \geq -1 + \frac{L_2 + \mu_2}{\mu_1} E_1(\frac{L_2}{\mu_1}) = \frac{\mu_2^2}{\mu_1(L_2 - \mu_2)}(-T_1) \geq 0$ , since  $E_{k+1}(\frac{L_2}{\mu_1})$  is increasing with  $k$ , thus the mixed terms can be dropped. Telescoping for  $k = 0, \dots, N-1$  and adding  $(B^k)$  with  $k = N$  gives the conclusion.  $\square$

**Theorem 3.** Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$ , with  $\mu_1 + \mu_2 > 0$  and  $L_2 + \mu_2 \leq 0$ . Then after  $N+1 \geq 2$  iterations of DCA starting from  $x^0$  it holds  $\frac{1}{2}\|x^N - x^{N+1}\|^2 \leq \frac{F(x^0) - F(x^{N+1})}{\mu_1 + \mu_2 + \mu_1 E_N(\frac{\mu_2}{\mu_1})}$ .

*Proof.* We take  $\beta_1 = E_{k+1}(\frac{\mu_2}{\mu_1})$  and  $\beta_2 = -\frac{\mu_2}{\mu_1}E_{k+1}(\frac{\mu_2}{\mu_1})$  in (8), where  $k = 0, \dots, N-1$ . After building the squares and simplifications we get:

$$\begin{aligned} \Delta F(x^k) \geq & -E_k(\frac{\mu_2}{\mu_1})\frac{\mu_1}{2}\|\Delta x^k\|^2 + E_{k+1}(\frac{\mu_2}{\mu_1})\frac{\mu_1}{2}\|\Delta x^{k+1}\|^2 + \frac{\|G^k - \mu_1 \Delta x^k\|^2}{2(L_1 - \mu_1)} + \\ & \frac{(L_1 + \mu_1)E_{k+1}(\frac{\mu_2}{\mu_1})\|G^{k+1} - \mu_1 \Delta x^{k+1}\|^2}{2\mu_1(L_1 - \mu_1)} + \frac{[1 - \frac{L_2 + \mu_2}{\mu_1}E_{k+1}(\frac{\mu_2}{\mu_1})]\|G^{k+1} - \mu_2 \Delta x^k\|^2}{2(L_2 - \mu_2)}. \end{aligned}$$

$E_{k+1}(\frac{\mu_2}{\mu_1})$  is positive and monotonically increasing with index  $k$  as  $\mu_1 + \mu_2 > 0$ . Then due to  $L_2 + \mu_2 \leq 0$ , we can drop all mixed squares. Telescoping for  $k = 0, \dots, N-1$  and adding  $(B^k)$  with  $k = N$  gives the conclusion.  $\square$

### 3.2 Conjectures for any number of iterations

*Conjecture 1.* Let  $f_1 \in \mathcal{F}_{\mu_1, L_1}$  and  $f_2 \in \mathcal{F}_{\mu_2, L_2}$ , with  $\mu_1 \geq 0$  and  $\mu_1 + \mu_2 > 0$ . Consider  $N+1 \geq 2$  iterations of DCA starting from  $x^0$ .

If  $\mu_2 < 0 < \mu_1 < L_2$ , thus  $F$  is nonconvex-nonconcave, then:

$$\frac{\mu_1}{2} \min_{0 \leq k \leq N} \{\|x^k - x^{k+1}\|^2\} \leq \frac{F(x^0) - F(x^{N+1})}{1 + \frac{\mu_2}{\mu_1} + \min\{P_N(\frac{L_2}{\mu_1}, \frac{\mu_2}{\mu_1}), E_N(\frac{\mu_2}{\mu_1})\}}, \quad (9)$$

where  $P_N(\eta, \rho) := \frac{(1+\eta)(1+\rho)}{\eta+\rho}(N + \frac{(1-\eta)(1-\rho)}{\eta-\rho} \sum_{k=0}^N [E_k(\eta) - E_k(\rho)]_+)$ .

If  $L_2 \leq \mu_1$  and  $\mu_1 > 0$ , thus  $F$  is (strongly) convex, then:

$$\frac{\mu_1}{2} \min_{0 \leq k \leq N} \{\|x^k - x^{k+1}\|^2\} \leq \frac{F(x^0) - F(x^{N+1})}{1 + \frac{\mu_2}{\mu_1} + \min\{E_N(\frac{L_2}{\mu_1}), E_N(\frac{\mu_2}{\mu_1})\}}. \quad (10)$$

Finally, if  $0 \leq \mu_1 < L_1 \leq \mu_2 < L_2$ , thus  $F$  is (strongly) concave, then:

$$\frac{\mu_2}{2} \min_{0 \leq k \leq N} \{\|x^k - x^{k+1}\|^2\} \leq \frac{F(x^0) - F(x^{N+1})}{1 + \frac{\mu_1}{\mu_2} + \min\{E_N(\frac{L_1}{\mu_2}), E_N(\frac{\mu_1}{\mu_2})\}}. \quad (11)$$

For  $N = 1$ , [Conjecture 1](#) includes the rates corresponding to regimes  $p_3$  and  $p_5$  in (9) when  $F$  is nonconvex, of regimes  $p_4$  and  $p_5$  in (10) when  $F$  is (strongly) convex and of regime  $p_6$  in (11) when  $F$  is concave. The rate in (10) includes the ones proved in [Theorem 2](#) (within the domain of regime  $p_4$ ) and in [Theorem 3](#). The expressions of the rates (9) and (10) are independent of  $L_1$ , hence they also hold when  $f_1$  is *nonsmooth*, i.e.,  $L_1 = \infty$ .

[Conjecture 1](#) is based on the exact rates developed for gradient descent (GD) derived in [18] and is confirmed by PEP numerical experiments. GD is a particular case of the proximal gradient descent (PGD), which is iteration equivalent with DCA when  $f_2$  is smooth.

## 4 Deriving rates for proximal gradient descent (PGD)

Assume  $F = \varphi + h$ , with  $\varphi \in \mathcal{F}_{\mu_\varphi, L_\varphi}$  smooth,  $L_\varphi \in (0, \infty)$ ,  $\mu_\varphi \in (-\infty, L_\varphi)$ , and  $h \in \mathcal{F}_{\mu_h, L_h}$  proper, closed, l.s.c. and convex, such that  $\mu_h \geq 0$  and  $L_h \in (0, \infty]$ .

The PGD iteration with stepsize  $\gamma > 0$ , starting from  $x^k$ , reads  $x^{k+1} = \operatorname{argmin}_{w \in \mathbb{R}^d} \{h(w) + \frac{1}{2\gamma}\|w - x^k + \gamma \nabla \varphi(x^k)\|^2\}$ . This iteration is exactly the DCA one applied to  $f_1 = h + \frac{\|\cdot\|^2}{2\gamma}$  and  $f_2 = \frac{\|\cdot\|^2}{2\gamma} - \varphi$ , as shown in [13, Section 3.3.4] and [19, Proposition 3]. Consequently, all the rates developed for DCA readily

translate to rates for PGD using the mapping  $\mu_1 = \gamma^{-1} + \mu_h$ ,  $L_1 = \gamma^{-1} + L_h$ ,  $\mu_2 = \gamma^{-1} - L_\varphi$ ,  $L_2 = \gamma^{-1} - \mu_\varphi$ . The condition  $\mu_1 + \mu_2 > 0$  translates to the standard upper limit of  $\gamma < \frac{2}{L_\varphi - \mu_h}$  and  $\mu_2 < 0$  corresponds to stepsizes  $\gamma > \frac{1}{L_\varphi}$ .

When  $\varphi$  is convex, [24] shows various tight rates employing different performance metrics, including gradient residual for stepsize  $\gamma = \frac{1}{L_\varphi}$ , but not the gradient mapping. In the gradient residual, for  $\varphi$  nonconvex, with  $\mu_\varphi = -L_\varphi$  and  $\mu_h = 0$ ,  $L_h = \infty$ , the tight rate for stepsizes shorter than  $\frac{1}{L_\varphi}$  is given in [1, Proposition 8.11]. Sublinear rates for the same criterion are given in [19].

The gradient descent (GD) is obtained by setting  $h = 0$ , hence  $\mu_h = L_h = 0$ . Convergence rates on the best gradient norm are proved in [18] for all curvatures of  $\varphi$ . These rates are a particular case of [Conjecture 1](#) and served as a base to conjecture the tight expressions, since the projection / proximal step is not modifying the behavior of the worst-case function.

To get exact rates from DCA, one just has to plug in the expressions of the curvatures  $\mu_i$ ,  $L_i$ ,  $i = \{1, 2\}$ , written with respect to curvatures of  $\varphi$  and  $h$ , in [Theorem 1](#) and [Conjecture 1](#). The convexity of  $h$  implies  $\mu_h \geq 0$ , thus  $\mu_1 > 0$ . Then regime  $p_1$  corresponds to taking stepsizes  $\gamma \leq \frac{1}{L_\varphi}$  for a splitting with  $\varphi$  and  $F$  nonconvex. Within the regimes  $p_3$  and  $p_4$ , the stepsize  $\gamma$  lies in the interval  $[\frac{1}{L_\varphi}, \bar{\gamma}^1(L_\varphi, \mu_\varphi, \mu_h)]$ , where  $\bar{\gamma}^1(L_\varphi, \mu_\varphi, \mu_h)$  is a threshold with an analytical expression obtained through the condition  $\mu_2 = \frac{-L_2\mu_1}{L_2 + \mu_1}$ . Regime  $p_3$  holds when  $\varphi$  and  $F$  are nonconvex. When  $\mu_h = 0$  and  $\mu_\varphi = -L_\varphi$ , then  $\bar{\gamma}^1 = \frac{\sqrt{3}}{L_\varphi}$ , as also obtained in [2] for the GD method. Regime  $p_4$  holds when  $F$  is strongly convex and  $\mu_\varphi + \mu_h \geq 0$ ; in particular, this includes the case of  $\varphi$  being strongly convex. For stepsizes  $\gamma \in (\bar{\gamma}^1(L_\varphi, \mu_\varphi, \mu_h), \frac{2}{L_\varphi + \mu_\varphi})$ , regime  $p_5$  is active if  $F$  is nonconcave. Regimes  $p_2$  and  $p_6$ , respectively, correspond to using stepsizes  $\gamma \leq \frac{1}{L_\varphi}$  for  $\varphi$  concave and  $F$  nonconcave or concave, respectively.

[Lemma 8](#) provides an example of convergence rate translation between PGD and DCA, corresponding to regime  $p_3$  from [Lemma 4](#), assuming the standard setup of PGD with  $\mu_h = 0$  and  $L_h = \infty$ .

**Lemma 8.** *Let  $\varphi \in \mathcal{F}_{\mu_\varphi, L_\varphi}$ ,  $h \in \mathcal{F}_{0, \infty}$  and  $\gamma \in [\frac{1}{L_\varphi}, \frac{2}{L_\varphi})$ . Then after two iterations of PGD with stepsize  $\gamma$  connecting  $x^k$ ,  $x^{k+1}$  and  $x^{k+2}$  it holds*

$$\Delta F(x^k) \geq \frac{[(2-L_\varphi)(2-\mu_\varphi)-1]}{\gamma(2-L_\varphi-\mu_\varphi)} \frac{1}{2} \|\Delta x^k\|^2 + \frac{1}{\gamma(2-L_\varphi-\mu_\varphi)} \frac{1}{2} \|\Delta x^{k+1}\|^2.$$

[Lemma 8](#) is the extension to PGD of the same result obtained for GD in [19, Lemma 4.3]. Therein, it is involved in deriving a stepsize schedule gradually increasing from  $\bar{\gamma}^1(L_\varphi, \mu_\varphi, 0)$  towards  $\frac{2}{L_\varphi + \mu_\varphi}$  (if  $\varphi$  is convex), which gives a better worst-case guarantee than the best constant stepsize. Based on [Lemma 8](#), the same schedule should give a similar performance for PGD. Further, we show how it may improve the DCA convergence.

### Curvature shifting and schedules for DCA

**Proposition 2.** *One PGD iteration with stepsize  $\gamma$  on the splitting  $F = \varphi + h$  produces the same iterate as one iteration with stepsize  $\tilde{\gamma}$  on  $F = \tilde{\varphi} + \tilde{h}$ , with  $\tilde{\varphi} = \varphi + \lambda \frac{\|\cdot\|^2}{2}$ ,  $\tilde{h} = h - \lambda \frac{\|\cdot\|^2}{2}$ ,  $\tilde{\gamma}^{-1} = \gamma^{-1} - \lambda$ , where  $\lambda < \gamma^{-1}$ .*

This relates to curvature shifting in DCA:  $f_i^\lambda := f_i - \frac{\lambda \|\cdot\|^2}{2}$ ,  $i = \{1, 2\}$ , where  $\lambda$  is a scalar. Based on a single-iteration analysis, [19] studies the best curvature shifting, observing that it usually reduces the convexity of both functions. The procedure can be improved, as (optimized) stepsize schedules  $\{\gamma^k(L_\varphi, \mu_\varphi)\}$  for PGD can be translated to DCA. Consider the standard setting of PGD with  $L_\varphi > 0$ ,  $\mu_h = 0$ ,  $L_h = \infty$ , assume  $\mu_1 > 0$  and choose  $\gamma = \mu_1^{-1}$ . Using Proposition 2, a curvature splitting schedule for DCA is defined as  $\{\lambda^k\} = \mu_1 - \{\gamma^k(\mu_1 - L_2, \mu_1 - \mu_2)\}$ .

**Acknowledgements** This research was funded within the framework of the Global PhD Partnership KU Leuven - UCLouvain.

## 5 Conclusion

We studied rates in the gradient mapping for DCA and PGD. Examining the behavior of DCA after two iterations, we identified six regimes partitioning the parameter space, and proved the corresponding six tight sublinear convergence rates. Three of those sublinear rates hold for any number of iterations and, for the other regimes, we conjectured their exact convergence rates. We also show how PGD rates follow directly from DCA ones. This connection suggests the potential for an optimized splitting of the objective function used in DCA, possibly varying at each iteration, inspired by stepsize schedules used for (proximal) gradient descent. We leave the practical testing of those as future work.

## References

1. H. Abbaszadehpeivasti. *Performance Analysis of Optimization Methods for Machine Learning*. CentER dissertation. CentER, Tilburg University, 2024.
2. Hadi Abbaszadehpeivasti, Etienne de Klerk, and Moslem Zamani. The exact worst-case convergence rate of the gradient method with fixed step lengths for  $l$ -smooth functions. *Optimization Letters*, November 2021.
3. Hadi Abbaszadehpeivasti, Etienne de Klerk, and Moslem Zamani. On the rate of convergence of the difference-of-convex algorithm (DCA). *Journal of Optimization Theory and Applications*, Mar 2023.
4. Heinz H. Bauschke, Walaa M. Moursi, and Xianfu Wang. Generalized monotone operators and their averaged resolvents. *Mathematical Programming*, 189(1):55–74, Sep 2021.
5. Damek Davis and Dmitriy Drusvyatskiy. Stochastic model-based minimization of weakly convex functions. *SIAM Journal on Optimization*, 29(1):207–239, 2019.
6. Tao Pham Dinh and Hoai An Le Thi. Convex analysis approach to d.c. programming: Theory, algorithm and applications. In *Acta Mathematica Vietnamica*, volume 22, 1997.
7. Yoel Drori and Marc Teboulle. Performance of first-order methods for smooth convex minimization: a novel approach. *Mathematical Programming*, 145(1-2):451–482, June 2014.
8. Oisín Faust, Hamza Fawzi, and James Saunderson. A Bregman divergence view on the Difference-of-Convex Algorithm. In *Proceedings of The 26th International Conference on Artificial Intelligence and Statistics*, volume 206 of *Proceedings of Machine Learning Research*, pages 3427–3439. PMLR, 25–27 Apr 2023.

9. Baptiste Goujaud, Céline Mouceur, François Glineur, Julien M. Hendrickx, Adrien B. Taylor, and Aymeric Dieuleveut. PEPit: computer-assisted worst-case analyses of first-order optimization methods in python. *Mathematical Programming Computation*, 16(3):337–367, Sep 2024.
10. R. Horst and N. V. Thoai. DC Programming: Overview. *Journal of Optimization Theory and Applications*, 103(1):1–43, October 1999.
11. Gert Lanckriet and Bharath K. Sriperumbudur. On the convergence of the concave-convex procedure. *Advances in Neural Information Processing Systems*, 22, 2009.
12. Hoai An Le Thi, Van Ngai Huynh, and Tao Pham Dinh. Convergence analysis of difference-of-convex algorithm with subanalytic data. *Journal of Optimization Theory and Applications*, 179(1):103–126, Oct 2018.
13. Hoai An Le Thi and Tao Pham Dinh. DC programming and DCA: thirty years of developments. *Mathematical Programming*, 169(1):5–68, May 2018.
14. Thomas Lipp and Stephen Boyd. Variations and extension of the convex–concave procedure. *Optimization and Engineering*, 17(2):263–287, Jun 2016.
15. Yurii Nesterov. Gradient methods for minimizing composite functions. *Mathematical Programming*, 140(1):125–161, Aug 2013.
16. Yurii Nesterov. *Lectures on Convex Optimization*. Springer Publishing Company, Incorporated, 2nd edition, 2018.
17. R. Tyrrell Rockafellar and Roger J.-B. Wets. *Variational Analysis*. Springer Verlag, Heidelberg, Berlin, New York, 1998.
18. Teodor Rotaru, François Glineur, and Panagiotis Patrinos. Exact worst-case convergence rates of gradient descent: a complete analysis for all constant stepsizes over nonconvex and convex functions. *arXiv preprint arXiv:2406.17506*, 2024.
19. Teodor Rotaru, Panagiotis Patrinos, and François Glineur. Tight analysis of difference-of-convex algorithm (DCA) improves convergence rates for proximal gradient descent. In *AISTats 2025 Proceedings*, 2025.
20. Pham Dinh Tao and Le Thi Hoai An. A D.C. optimization algorithm for solving the trust-region subproblem. *SIAM Journal on Optimization*, 8(2):476–505, 1998.
21. Adrien Taylor, Julien Teboulle Hendrickx, and François Glineur. Smooth strongly convex interpolation and exact worst-case performance of first-order methods. *Mathematical Programming*, 161(1-2):307–345, January 2017.
22. Adrien B. Taylor, Julien M. Hendrickx, and François Glineur. Exact worst-case performance of first-order methods for composite convex optimization. *SIAM Journal on Optimization*, 27(3):1283–1313, 2017.
23. Adrien B. Taylor, Julien M. Hendrickx, and François Glineur. Performance estimation toolbox (PESTO): Automated worst-case analysis of first-order optimization methods. In *2017 IEEE 56th Annual Conference on Decision and Control (CDC)*, pages 1278–1283, December 2017.
24. Adrien B. Taylor, Julien M. Hendrickx, and François Glineur. Exact worst-case convergence rates of the proximal gradient method for composite convex minimization. *Journal of Optimization Theory and Applications*, 178(2):455–476, 2018.
25. J. F. Toland. A duality principle for non-convex optimisation and the calculus of variations. *Archive for Rational Mechanics and Analysis*, 71(1):41–61, May 1979.
26. Chaorui Yao and Xin Jiang. A globally convergent difference-of-convex algorithmic framework and application to log-determinant optimization problems, 2023.
27. Alan L Yuille and Anand Rangarajan. The concave-convex procedure (CCCP). In T. Dietterich, S. Becker, and Z. Ghahramani, editors, *Advances in Neural Information Processing Systems*, volume 14. MIT Press, 2001.
28. Alp Yurtsever and Suvrit Sra. CCCP is Frank-Wolfe in disguise. In *Advances in Neural Information Processing Systems*, volume 35, pages 35352–35364, 2022.