

Molecular Physiology of Water Balance

TO THE EDITOR: In their review article on water balance, Knepper et al. (April 2 issue)¹ discuss water channels (aquaporins) in renal tubular cells. They omit mention of the critical role played by aquaporin-1 in microvascular endothelia.

In the renal microvasculature, endothelial aquaporin-1 mediates the osmotic water efflux across descending vasa recta and is required for regulation of medullary blood flow, maintenance of the medullary interstitial gradient, and urinary concentrating ability.²

Aquaporin-1 is also highly expressed in the endothelium lining peritoneal capillaries, where it facilitates osmotically driven water transport during peritoneal dialysis. Peritoneal dialysis is a technique of renal-replacement therapy that is increasingly used worldwide to restore homeostasis, including water balance, in patients with end-stage renal disease.³ Studies in experimental models have shown that aquaporin-1 mediates up to 50% of ultrafiltration during peritoneal dialysis with crystalloid (glucose-based) solutions.⁴ In these models, pharmacologic regulation of aquaporin-1, either through induction with glucocorticoids⁴ or gating with furosemide-based compounds,⁵ enhances aquaporin-1-mediated water transport across the peritoneal membrane. These data show the important role of endothelial water channels in water homeostasis.

Johann Morelle, M.D.

Catholic University of Louvain Medical School
Brussels, Belgium

Eric Goffin, M.D.

Saint-Luc Academic Hospital
Brussels, Belgium

Olivier Devuyst, M.D., Ph.D.

University of Zurich
Zurich, Switzerland
olivier.devuyst@uzh.ch

No potential conflict of interest relevant to this letter was reported.

1. Knepper MA, Kwon TH, Nielsen S. Molecular physiology of water balance. *N Engl J Med* 2015;372:1349-58.
2. Pallone TL, Edwards A, Ma T, Silldorff EP, Verkman AS. Requirement of aquaporin-1 for NaCl-driven water transport across descending vasa recta. *J Clin Invest* 2000;105:215-22.

3. Pastan S, Bailey J. Dialysis therapy. *N Engl J Med* 1998;338:1428-37.

4. Devuyst O, Rippe B. Water transport across the peritoneal membrane. *Kidney Int* 2014;85:750-8.

5. Yool AJ, Morelle J, Cnops Y, et al. AqF026 is a pharmacologic agonist of the water channel aquaporin-1. *J Am Soc Nephrol* 2013;24:1045-52.

DOI: 10.1056/NEJMc1505505

THE AUTHORS REPLY: Morelle and colleagues point out several key roles of aquaporin-1 in microvascular endothelia that are relevant to our article. We agree with their points and add the following ones. Aquaporin-1 is strongly expressed in the endothelium of the descending vasa recta of the renal medulla.¹ The resulting high water permeability of the descending vasa recta² contributes to the countercurrent exchange process that permits perfusion of the renal medulla without dissipation of medullary solute gradients.³ Therefore, aquaporin-1 expression in the vasa recta is crucial to the ability of the kidneys to conserve water.

Mark A. Knepper, M.D., Ph.D.

National Heart, Lung, and Blood Institute
Bethesda, MD
knepp@helix.nih.gov

Tae-Hwan Kwon, M.D., Ph.D.

Kyungpook National University
Taegu, South Korea

Soren Nielsen, M.D., Ph.D.

Aalborg University
Aalborg, Denmark

Since publication of their article, the authors report no further potential conflict of interest.

1. Nielsen S, Pallone T, Smith BL, Christensen EI, Agre P, Maunsbach AB. Aquaporin-1 water channels in short and long loop descending thin limbs and in descending vasa recta in rat kidney. *Am J Physiol* 1995;268:F1023-F1037.

2. Pallone TL, Kishore BK, Nielsen S, Agre P, Knepper MA. Evidence that aquaporin-1 mediates NaCl-induced water flux across descending vasa recta. *Am J Physiol* 1997;272:F587-96.

3. Zimmerhackl BL, Robertson CR, Jamison RL. The medullary microcirculation. *Kidney Int* 1987;31:641-7.

DOI: 10.1056/NEJMc1505505