

Journal Pre-proof

Higher buccal mtDNA content is associated with residential surrounding green in a panel study of primary school children

Pauline Hautekiet, Nelly D. Saenen, Raf Aerts, Dries S. Martens, Harry A. Roels, Esmée M. Bijmens, Tim S. Nawrot

PII: S0013-9351(22)00878-7

DOI: <https://doi.org/10.1016/j.envres.2022.113551>

Reference: YENRS 113551

To appear in: *Environmental Research*

Received Date: 24 February 2022

Revised Date: 18 May 2022

Accepted Date: 19 May 2022



Please cite this article as: Hautekiet, P., Saenen, N.D., Aerts, R., Martens, D.S., Roels, H.A., Bijmens, Esmée.M., Nawrot, T.S., Higher buccal mtDNA content is associated with residential surrounding green in a panel study of primary school children, *Environmental Research* (2022), doi: <https://doi.org/10.1016/j.envres.2022.113551>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Published by Elsevier Inc.

Higher buccal mtDNA content is associated with residential surrounding green in a panel study of primary school children

Pauline Hautekiet^{a,b*}, Nelly D. Saenen^{a,b*}, Raf Aerts^{a,b,c}, Dries S. Martens^a, Harry A. Roels^{a,d}, Esmée M. Bijmens^{a,e}, Tim S. Nawrot^{a,f}

^a Centre for Environmental Sciences, Hasselt University, Agoralaan gebouw D, BE-3590 Hasselt, Belgium

^b Risk and Health Impact Assessment, Sciensano (Belgian Institute of Health), Juliette Wytsmanstraat 14, BE-1050 Brussels, Belgium

^c Department of Ecology, Evolution and Biodiversity Conservation, University of Leuven (KU Leuven), Kasteelpark Arenberg 31-2435, BE-3001 Leuven, Belgium

^d Louvain Centre for Toxicology and Applied Pharmacology, Université catholique de Louvain, Av. Hippocrate 57, BE-1200 Woluwe-Saint-Lambert, Belgium

^e Department of Human Structure and Repair, Ghent University Hospital, Corneel Heymanslaan 10, BE-9000 Ghent, Belgium

^f Department of Public Health & Primary Care, University of Leuven (KU Leuven), O&N I Herestraat 49 - bus 706, BE-3000 Leuven, Belgium

Corresponding author: Tim Nawrot, Hasselt University, Centre for Environmental Sciences, Agoralaan gebouw D, BE-3590 Hasselt, Belgium. E-mail address: tim.nawrot@uhasselt.be.

*These authors contributed equally to this work

Abstract

Background: Mitochondria are known to respond to environmental stressors but whether green space is associated with mitochondrial abundance is unexplored. Furthermore, as exposures may affect health from early life onwards, we here evaluate if residential green space is associated with mitochondria DNA content (mtDNAC) in children.

Methods: In primary schoolchildren (COGNAC study), between 2012 and 2014, buccal mtDNAC was repeatedly (three times) assessed using qPCR. Surrounding low (<3m), high (≥3m) and total (sum of low and high) green space within different radii (100m – 1000m) from the residence and distance to the nearest large green space (> 0.5ha) were estimated using a remote sensing derived map. Given the repeated measures design, we applied a mixed-effects model with school and subject as random effect while adjusting for *a priori* chosen fixed covariates.

Results: mtDNAC was assessed in 246 children with a total of 436 measurements (mean age 10.3 years). Within a 1000m radius around the residential address, an IQR increment in low (11.0%), high (9.5%), and total (13.9%) green space was associated with a respectively 15.2% (95% CI: 7.2% to 23.7%), 10.8% (95% CI: 4.5% to 17.5%), and 13.4% (95% CI: 7.4% to 19.7%) higher mtDNAC. Conversely, an IQR increment (11.6%) in agricultural area in the same radius was associated with a -3.4% (95% CI: -6.7% to -0.1%) lower mtDNAC. Finally, a doubling in distance to large green space was associated with a -5.2% (95% CI: -7.9 to -2.4%) lower mtDNAC.

Conclusion: To our knowledge, this is the first study evaluating associations between residential surrounding green space and mtDNAC in children. Our results showed that green space was associated with a higher mtDNAC in children, which indicates the importance of the early life environment. To what extent these findings contribute to later life health effects should be further examined.

Keywords

Green space, agriculture, mitochondrial DNA content, mtDNA, children

Source of funding

The COGNAC study was supported by the European Research Council [ERC-2012-StG310898] and the Flemish Scientific Fund [FWO, G073315N]. Esmée Bijmens holds a fellow-ship from the Marguerite-Marie Delacroix foundation. Dries Martens is a postdoctoral fellow of the Research Foundation - Flanders (FWO 12X9620N). The study complied with the Helsinki declaration and was approved by the ethics committees of Hasselt University and the Eastern-Limburg Hospital, Belgium (11/065U).

1. Introduction¹

Mitochondria are essential for the function and homeostasis of the cell due to their role in energy (ATP) production, apoptosis, the control of cytosolic calcium levels, and cell signalling (Shaughnessy et al. 2014). They have their own double-stranded genome, called mitochondrial DNA. Being the powerhouse of the cell, mitochondrial DNA content (mtDNAC) in cells and tissues must be adjusted according to the metabolic need (Filograna et al. 2021). As a proxy for mitochondrial function (Malik and Czajka 2013), mtDNAC is used to assess the role of mitochondria in disease (Longchamps et al. 2020). As such, a higher mtDNAC (measured in peripheral blood, buccal cells and placenta tissue, respectively) has been associated with cancer (Zhou et al. 2014) in adults and heart rate variability (Saenen et al. 2019) and intelligence (Bijnens et al. 2019) in children. On the other hand, a lower mtDNAC (measured in peripheral blood and leucocytes, respectively) has been associated with health outcomes such as Parkinson's disease (Pyle et al. 2016), and schizophrenia (Shivakumar et al. 2020) in adults.

Literature suggests that environmental exposures might affect mtDNAC as mitochondria coordinate cellular adaptation to stressors like oxidative stress and DNA damage (Vyas et al. 2016). In adults, a decreased mtDNAC has been associated with particulate matter air pollution (Wang et al. 2020) and PAH exposure (Pavanello et al. 2013). Although limited, this effect is also observed during childhood. For example, exposure to particulate matter has been associated with a lower buccal mtDNAC (Hautekiet et al. 2020; Saenen et al. 2019) and exposure to lead was associated with a lower leukocyte mtDNAC (Alegría-Torres et al. 2020). What has, to our knowledge, not been evaluated in literature is the effect of green space on mtDNAC. Assessing green space in early life is important as this might affect health outcomes later in life. For example, reduced exposure to residential green space during early life has

¹ **Abbreviations:** Cq, cycle quantification value; mtDNAC, mitochondrial DNA content; IQR, interquartile range; GIS, Geographic Information System

been associated with higher blood pressure during adulthood (Bijnens et al. 2017). Also, children living in the lowest decile of green space showed a 15% to 55% higher risk for mental illnesses later in life compared to those living in the highest decile of green space (Engemann et al. 2019). Hence, in this panel study, we examined whether residential surrounding green space and distance to green space were associated with mtDNAc in children between 9 and 12 years old. We hypothesize that children living in a neighbourhood with more green space or living close to green space have higher mtDNAc.

2. Materials and methods

2.1. Study population

This study was part of the COGNAC (COGNition and Air pollution in Children) study and included repeated measurements of children between 9 and 12 years old (Saenen et al. 2016). The study was executed between January 2012 and February 2014 in three different schools, each in another Belgian municipality (Tienen, Zonhoven, and Hasselt-Kiewit). 770 children were invited, of which 334 (43.4%) agreed to participate. In the present study, a final study population of 246 children (73.7%), with a total of 436 measurements were included after exclusion of samples not meeting the quality control of the molecular measurements.

Parents of the children signed an informed consent and completed a questionnaire to provide information on the demographic and lifestyle characteristics of the child and its family. Socioeconomic status was based on the highest educational level of either parent (up to high school diploma; college or university diploma). Children were considered to be exposed to passive smoking if one or more family members smoked inside the house. Physical activity like for example soccer, basketball, gymnastics, swimming, or horseback riding was calculated in hours per week. The study complied with the Helsinki declaration and was approved by the ethics committees of Hasselt University and the Eastern-Limburg Hospital, Belgium (11/065U).

2.2. Sample collection

Buccal cell swabs (SK2, Isohelix, Kent, UK) were sampled during three different examinations at the school. Per examination, we collected two buccal samples per child. Sampling for each examination was carried out at the same time during the day and on the same day of the week. Before sampling, the children did not eat or drink for at least 30 minutes and rinsed their mouths three times with water. Each swab was rubbed inside the cheek for one minute. Afterwards the samples were kept on ice until storage at -80°C. The mean (SD) time interval between two consecutive examinations was 46 (20) days. Of the 246 children included, 105 (42.7%) provided a buccal swab at one time point, 92 (37.4%) at two time points, and 49 (19.9%) at three time points, resulting in a total number of 436 samples.

2.3. Mitochondrial DNA assay

DNA was extracted from both buccal swabs at once using the QIAgen Micro Kit (Qiagen, N.V. Venlo, the Netherlands). The quantity and purity of the sample were measured with a Nanodrop spectrophotometer (ND-1000; Ismgen Life Science, De Meern, the Netherlands). To ensure a uniform DNA input for each qPCR reaction, samples were diluted and checked using the Quant-iT™ PicoGreen® dsDNA Assay Kit (Life Technologies, Europe). Extracted DNA was stored at -80°C until analysis.

Average mtDNA_C was determined by averaging the ratios of two mitochondrial gene copy numbers (MTF3212/R3319 and MT-ND1) (M) to one reference gene (36B4) (S). Samples were measured in triplicate using a previously described quantitative real-time PCR (qPCR) assay with minor modifications (Janssen et al. 2012). Briefly, the reaction mixtures contained Qiagen 1x QuantiTect SYBR Green Mastermix, forward (300 nM) and reverse (300 nM) primer, and 12.5 ng DNA. The forward and reverse primers for the mitochondrial genes were respectively 5'-CACCCAAGAACAGGGTTTGT-3' and 5'-TGGCCATGGGTATGTTGTTAA-3' for MTF3212/3319, and 5'-ATGGCCAACC TCCTACTCCT-3' and 5'-

CTACAACGTTGGGGCCTTT-3' for MT-ND1. For the reference gene, the forward and reverse primers were 5'-GGAATGTGGGCTTTGTGTTC-3' and 5'-CCCAATTGTCCCCTTACCTT-3'.

All PCR-reactions were performed on a 7900HT Fast Real-Time PCR System (Applied Biosystems, Foster City, CA, USA) using the following thermal cycling profile: 'M' – 1 cycle of 10 min at 95°C, followed by 30 cycles of 15 s at 95°C and 1 min 10 s at 58°C; 'S' – 1 cycle of 10 min at 95°C, followed by 35 cycles of 15 s at 95°C and 1 min 10 s at 58°C. Six inter-run calibrators (IRCs) were used to account for inter-run variability. Also, non-template controls were used in each run. Raw data were processed and normalized to the reference gene using qBase-plus software (Biogazelle, Zwijnaarde, Belgium), taking into account run-to-run differences. Quality control measures were taken into account for data pre-processing after qPCR: (1) absolute difference < 0.3 of triplicate Ct (cycle threshold) values and (2) maximum 1 absolute Ct value deviation of the average Ct of each plate. The intra-plate ICC (95% CI) of M/S ratios (MTF3212/3319 and MT-ND1), mtDNAC runs (MTF3212/3319 and MT-ND1) and single-copy gene runs were respectively 0.953 (0.945, 0.960), 0.962 (0.955, 0.968), 0.975 (0.970, 0.978), 0.957 (0.950, 0.963), and 0.975 (0.971, 0.979) (Telomere research network 2020).

2.4. Residential green space exposure

Residential addresses of the children were geocoded. We characterized two indicators of exposure to green space: 1) residential surrounding green space (%) and 2) residential distance to green space (m). To characterize green space we used the Green Map of Flanders 2012, generated by the Agency for Geographic Information Flanders (AGIV) (Geopunt Vlaanderen 2012). This 1 m² resolution raster geodataset derived from orthorectified aerial imagery classifies every pixel in the entire region as: not green, agricultural area, low green (vegetation height < 3 m), and high green (vegetation height ≥ 3 m) areas. The percentage of agricultural area and low, high, and total (sum of low and high green space) green space was

calculated for different radii (100 m, 300 m, 500 m, and 1000 m) around each child's residence. To evaluate scale effects and the consistency and robustness of our results, we chose different radii. The 100 m, 300 m, and 500 m radius were used to indicate the immediate, intermediate, and neighbourhood surrounding green space, respectively, whereas 1000 m indicates the common walking distance to places nearby (Dadvand et al. 2017; Stockton et al. 2016). Finally, to characterize residential distance to green space, we calculated the distance (m) from the child's residence to the nearest large green space (≥ 0.5 ha) (Ludlow et al. 2003). For children with more than one residential address at the moment of the study ($n = 38$; 15.5%), we calculated a weighted average using the proportion of time spent at each location. ArcGIS 10 (ESRI Inc, Redmonds, CA, US) software was used for all GIS analyses.

2.5. Statistical analysis

Statistical analyses were performed using SAS software (version 9.4; SAS Institute Inc., Cary, NC, USA). To reduce skewness and to approximate a normal distribution, we $\log_{10}$ -transformed the mtDNAc data, the distance to major roads, and the distance to green space. As our study design included repeated measurements of mtDNAc, we used a mixed-effects model (unstructured covariance matrix) with school (representing the sampling framework) and subject as random effects to assess the associations between surrounding green space exposure or distance to green space and mtDNAc as the outcome variable. *A priori* selected covariates included age (continuous), sex (boy; girl), BMI (continuous), passive smoking (yes; no), season of examination (winter; summer; autumn; spring), highest educational level of either parent (up to high school; college or university), date of examination (continuous) and distance to major road (continuous). Distance to major road represents the distance from the child's residence to the nearest highway or other national road and was calculated with ArcGIS 10 (ESRI Inc, Redmonds, CA, US) software. We chose these covariates as socio-economic status has been associated with greens space (P. Dadvand et al. 2012) whereas the others

have been associated with mtDNAC (Hou et al. 2019; Janssen et al. 2012; Knez et al. 2016; Skuratovskaia et al. 2019; Wu et al. 2019).

All covariates were measured at baseline except season, which was accounted for at each examination. Results of the analyses were presented as the percentage difference in relative average mtDNAC associated with an interquartile range (IQR) increment in residential green space or with a doubling in distance to green space. Because of the skewedness of the data for the 100 m radius of agriculture, we expressed the effect size for a dichotomized variable comparing the presence to the absence of agriculture within this radius.

Bivariate associations between the fixed covariates and the outcome were assessed using the unadjusted model, but taking into account the random effect of the subjects and school. Unadjusted Spearman's correlations were used to assess the associations between the different types of green space within each of the radii.

2.6. Sensitivity analysis

In a sensitivity analysis, we additionally adjusted the model separately and combined for residential PM_{2.5} exposure in the year before examination (long-term exposure), the average ultrafine particle (UFP) < 300 nm exposure at school measured between 9 and 12 O'clock on the day of examination (short-term exposure), and physical activity, based on out-of-school sport activities (hours/week).

Daily residential exposure ($\mu\text{g}/\text{m}^3$) to particulate matter $\leq 2.5\mu\text{m}$ (PM_{2.5}) at the child's residence was modelled using a spatiotemporal interpolation model, designed for Belgium (Janssen et al. 2008; Lefebvre et al. 2013; Maiheu et al. 2013), as described before (Hautekiet et al. 2020). We used the daily modelled exposure values obtained by this model to assign the residential exposure during the year before each examination. For children with more than one residential address at the moment of the study, we calculated a weighted average using the proportion of time spent at each location. Exposure to ultrafine particles (UFP) < 300 nm at school was

measured continuously between 9 and 12 O'clock with a nanotracer (Aerasense, Best, The Netherlands) on the day of the examination.

Finally, to evaluate the effect of the repeated measures design, we also made the association between green space and mtDNAC when only the first measurement of each child was included. Mixed models as for the main analysis were used but without the random statement.

3. Results

3.1. Population characteristics

The characteristics of the 251 children included in this study are presented in Table 1. Children comprising of 46.8% girls had an average age and BMI (SD) of 10.3 (1.3) years and 17.4 (2.8) kg/m², respectively. The most common highest category of education of either parent was college or university (69.5%). 15.9% of the children were exposed to passive smoking at home. The average distance (SD) to major road was 576.2 (559.0) m. The children's geometric mean of relative mtDNAC (95% CI) was 0.98 (0.94 – 1.01). The mtDNAC results of the three examinations were positively correlated with each other: the Pearson correlation coefficients between examinations 1 and 2, 1 and 3, and 2 and 3 were respectively 0.44 ($p < 0.0001$), 0.30 ($p = 0.018$), and 0.27 ($p = 0.023$) (Suppl. figure 1). The average number of days between examination 1 and 2, 1 and 3 and 2 and 3 was respectively 55, 95 and 27 days.

Bivariate analysis showed that girls tended to have a -6.8% (95% CI: -14.2% to 1.3%, $p = 0.095$) lower mtDNAC compared to boys. A one-year increase in age was associated with a 3.9% (95% CI: 0.5 to 7.3, $p = 0.023$) higher mtDNAC. No significant association was found between mtDNAC and BMI ($p = 0.57$), exposure to passive smoking ($p = 0.16$), education ($p = 0.54$), or physical activity ($p = 0.80$).

3.2. Green space exposure characteristics

Mean exposure (IQR) to low green, high green, total green and agriculture in a 1000 m radius was respectively 28.7 (11.0)%, 20.9 (9.5)%, 49.6 (13.9)%, and 17.0 (13.9)% (Table 2). The median distance (IQR) to large green space was 123.7 (144.6) m (Table 2). The different types of green space and the different radii were highly correlated (Suppl. table 1). Agricultural area was negatively correlated with low and total green, with the strongest association between low green and agriculture in a 1000 m radius (Spearman's correlation = -0.54, $p < 0.001$).

3.3. Residential surrounding green space and distance to green space in association with mtDNAC

After adjustment for the *a priori* selected covariates, residential surrounding green space was positively associated with mtDNAC in all buffers except for the 100 m radius (Fig. 1 & Suppl. table 2). The strongest associations were found for the 500 m and 1000 m radii. An IQR increment in low (11.0%), high (9.5%), and total (13.9%) green space in the 1000 m radius was associated with respectively a 15.2% (95% CI: 7.2% to 23.7%, $p = 0.00014$), 10.8% (95% CI: 4.5% to 17.5%, $p = 0.00070$), and 13.4% (95% CI: 7.4% to 19.7%, $p < 0.0001$) higher mtDNAC (Fig. 1). On the contrary, the opposite effect was found for agricultural area where an IQR increment (11.9%) in agricultural area in a 1000 m radius was associated with a -3.4% (95% CI: -6.7% to -0.1%, $p = 0.047$) lower mtDNAC (Fig. 1).

In addition, mtDNAC was inversely correlated with distance to green space (area with more than 0.5 ha of green). After adjustment, a doubling in distance was significantly associated with a -5.2% (95% CI: -7.9 to -2.4%, $p = 0.00040$) lower mtDNAC (Fig. 2).

3.4. Sensitivity analysis

Average (SD) PM_{2.5} exposure during the year before examination was 14.7 (1.1) µg/m³. Average UFP (SD) exposure measured at school on the day of the examination was 23,604 (15,272.2) particles/cm³. Additional adjustment of the main model for PM_{2.5} exposure, UFP exposure and physical activity, separately or combined are presented in table 3 for the 1000 m radius and in and supplementary table 3-5 for the other radii. Results did not show a notable change in the effect estimates of the aforementioned associations. However, after adjusting for physical activity, the results for agriculture showed slightly higher *p*-values. Also, only including the first measurement of each child did not alter the results (Suppl. table 6).

4. Discussion

In 9 to 12 year old children, we investigated whether residential surrounding green space and distance to green space were associated with mitochondrial DNA content. We found that an increase in surrounding green space was associated with a higher mtDNA_c, whereas the opposite was found for agricultural area. Secondly, our results showed a higher mtDNA_c for children living closer to green space.

4.1. Interpretation of the findings

To our knowledge, this is the first study to evaluate the association between green space exposure and mtDNA_c. Although we cannot compare with other studies, our results are in line with several previous observations. More time spent in green spaces has been associated with lower stress, better mental health, and improved social contacts (Beyer et al. 2014; Brown et al. 2018; Dadvand et al. 2019), which in turn, could be associated with higher mtDNA_c (Brown et al. 2018; Orban et al. 2017; Pun et al. 2018).

Furthermore, we found that green space within a 500 m and 1000 m radius had the strongest effect on mtDNAC. This is consistent with previous evidence where larger radii of green space were more strongly associated with health outcomes. For example, the 500 m radius compared with smaller radii was more strongly associated with birth weight and head circumference (Payam Dadvand et al. 2012). Also, greenness within 3 km was associated with mental health whereas no association was found for 1 km (Bos et al. 2016). Finally, built greenness radii between 100 m and 2000 m showed stronger associations for larger radii and intelligence quotient in children (Lee et al. 2021).

However, two studies also found associations with health outcomes for smaller radii. A higher birthweight was found for exposure to green space in a 50 m and 100 m radius (Laurent et al. 2013) and the strongest association between green space and mental and self-perceived health was found for the 100 m and 250 m radius, respectively (Su et al. 2019). This indicates the possibility of different underlying mechanisms. Dadvand and colleagues hypothesized that visual access and heat reduction might be important mechanisms of immediate surrounding green space, whereas larger radii might be associated with other mechanisms like social health (Dadvand et al. 2016) and physical activity (Dadvand et al. 2014). Indeed, research showed that larger radii are more strongly associated with social support (Dadvand et al. 2016) and physical activity (Su et al. 2019), and that they better predict physical activity (Browning and Lee 2017), which has been associated with higher mtDNAC (Steiner et al. 2011). Finally, not only surrounding green space but also access to a large green space is important in the context of physical activity (McCormack et al. 2010). The association of residential green space with mtDNAC is confirmed by our finding that living closer to green space was associated with a higher mtDNAC.

On the other hand, we found an inverse association between surrounding agricultural area and mtDNAC. One possible hypothesis could be that children living in an environment with more agricultural areas are exposed to a lower percentage of surrounding green space and therefore might experience less positive effects of green space. Our findings showed that with

the green map of Flanders, green space was inversely associated with agriculture. Another explanation from literature could be that agricultural activities are associated with emissions of primary particulate matter (PM), nitrogen oxides (NO_x), and ammonia, which in turn give rise to the formation of secondary PM. These agricultural emissions have been associated with adverse health effects (Pozzer et al. 2017), and pesticides have been linked with oxidative stress and DNA damage (Kisby et al. 2009). However, additionally adjusting for long-term PM_{2.5} in the sensitivity analysis did not alter the observed effect. Future studies should confirm the association between agriculture and mtDNAc and further elucidate underlying pathways.

4.2. *Biological processes*

The biological pathways underlying the association between green space and mtDNAc are not yet known. One hypothesis is oxidative stress and inflammation. Higher exposure to green space has been associated with higher levels of physical activity (Steiner et al. 2011), lower levels of obesity (Knobel et al. 2020), and lower levels of psychological stress (Beyer et al. 2014). All three factors have been associated with a decrease in oxidative stress and inflammation (Eick et al. 2018; McMurray et al. 2016; Simioni et al. 2018), which in turn has been associated with higher levels of mtDNAc (Castellani et al. 2020). Furthermore, as mtDNAc is strongly correlated with telomere length (Martens and Nawrot 2016), our results support the research by Miri et al (2020) which showed a positive effect between green space and telomere length in children (Miri et al. 2020).

Even though mtDNAc is often evaluated in adults and elderly, we believe that it is important to also assess these mtDNAc processes in early life because environmental exposures might already affect mtDNAc *in utero* as shown for multiple toxic metals by Vriens et al. (2017) and during childhood as shown for air pollution in our previous study (Hautekiet et al. 2020). The Developmental Origins of Health and Disease (DOHaD) hypothesis states that adverse influences of the early-life environment can result in permanent changes in adulthood (De Boo

and Harding 2006). Accordingly, the positive effect of green space in early life has been shown to be associated with health outcomes later in life such as blood pressure (Bijnens et al. 2017) and mental health (Engemann et al. 2019). Therefore, increasing exposure to beneficial environments, like green space, during early life might promote health in later life.

4.3. *Strengths and limitations*

The main feature of our study is that it is the first to explore the associations of surrounding green space with mtDNAC in children. Other strengths are the use of a repeated measures design and the fact that we were able to estimate surrounding green space using a specific high-resolution map. However, the latter makes it less evident to compare our results with other studies which often use the normalized difference vegetation index (NDVI, which is an indicator of greenness) or the Corine land cover (which is a coarse land cover map in which smaller green spaces are dissolved in mixed pixels). We chose to use the green map of Flanders in this study as it captures green space on a high-resolution (1m x 1m), whereas CORINE land-cover data does only include green areas that are at least 25 ha and NDVI often fails to distinguish between natural and agricultural areas.

A limitation of our study is the use of buccal swabs. Because children are the subjects of our study, we used buccal swabs to sample DNA as this is a common, non-invasive method compared to the drawing of blood. However, there are three disadvantages to the use of buccal swabs. Firstly, the quality of the obtained DNA is lower (Hansen et al. 2007), which resulted in a loss of data. Secondly, we were not able to compare the mtDNAC of buccal swabs with other somatic cells. Therefore, our results should be interpreted with caution. Finally, the oral cell composition might be influenced by poor oral hygiene or infection (Shalev 2012). However, to avoid this, we used a standardized protocol in which the children rinsed their mouth three times with water and were not allowed to eat or drink 30 minutes before sampling. Lastly, the study design and statistical approach including the repeated measurements caused

a lower risk of reverse causality. However, as for multiple children only one measurement was available, the power of the study design declines so caution is advised when interpreting the causality of these results.

5. Conclusion

In the present study, we found a positive association of buccal mtDNAc with residential surrounding green space and an inverse association with agricultural area. We also showed that living closer to large green space of > 0.5 ha was positively associated with mtDNAc. To what extent higher mtDNAc due to green space contributes to human health effects should be further elucidated.

Author contributions

Pauline Hautekiet: Methodology, Formal analysis, Writing – Original draft, Visualization;
Nelly D. Saenen: Conceptualization, Investigation, Methodology, Writing – Review & Editing,
Data curation, Project administration; **Raf Aerts:** Writing – Review & Editing; **Dries S.**
Martens: Methodology, Writing – Review & Editing; **Esmée M. Bijmens:** Methodology, Writing
– Review & Editing; **Harry A. Roels:** Writing – Review & Editing; **Tim S. Nawrot:**
Conceptualization, Methodology, Resources, Writing – Review & Editing, Supervision,
Validation, Funding acquisition, Project administration.

Competing interests

All authors declare that they have no competing interests.

Funding

The COGNAC study was supported by the European Research Council [ERC-2012-StG310898]; and the Flemish Scientific Fund [FWO, G073315N]. Esmée Bijmens holds a fellow-ship from the Marguerite-Marie Delacroix foundation. Dries Martens is a postdoctoral fellow of the Research Foundation - Flanders (FWO 12X9620N).

Acknowledgments

We are grateful to the children, their parents, and the school directions for participating in this study.

References

- Alegría-Torres J.A., Pérez-Rodríguez R.Y., García-Torres L., Costilla-Salazar R., Rocha-Amador D. Exposure to arsenic and lead in children from salamanca méxico, effects on telomeric lengthening and mitochondrial DNA. *Environmental Science and Pollution Research* 2020;27(6):6420-6428 <https://doi.org/10.1007/s11356-019-07108-4>
- Beyer K.M., Kaltenbach A., Szabo A., Bogar S., Nieto F.J., Malecki K.M. Exposure to neighborhood green space and mental health: Evidence from the survey of the health of wisconsin. *International Journal of Environmental Research and Public Health* 2014;11(3):3453-3472 <http://doi.org/10.3390/ijerph110303453>
- Bijnens E.M., Nawrot T.S., Loos R.J.F., Gielen M., Vlietinck R., Derom C., et al. Blood pressure in young adulthood and residential greenness in the early-life environment of twins. *Environmental Health* 2017;16(1):53 <https://doi.org/10.1186/s12940-017-0266-9>
- Bijnens E.M., Derom C., Weyers S., Janssen B.G., Thiery E., Nawrot T.S. Placental mitochondrial DNA content is associated with childhood intelligence. *Journal of Translational Medicine* 2019;17(1):361 <https://doi.org/10.1186/s12967-019-2105-y>
- Bos E.H., van der Meulen L., Wichers M., Jeronimus B.F. A primrose path? Moderating effects of age and gender in the association between green space and mental health. *International Journal of Environmental Research and Public Health* 2016;13(5) <https://doi.org/10.3390/ijerph13050492>
- Brown S.C., Perrino T., Lombard J., Wang K., Toro M., Rundek T., et al. Health disparities in the relationship of neighborhood greenness to mental health outcomes in 249,405 u.S. Medicare beneficiaries. *International Journal of Environmental Research and Public Health* 2018;15(3):430 <http://doi.org/10.3390/ijerph15030430>
- Browning M., Lee K. Within what distance does "greenness" best predict physical health? A systematic review of articles with gis buffer analyses across the lifespan. *International Journal of Environmental Research and Public Health* 2017;14(7):675 <http://doi.org/10.3390/ijerph14070675>
- Castellani C.A., Longchamps R.J., Sun J., Guallar E., Arking D.E. Thinking outside the nucleus: Mitochondrial DNA copy number in health and disease. *Mitochondrion* 2020;53:214-223 <https://doi.org/10.1016/j.mito.2020.06.004>
- Dadvand P., de Nazelle A., Figueras F., Basagana X., Su J., Amoly E., et al. Green space, health inequality and pregnancy. *Environment International* 2012;40:110-115 <http://doi.org/10.1016/j.envint.2011.07.004>
- Dadvand P., Sunyer J., Basagaña X., Ballester F., Lertxundi A., Fernandez-Somoano A., et al. Surrounding greenness and pregnancy outcomes in four spanish birth cohorts. *Environmental Health Perspectives* 2012;120(10):1481-1487 <https://doi.org/10.1289/ehp.1205244>

- Dadvand P., Wright J., Martinez D., Basagaña X., McEachan R.R., Cirach M., et al. Inequality, green spaces, and pregnant women: Roles of ethnicity and individual and neighbourhood socioeconomic status. *Environment International* 2014;71:101-108 <https://doi.org/10.1016/j.envint.2014.06.010>
- Dadvand P., Bartoll X., Basagaña X., Dalmau-Bueno A., Martinez D., Ambros A., et al. Green spaces and general health: Roles of mental health status, social support, and physical activity. *Environment International* 2016;91:161-167 <https://doi.org/10.1016/j.envint.2016.02.029>
- Dadvand P., Tischer C., Estarlich M., Llop S., Dalmau-Bueno A., López-Vicente M., et al. Lifelong residential exposure to green space and attention: A population-based prospective study. *Environmental Health Perspectives* 2017;125(9):097016 <https://doi.org/10.1289/EHP694>
- Dadvand P., Hariri S., Abbasi B., Heshmat R., Qorbani M., Motlagh M.E., et al. Use of green spaces, self-satisfaction and social contacts in adolescents: A population-based caspian-v study. *Environmental Research* 2019;168:171-177 <http://doi.org/10.1016/j.envres.2018.09.033>
- De Boo H.A., Harding J.E. The developmental origins of adult disease (barker) hypothesis. *Australian and New Zealand Journal of Obstetrics and Gynaecology* 2006;46(1):4-14 <https://doi.org/10.1111/j.1479-828X.2006.00506.x>
- Eick S.M., Barrett E.S., van 't Erve T.J., Nguyen R.H., Bush N.R., Milne G., et al. Association between prenatal psychological stress and oxidative stress during pregnancy. *Paediatric and Perinatal Epidemiology* 2018;32(4):318-326 <https://doi.org/10.1111/ppe.12465>
- Engemann K., Pedersen C.B., Arge L., Tsirogiannis C., Mortensen P.B., Svenning J.-C. Residential green space in childhood is associated with lower risk of psychiatric disorders from adolescence into adulthood. *Proceedings of the National Academy of Sciences* 2019;116(11):5188-5193 <https://doi.org/10.1073/pnas.1807504116>
- Filograna R., Mennuni M., Alsina D., Larsson N.G. Mitochondrial DNA copy number in human disease: The more the better? *FEBS Letters* 2021;595(8):976-1002 <https://doi.org/10.1002/1873-3468.14021>
- Geopunt Vlaanderen. 2012. Geopunt-kaart. Available: <https://www.geopunt.be/> [accessed 17/10/2019].
- Hansen T.v.O., Simonsen M.K., Nielsen F.C., Hundrup Y.A. Collection of blood, saliva, and buccal cell samples in a pilot study on the danish nurse cohort: Comparison of the response rate and quality of genomic DNA. *Cancer Epidemiology and Prevention Biomarkers* 2007;16(10):2072-2076 <https://doi.org/10.1158/1055-9965.EPI-07-0611>
- Hautekiet P., Nawrot T.S., Janssen B.G., Martens D.S., De Clercq E.M., Dadvand P., et al. Child buccal telomere length and mitochondrial DNA content as biomolecular markers of ageing in association with air pollution. *Environment International* 2020;147:106332 <https://doi.org/10.1016/j.envint.2020.106332>

- 525
 526 Hou J., Yin W., Li P., Hu C., Zhang Y., Wang X., et al. Seasonal modification of the associations of
 527 exposure to polycyclic aromatic hydrocarbons or phthalates of cellular aging. *Ecotoxicology and*
 528 *Environmental Safety* 2019;182:109384 <https://doi.org/10.1016/j.ecoenv.2019.109384>
- 529
 530 Janssen B.G., Munters E., Pieters N., Smeets K., Cox B., Cuypers A., et al. Placental mitochondrial
 531 DNA content and particulate air pollution during in utero life. *Environmental Health Perspectives*
 532 2012;120(9):1346-1352 <https://doi.org/10.1289/ehp.1104458>
- 533
 534 Janssen S., Dumont G., Fierens F., Mensink C. Spatial interpolation of air pollution measurements
 535 using corine land cover data. *Atmospheric Environment* 2008;42(20):4884-4903
 536 <https://doi.org/10.1016/j.atmosenv.2008.02.043>
- 537
 538 Kisby G.E., Muniz J.F., Scherer J., Lasarev M.R., Koshy M., Kow Y.W., et al. Oxidative stress and DNA
 539 damage in agricultural workers. *Journal of Agromedicine* 2009;14(2):206-214
 540 <https://doi.org/10.1080/10599240902824042>
- 541
 542 Knez J., Winckelmans E., Plusquin M., Thijs L., Cauwenberghs N., Gu Y., et al. Correlates of peripheral
 543 blood mitochondrial DNA content in a general population. *American Journal of Epidemiology*
 544 2016;183(2):138-146 <http://doi.org/10.1093/aje/kwv175>
- 545
 546 Knobel P., Maneja R., Bartoll X., Alonso L., Bauwelinck M., Valentin A., et al. Quality of urban green
 547 spaces influences residents' use of these spaces, physical activity, and overweight/obesity.
 548 *Environmental Pollution* 2020;271:116393 <https://doi.org/10.1016/j.envpol.2020.116393>
- 549
 550 Laurent O., Wu J., Li L., Milesi C. Green spaces and pregnancy outcomes in southern california.
 551 *Health & Place* 2013;24:190-195 <https://doi.org/10.1016/j.healthplace.2013.09.016>
- 552
 553 Lee K.S., Kim B.N., Cho J., Jang Y.Y., Choi Y.J., Lee W.S., et al. Associations between surrounding
 554 residential greenness and intelligence quotient in 6-year-old children. *Science of the Total*
 555 *Environment* 2021;759:143561 <https://doi.org/10.1016/j.scitotenv.2020.143561>
- 556
 557 Lefebvre W., Degrawe B., Beckx C., Vanhulsel M., Kochan B., Bellemans T., et al. Presentation and
 558 evaluation of an integrated model chain to respond to traffic-and health-related policy questions.
 559 *Environmental Modelling & Software* 2013;40:160-170
 560 <https://doi.org/10.1016/j.envsoft.2012.09.003>
- 561
 562 Longchamps R.J., Castellani C.A., Yang S.Y., Newcomb C.E., Sumpter J.A., Lane J., et al. Evaluation of
 563 mitochondrial DNA copy number estimation techniques. *PLoS One* 2020;15(1):e0228166
 564 <https://doi.org/10.1371/journal.pone.0228166>
- 565
 566 Ludlow D., Mitchell C., Webster M. 2003. European common indicators (eci). Towards a local
 567 sustainability profile. Milano, Italia:Ambiente Italia Research Institute.
- 568

- Maiheu B., Veldeman B., Viaene P., De Ridder K., Lauwaet D., Smeets N., et al. 2013. Identifying the best available large-scale concentration maps for air quality in Belgium. (). Mechelen, Belgium:Flemish Institute for Technological Research (VITO).
- Malik A.N., Czajka A. Is mitochondrial DNA content a potential biomarker of mitochondrial dysfunction? *Mitochondrion* 2013;13(5):481-492 <https://doi.org/10.1016/j.mito.2012.10.011>
- Martens D.S., Nawrot T.S. Air pollution stress and the aging phenotype: The telomere connection. *Current Environmental Health Reports* 2016;3(3):258-269 <https://doi.org/10.1007/s40572-016-0098-8>
- McCormack G.R., Rock M., Toohey A.M., Hignell D. Characteristics of urban parks associated with park use and physical activity: A review of qualitative research. *Health Place* 2010;16(4):712-726 <http://doi.org/10.1016/j.healthplace.2010.03.003>
- McMurray F., Patten D.A., Harper M.E. Reactive oxygen species and oxidative stress in obesity - recent findings and empirical approaches. *Obesity* 2016;24(11):2301-2310 <https://doi.org/10.1002/oby.21654>
- Miri M., de Prado-Bert P., Alahabadi A., Najafi M.L., Rad A., Moslem A., et al. Association of greenspace exposure with telomere length in preschool children. *Environmental Pollution* 2020;266:115228 <https://doi.org/10.1016/j.envpol.2020.115228>
- Orban E., Sutcliffe R., Dragano N., Jökel K.H., Moebus S. Residential surrounding greenness, self-rated health and interrelations with aspects of neighborhood environment and social relations. *Journal of Urban Health* 2017;94(2):158-169 <https://doi.org/10.1007/s11524-016-0112-3>
- Pavanello S., Dioni L., Hoxha M., Fedeli U., Mielzynska-Svach D., Baccarelli A.A. Mitochondrial DNA copy number and exposure to polycyclic aromatic hydrocarbons. *Cancer Epidemiol Biomarkers Prev* 2013;22(10):1722-1729 <https://doi.org/10.1158/1055-9965.Epi-13-0118>
- Pozzer A., Tsimpidi A.P., Karydis V.A., De Meij A., Lelieveld J. Impact of agricultural emission reductions on fine-particulate matter and public health. *Atmospheric Chemistry and Physics* 2017;17(20):12813-12826 <https://doi.org/10.5194/acp-17-12813-2017>
- Pun V.C., Manjourides J., Suh H.H. Association of neighborhood greenness with self-perceived stress, depression and anxiety symptoms in older u.S adults. *Environmental Health* 2018;17(1):39 <http://doi.org/10.1186/s12940-018-0381-2>
- Pyle A., Anugraha H., Kurzawa-Akanbi M., Yarnall A., Burn D., Hudson G. Reduced mitochondrial DNA copy number is a biomarker of parkinson's disease. *Neurobiology of Aging* 2016;38:216.e217-216.e210 <https://doi.org/10.1016/j.neurobiolaging.2015.10.033>
- Saenen N.D., Provost E.B., Viaene M.K., Vanpoucke C., Lefebvre W., Vrijens K., et al. Recent versus chronic exposure to particulate matter air pollution in association with neurobehavioral

performance in a panel study of primary schoolchildren. *Environment International* 2016;95:112-119
<https://doi.org/10.1016/j.envint.2016.07.014>

Saenen N.D., Provost E.B., Cuypers A., Kicinski M., Pieters N., Plusquin M., et al. Child's buccal cell mitochondrial DNA content modifies the association between heart rate variability and recent air pollution exposure at school. *Environment International* 2019;123:39-49
<http://doi.org/10.1016/j.envint.2018.11.028>

Shalev I. Early life stress and telomere length: Investigating the connection and possible mechanisms: A critical survey of the evidence base, research methodology and basic biology. *BioEssays* 2012;34(11):943-952 <https://doi.org/10.1002/bies.201200084>

Shaughnessy D.T., McAllister K., Worth L., Haugen A.C., Meyer J.N., Domann F.E., et al. Mitochondria, energetics, epigenetics, and cellular responses to stress. *Environmental Health Perspectives* 2014;122(12):1271-1278 <https://doi.org/10.1289/ehp.1408418>

Shivakumar V., Rajasekaran A., Subbanna M., Kalmady S.V., Venugopal D., Agrawal R., et al. Leukocyte mitochondrial DNA copy number in schizophrenia. *Asian Journal of Psychiatry* 2020;53:102193 <https://doi.org/10.1016/j.ajp.2020.102193>

Simioni C., Zauli G., Martelli A.M., Vitale M., Sacchetti G., Gonelli A., et al. Oxidative stress: Role of physical exercise and antioxidant nutraceuticals in adulthood and aging. *Oncotarget* 2018;9(24):17181-17198 <https://doi.org/10.18632/oncotarget.24729>

Skuratovskaia D., Litvinova L., Vulf M., Zatolokin P., Popadin K., Mazunin I. From normal to obesity and back: The associations between mitochondrial DNA copy number, gender, and body mass index. *Cells* 2019;8(5):430 <https://doi.org/10.3390/cells8050430>

Steiner J.L., Murphy E.A., McClellan J.L., Carmichael M.D., Davis J.M. Exercise training increases mitochondrial biogenesis in the brain. *Journal of Applied Physiology* 2011;111(4):1066-1071
<http://doi.org/10.1152/japplphysiol.00343.2011>

Stockton J.C., Duke-Williams O., Stamatakis E., Mindell J.S., Brunner E.J., Shelton N.J. Development of a novel walkability index for london, united kingdom: Cross-sectional application to the whitehall ii study. *BMC Public Health* 2016;16:416 <http://doi.org/10.1186/s12889-016-3012-2>

Su J.G., Dadvand P., Nieuwenhuijsen M.J., Bartoll X., Jerrett M. Associations of green space metrics with health and behavior outcomes at different buffer sizes and remote sensing sensor resolutions. *Environment International* 2019;126:162-170 <http://doi.org/10.1016/j.envint.2019.02.008>

Telomere research network. 2020. Study design and analysis. Available:
<https://trn.tulane.edu/resources/study-design-analysis/> [accessed 21/02/2022].

- Vriens A., Nawrot T.S., Baeyens W., Den Hond E., Bruckers L., Covaci A., et al. Neonatal exposure to environmental pollutants and placental mitochondrial DNA content: A multi-pollutant approach. *Environment International* 2017;106:60-68 <https://doi.org/10.1016/j.envint.2017.05.022>
- Vyas S., Zaganjor E., Haigis M.C. Mitochondria and cancer. *Cell* 2016;166(3):555-566 <https://doi.org/10.1016/j.cell.2016.07.002>
- Wang X., Hart J.E., Liu Q., Wu S., Nan H., Laden F. Association of particulate matter air pollution with leukocyte mitochondrial DNA copy number. *Environment International* 2020;141:105761 <https://doi.org/10.1016/j.envint.2020.105761>
- Wu S., Li X., Meng S., Fung T., Chan A.T., Liang G., et al. Fruit and vegetable consumption, cigarette smoke, and leukocyte mitochondrial DNA copy number. *The American Journal of Clinical Nutrition* 2019;109(2):424-432 <https://doi.org/10.1093/ajcn/nqy286>
- Zhou W., Zhu M., Gui M., Huang L., Long Z., Wang L., et al. Peripheral blood mitochondrial DNA copy number is associated with prostate cancer risk and tumor burden. *PloS One* 2014;9(10):e109470 <https://doi.org/10.1371/journal.pone.0109470>

Table 1: Characteristics of the study population (n = 246)

Anthropometric characteristics	Mean (SD) or n (%)
Girls	115 (46.8%)
Age, years	10.3 (1.3)
Body Mass Index (BMI) (kg/m ²)	17.4 (2.8)
Lifestyle characteristics	
Highest category of education of either parent	
Up to high school diploma	75 (30.5%)
College or university diploma	171 (69.5%)
Exposed to passive tobacco smoke	39 (15.9%)
School	
Tienen	56 (22.8%)
Zonhoven	122 (49.6%)
Hasselt-Kiewit	68 (27.6%)
Distance to major road (m)	576.2 (559.0)
Physical activity (hours/week) ^a	3.4 (2.4)

^a Data available for 240 children.

Table 2: Residential exposure characteristics (n = 246).

	Mean	Median	IQR
Low green (%)			
100 m radius	34.2	35.0	18.5
300 m radius	31.6	32.4	12.8
500 m radius	30.4	31.2	11.3
1000 m radius	28.7	31.5	11.0
High green (%)			
100 m radius	16.1	13.7	11.7
300 m radius	19.1	17.5	8.9
500 m radius	20.2	19.2	9.0
1000 m radius	20.9	20.2	9.5
Total green (%)			
100 m radius	50.2	52.7	22.0
300 m radius	50.7	52.7	18.6
500 m radius	50.6	52.9	18.8
1000 m radius	49.6	53.5	13.9
Agriculture (%)			
100 m radius	6.1	0.0	8.3
300 m radius	11.5	6.6	15.0
500 m radius	13.7	9.1	12.7
1000 m radius	17.0	13.5	11.6
Distance to green space > 0.5 ha (m)	210.8	123.7	144.6

IQR: interquartile range; Low green: <3m; High green: ≥3m; Total green: sum of low and high green


Table 3. Sensitivity analysis for the 1000 m radius and distance to green space.


	n	Mitochondrial DNA content (mtDNAC)		
		% Difference	95% CI	p-value
Low green	436	15.2	7.2 to 23.7	0.00014
+ Physical activity	424	14.4	6.1 to 23.3	0.00051
+ PM _{2.5} exposure	436	14.2	6.3 to 22.8	0.00037
+ UFP exposure	436	15.2	7.2 to 23.8	0.00014
+ Physical activity, PM _{2.5} and UFP	424	13.6	5.4 to 22.5	0.0010
High green	436	10.8	4.5 to 17.5	0.00070
+ Physical activity	424	10.2	3.7 to 17.1	0.0018
+ PM _{2.5} exposure	436	9.9	3.4 to 16.7	0.0024
+ UFP exposure	436	10.7	4.4 to 17.4	0.00079
+ Physical activity, PM _{2.5} and UFP	424	9.3	2.7 to 16.3	0.0051
Total green	436	13.4	7.4 to 19.7	< 0.0001
+ Physical activity	424	13.0	6.7 to 19.6	< 0.0001
+ PM _{2.5} exposure	436	12.7	6.6 to 19.2	< 0.0001
+ UFP exposure	436	13.3	7.4 to 19.7	< 0.0001
+ Physical activity, PM _{2.5} and UFP	424	12.4	6.0 to 19.1	0.00012
Agriculture	436	-3.4	-6.7 to -0.1	0.047
+ Physical activity	424	-3.0	-6.5 to 0.7	0.11
+ PM _{2.5} exposure	436	-3.4	-6.7 to 0.1	0.055
+ UFP exposure	436	-3.4	-6.8 to 0.03	0.052
+ Physical activity, PM _{2.5} and UFP	424	-2.9	-6.4 to 0.8	0.12
Distance to green space > 0.5 ha	436	-5.2	-7.9 to -2.4	0.00040
+ Physical activity	424	-4.9	-7.6 to -2.0	0.00098
+ PM _{2.5}	436	-5.0	-7.7 to -2.2	0.00057
+ UFP exposure	436	-5.2	-7.9 to -2.4	0.00036
+ Physical activity, PM _{2.5} and UFP	424	-4.8	-7.5 to -1.9	0.0012

Percentage difference (95% CI) in relative average mitochondrial DNA content (mtDNAC) for an IQR increment in residential green space or a doubling in distance to green space > 0.5 ha. The main model was adjusted for sex, age, BMI, highest educational level of either parent, passive smoking, season, date of examination, distance to major road, and the random effect of school and subject. In the sensitivity analysis, the models were additionally adjusted for physical activity, PM_{2.5} exposure during the year before the examination, and ultrafine particle (UFP) < 300 nm exposure at school separately and combined.

Figure 1: Difference (%) in average relative mitochondrial DNA content (mtDNAC) (with 95% CI) in association with an IQR increment in residential green space (high green, low green, total green, and agriculture). Because of the skewedness of the data for the 100 m radius of agriculture, we expressed the effect size for a dichotomized variable comparing the presence to the absence of agriculture within this radius. Estimates were adjusted for sex, age, BMI, highest educational level of either parent, passive smoking, season, date of examination, distance to major road, and the random effect of school and subject. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Figure 2: Scatterplot showing the association between distance to green space > 0.5 ha (area with more than 0.5 ha of green) and average residualized mtDNAC per child. MtDNAC was residualized by regressing on the covariates (age, sex, BMI, passive smoking, season of examination, highest educational level of either parent, date of examination and distance to major road).





Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: