

Editorial

COVID-19: Lessons on malnutrition, nutritional care and public health from the ESPEN-WHO Europe call for papers

S U M M A R Y

Keywords:
 COVID-19
 Sars-CoV-2
 Malnutrition
 Nutritional care
 Public health

With prolonged pandemic conditions, and emerging evidence but persisting low awareness of the importance of nutritional derangements, ESPEN has promoted in close collaboration with World Health Organization-Europe a call for papers on all aspects relating COVID-19 and nutrition as well as nutritional care, in the Society Journals Clinical Nutrition and Clinical Nutrition ESPEN. Although more COVID-related papers are being submitted and continue to be evaluated, ESPEN and WHO present the current editorial to summarize the many published findings supporting major interactions between nutritional status and COVID-19. These include 1) high risk of developing the disease and high risk of severe disease in the presence of pre-existing undernutrition (malnutrition) including micronutrient deficiencies; 2) high risk of developing malnutrition during the course of COVID-19, with substantial impact on long-term sequelae and risk of long COVID; 3) persons with obesity are also prone to develop or worsen malnutrition and its negative consequences during the course of COVID-19; 4) malnutrition screening and implementation of nutritional care may improve disease outcomes; 5) social and public health determinants contribute to the interaction between nutritional status and COVID-19, including negative impact of lockdown and social limitations on nutrition quality and nutritional status. We believe the evidence supports the need to consider COVID-19 as (also) a case of malnutrition-enhanced disease and disease-related malnutrition, with added risk for persons both with and without obesity. Similarities with many other disease conditions further support recommendations to implement standard nutritional screening and care in COVID-19 patients, and they underscore the relevance of appropriate nutritional and lifestyle prevention policies to limit infection risk and mitigate the negative health impact of acute pandemic bouts.

© 2022 Elsevier Ltd and European Society for Clinical Nutrition and Metabolism. All rights reserved.

1. Introduction

The COVID-19 pandemic continues to impose substantial global threats for individual health and public health systems. A strong association between nutritional derangements and COVID-19 has been hypothesized very early in the pandemic course, with potential substantial impact of malnutrition on infection risk, disease severity and outcome. ESPEN has therefore rapidly issued guidance documents for nutritional management of COVID-19 patients, at the onset of the pandemic and later in 2021 to focus on the emerging nutritional risk in COVID-19 patients with obesity [1,2]. With prolonged pandemic conditions and emerging evidence, but persisting low awareness of the importance of nutritional derangements, ESPEN has promoted in close collaboration with World Health Organization (WHO)-Europe a call for papers on all aspects relating COVID-19 and nutrition as well as nutritional care, in the Society Journals Clinical Nutrition and Clinical Nutrition ESPEN. ESPEN and WHO-Europe present the current paper to summarize the many findings published as a result of this call in ESPEN journals. In particular, the paper was prepared by collecting and summarizing all original

articles accepted and published in the ESPEN journals Clinical Nutrition and Clinical Nutrition ESPEN on the topic of COVID-19 until April 2022. Due to the nature of this work, that is not a formal systematic review, there were no explicitly defined inclusion and exclusion criteria related to study type, exposure, outcome and population group. A comprehensive review of papers published in other journals was also outside the scope of the current work, which is focused on the ESPEN-WHO Europe call for papers. The authors then identified five main topics under which published papers were sub-divided, based on authors judgment, into general fields of interaction between disease and nutritional status: 1) negative impact of pre-existing undernutrition (hereafter indicated as malnutrition according to clinical terminology in the clinical nutrition community) and micronutrient deficiencies on COVID-19 risk and severity; 2) COVID-19-related malnutrition as a case of disease-related malnutrition (DRM), and its potential involvement in the more recently-defined long COVID; 3) association between obesity and malnutrition in COVID-19 and their clinical impact; 4) initial evidence supporting implementation of nutritional treatment to improve COVID-19 outcomes; 5) lifestyle and public health nutritional implications.

2. Negative impact of pre-existing malnutrition and micronutrient deficiencies on COVID-19 risk and severity

COVID-19 is a respiratory disease caused by SARS-CoV-2 infection that can lead to respiratory insufficiency with potentially fatal complications [3–7]. Patient groups with pre-existing comorbidities as well as older age have been soon recognized to have higher risk for severe disease and mortality [3–7]. Also importantly, both older age and polymorbidity are per se associated with high risk and prevalence of malnutrition [8,9] due to combined catabolic derangements, low food intake and low physical activity, resulting in loss of body and skeletal muscle mass and function [8–10]. Malnutrition is conversely an independent cause of morbidity and mortality in most disease conditions, through mechanisms including negative impact on immune function and high risk of infections or superinfection [11,12] as well as respiratory and cardiac muscle dysfunction [13,14]. Micronutrient deficiencies are also common in high-risk and older individuals, and they are well-recognized risk factors for immune dysfunction and consequent infection risk [2]. Malnutrition characterized by loss of skeletal muscle mass may also notably worsen quality of life, disability and morbidities in disease survivors long after infection resolution and hospital discharge [8–10,15–17].

2.1. Malnutrition: prevalence

Consistent with these concepts, evidence has emerged that risk of malnutrition, and malnutrition per se, are highly prevalent in COVID-19 patients in various clinical settings [18–24], fully consistent with high risk of COVID-19 infection in malnourished individuals. The ESPEN-WHO Europe call has contributed to further establish and expand available information. Malnutrition was confirmed to be common in both non-ICU and ICU hospitalized cohorts and in various age groups, with combined prevalence of malnutrition and malnutrition risk consistently above 40% [24–30], with a systematic review of 27 studies identifying average prevalence at 49% [31]. The recent GLIM approach for malnutrition diagnosis was also employed, and detected malnutrition with comparable prevalence [29,32], while reports from several Countries and world regions confirmed the global negative role of malnutrition in the COVID-19 pandemic [29,32].

2.2. Malnutrition: negative clinical impact

Similar to virtually any other disease condition [10], malnutrition risk and malnutrition per se were associated with higher disease severity and worse outcomes in early reports on COVID-19 [18–21,23,24,27]. Several studies published in the ESPEN-WHO Europe call have further confirmed and dissected this association. In non-ICU patients with COVID-19, poor nutritional state was associated with higher mortality in a systematic review [31] and with higher risk of transfer to ICU from hospital wards [30] as well as longer length of hospital stay [30,33], active complications at discharge [34], and ultimately negative outcomes and mortality in several studies and databases [29,30,34,35]. Specifically in geriatric settings, malnutrition risk at admission predicted hospital length of stay [27] while MNA-assessed malnutrition also predicted in-hospital mortality in a large population cohort [36]. Despite ongoing discussions on its suitability for assessment of nutritional status in ICU, the NUTRIC score also predicted mortality in critically ill patients [24,37]. Independently of other nutritional parameters, self-reported reduction of food intake before admission, and physician judgment of expected low food intake after admission were also reported to predict admission to ICU and mortality in non-ICU patients [38]. This finding is notably consistent

with observations that recent pre-admission weight-loss is an independent risk factor for mortality in all-cause hospitalized patients from the nDay database [39].

2.3. Malnutrition: role of muscle mass

Besides changes in body mass, altered body composition with selective reduction in skeletal muscle mass has clearly emerged as a major determinant of malnutrition-induced acute and chronic complications, particularly in disease-related malnutrition [10]. New evidence from the ESPEN-WHO Europe call has unveiled that altered body composition per se, with low measured or estimated muscle and lean body mass, are also predictors of poor clinical outcomes in COVID-19. In non-ICU patients [40] phase-angle measured with bioelectrical impedance, a surrogate of lean body mass, was associated with 90-day mortality as well as longer hospital stay and inflammation markers, such as low albumin and elevated C-reactive protein. Phase angle also predicted COVID-19 severity with a composite score of morbidity, ICU admission and mortality in a mixed non-ICU and ICU population [41,42]. In several studies, low muscle mass measured by thoracic CT scans in hospitalized COVID-19 patients was associated with clinical parameters ranging from risk of ICU admission to successful ventilator weaning, length of stay and in-hospital mortality [43]. Consistent with these findings, a meta-analysis of nine studies confirmed associations between low skeletal muscle mass and COVID-19 severity and mortality [44]. Low muscle quality, particularly myosteatosis with high muscle lipid content, is an additional component of sarcopenia closely associated with malnutrition and potentially low muscle mass [10]. CT-assessed thoracic myosteatosis was accordingly also independently associated with negative ICU outcomes [43] or with transition to severe disease [45] in ICU patient cohorts.

2.4. Micronutrient deficiencies: prevalence and impact

Several vitamins (vitamin A, B₆, B₁₂, folate, C, D, E) and trace elements (Zinc, Selenium, Copper) are known to play important roles in regulating both innate and adaptive immune responses, with putative negative roles of micronutrient deficiencies to enhance susceptibility to infectious diseases including, but not limited to, viral respiratory tract infections [46]. Vitamin D is the most commonly measured micronutrient, and vitamin D deficiencies are common in many high-risk conditions including older age and obesity [2], being associated with viral infections including influenza. An early association between lower plasma vitamin D and risk of Sars-CoV2 infection was reported early in the course of the pandemic [47], and studies from the ESPEN-WHO Europe call have addressed and expanded this concept. Most reports confirmed the association between vitamin D deficiency and COVID-19 infection in the general population both in Europe and Asia [48], as well as in hospitalized patients with COVID-19 [49,50], with highest prevalence in older patient groups with obesity [51,52]. Low 25OH vitamin D was reported in ICU patients with COVID-19 and associated with higher systemic inflammation markers in one study [53,54]. Another study further suggested that lower 25OH vitamin D may predict risk of death, particularly in older COVID-19 ICU patients [55]. The association between low 25OH vitamin D and poor clinical outcomes was however not confirmed in a larger study that actually suggested a paradoxical association between high 25OH vitamin D and mortality [56]. Inconsistent findings could at least in part be explained by heterogeneous measurements, since another ICU study reported no clinical impact of low 25OH vitamin D but higher disease severity and need for mechanical ventilation in patients with low 1,25OH dihydroxylated form [57]. In additional studies, deficient zinc and

iron status were also reported to be common in patients with COVID-19 [26,49,58]. Hypozincemia was reported in early disease stages and associated with risk of hospitalization [58], whereas low iron status was associated with severe disease in hospitalized patients [59].

Altogether the above-summarized reports provide strong support for a causal link between malnutrition, micronutrient deficiencies and risk of COVID-19 infection and severity (Fig. 1).

3. COVID-19: a case of disease-related malnutrition (DRM) - impact on long covid

The ESPEN guidelines on definitions and terminology in clinical nutrition define DRM as a “specific type of malnutrition caused by a concomitant disease”. Inflammation is common in most acute and chronic disease conditions, representing a key determinant of DRM [10,11,16]. While factors triggering inflammation may vary in different diseases, metabolic and nutritional consequences of the inflammatory response are common and include skeletal muscle catabolism with muscle loss as well as anorexia and reduced food intake. Reduced physical activity or bedrest may also contribute to muscle catabolism. The same ESPEN guideline also generally defined two levels of inflammation intensity, with milder inflammatory response being more common in chronic disease and stronger inflammation in acute disease- or injury-related malnutrition [10].

COVID-19 is an acute infectious disease with potentially strong inflammatory response also at systemic level [60]. In addition, gastrointestinal symptoms, ageusia and anosmia are common and may contribute to reduce food intake. Pre-existing comorbidities may also occur, particularly in older patients, thereby synergistically enhancing catabolic responses. The combined catabolic impact of inflammation, bedrest and lack of spontaneous food intake may be strongest in intensive care COVID-19 patients [16]. Again the ESPEN terminology guideline [10] indicates that high pro-inflammatory cytokines, increased stress hormones as well as treatment, resistance to insulin, bed rest and no or reduced food intake “pave the way for a fast decline of body energy and nutrient stores”.

Evidence of malnutrition ensuing or being exacerbated during the course of COVID-19 has accumulated in the past two years and the ESPEN-WHO Europe call to action has contributed to clarify nutritional defects and their clinical consequences. Importantly, acute impairment of nutritional state may have prolonged consequences on patient health, by favoring the onset of complications and comorbidities and by impairing functional status, thereby directly causing and sustaining the long-term clinical picture that is currently defined as “long COVID”. COVID-induced malnutrition, with potential muscle loss and sarcopenia, may indeed directly contribute to major symptoms of long COVID such as fatigue and

weakness, which may delay recovery and cause prolonged disability in a large fraction of COVID patients following acute infection.

3.1. COVID-19 causes malnutrition

Several studies reported high prevalence of moderate to severe weight loss (>5 or 10% of body weight) during the course of COVID-19 in non-ICU [61–66] and ICU hospitalized patients [63,67,68], with malnutrition prevalence at discharge reaching >80% in ICU survivors [67]. One study also reported 21% prevalence of weight loss above 5% also in non-hospitalized patients [69]. Also important, one study investigated changes in body composition measured by body impedance analysis in a small non-ICU patient cohort [70], reporting stable body fat and water but significant reduction in phase angle during two-week hospital stays. Causes of negative energy balance appeared to include lower food intake also before admission [65,71], gastrointestinal symptoms with reduced appetite, feeling of gastric and abdominal fullness, altered taste and smell [66,71]. Higher inflammation as assessed by C-reactive protein was also reported in weight-losing compared to weight-stable individuals [69]. In post-intubation ICU patients, but also in the non-ICU setting, a high prevalence of dysphagia was reported [64,72]. Also relevant for ICU patients, marked and prolonged hyper-metabolism was consistently described by different authors through indirect calorimetry [73–76]; from a methodological standpoint, poor predictive value of common equations to estimate energy expenditure was reported by one group [75]. A pilot study also addressed the issue of use of protective masks while implementing indirect calorimetry [77] suggesting superimposable results with or without protective device. In full agreement with the above observations, increased prevalence of malnutrition according to GLIM criteria was documented during hospital stay from 18 to 79% in one study, with 55% characterized as severe [68], and up to 80% malnutrition prevalence was reported at discharge from ICU also in another study [67].

3.2. COVID-19-INDUCED malnutrition: clinical impact and long COVID

COVID-19-induced impairment of nutritional status was associated with negative impact on patient clinical outcomes. Weight loss was associated with longer disease duration and hospital stay [63,64,69] as well as risk of ICU transfer [66]. Impaired functional status was reported at discharge based on Barthel score in a non-ICU group [64] and lower mobility and ability to perform usual daily activities were observed at discharge in 70% of ICU patients [67]. A few studies documented follow-up of nutritional parameters for up to six months. Low appetite and gastro-intestinal complaints were reported in approximately one third of patients up to 5 months after discharge with parallel continuing weight loss




Fig. 1. Schematic representation of the potential causal link between malnutrition and Covid-19.

in 28% of patients [66]. Malnutrition according to GLIM criteria was also reported to be persistently high with more than 50% prevalence and approximately one third severe cases three months after ICU discharge [68]. Impaired functional status was also not completely recovered six months after discharge in a general hospital cohort particularly in the presence of in-hospital dysphagia [64].

Taken together, the above summarized reports provide support for COVID-19 infection as a strong risk factor for the deterioration of the nutritional status and the development of malnutrition (Fig. 2).

4. Obesity and malnutrition in COVID-19 and their clinical impact

Early evidence from COVID-19 cohorts has shown that persons with obesity are at high risk for severe disease and mortality [78–83]. Obesity-induced metabolic derangements including hypertension, insulin resistance and hyperglycemia and the metabolic syndrome (MS) have been reported to worsen COVID-19 severity and prognosis and may partly explain obesity-associated clinical risk [84–86]. Pre-existing systemic and tissue low-grade inflammation may also contribute to impaired immune function in persons with obesity with enhanced risk for organ damage and failure in the presence of COVID-19 [87]. From the nutritional standpoint, obesity with metabolic complications, comorbidities and older age also increased risk for loss of skeletal muscle mass and function, which may lead to a double burden of over-nutrition and malnutrition [88–92]. Also importantly, obesity is a strong risk factor for many diseases and for chronic and acute organ failures [93–95], with additional independent negative impact on muscle mass and nutritional status [96–99]. It should be further pointed out that obesity is commonly associated with micronutrient deficiencies that may negatively affect both skeletal muscle protein metabolism and immune function [100,101], thereby enhancing risk of infection and loss of skeletal muscle mass. The concept of the double burden of obesity and malnutrition is still underappreciated, but efforts are being made to enhance awareness on this topic. The recent GLIM criteria for malnutrition diagnosis allow for practical detection of malnutrition when involuntary weight loss or low muscle mass are observed at any level of body mass index (BMI), provided that low food intake or disease are also present (GLIM). Most importantly, when malnutrition or malnutrition risk are detected, nutritional care should be implemented also in persons with obesity. ESPEN has issued in 2021 a guidance paper for nutritional management of persons with obesity and COVID-19 endorsed by the ESPEN Council, suggesting malnutrition screening for all patients with obesity and risk factors for malnutrition and muscle loss, with particular regard to comorbidities and older age [1].

The ESPEN WHO Europe call has enhanced available evidence for a negative clinical impact of concomitant malnutrition in patients with obesity and COVID-19. Indeed, several studies on the

prevalence of malnutrition and micronutrient deficiencies reported on clinical cohorts with overall average BMI in the overweight and obesity range, that were also affected by malnutrition or malnutrition risk [25,32–34,37,51]. These findings confirm that malnutrition is also a risk factor for infection and severe COVID-19 in persons with obesity. Also important, most studies on COVID-19-induced malnutrition similarly include overweight and obese subgroups, representing up to more than 90% of the study cohort [63,64,66–68], confirming that persons with obesity are prone to the negative nutritional impact of COVID-19. Moreover, studies on altered body composition showing a specific negative impact on lean body mass also reported on patient groups with average BMI in the obesity range [41,70], supporting the concept that loss of skeletal muscle mass may be the most prominent feature of malnutrition in persons with obesity. Additional papers reported on negative outcomes in COVID-19 patients with obesity and metabolic syndrome or elevated inflammation markers, that may also impair nutritional status [102–104]. Patients with high visceral adiposity, also a major risk factor for inflammation and low muscle mass, were found to also have higher risk for severe COVID-19 in a systematic review and meta-analysis [105].

Taken together, these findings provided strong support for a close nutritional screening and assessment of COVID-19 patients with overweight and obesity for early detection of malnutrition which may enhance obesity-associated risk of poor COVID-19 outcomes and requires nutritional treatment (Fig. 3).

5. Nutritional treatment to improve COVID-19 outcomes: benefits and hurdles

As malnutrition has been gradually established as both a consequence of COVID-19 infection and a risk factor for COVID-19 infection and severity in the past two years, studies on the clinical impact of nutritional treatment to improve COVID-19 outcomes are less numerous. Evidence is however increasingly available to indicate beneficial clinical effects of medical nutrition.

5.1. Benefits: acute setting (hospital and ICU)

Several papers on from the ESPEN-WHO Europe call for COVID-19 papers have reported on the impact medical nutrition in critically ill patients that are at highest nutritional risk. Earlier implementation of nutritional care within 48 h of ICU admission was associated with survival in two studies [26,106]. In another study [71], ICU patients reaching satisfactory caloric intake through any route similarly had lower mortality compared to those not reaching nutritional targets. In a Brazilian ICU cohort, protein provision above 0.8 g/Ideal Body Weight/day was also associated with lower mortality [107]. A protective impact was also reported in a mixed cohort of non-ICU and ICU patients [108] for reaching overall adequacy of calorie and protein provision at 25 kcal and 1.2–1.5 g per kg/day,




Fig. 2. Schematic representation of the link between Covid-19 and malnutrition development.




Fig. 3. Schematic representation of the proposed nutritional screening and treatment algorithm in Covid-19 patients.

respectively. Conversely, in a study from the United States, intolerance to enteral nutrition in ICU was associated with nutritional complications including higher risk of malnutrition, higher risk of organ failure and overall mortality [109]. Also in the ICU setting [110], parenteral nutrition in patients receiving non-invasive ventilation was per se associated with lower odds of oxygen escalation. Consistent with high prevalence of malnutrition in COVID-19 patients with overweight and obesity, and with high risk of developing malnutrition during the course of COVID-19 [2], most of the above reports on nutritional treatment included large subgroups of patients with overweight and obesity.

It should be pointed out that evidence on potential beneficial effects of micronutrient supplementation is less conclusive. With regard to oral supplements, a systematic review of 12 randomized controlled trials on use of dietary herbal supplements to increase micronutrient provision [111] showed that high risk of bias makes it difficult to interpret results; the authors conclude that some promising effects were suggested in improving recovery, and zinc supplements could lead to faster recovery from olfactory dysfunction. A meta-analysis focusing on vitamin C, vitamin D and zinc reported beneficial effects of vitamin D supplementation in reducing risk of intubation during COVID-19 illness, whereas no significant impact was detected for vitamin C and zinc on mortality, intubation risk and length of hospital stay [112].

5.2. Benefits: post discharge

Evidence of nutritional derangements and high risk of malnutrition following COVID-19 infection strongly support continuing

nutritional follow-up and treatment for several months after infection resolution, to potentially prevent or treat nutritional components of long COVID. Very few reports are however available on sustained nutritional treatment in this setting. In the setting of the ESPEN-WHO call for papers, one study from Italy in non-ICU patients reporting post-COVID fatigue investigated the impact of a commercial oral supplement including vitamins, minerals, amino acids and plant extracts [113]. Questionnaire-based improvements of quality of life, fatigue and global health status were reported after 14 and 28 days [113]. Another study described the effects of a multidisciplinary rehabilitation program for patients discharged from acute COVID hospitals in northern Italy after worsening of nutritional status [62]. 61% of patients had developed malnutrition during hospitalization according to GLIM criteria, despite average BMI in the overweight range [62]. Four-week interventions including diet optimization (27–30 kcal and 1–1.3 g protein per kg/day) and controlled aerobic and resistance exercise were reported to improve body composition with increased phase angle and muscle strength [62]. Further studies are clearly needed to assess the potential positive clinical impact of rigorous comprehensive nutritional intervention programmes on nutritional sequelae of COVID-19.

5.3. Hurdles: awareness

Since the start of the pandemic, ESPEN and other scientific Societies have issued guidance papers on implementation of nutritional care for prevention and treatment of malnutrition in patients with COVID-19 [1,2], aimed at reducing morbidity and

mortality as well as infection risk particularly in high-risk patients with older age and pre-existing comorbidities [1,2]. ESPEN also specifically addressed the issue of obesity [2], with largely similar recommendations. Guidance for pediatric patients is also available [114] for specific needs of this age group [115]. Despite early availability of guidance documents, several hurdles may however impair implementation of nutritional care in routine clinical practice. Low awareness of the negative impact of malnutrition and the importance of nutritional care in virtually any acute disease conditions is a pre-existing hurdle [116–120]. Even in tertiary care centers, inadequate implementation of nutritional care in patients with COVID-19 has been documented when compared to malnutrition prevalence [28,67]. In surveys from South-East Asia, limitations were confirmed in implementation of nutrition guidelines in COVID-19 patients, with reasons including lack of confidence as well as budget limitations [121]. Similarly, insufficient knowledge of nutrition management was self-reported by ICU personnel in a Swiss survey [122]. Importantly, low adherence to nutritional guidelines was associated with negative clinical outcomes including higher length of stay in ICU and mortality [121]. On the other hand, studies reporting adequate implementation of nutritional care in ICU to reach nutritional targets also reported good access to healthcare professionals such as nutritionists and dietitians [123–126]. The ESPEN-WHO Europe call for papers on COVID-19 therefore also contributes to highlight the need for action to increase awareness on the importance of nutritional care in this setting, both for healthcare professionals and policy-makers.

5.4. Hurdles: practical COVID-related issues

The onset of the COVID-19 pandemic and pandemic waves at different times around the world have also posed huge practical hurdles to implementation of nutritional care, as priorities and resources for healthcare systems had to shift towards COVID-19. Negative consequences included limitation of workforce, need for changes in hospital organization and limited physical access to patients for examination and monitoring. The ESPEN-WHO Europe call for papers has also allowed to collect reports on potential approaches to mitigate practical hurdles to nutritional care. Simplified organization of nutrition protocols for ICU healthcare providers to COVID-19 patient has been proposed and implemented [122], with higher adherence to nutritional targets reported in a subsequent study in the same setting [127]. Reports on re-organizations of nutrition support activities, and the specific role of different professional components of the nutritional team have been also provided [128]. Limited or no patient contact has been also addressed both in the hospital and home care settings. Recommendations have been proposed for remote evaluation of the anthropometric component of nutritional assessment in ambulatory settings [129], and a pilot study reported good performance of an abridged format of Subjective Global Assessment without physical examination [130]. Results of the latter study appeared to support the use of the abridged format, with good comparison in detection of malnutrition with the complete form [130]. A remote nutritional screening tool suitable for app implementation has been also proposed [131]. The ESPEN Special Interest group for Home Artificial Nutrition issued early during the pandemic a position paper with considerations on the potentially problematic topic of management of home parenteral nutrition in the setting of pandemic-related limitations [132], with additional statements coming from a network of home nutrition centers in Poland [133]. Altogether, the above reports have started to provide guidance and practical examples for effective implementation of nutritional care in COVID-19 patients.

6. Lifestyle and public health nutritional implications

The COVID-19 pandemic also created fundamental challenges through profound limitations in social interactions with potential negative impact on individual lifestyle habits and nutrition. Variable reduction of economic activities with loss of income may have also directly and negatively affected food quality and security; to this regard the pandemic also exacerbated or unveiled social and regional inequalities in terms of food habits and access to balanced and adequate nutrition. The ESPEN-WHO Europe call for papers also addressed these topics, and several studies have been published on these important issues.

6.1. Lifestyle

Measures to limit the spread of the COVID-19 pandemic have included strong limitations to usual activities and social confinement, up to general lockdowns. Changes in dietary habits and their impact on body weight have been investigated in original studies and systematic reviews. Different patterns of dietary changes have been reported, indicating heterogeneous responses to lockdowns in the general population. In a large sample of >22,000 UK adults, 64% of participants reported no modifications in nutrition quantity [134], whereas two alternative minority trajectories emerged, with reductions or increments in food intake. Other studies reported a majority of individuals with overall increasing food intake [135,136]. Interestingly, emotional eating defined as food consumption in response to stress was reported to be higher under lockdown conditions in one survey, with more common emotional eating in persons with obesity and overweight [137]. A systematic review of 32 studies also reported increments in snacking and alcohol consumption (10–51%) [138], and number of snacks also increased in an Iranian cohort in all body weight categories [135]. A sizable fraction of studied populations (15–41%) intriguingly reported reduced consumption of fast food and ordered food, with a counterintuitive adaptation to lockdown conditions that could favor consumption of more healthy home-prepared meals, also suggested by a subsequent report [138]. Also interestingly, an urban household survey suggested that consumption of fresh, minimally processed food is associated with lower prevalence of anxiety or depression compared to ultra-processed food [139]. More studies are needed to further investigate the potential role of healthy food consumption to prevent nutritional and mental negative consequences during lockdown conditions. Body weight changes during lockdowns were investigated in a systematic review of 36 studies that reported an overall trend for increments (11–72%) although this was accompanied by a sizable fraction of the overall population reporting weight loss (7–52% in different studies) [140]. Interestingly, the only study from this review reporting overall weight loss was characterized by older age >60 [141], suggesting a potential negative impact of aging on weight maintenance during the COVID-19 pandemic. Another study in older adults did not report lockdown-induced BMI changes, but it confirmed a reduction in physical activity, associated with worsening of mental health parameters [142]. Changes in body composition should be directly investigated in future studies, since reduction of physical activity during lockdowns could favor reduction in muscle mass and function, with ensuing malnutrition and sarcopenia [143]. Since low physical activity also potentially affects intestinal transit and bowel rhythm, these parameters were investigated in a specific survey in an adult cohort; while a majority of responders reported adequate transit, approximately one quarter reported altered rhythm with either faster or slower transit [144]. Patients with eating disorders (ED) represent an additional at-risk group with potential more profound negative impact of lockdown

conditions on eating behavior patterns. A meta-analysis focusing on this patient group confirmed a high risk of worsening of ED symptoms and mental health, with potential negative impact on nutritional status during the pandemic [145]. The impact of lockdowns was also studied on additional behavioral and lifestyle components. One report from Peru found increments in body weight in a cohort of young university students [146], and this change was associated with reduced physical activity, light exposure and sleep duration. Other studies however suggested that changes in sleep duration may vary in different populations with reported no change or increase in up to 90% of participants in a study from Turkey [137].

Taken together, the above reports from the ESPEN-WHO Europe call for papers contribute to identify potential negative changes in eating behavior that may enhance the risk to develop or worsen overweight and obesity, with potential for associated reduction in muscle mass that should be specifically investigated.

6.2. Public health nutrition

Nutrition patterns and diet quality, as well as levels of food security may on one hand influence the risk of developing severe infections including COVID-19 by modulating nutritional state and immune function. On the other hand, the COVID-19 pandemic may worsen economic conditions, individual income and food security with synergistic impact on infection risk. Selected reports from the ESPEN-WHO Europe call have addressed these important issues. One study analyzed the impact of food supply available in 170 different Countries from Food and Agriculture Organization of the United Nations data on rates of recovery from COVID-19 [147]. The study demonstrated that recovery rates were highest in countries stratified as having eradicated hunger, and lowest in those with moderate or high global hunger index [147], although the potential impact of healthcare quality and accessibility should also be considered since this is a relevant potential factor that could affect study results. Major benefits from individual nutrients were attributed to animal-derived food while sources of immune-modulating micronutrients were also beneficial [147]. The potential relevance of habitual dietary patterns was further suggested in a study from South Korea showing that population groups stratified as having high risk to develop COVID-19 also had higher risk of having nutritional deficiencies, with particular regard to multiple vitamins [148]. On the other hand, higher adherence to diet rich in antioxidants, anti-inflammatory and micronutrients such as the Mediterranean diet was associated with lower risk to develop COVID-19 in a Spanish cohort [149]. Two additional studies addressed the impact of COVID-19 on food security. In one report from Jordan using a web-based validated questionnaire, 60% of participants were identified as having severe or moderate food insecurity, associated with reduced monthly income and number of family members [150]. In another paper, a systematic review was performed on the impact of COVID-19 on food systems and security [151]. Findings from 35 studies confirmed adverse effects of the pandemic that were modulated by timing, intensity and duration of social restrictions as well as by existence of social and government safety nets [151]. Inequalities in negative impact also emerged along lines of female gender, age and low socio-economic status without stable employment [151]. Concern for high risk categories was further expressed in additional papers focusing on high risk of food insecurity in children [152] and on pregnant women with diabetes mellitus [153].

Overall, the above reports from the ESPEN-WHO call contributed to confirm the extended COVID-19 challenge for public health through potentially deleterious changes in lifestyle and food

security, with a strong risk for enhancement of inequalities within and among countries and world regions.

7. Conclusions

ESPEN and WHO Europe have strongly supported the collection of scientific evidence on the role of nutrition and nutritional care to effectively address the challenges and consequences of the COVID-19 pandemic. At the time of submission of this paper more than 260,000 publications are retrievable in PubMed using COVID as keyword. The total number of publications is however substantially lower (5705 - approximately 2%) when COVID and nutrition are both used for publications search. When COVID and malnutrition are used for publications search, only 898 publications are retrieved (less than 0.4%). We are pleased to offer more than 120 publications in ESPEN journals on these important topics and hope that these findings will contribute to enhance awareness on the importance of nutrition in prevention, treatment and post-infection care of COVID-19. As we join the universal hope that the impact of the COVID-19 pandemic will decline as soon as possible through vaccinations and potential new treatments, we believe and advocate that optimal nutritional care may provide important contributions to reach this goal.

Funding

No funding was sought or received for preparation of the manuscript.

Conflict of interest

The authors declare that they have no competing interests for the content of this paper.

Covid-19 call editorial board author list

Osman Abbasoglu, Turkey
 Judith Beurskens Meijerink, Netherlands
 Stephan Bischoff, Germany
 Rosa Burgos Pelaez, Spain
 Diana Cardenas, France
 Tommy Cederholm, Sweden
 Emanuele Cereda, Italy
 Michael Chourdakis, Greece
 Maria Isabel Toulson Davisson Correia, Brazil
 Marian de van der Schuren, Netherlands
 Nathalie Delzenne, Belgium
 Evelyn Frias-Toral, Ecuador
 Laurence Genton, Switzerland
 Gianluca Gortan Cappellari, Italy
 Burcu Kelleci Cakir, Turkey
 Stanislaw Klek, Poland
 Zeljko Krznaric, Croatia
 Alessandro Laviano, Italy
 Dileep Lobo, United Kingdom
 Maurizio Muscaritoli, Italy
 Johann Ockenga, Germany
 Matthias Pirlich, Germany
 Mireille JM Serlie, Netherlands
 Han Ping Shi, China
 Pierre Singer, Israel
 Mattias Soop, Sweden
 Stephane Walrand, France
 Arved Weimann, Germany

References

- [1] Barazzoni R, Bischoff SC, Breda J, Wickramasinghe K, Krznaric Z, Nitzan D, et al. ESPEN expert statements and practical guidance for nutritional management of individuals with SARS-CoV-2 infection. *Clin Nutr* 2020;39(6):1631–8.
- [2] Barazzoni R, Bischoff SC, Busetto L, Cederholm T, Chourdakis M, Cuerda C, et al. Nutritional management of individuals with obesity and COVID-19: ESPEN expert statements and practical guidance. Edinburgh, Scotland: Clinical nutrition; 2021.
- [3] Bouadma L, Lescure FX, Lucet JC, Yazdanpanah Y, Timsit JF. Severe SARS-CoV-2 infections: practical considerations and management strategy for intensivists. *Intensive Care Med* 2020.
- [4] Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;395(10223):507–13.
- [5] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395(10223):497–506.
- [6] Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020.
- [7] Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020;382(8):727–33.
- [8] Gomes F, Schuetz P, Bounoure L, Austin P, Ballesteros-Pomar M, Cederholm T, et al. ESPEN guidelines on nutritional support for polymorbid internal medicine patients. *Clin Nutr* 2018;37(1):336–53.
- [9] Volkert D, Beck AM, Cederholm T, Cruz-Jentoft A, Goisser S, Hooper L, et al. ESPEN guideline on clinical nutrition and hydration in geriatrics. *Clin Nutr* 2019;38(1):10–47.
- [10] Cederholm T, Barazzoni R, Austin P, Ballmer P, Biolo G, Bischoff SC, et al. ESPEN guidelines on definitions and terminology of clinical nutrition. *Clin Nutr* 2017;36(1):49–64.
- [11] Calder PC. Feeding the immune system. *Proc Nutr Soc* 2013;72(3):299–309.
- [12] Schneider SM, Veyres P, Pivot X, Soummer AM, Jambou P, Filippi J, et al. Malnutrition is an independent factor associated with nosocomial infections. *Br J Nutr* 2004;92(1):105–11.
- [13] Arora NS, Rochester DF. Respiratory muscle strength and maximal voluntary ventilation in undernourished patients. *Am Rev Respir Dis* 1982;126(1):5–8.
- [14] Heymsfield SB, Bethel RA, Ansley JD, Gibbs DM, Felner JM, Nutter DO. Cardiac abnormalities in cachectic patients before and during nutritional repletion. *Am Heart J* 1978;95(5):584–94.
- [15] Hiesmayr M, Frantal S, Schindler K, Themessl-Huber M, Mouhieddine M, Schuh C, et al. The patient- and nutrition-derived outcome risk assessment score (PANDORA): development of a simple predictive risk score for 30-day in-hospital mortality based on demographics, clinical observation, and nutrition. *PLoS One* 2015;10(5):e0127316.
- [16] Singer P, Blaser AR, Berger MM, Alhazzani W, Calder PC, Casaer MP, et al. ESPEN guideline on clinical nutrition in the intensive care unit. *Clin Nutr* 2019;38(1):48–79.
- [17] Souza TT, Sturion CJ, Faintuch J. Is the skeleton still in the hospital closet? A review of hospital malnutrition emphasizing health economic aspects. *Clin Nutr* 2015;34(6):1088–92.
- [18] Allard L, Ouedraogo E, Molleville J, Bihan H, Giroux-Leprieur B, Sutton A, et al. Malnutrition: percentage and association with prognosis in patients hospitalized for coronavirus disease 2019. *Nutrients* 2020;12(12).
- [19] Hu X, Deng H, Wang Y, Chen L, Gu X, Wang X. Predictive value of the prognostic nutritional index for the severity of coronavirus disease 2019. *Nutrition* 2020;84:111123.
- [20] Li G, Zhou CL, Ba YM, Wang YM, Song B, Cheng XB, et al. Nutritional risk and therapy for severe and critical COVID-19 patients: a multicenter retrospective observational study. Edinburgh, Scotland: Clinical nutrition; 2020.
- [21] Li T, Zhang Y, Gong C, Wang J, Liu B, Shi L, et al. Prevalence of malnutrition and analysis of related factors in elderly patients with COVID-19 in Wuhan, China. *Eur J Clin Nutr* 2020;74(6):871–5.
- [22] Pironi L, Sasdelli AS, Ravaoli F, Baracco B, Battaiola C, Bocedi G, et al. Malnutrition and nutritional therapy in patients with SARS-CoV-2 disease. Edinburgh, Scotland: Clinical nutrition; 2020.
- [23] Yu Y, Ye J, Chen M, Jiang C, Lin W, Lu Y, et al. Malnutrition prolongs the hospitalization of patients with COVID-19 infection: a clinical epidemiological analysis. *J Nutr Health Aging* 2021;25(3):369–73.
- [24] Zhang P, He Z, Yu G, Peng D, Feng Y, Ling J, et al. The modified NUTRIC score can be used for nutritional risk assessment as well as prognosis prediction in critically ill COVID-19 patients. *Clin Nutr* 2021;40(2):534–41.
- [25] Larrazabal Jr RB, Perez BMB, Masamayor EMI, Chiu HHC, Palileo-Villanueva LAM. The prevalence of malnutrition and analysis of related factors among adult patients with the Coronavirus Disease 2019 (COVID 19) in a tertiary government hospital: the MalnutriCoV study. *Clinical nutrition ESPEN* 2021;42:98–104.
- [26] Li G, Zhou CL, Ba YM, Wang YM, Song B, Cheng XB, et al. Nutritional risk and therapy for severe and critical COVID-19 patients: a multicenter retrospective observational study. *Clin Nutr* 2021;40(4):2154–61.
- [27] Mendes A, Serratrice C, Herrmann FR, Gold G, Graf CE, Zekry D, et al. Nutritional risk at hospital admission is associated with prolonged length of hospital stay in old patients with COVID-19. Edinburgh, Scotland: Clinical nutrition; 2021.
- [28] Pironi L, Sasdelli AS, Ravaoli F, Baracco B, Battaiola C, Bocedi G, et al. Malnutrition and nutritional therapy in patients with SARS-CoV-2 disease. *Clin Nutr* 2021;40(3):1330–7.
- [29] Shahbazi S, Hajimohammadebrahim-Ketabforoush M, Vahdat Shariatpanahi M, Shahbazi E, Vahdat Shariatpanahi Z. The validity of the global leadership initiative on malnutrition criteria for diagnosing malnutrition in critically ill patients with COVID-19: a prospective cohort study. *Clinical nutrition ESPEN* 2021;43:377–82.
- [30] Nicolau J, Ayala L, Sanchis P, Olivares J, Dotres K, Soler AG, et al. Influence of nutritional status on clinical outcomes among hospitalized patients with COVID-19. *Clinical nutrition ESPEN* 2021;43:223–9.
- [31] Abate SM, Chekole YA, Estifanos MB, Abate KH, Kabthymmer RH. Prevalence and outcomes of malnutrition among hospitalized COVID-19 patients: a systematic review and meta-analysis. *Clinical nutrition ESPEN* 2021;43:174–83.
- [32] Bedock D, Bel Lassen P, Mathian A, Moreau P, Couffignal J, Ciangura C, et al. Prevalence and severity of malnutrition in hospitalized COVID-19 patients. *Clinical nutrition ESPEN* 2020;40:214–9.
- [33] Larrazabal Jr RB, Chiu HHC, Palileo-Villanueva LAM. Outcomes of nutritionally at-risk Coronavirus Disease 2019 (COVID 19) patients admitted in a tertiary government hospital: a follow-up study of the MalnutriCoV study. *Clinical nutrition ESPEN* 2021;43:239–44.
- [34] Stefano M, Andrea B, Daniela C, Emanuela M, Lorena P, Daniela D, et al. Malnutrition risk as a negative prognostic factor in COVID-19 patients. *Clinical nutrition ESPEN* 2021;45:369–73.
- [35] Bouziotis J, Arvanitakis M, Preiser JC. Association of body mass index with COVID-19 related in-hospital death. Edinburgh, Scotland: Clinical nutrition; 2022.
- [36] Kananen L, Eriksdotter M, Boström AM, Kivipelto M, Annetorp M, Metzner C, et al. Body mass index and Mini Nutritional Assessment-Short Form as predictors of in-geriatric hospital mortality in older adults with COVID-19. Edinburgh, Scotland: Clinical nutrition; 2021.
- [37] Liberti A, Piacentino E, Umbrello M, Muttini S. Comparison between Nutric Score and modified nutric score to assess ICU mortality in critically ill patients with COVID-19. *Clinical nutrition ESPEN* 2021;44:479–82.
- [38] Caccialanza R, Formisano E, Klersy C, Ferretti V, Ferrari A, Demontis S, et al. Nutritional parameters associated with prognosis in non-critically ill hospitalized COVID-19 patients: the NUTRI-COVID19 study. Edinburgh, Scotland: Clinical nutrition; 2021.
- [39] Barazzoni R, Sulz I, Schindler K, Bischoff SC, Gortan Cappellari G, Hiesmayr M, et al. A negative impact of recent weight loss on in-hospital mortality is not modified by overweight and obesity. *Clin Nutr* 2020;39(8):2510–6.
- [40] Cornejo-Pareja I, Vegas-Aguilar IM, García-Almeida JM, Bellido-Guerrero D, Talluri A, Lukaski H, et al. Phase angle and standardized phase angle from bioelectrical impedance measurements as a prognostic factor for mortality at 90 days in patients with COVID-19: a longitudinal cohort study. Edinburgh, Scotland: Clinical nutrition; 2021.
- [41] Moonen H, van Zanten FJL, Driessen L, de Smet V, Slingerland-Boot R, Mensink M, et al. Association of bioelectric impedance analysis body composition and disease severity in COVID-19 hospital ward and ICU patients: the BIAC-19 study. *Clin Nutr* 2021;40(4):2328–36.
- [42] Moonen HP, Bos AE, Hermans AJ, Stikkelman E, van Zanten FJ, van Zanten AR. Bioelectric impedance body composition and phase angle in relation to 90-day adverse outcome in hospitalized COVID-19 ward and ICU patients: the prospective BIAC-19 study. *Clinical nutrition ESPEN* 2021;46:185–92.
- [43] Damanti S, Cristel G, Ramirez GA, Bozzolo EP, Da Prat V, Gobbi A, et al. Influence of reduced muscle mass and quality on ventilator weaning and complications during intensive care unit stay in COVID-19 patients. Edinburgh, Scotland: Clinical nutrition; 2021.
- [44] Siahaan YMT, Hartoyo V, Hariyanto TI, Kurniawan A. Coronavirus disease 2019 (Covid-19) outcomes in patients with sarcopenia: a meta-analysis and meta-regression. *Clinical nutrition ESPEN* 2022;48:158–66.
- [45] Yi X, Liu H, Zhu L, Wang D, Xie F, Shi L, et al. Myosteatosis predicting risk of transition to severe COVID-19 infection. Edinburgh, Scotland: Clinical nutrition; 2021.
- [46] Calder PC, Carr AC, Gombart AF, Eggensdorfer M. Optimal nutritional status for a well-functioning immune system is an important factor to protect against viral infections. *Nutrients* 2020;12(4).
- [47] Meltzer DO, Best TJ, Zhang H, Vokes T, Arora V, Solway J. Association of vitamin D status and other clinical characteristics with COVID-19 test results. *JAMA Netw Open* 2020;3(9):e2019722.
- [48] Sooriyaarachchi P, Jeyakumar DT, King N, Jayawardena R. Impact of vitamin D deficiency on COVID-19. *Clinical nutrition ESPEN* 2021;44:372–8.
- [49] Elham AS, Azam J, Mostafa L, Nasrin B, Marzieh N. Serum vitamin D, calcium, and zinc levels in patients with COVID-19. *Clinical nutrition ESPEN* 2021;43:276–82.
- [50] Tehrani S, Khabiri N, Moradi H, Mosavat MS, Khabiri SS. Evaluation of vitamin D levels in COVID-19 patients referred to Labafinejad hospital in Tehran and its relationship with disease severity and mortality. *Clinical nutrition ESPEN* 2021;42:313–7.
- [51] Gonçalves TJM, Gonçalves S, Guarnieri A, Risegato RC, Guimarães MP, de Freitas DC, et al. Prevalence of obesity and hypovitaminosis D in elderly with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Clinical nutrition ESPEN* 2020;40:110–4.

- [52] Biesalski HK. Obesity, vitamin D deficiency and old age a serious combination with respect to coronavirus disease-2019 severity and outcome. *Curr Opin Clin Nutr Metab Care* 2021;24(1):18–24.
- [53] Pimentel GD, Dela Vega MCM, Laviano A. High neutrophil to lymphocyte ratio as a prognostic marker in COVID-19 patients. *Clinical nutrition ESPEN* 2020;40:101–2.
- [54] Pimentel GD, Dela Vega MCM, Pichard C. Low vitamin D levels and increased neutrophil in patients admitted at ICU with COVID-19. *Clinical nutrition ESPEN* 2021;44:466–8.
- [55] Jenei T, Jenei S, Tamás LT, Putics Á, Knausz M, Hegedűs I, et al. COVID-19 mortality is associated with low vitamin D levels in patients with risk factors and/or advanced age. *Clinical nutrition ESPEN* 2022;47:410–3.
- [56] Cereda E, Bogliolo L, Klersy C, Lobascio F, Masi S, Crotti S, et al. Vitamin D 25OH deficiency in COVID-19 patients admitted to a tertiary referral hospital. *Clin Nutr* 2021;40(4):2469–72.
- [57] Notz Q, Herrmann J, Schlesinger T, Kranke P, Sitter M, Helmer P, et al. Vitamin D deficiency in critically ill COVID-19 ARDS patients. *Edinburgh, Scotland: Clinical nutrition*; 2021.
- [58] Fromont J, Gette M, Ben Lassoued A, Guéant JL, Guéant-Rodriguez RM, Guieu R. Hypozincemia in the early stage of COVID-19 is associated with an increased risk of severe COVID-19. *Edinburgh, Scotland: Clinical nutrition*; 2021.
- [59] Lv Y, Chen L, Liang X, Liu X, Gao M, Wang Q, et al. Association between iron status and the risk of adverse outcomes in COVID-19. *Clin Nutr* 2021;40(5):3462–9.
- [60] Reddy K, Rogers AJ, McAuley DF. Delving beneath the surface of hyperinflammation in COVID-19. *The Lancet Rheumatology* 2020;2(10):e578–9.
- [61] Fiorindi C, Campani F, Rasero L, Campani C, Livi L, Giovannoni L, et al. Prevalence of nutritional risk and malnutrition during and after hospitalization for COVID-19 infection: preliminary results of a single-centre experience. *Clinical nutrition ESPEN* 2021;45:351–5.
- [62] Gobbi M, Brunani A, Arregghini M, Baccalaro G, Dellepiane D, La Vela V, et al. Nutritional status in post SARS-CoV2 rehabilitation patients. *Edinburgh, Scotland: Clinical nutrition*; 2021.
- [63] Haraj NE, El Aziz S, Chadli A, Dafir A, Mjabber A, Aissaoui O, et al. Nutritional status assessment in patients with Covid-19 after discharge from the intensive care unit. *Clinical nutrition ESPEN* 2021;41:423–8.
- [64] Martín-Martínez A, Ortega O, Viñas P, Arreola V, Nascimento W, Costa A, et al. COVID-19 is associated with oropharyngeal dysphagia and malnutrition in hospitalized patients during the spring 2020 wave of the pandemic. *Edinburgh, Scotland: Clinical nutrition*; 2021.
- [65] Ramos A, Joaquín C, Ros M, Martín M, Cachero M, Sospedra M, et al. Impact of COVID-19 on nutritional status during the first wave of the pandemic. *Edinburgh, Scotland: Clinical nutrition*; 2021.
- [66] Wierdsma NJ, Kruizenga HM, Konings LA, Krebbers D, Jorissen JR, Joosten MI, et al. Poor nutritional status, risk of sarcopenia and nutrition related complaints are prevalent in COVID-19 patients during and after hospital admission. *Clinical nutrition ESPEN* 2021;43:369–76.
- [67] Cuerda C, Sánchez López I, Gil Martínez C, Merino Viveros M, Velasco C, Cevallos Peñafiel V, et al. Impact of COVID-19 in nutritional and functional status of survivors admitted in intensive care units during the first outbreak. Preliminary results of the NUTRICOVID study. *Edinburgh, Scotland: Clinical nutrition*; 2021.
- [68] Rives-Lange C, Zimmer A, Merazka A, Carette C, Martins-Bexinga A, Hauw-Berlemont C, et al. Evolution of the nutritional status of COVID-19 critically-ill patients: a prospective observational study from ICU admission to three months after ICU discharge. *Edinburgh, Scotland: Clinical nutrition*; 2021.
- [69] Di Filippo L, De Lorenzo R, D'Amico M, Sofia V, Roveri L, Mele R, et al. COVID-19 is associated with clinically significant weight loss and risk of malnutrition, independent of hospitalisation: a post-hoc analysis of a prospective cohort study. *Clin Nutr* 2021;40(4):2420–6.
- [70] Kellnar A, Hoppe JM, Brunner S, Stremmel C. Hospitalization for COVID-19 is associated with significant changes in body composition. *Clinical nutrition ESPEN* 2021;45:499–502.
- [71] Cereda E, Guzzardella A, Klersy C, Belliati M, Pellegrini A, Sciutti F, et al. Early caloric deficit is associated with a higher risk of death in invasive ventilated COVID-19 patients. *Edinburgh, Scotland: Clinical nutrition*; 2021.
- [72] Bordejé Laguna L, Marcos-Neira P, de Lagrán Zurbano IM, Marco EM, Guisasaola CP, Viñas Soria CD, et al. Dysphagia and mechanical ventilation in SARS-COV-2 pneumonia: it's real. *Edinburgh, Scotland: Clinical nutrition*; 2021.
- [73] Karayiannis D, Maragkouti A, Mikropoulos T, Sarri A, Kanavou A, Katsagoni C, et al. Neuromuscular blockade administration is associated with altered energy expenditure in critically ill intubated patients with COVID-19. *Edinburgh, Scotland: Clinical nutrition*; 2021.
- [74] Lakenman PLM, van der Hoven B, Schuijs JM, Eveleens RD, van Bommel J, Olieman JF, et al. Energy expenditure and feeding practices and tolerance during the acute and late phase of critically ill COVID-19 patients. *Clinical nutrition ESPEN* 2021;43:383–9.
- [75] Niederer LE, Miller H, Haines KL, Molinger J, Whittle J, MacLeod DB, et al. Prolonged progressive hypermetabolism during COVID-19 hospitalization undetected by common predictive energy equations. *Clinical nutrition ESPEN* 2021;45:341–50.
- [76] von Renesse J, von Bonin S, Held HC, Schneider R, Seifert AM, Seifert L, et al. Energy requirements of long-term ventilated COVID-19 patients with resolved SARS-CoV-2 infection. *Clinical nutrition ESPEN* 2021;44:211–7.
- [77] Bertoli S, Leone A, De Amicis R, Foppiani A, Osio D, Battezzati A. Effects of wearing a FFP2 mask on indirect calorimetry measurements: a pilot study. *Clinical nutrition ESPEN* 2021;41:443–6.
- [78] Foldi M, Farkas N, Kiss S, Dembrowsky F, Szakacs Z, Balasko M, et al. Visceral adiposity elevates the risk of critical condition in COVID-19: a systematic review and meta-analysis. *Obesity* 2021;29(3):521–8.
- [79] Huang Y, Lu Y, Huang YM, Wang M, Ling W, Sui Y, et al. Obesity in patients with COVID-19: a systematic review and meta-analysis. *Metab Clin Exp* 2020;113:154378.
- [80] Muscogiuri G, Pugliese G, Barrea L, Savastano S, Colao A. Commentary: obesity: the "achilles heel" for COVID-19? *Metabolism. clinical and experimental* 2020;108:154251.
- [81] Sharma A, Garg A, Rout A, Lavie CJ. Association of obesity with more critical illness in COVID-19. *Mayo Clin Proc* 2020;95(9):2040–2.
- [82] Simonnet A, Chetboun M, Poissy J, Raverdy V, Noulette J, Duhamel A, et al. High prevalence of obesity in severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) requiring invasive mechanical ventilation. *Obesity* 2020;28(7):1195–9.
- [83] Singh S, Bilal M, Pakhchanian H, Raiker R, Kochhar GS, Thompson CC. Impact of obesity on outcomes of patients with coronavirus disease 2019 in the United States: a multicenter electronic health records network study. *Gastroenterology* 2020;159(6):2221–2225 e6.
- [84] Collard D, Nurmohamed NS, Kaiser Y, Reeskamp LF, Dormans T, Moeniralam H, et al. Cardiovascular risk factors and COVID-19 outcomes in hospitalised patients: a prospective cohort study. *BMJ Open* 2021;11(2):e045482.
- [85] Coppelli A, Giannarelli R, Aragona M, Penno G, Falcone M, Tiseo G, et al. Hyperglycemia at hospital admission is associated with severity of the prognosis in patients hospitalized for COVID-19: the pisa COVID-19 study. *Diabetes Care* 2020;43(10):2345–8.
- [86] Racine-Brzostek SE, Yang HS, Jack GA, Chen Z, Chaddburn A, Ketts TJ, et al. Post-bronchovaccinal SARS-CoV-2 IgG and neutralizing antibodies are elevated in individuals with poor metabolic health. *J Clin Endocrinol Metab* 2021.
- [87] Huizinga GP, Singer BH, Singer K. The collision of meta-inflammation and SARS-CoV-2 pandemic infection. *Endocrinology* 2020;161(11).
- [88] Barazzoni R, Bischoff SC, Boirie Y, Busetto L, Cederholm T, Dicker D, et al. Sarcopenic obesity: time to meet the challenge. *Clin Nutr* 2018;37(6 Pt A):1787–93.
- [89] Guillet C, Delcourt I, Rance M, Giraudet C, Walrand S, Bedu M, et al. Changes in basal and insulin and amino acid response of whole body and skeletal muscle proteins in obese men. *J Clin Endocrinol Metab* 2009;94(8):3044–50.
- [90] Guillet C, Masgrau A, Walrand S, Boirie Y. Impaired protein metabolism: interlinks between obesity, insulin resistance and inflammation. *Obes Rev* : an official journal of the International Association for the Study of Obesity 2012;13(Suppl 2):51–7.
- [91] Tardif N, Salles J, Guillet C, Tordjman J, Reggion S, Landrier JF, et al. Muscle ectopic fat deposition contributes to anabolic resistance in obese sarcopenic old rats through eIF2alpha activation. *Ageing Cell* 2014;13(6):1001–11.
- [92] Wall BT, Cruz AM, Otten B, Dunlop MV, Fulford J, Porter C, et al. The impact of disuse and high-fat overfeeding on forearm muscle amino acid metabolism in humans. *J Clin Endocrinol Metab* 2020;105(7).
- [93] Martín-Del-Campo F, Ruvalcaba-Contreras N, Velázquez-Vidaurre AL, Cueto-Manzano AM, Rojas-Campos E, Cortés-Sanabria L, et al. Morbid obesity is associated with mortality and acute kidney injury in hospitalized patients with COVID-19. *Clinical nutrition ESPEN* 2021;45:200–5.
- [94] Pinhel MAS, Watanabe LM, Noronha NY, Junior WS, Nonino CB. The intersection between COVID-19 and obesity in the context of an emerging country. *Clinical nutrition ESPEN* 2021;44:472–4.
- [95] Bischoff SC, Boirie Y, Cederholm T, Chourdakis M, Cuerda C, Delzenne NM, et al. Towards a multidisciplinary approach to understand and manage obesity and related diseases. *Clin Nutr* 2017;36(4):917–38.
- [96] Atkins JL, Wannamethee SG. Sarcopenic obesity in ageing: cardiovascular outcomes and mortality. *Br J Nutr* 2020;124(10):1102–13.
- [97] Montano-Loza AJ, Angulo P, Meza-Junco J, Prado CM, Sawyer MB, Beaumont C, et al. Sarcopenic obesity and myosteatosis are associated with higher mortality in patients with cirrhosis. *Journal of cachexia, sarcopenia and muscle* 2016;7(2):126–35.
- [98] Van Aller C, Lara J, Stephan BCM, Donini LM, Heymsfield S, Katzmarzyk PT, et al. Sarcopenic obesity and overall mortality: results from the application of novel models of body composition phenotypes to the National Health and Nutrition Examination Survey 1999–2004. *Clin Nutr* 2019;38(1):264–70.
- [99] Zhang X, Xie X, Dou Q, Liu C, Zhang W, Yang Y, et al. Association of sarcopenic obesity with the risk of all-cause mortality among adults over a broad range of different settings: a updated meta-analysis. *BMC Geriatr* 2019;19(1):183.
- [100] Fedele D, De Francesco A, Riso S, Collo A. Obesity, malnutrition, and trace element deficiency in the coronavirus disease (COVID-19) pandemic: an overview. *Nutrition* 2021;81:111016.
- [101] Sahin E, Orhan C, Uckun FM, Sahin K. Clinical impact potential of supplemental nutrients as adjuncts of therapy in high-risk COVID-19 for obese patients. *Front Nutr* 2020;7:580504.
- [102] Alcántara-Alonso E, Molinar-Ramos F, González-López JA, Alcántara-Alonso V, Muñoz-Pérez MA, Lozano-Nuevo JJ, et al. High triglyceride to HDL-cholesterol ratio as a biochemical marker of severe outcomes in COVID-19 patients. *Clinical nutrition ESPEN* 2021;44:437–44.
- [103] Nicolau J, Ayala L, Sanchis P, Rodríguez I, Romano A, Dotres K, et al. The coexistence of low albumin levels and obesity worsens clinical outcomes among

- subjects admitted for sars-cov-2 infection. *Clinical nutrition ESPEN* 2021;46:434–8.
- [104] Soetedjo NNM, Iryaningrum MR, Damara FA, Permadhi I, Sutanto LB, Hartono H, et al. Prognostic properties of hypoalbuminemia in COVID-19 patients: a systematic review and diagnostic meta-analysis. *Clinical nutrition ESPEN* 2021;45:120–6.
- [105] Pranata R, Lim MA, Huang I, Yonas E, Henrina J, Vania R, et al. Visceral adiposity, subcutaneous adiposity, and severe coronavirus disease-2019 (COVID-19): systematic review and meta-analysis. *Clinical nutrition ESPEN* 2021;43:163–8.
- [106] Alves T, Guimarães RS, Souza SF, Brandão NA, Daltro C, Conceição-Machado MEP, et al. Influence of nutritional assistance on mortality by COVID-19 in critically ill patients. *Clinical nutrition ESPEN* 2021;44:469–71.
- [107] Silvah JH, de Lima CMM, Nicoletti CF, Barbosa AC, Junqueira GP, da Cunha SFC, et al. Protein provision and lower mortality in critically ill patients with COVID-19. *Clinical nutrition ESPEN* 2021;45:507–10.
- [108] Chada RR, Chidrawar S, Siddiqua A, Medanki R, Omer SA, Nagalla B. Tailoring nutrition therapy amid the COVID-19 pandemic: does it work? *Clinical nutrition ESPEN* 2021;45:381–8.
- [109] Liu R, Paz M, Siraj L, Boyd T, Salamone S, Lite TV, et al. Feeding intolerance in critically ill patients with COVID-19. Edinburgh, Scotland: *Clinical nutrition*; 2021.
- [110] Subramanian K, Solomon N, Faillace R, Menon V, Raiszadeh F, Brandeis G. Effect of parenteral nutrition in oxygen escalation/de-escalation in SARS-CoV-2 infected patients who are pre-intubation: a multicenter, observational study. *Clinical nutrition ESPEN* 2021;46:206–9.
- [111] Feng Z, Yang J, Xu M, Lin R, Yang H, Lai L, et al. Dietary supplements and herbal medicine for COVID-19: a systematic review of randomized control trials. *Clinical nutrition ESPEN* 2021;44:50–60.
- [112] Beran A, Mhanna M, Srour O, Ayesh H, Stewart JM, Hjouj M, et al. Clinical significance of micronutrient supplements in patients with coronavirus disease 2019: a comprehensive systematic review and meta-analysis. *Clinical nutrition ESPEN* 2022;48:167–77.
- [113] Rossato MS, Brilli E, Ferri N, Giordano G, Tarantino G. Observational study on the benefit of a nutritional supplement, supporting immune function and energy metabolism, on chronic fatigue associated with the SARS-CoV-2 post-infection progress. *Clinical nutrition ESPEN* 2021;46:510–8.
- [114] Marino LV, Valla FV, Tume LN, Jotterand-Chaparro C, Moulet C, Latten L, et al. Considerations for nutrition support in critically ill children with COVID-19 and paediatric inflammatory multisystem syndrome temporally associated with COVID-19. *Clin Nutr* 2021;40(3):895–900.
- [115] Karakaya Molla G, Ünal Uzun Ö, Koç N, Özen Yeşil B, Bayhan G. Evaluation of nutritional status in pediatric patients diagnosed with Covid-19 infection. *Clinical nutrition ESPEN* 2021;44:424–8.
- [116] Cuerda C, Muscaritoli M, Donini LM, Baque P, Barazzoni R, Gaudio E, et al. Nutrition education in medical schools (NEMS). An ESPEN position paper. *Clin Nutr* 2020;39(9):2938–9.
- [117] Cuerda C, Muscaritoli M, Donini LM, Baque P, Barazzoni R, Gaudio E, et al. Nutrition education in medical schools (NEMS). An ESPEN position paper. *Clin Nutr* 2019;38(3):969–74.
- [118] Cuerda C, Muscaritoli M, Krznaric Z, Pirlich M, Van Gossum A, Schneider S, et al. Nutrition education in medical schools (NEMS) project: joining ESPEN and university point of view. *Clin Nutr* 2021;40(5):2754–61.
- [119] Cardenas D, Correia M, Ochoa JB, Hardy G, Rodríguez-Ventimilla D, Bermudez CE, et al. Clinical nutrition and human rights. An international position paper. *Clin Nutr* 2021;40(6):4029–36.
- [120] Cardenas D, Bermudez C, Barazzoni R. The Cartagena Declaration: a call for global commitment to fight for the right to nutritional care. *Clin Nutr* 2019;38(5):2458–9.
- [121] Ho DKN, Nguyen HS, Irnandi DF, Faradina A, Dang TD, Wiratama BS, et al. Adherence to COVID-19 nutritional guidelines and their impact on the clinical outcomes of hospitalized COVID-19 patients. *Clinical nutrition ESPEN* 2021;46:491–8.
- [122] de Watteville A, Genton L, Barcelos GK, Pugin J, Pichard C, Heidegger CP. Easy-to-prescribe nutrition support in the intensive care in the era of COVID-19. *Clinical nutrition ESPEN* 2020;39:74–8.
- [123] Miguélez M, Velasco C, Cambor M, Cedeño J, Serrano C, Bretón I, et al. Nutritional management and clinical outcome of critically ill patients with COVID-19: a retrospective study in a tertiary hospital. Edinburgh, Scotland: *Clinical nutrition*; 2021.
- [124] Nakamura K, Liu K, Katsukawa H, Nydahl P, Ely EW, Kudchadkar SR, et al. Nutrition therapy in the intensive care unit during the COVID-19 pandemic: findings from the ISIC point prevalence study. Edinburgh, Scotland: *Clinical nutrition*; 2021.
- [125] Osuna-Padilla I, Rodríguez-Moguel NC, Aguilar-Vargas A, Rodríguez-Llamazares S. Safety and tolerance of enteral nutrition in COVID-19 critically ill patients, a retrospective study. *Clinical nutrition ESPEN* 2021;43:495–500.
- [126] Salazar E, Cheah MCC. Prescribing parenteral nutrition in acute hospital setting during COVID-19: the Singapore experience. *Clin Nutr* 2020;39(7):2321.
- [127] de Watteville A, Montalbano F, Wozniak H, Collet TH, Jaksic C, Le Terrier C, et al. Impact of nutritional therapy during the first wave of the COVID-19 pandemic in intensive care patients: a retrospective observational study. Edinburgh, Scotland: *Clinical nutrition*; 2021.
- [128] Turri Quarenghi R, Bertolotti D, Gavazzoni M, Ossola N, Quarenghi M. The reorganisation of nutritional assessment and care in a hospital that became a COVID centre during the COVID-19 pandemic. *Clinical nutrition ESPEN* 2021;46:142–6.
- [129] Viana Bagni U, da Silva Ribeiro KD, Soares Bezerra D, Cavalcante de Barros D, de Magalhães Fittipaldi AL, Pimenta da Silva Araújo RG, et al. Anthropometric assessment in ambulatory nutrition amid the COVID-19 pandemic: possibilities for the remote and in-person care. *Clinical nutrition ESPEN* 2021;41:186–92.
- [130] Vivanti A. Pilot investigation shows high sensitivity and specificity using an Abridged Subjective Global Assessment without physical examination in a tertiary hospital; pertinence for use amongst those without COVID-19 when physical distancing required during the pandemic. *Clinical nutrition ESPEN* 2021;44:463–5.
- [131] Ž Krznarić, Bender DV, Laviano A, Cuerda C, Landi F, Monteiro R, et al. A simple remote nutritional screening tool and practical guidance for nutritional care in primary practice during the COVID-19 pandemic. *Clin Nutr* 2020;39(7):1983–7.
- [132] Lai S, Van Gossum A, Joly F, Bozzetti F, Cuerda C, Lamprecht G, et al. Considerations for the management of home parenteral nutrition during the SARS-CoV-2 pandemic: a position paper from the home artificial nutrition and chronic intestinal failure special interest group of ESPEN. *Clin Nutr* 2020;39(7):1988–91.
- [133] Matras P, Klek S, Folwarski M, Zmarzly A, Bartoszewski L, Cebulski W, et al. Home medical nutrition during SARS-CoV-2 pandemic – a position paper. *Clinical nutrition ESPEN* 2020;38:196–200.
- [134] Herle M, Smith AD, Bu F, Steptoe A, Fancourt D. Trajectories of eating behavior during COVID-19 lockdown: longitudinal analyses of 22,374 adults. *Clinical nutrition ESPEN* 2021;42:158–65.
- [135] Al-Domi H, Al-Dalaeen A, Al-Rosan S, Batarseh N, Nawaiseh H. Healthy nutritional behavior during COVID-19 lockdown: a cross-sectional study. *Clinical nutrition ESPEN* 2021;42:132–7.
- [136] Moradi F, Heshmati J, Daneshzad E, Ahmadi A, Jafari T, Persad E, et al. Association between dietary satisfaction and depression, anxiety and stress in obese and overweight patients during the coronavirus pandemic. *Clinical nutrition ESPEN* 2021;45:399–403.
- [137] Madalı B, Alkan Ş B, Örs ED, Ayrançı M, Taşkın H, Kara HH. Emotional eating behaviors during the COVID-19 pandemic: a cross-sectional study. *Clinical nutrition ESPEN* 2021;46:264–70.
- [138] Bakaloudi DR, Jeyakumar DT, Jayawardena R, Chourdakis M. The impact of COVID-19 lockdown on snacking habits, fast-food and alcohol consumption: a systematic review of the evidence. Edinburgh, Scotland: *Clinical nutrition*; 2021.
- [139] Coletro HN, Mendonça RD, Meireles AL, Machado-Coelho GLL, Menezes MC. Ultra-processed and fresh food consumption and symptoms of anxiety and depression during the COVID - 19 pandemic: COVID Inconfidentes. *Clinical nutrition ESPEN* 2022;47:206–14.
- [140] Bakaloudi DR, Barazzoni R, Bischoff SC, Breda J, Wickramasinghe K, Chourdakis M. Impact of the first COVID-19 lockdown on body weight: a combined systematic review and a meta-analysis. Edinburgh, Scotland: *Clinical nutrition*; 2021.
- [141] Di Santo SG, Franchini F, Filippini B, Martone A, Sannino S. The effects of COVID-19 and quarantine measures on the lifestyles and mental health of people over 60 at increased risk of dementia. *Front Psychiatr* 2020;11:578628.
- [142] Stival C, Lugo A, Bosetti C, Amerio A, Serafini G, Cavaliere d'Oro L, et al. COVID-19 confinement impact on weight gain and physical activity in the older adult population: data from the LOST in Lombardia study. *Clinical nutrition ESPEN* 2022;48:329–35.
- [143] Donini LM, Busetto L, Bischoff SC, Cederholm T, Ballesteros-Pomar MD, Batsis JA, et al. Definition and diagnostic criteria for sarcopenic obesity: ESPEN and EASO consensus statement. *Clin Nutr* 2022;41(4):990–1000.
- [144] Hamid A, Dos Santos C, da Silva Schlickmann D, Gama da Silva T, Molz P, Brand C, et al. Intestinal transit rhythm and associated factors during the COVID-19 pandemic: a pilot study. *Clinical nutrition ESPEN* 2022;48:220–6.
- [145] Khraisat BR, Al-Jeady AM, Alqatawneh DA, Toubasi AA, AlRyalat SA. The prevalence of mental health outcomes among eating disorder patients during the COVID-19 pandemic: a meta-analysis. *Clinical nutrition ESPEN* 2022;48:141–7.
- [146] Baquerizo-Sedano L, Chaquila JA, Aguilar L, Ordóñez JM, González-Muniesa P, Garaulet M. Anti-COVID-19 measures threaten our healthy body weight: changes in sleep and external synchronizers of circadian clocks during confinement. Edinburgh, Scotland: *Clinical nutrition*; 2021.
- [147] Cobre AF, Surek M, Vilhena RO, Böger B, Fachi MM, Momade DR, et al. Influence of foods and nutrients on COVID-19 recovery: a multivariate analysis of data from 170 countries using a generalized linear model. Edinburgh, Scotland: *Clinical nutrition*; 2021.
- [148] Baik I. Region-specific COVID-19 risk scores and nutritional status of a high-risk population based on individual vulnerability assessment in the national survey data. Edinburgh, Scotland: *Clinical nutrition*; 2021.
- [149] Perez-Araluce R, Martinez-Gonzalez MA, Fernández-Lázaro CI, Bes-Rastrollo M, Gea A, Carlos S. Mediterranean diet and the risk of COVID-19 in the 'Seguimiento Universidad de Navarra' cohort. Edinburgh, Scotland: *Clinical nutrition*; 2021.

- [150] Elshahry N, Al-Sayyed H, Odeh M, McGrattan A, Hammad F. Effect of Covid-19 on food security: a cross-sectional survey. *Clinical nutrition ESPEN* 2020;40:171–8.
- [151] Piccioni F, Goulao LF, Roberfroid D. The impact of COVID-19 on diet quality, food security and nutrition in low and middle income countries: a systematic review of the evidence. Edinburgh, Scotland: *Clinical nutrition*; 2021.
- [152] Kabir M, Saqib MAN, Zaid M, Ahmed H, Afzal MS. COVID-19, economic impact and child mortality: a global concern. *Clin Nutr* 2020;39(7):2322–3.
- [153] Amataiti TA, Hood F, Krebs JD, Weatherall M, Hall RM. The Impact of COVID-19 on diet and lifestyle behaviours for pregnant women with diabetes. *Clinical nutrition ESPEN* 2021;45:404–11.

Rocco Barazzoni*

Department of Medical, Surgical and Health Sciences, University of Trieste, Italy

Joao Breda

World Health Organization (WHO) Regional Office for Europe, Athens, Greece

Cristina Cuerda

Departamento de Medicina, Universidad Complutense de Madrid, Nutrition Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain

Stephane Schneider
Gastroenterology and Nutrition, Nice University Hospital, Université Côte d'Azur, Nice, France

Nicolaas E. Deutz
Center for Translational Research in Aging and Longevity, Texas A&M University, College Station, TX, USA

Kremlin Wickramasinghe
World Health Organization European Office for the Prevention and Control of Noncommunicable Diseases, Moscow, Russian Federation

the COVID-19 Call Editorial Board

* Corresponding author. Department of Medical, Surgical and Health Sciences, Cattinara University Hospital, Strada di Fiume 447, 34149 Trieste, Italy.
E-mail address: barazzon@units.it (R. Barazzoni).

20 July 2022