

Assessment of cognition in aphasia: Perspectives from clinicians and researchers

Bruna Tessaro^{1,2,3}, Sonia L.E. Brownsett^{4,5,6,7}, Solène Hameau^{1,8}, Tijana Simic^{6,9,10}, Lyndsey A. Nickels¹, Natalie Gilmore^{6,11}, Claudia Peñaloza^{6,12,13,14}, Holly Robson^{6,15}, Christos Salis^{2,3}.

¹ School of Psychological Sciences, Macquarie University, Sydney, Australia

² Speech & Language Sciences, Newcastle University, Newcastle upon Tyne, UK

³ International Doctorate for Experimental Approaches to Language & Brain (IDEALAB), Newcastle University (UK), Macquarie University (AU), Potsdam University (DE), Groningen University (NL)

⁴ Queensland Aphasia Research Centre, School of Health & Rehabilitation Sciences, University of Queensland, Brisbane, Australia

⁵ NHMRC Centre for research excellence in Aphasia Recovery & Rehabilitation, La Trobe University, Melbourne, Australia

⁶ Collaboration of Aphasia Trialists

⁷ STARS Education and Research Alliance, Surgical Treatment and Rehabilitation Service (STARS), The University of Queensland and Metro North Health, Brisbane, Queensland, Australia.

⁸ Psychological Sciences Research Institute and Institute of Neuroscience, Université Catholique de Louvain, Louvain-la-Neuve, Belgium

⁹ Department of Speech-Language Pathology, University of Toronto, Toronto, Canada

¹⁰ KITE-Toronto Rehabilitation Institute, University Health Network, Toronto, Canada

¹¹ Laboratory of NeuroImaging for Coma & Consciousness, Department of Neurology, Massachusetts General Hospital, Boston, USA

¹² Department of Cognition, Development & Educational Psychology, University of Barcelona, Barcelona, Spain

¹³ Institute of Neurosciences, University of Barcelona, Barcelona, Spain

¹⁴ Cognition and Brain Plasticity Unit, Bellvitge Biomedical Research Institute-IDIBELL, L'Hospitalet de Llobregat, Barcelona, Spain

¹⁵ Division of Psychology & Language Sciences, University College London, London, UK

Corresponding author: Dr Sonia Brownsett s.brownsett@uq.edu.au

Funding sources

This work was supported by the NHMRC-funded Centre of Research Excellence in Aphasia Recovery and Rehabilitation (#1153236). BT was supported by an International Macquarie University Research Excellence Scholarship (iMQRES, Allocation number 20191485) and a Newcastle University Postgraduate Scholarship. Claudia Peñaloza is supported by grant RYC2021-034561-I funded by MCIN/AEI/10.13039/501100011033 and the European Union NextGenerationEU/PRTR. We would like to acknowledge the Collaboration of Aphasia Trialists (IS1208) EU Cooperation in Science and Technology and The Tavistock Trust for Aphasia provided important support to develop this research.

Word count (including introduction, methods, results and discussion): 9542

Word count (including abstract, introduction, methods, results and discussion, tables and figures and references): 13,717

Pages: 52

Tables:3

Figures:5

Supplementary material: 9 pages

Abstract

People with aphasia may present with cognitive impairments beyond language. In this population, reliance on language-based assessments of cognition may lead to unreliable metrics of cognitive performance across clinical and research settings. We investigated the underlying philosophy and practice of assessing cognition in people with aphasia.

Methods: An e-survey was developed for multidisciplinary clinicians and researchers. Snowball sampling was used to recruit international participants. The e-survey comprised 29 items (a mix of multiple-choice and open-ended items) addressing definitions of cognition, assessment of cognition, tools used to assess cognition and participant demographics. Data were analysed using descriptive statistics and thematic analysis.

Results: 291 respondents participated from a range of disciplines and countries. Over 80% of respondents considered it important to assess attention, executive functions, learning and memory. The main barrier to assessment was the lack of appropriate tools available for people with aphasia. Responses indicated that whilst professionals felt that understanding the interaction between language and cognition in aphasia was important for providing optimal care.

Discussion: This study highlights the need for better awareness and training in the assessment of cognition in people with aphasia, and for psychometrically robust assessments, appropriate for the assessment of cognition in the presence of aphasia.

Key words: cognition, aphasia, assessment, survey, multidisciplinary

Introduction

In addition to impaired linguistic abilities, people with aphasia may present with non-linguistic cognitive difficulties, such as those affecting memory, attention or executive functioning (Fucetola et al., 2009; Gilmore et al., 2019; Mayer & Murray, 2012; Murray, 2012). As such, comprehensive cognitive assessment in people with aphasia is recommended in research settings (Murray, 2012; Murray, 2017; Kuzmina & Weekes, 2017). The need for such testing, either directly or indirectly, is also included in the most recent clinical guidelines for healthcare professionals' (e.g., American Speech-Language Hearing Association, 2024; Australian Aphasia Rehabilitation Pathway, 2024; Australian and New Zealand Clinical Guidelines for Stroke Management, 2024; Canadian Stroke Best Practice, 2020; National Clinical Guideline for Stroke for the UK and Ireland, 2023). Disentangling linguistic and other cognitive impairments and residual abilities is essential for understanding the extent of the non-linguistic cognitive difficulties people with aphasia may face, how these interact with their linguistic impairment, and how they might impact intervention planning and prognosis (Brownsett et al., 2014; Diedrichs et al., 2022; Fonseca et al., 2019; Simic et al., 2020). The influence of non-linguistic cognitive ability on a person's response to therapy is also not understood. Whilst some studies have reported an absence of a relationship between residual cognitive abilities and therapy response (e.g., executive control: van de Sandt-Koenderman et al., 2007; for a review see Diedrichs et al., 2022), others have drawn attention to the relevance of cognitive abilities, beyond language, in aphasia research and clinical practice (Brownsett et al., 2014; Lacroix et al., 2021; Schumacher et al., 2019; Varkanitsa et al., 2023), and have demonstrated that better cognitive abilities are associated with a greater response to therapy (e.g., executive control: Simic et al., 2020).

Despite the importance of understanding the role of non-linguistic cognitive abilities in aphasia, aphasia research does not consistently test or report on cognitive abilities. Indeed, our review of published data between 2010-2020 showed that nearly half (47.5%) of studies in the aphasia literature did not report assessments of cognition (Tessaro et al., 2022). Additionally, despite aphasia care typically involving a range of healthcare professionals with varying skills and experience in neuropsychological testing (Block et al., 2017), the extent to which cognitive assessment is carried out in clinical settings, and by whom, is unclear.

There are a few reasons why the reporting of cognitive testing and abilities might be limited. First, clinicians may be unsure which non-linguistic abilities are important to assess in aphasia. For example, there is little consensus regarding which cognitive functions should be evaluated when establishing decision making capacity in aphasia (see Kim et al., 2020). Clinicians

and researchers may also lack clear guidelines on the most suitable assessment tools available (Fonseca et al., 2016; Simic et al., 2019), and who, within the multidisciplinary team, should be responsible for conducting and interpreting these assessments.

If clinicians and researchers are able to reach consensus on how best to assess cognition in people living with aphasia, this will guide clinical decision-making and management to ensure appropriate diagnosis, and optimised management. In addition, it will support researchers working across these domains to better understand the concomitant contribution of different impairments, how these might interact, differential diagnosis, and recovery and rehabilitation. Without a clear understanding of how cognition and language interact, and dissociate, clinical and research disciplines are unable to tease out the different essential ingredients of therapeutic targets. If consensus is unable to be obtained using existing tools, this will serve as an important motivator for the research and clinical communities to collaboratively develop an optimised tool, or series of tools, that specifically address this clinical problem.

The present study aimed to identify current practice, including clinical and research requirements, relating to assessment of cognition in aphasia. This was achieved by surveying healthcare professionals and researchers from diverse disciplines on their philosophy and practice of assessing cognition in people with aphasia. To establish a clear working definition of cognition in this context, we first surveyed respondents regarding their understanding of cognition. We then probed their opinions about the importance of assessing cognition in aphasia, and which cognitive constructs they felt needed to be assessed. Furthermore, given the lack of consensus as to which assessment tools should be used (Fonseca et al., 2016), we probed which tools were being used, how they were selected, what aspects of cognition they aimed to assess, and any barriers faced when assessing cognition in aphasia. Finally, we also asked respondents' opinions about whose role it is to assess cognition and convey assessment results to people with aphasia and their carers and/or family members.

Materials and Methods

This research was approved by the Research Ethics Committee of Newcastle University, School of Education, Communication & Language Sciences. The study used a cross-sectional online survey design. To ensure best practice in survey methodology, the Checklist for Reporting Results of Internet E-Surveys (CHERRIES) was used as a guideline (Eysenbach, 2004). LimeSurvey, a free and open-source on-line survey tool, was used to design and host the survey. The survey was open, voluntary and accessible via a web link. The survey began with a brief introduction of its purpose. Respondents could read the full participant information sheet by

clicking on a hyperlink that opened a PDF containing detailed information about the study, the investigators, the purpose of the study and expected duration of the study. Consent was provided by informing participants that continuing with the survey (i.e., clicking “next”) they agreed to take part. Responses were timed and anonymised (responses did not have the IP address logged), and participants could save a partially finished survey and return to it at a later date. Participants were not able to go back to earlier pages and responding to all questions was not mandatory. All data were collected from April to July 2021. No incentives were offered to complete the e-survey.

Survey Structure and Content

The survey comprised 29 questions organised into four sections: Section 1 - definitions of cognition in aphasia (2 questions), Section 2 - assessment of cognition in aphasia (10 questions), Section 3 - tools to assess cognition in aphasia (7 questions), and Section 4 - participant demographics (10 questions) (for specific questions see supplementary table 1) . Multiple choice, yes/no, open-ended (free text), and Likert scales response types were used. Following the introduction and consent page, the survey contained 5 pages with all questions from the same section on the same page, except that the section “Definition of cognition in aphasia” was split over two pages.

Given ambiguity around the extent to which language is considered a component of cognition, with language often excluded from the operationalisation of cognition, we wanted to understand respondents’ definitions of cognition, prior to asking more specific questions about the assessment of cognition. Therefore, the first section included two questions regarding the definition of cognition in aphasia. An open-ended question ‘What does the term cognition mean to you?’ (Question 1) was first asked, then a second question asked respondents to indicate, from a list provided, the domains that they considered fell under their definition of cognition (Question 2).

The second section included ten questions that aimed to gather respondents’ opinions about assessment practices in the context of aphasia. Six questions required forced choice answers or ratings, and four questions free text responses. Questions 3, 6, 7 required respondents to select options from a list provided, with question 3a, 4a, and 5a providing an opportunity to expand using free text responses. Questions 4 and 5 required a Likert scale response, asking to rate the importance of assessment of cognition, and the domains of cognition that should be assessed respectively. Question 8 required yes/no responses, and 8a asked respondents to ‘list in rank order list any barriers you have experienced find in assessing cognition in aphasia’. If respondents

indicated in Section 2 that they never assessed cognition in aphasia, then they were instructed to skip Section 3 and move to Section 4.

The third section included seven questions regarding the tools used by clinicians and researchers to assess cognition in aphasia (free text Questions 9 and 10), what the difficulties were (if any) in this assessment (Questions 10 (open-ended), 11 (yes/no), 11a (open-ended)), and if they changed their approach when assessing cognition in aphasia as compared to individuals without aphasia (Questions 12 (yes/no), 12a and 13 (open-ended)).

The fourth section comprised demographic questions about respondents' country of practice, age group, professional identity, work setting, and experience of working with people with aphasia.

Participant Recruitment

The survey was disseminated through mailing lists (aphasia associations, research groups, professional societies, university departments), newsletters, and/or posting on social media pages. A snowball sampling approach was used with e-mail invitations sent to professional contacts of the authors, asking that they both complete the survey and forward it to relevant people and/or mailing lists. To increase participation from non-English speaking countries, whilst the questionnaire was only available in English, we allowed responses in languages other than English. Responses that were not in English (i.e., French, Spanish and German) were translated by speakers of the respective languages (SH, SB, CP).

Pilot survey

The survey was developed by iterative discussion by the author team. Eight professionals working with people with aphasia were invited to complete a pilot version of the survey. As we aimed to survey a multidisciplinary team, given that the initial survey was designed by speech and language therapists, we ensured that we selected other professionals to complete the pilot. Specifically, respondents were two neuropsychologists, three occupational therapists, one physiotherapist, and two neurologists. Of these, two identified as clinicians, two as researchers and four as both clinicians and researchers. Two respondents had more than 10 years of experience working with people with aphasia and six respondents had between four to 10 years of experience. They provided feedback on the survey structure. Following their feedback, changes were made to reduce the length of the Introduction and consent page. Given that no changes were recommended to the survey items themselves, the pilot data were included in the final analysis.

Data Analyses

Survey items were analysed based on response type. Accordingly, quantitative methods were used to analyse data from multiple-choice, yes/no, and Likert scale responses (Questions 2-5, 6-9, 11-12). Qualitative methods were used to analyse data from open-ended questions (Questions 1, 3-5, 8, 11-12).

Descriptive statistics were computed in R Core Team version 4.2.1 (2022) for multiple-choice questions (i.e., Questions 2, 6-7, 9). A qualitative approach, described below, was used to analyse the open-ended question (i.e., Question 1). Mixed qualitative and quantitative analyses were performed for yes/no, multiple choice, and Likert scale questions that were combined with an open-ended question (i.e., Questions 3-5, 8, 11-12).

Qualitative Analysis

Thematic analysis (Braun et al., 2019; Braun & Clarke, 2006; Clarke & Braun, 2017) was used to analyse the text data from the survey. Codes were created by identifying and labelling discrete segments of data that captured a single idea of interest relevant to the research question (Braun et al., 2009). Themes were then created by identifying an idea or concept that summarised recurring patterns of codes across the data (Braun & Clarke, 2009).

There were three steps to this analysis:

Step 1 - identification: one author (BT) created codes based on the responses to each question and identified possible themes. Each code was revised by checking the responses assigned to it and, if needed, the name of the code was refined to better reflect the responses under that code. At this stage, some codes were either combined into a single code (if sufficiently similar to the same code) or divided into more than one code (if sufficiently different to justify the creation of one or more additional codes). Responses were re-checked against the assigned codes.

Step 2 – codebook revision: one author (SB), who also participated in the pilot stage, reviewed and revised the codebook, indicating if they agreed with the name of the codes, the code structures (relationships between higher – e.g., “Materials” in the responses to the question targeting barriers to assessing cognition in aphasia, and lower-level codes – e.g., “Access to assessments and resources” for the same question), and the examples for each code. Suggested revisions were reviewed (BT, CS, SH), via consensus, to determine which should be implemented. The data were recoded using the revised codebook.

Step 3 – interrater reliability: an interrater reliability measure to assess the rigour and transparency of the analysis was calculated following the guidelines of O’Connor and Joffe (2020).

Following initial coding (BT), second rater (TS), blinded to the initial coding, coded 25% of the dataset (25% of the responses to each question) according to the revised codebook. Interrater reliability was calculated in two ways: percentage agreement (PA), the percentage of the content where the two raters agreed on the coding, and Cohen's Kappa coefficient, a statistical measure of inter-rater reliability which considers the amount of agreement that could be expected to occur through chance. Kappa values below 0.40 indicate poor agreement, values between 0.40-0.60 indicate moderate agreement, values between 0.61-0.80 indicate substantial agreement and values over 0.81 indicate almost perfect agreement (Landis & Koch, 1977).

Reliability

The inter-rater reliability of the final coding (see Supplementary Table 3) showed very high (>97%) percentage agreement on all questions, and Cohen's Kappa co-efficient showed moderate to almost perfect agreement between raters. As such, no further coding revisions were considered necessary.

Results

504 survey attempts were made, and, of these, 205 surveys were completed, providing a completion ratio of 0.41. One attempt was excluded because the respondent identified as a person with aphasia, for a total of 204 survey respondents. The average time spent responding to the survey was 32.6 minutes (SD = 172.2, median = 12.5), including pauses.

Respondents' demographic characteristics

Most respondents (52%) were located in European countries, followed by North America (28%), Oceania (15%), Asia (3.9%), Africa (0.5%), and South America (0.5%). Most of the sample (62.6%) were aged between 18 and 44 years, with 37.4% aged 45 or older (see Supplementary Table 2).

Most respondents indicated they were Speech and Language Pathologists (SLPs) (64%), followed by neuropsychologists (13%) and occupational therapists (8.3%). About half of respondents identified as clinicians (51%), followed by those that identified as both clinician and researchers (35%), and researchers (14%).

About half the sample (53.4%) indicated that they had more than 10 years of experience working with people with aphasia, and 35.3% reported having between 4 to 10 years of experience. Most of these professionals reported working with people with aphasia three or more days per week (58%), with 7.5% reporting variability (e.g., 'depending on the research project I am engaged in'). 5% indicated they were not currently working with people living with aphasia (PWA).

Qualitative Analysis

First, 294 codes encompassed by the responses were identified and created. Next, 222 of 244 suggested revisions to the codebook reached consensus for revision and were incorporated in the revised codebook. The following sections discuss each question in turn and summarise the main themes identified from the codes, providing some illustrative responses. Quantitative and qualitative results for each survey question are summarized, in turn below.

Section 1: Defining cognition in the context of aphasia.

The first question asked respondents to define cognition. Three main themes were identified in the responses: (1) how the concept of cognition was referred to, (2) the processes that cognition allows, and (3) cognitive domains listed as part of cognition. For example:

‘The (thought processes) [1] that allow us (to understand and interact with our environment) [2].’

‘Cognition is a broad term comprising of (memory, language, attention, thinking, reasoning, executive functions) [3] which are important in daily functioning.’

Most respondents defined cognition as a set of skills, mental processes and brain functions that allow individuals to think, problem solve, learn, plus process and integrate information. They also indicated that cognition allows individuals to perceive, navigate and interact with the environment, and to reason and plan. Most responses listed domains that the respondents would include as part of cognition (e.g., memory, attention, executive functions, language).

In Question 2, respondents were given a list of several cognitive domains and were asked which fell under their definition of cognition. Most respondents indicated they would include attention, memory, executive functions, learning, processing speed, social cognition, visuo-perceptual and visuo-constructional abilities, and language (see Figure 1). Whilst only 14 people (11%) included motor function and praxis in their definition of cognition in Question 1, on this second question 160 respondents (54%) indicated that they considered praxis to be part of cognition. Specifically, 148 respondents included praxis in their multiple-choice definition of cognition, but not in their open-ended response to Question 1. Thirty-one respondents (24%) commented that constructs such as praxis and social cognition were composite abilities reliant on other abilities such as language, attention and memory.

<insert figure 1>

When asked to define cognition (Q1, 260 respondents), or comment on the language domain (Q2, optional, 35 respondents), 16% of respondents (47/295 responses) mentioned that language and cognition possibly interact. Nine respondents indicated they did not consider language to be part of cognition, and two respondents (0.7%) mentioned that they believed cognition was not dependent on language. Two further respondents commented that they were aware that language and cognition can be selectively impaired:

‘This [language] is hard to tease out from cognition, but there are people who are profoundly affected in their ability to receive and convey symbols who nevertheless are functional otherwise.’

Twenty-six respondents (9%) suggested that while both domains are inextricably linked, they may still be considered discrete functions. Finally, eight respondents shared that they considered language to be part of cognition, and how this is reflected in clinical practice:

'I find it unfortunate that clinical tools, especially assessments, often dichotomize language and cognition.'

Section 2: Current practice in the assessment of cognition in aphasia

This section comprised questions that sought to gauge current practices regarding the assessment of cognition in aphasia, focusing on the reasons that influence the decision to assess cognition in people with aphasia (Question 3, 3a, 4, 4a), which aspects of cognition should be assessed (Question 5, 5a), which professionals should be responsible for assessing cognition and discussing assessment findings (Question 6), and identifying any barriers to assessment (Question 8, 8a).

Question 3: What considerations influence your decision to assess cognition in PWA?

This question asked which factors influenced the decision to assess cognition, but as it was a multiple-choice question, respondents did not indicate how the decision was influenced. As shown in Figure 2, the greatest numbers of respondents indicated that their decision to assess cognition in aphasia was influenced by reports of cognitive deficits by the person with aphasia or their caregiver (69.9% of 266 respondents), and/or when caregivers reported concern about the cognition of people with aphasia (63.9%).

<insert figure 2>

Half the respondents indicated that age did not influence whether they would assess cognition (50.4%), with relatively small numbers indicating that older (7.52%) and younger (6.8%) age would influence their decision to assess cognition. Nearly half of respondents indicated that stroke severity influenced their decision to assess cognition, with similar numbers for severe (45%), moderate (44.7%) and mild (40.2%).

Language impairment was also considered by slightly less than half of the respondents: 45.5% responded that they assessed cognition for people with mild, 44.3% with moderate, and 43.2% with severe language impairment. Twenty percent of respondents reported that they assessed cognition in aphasia for research purposes. Only 1.1% (2 respondents) indicated that they

did not assess or refer people with aphasia for cognitive assessment, noting that they lacked training in completing cognitive assessments, and a lack of appropriate time and resources.

Question 4: How important is the assessment of cognition for specific purposes?

Respondents were asked to rate the importance of each of a list of potential reasons for assessing cognition in aphasia (Figure 3). Overall, most respondents (>60%) considered it important, or very important, to assess cognition for treatment planning and rehabilitation, to facilitate return to employment, to provide education about cognitive function and impairment, to recommend strategies for patients and/or caregivers, to determine legal capacity, and to determine the level of cognitive impairment beyond language. Assessing cognition for insurance purposes applied to the practice of about half the respondents, with 15% of these respondents considering this important, or very important.

<insert figure 3>

In response to an open-ended optional follow-up to Question 4 (Question 4a), 66 participants listed other reasons to assess cognition not listed in the response options. Three themes emerged from these responses (Supplementary table 4): discharge and intervention planning, understanding of cognition in aphasia, and determining cognitive status.

Comments related to assessing cognition for discharge and intervention planning mostly referred to returning to activities of daily living, such as driving and returning to work. Respondents reported assessing cognition to help determine the best treatment options for people with aphasia, and how to provide better multidisciplinary support. Other reported purposes of cognitive assessments included to determine the impact of language and cognition on activities of daily living, provide information on cognitive status, monitor change, plan neurosurgery, and determine research inclusion criteria.

Question 5: Which aspects of cognition should be assessed in aphasia?

This question aimed to gauge the perceived importance (not important, slightly important, moderately important, important, very important) of assessing each aspect of cognition. Participants were asked the level of importance of assessing nine different cognitive domains: attention, executive functions, learning, memory, praxis, processing speed, social cognition, visuoconstructional, and visuoperceptual processing.

All domains were considered at least moderately important by at least half of the respondents. The domains of attention, executive functions, learning and memory were considered important or very important by over 80% of respondents. Over half of the respondents indicated that attention (55%) and executive functions (51.2%) were very important. These were followed by about 60% of respondents indicating that the assessment of praxis, processing speed, social cognition and visuo-perceptual abilities were important or very important. Half of the respondents considered visuoconstructional abilities important, or very important to assess.

A follow-up optional question asked respondents to indicate any other aspects of cognition they believed should be assessed in aphasia. There were 42 responses to this question, but with very little consensus on the additional areas including: insight and awareness (n = 4), mathematical skills (n = 3), self-monitoring and/or self-correction (n = 5), emotional issues and/or mood (n = 3), non-verbal cognition (n = 2), visual processing (n = 2), working memory (n = 2), abstraction (n = 1), flexibility (n = 1), inhibition (n = 1), orientation (n = 1), processing speed (n = 1), reasoning (n = 1), set-shifting (n = 1), and social ability (n = 1).

Eleven respondents indicated the importance of individually determining assessments:

'It depends on why the person is tested. I seldom test learning and processing speed because these functions normally are very reduced in my patients. Instead, I inform patients and relatives about how to take this into account. If the question is whether the patient can drive a car again, I test processing speed.'

One respondent highlighted the important point that an overview of all cognitive abilities of the person with aphasia is crucial for both diagnosis and treatment:

'I think it is ill-considered to treat aphasia in isolation, as if it is 'just' a language deficit. If you don't have a big picture overview of cognitive functioning, you cannot properly treat or understand the aphasic patient.'

Questions 6 and 7: Who should assess cognition in aphasia and discuss the results?

In Questions 6 and 7, participants were asked about whose role it was to assess cognition and discuss results with caregivers and/or family members of people with aphasia (Figure 4). Respondents could choose as many professions as they felt were applicable, and hence the responses do not reflect the opinion that only a single professional group should assess cognition

in people with aphasia. Around 80% of respondents indicated that neuropsychologists should assess and discuss cognitive functioning in people with aphasia, and a similar percentage of respondents indicated the same for SLPs. This was followed by around 56% of respondents indicating that it could be the role of occupational therapists. A third of respondents (33%) felt that neurologists should be responsible for assessing cognition in aphasia, but half suggested that neurologists could take the role of discussing the results.

<insert figure 4>

Respondents were also given the opportunity to comment further and specify additional professionals. Most of these comments highlighted that both the assessment of cognition and discussion of results should be carried out by a multidisciplinary team. Others noted that the professional carrying out the assessment would vary across cases, and that the professional discussing the results should either have performed the assessment or be involved in their rehabilitation. Some respondents stated that both assessment and discussion should be carried out by professionals trained in the assessment of cognition, and that assessments also required a professional who had specific experience working with people with aphasia. Two respondents indicated that the discussion should have SLP support.

Question 8 and 8a: What are the barriers to assessing cognition in aphasia?

Question 8 (yes/no question) aimed to gain insights into any barriers experienced when assessing cognition in aphasia. Some respondents indicated that this was not applicable to their professional role (6.8%) and/or workplace (10.4%). Of the remaining respondents, over half had experienced barriers in their professional role (68.9%) and/or in their workplace (64.6%). 158 respondents specified these barriers in open-ended responses to Question 8a (431 barriers), which were classified into five themes: barriers related to materials (39.2%), professional role (26.9%), professionals' knowledge and skills (17.2%), the people with aphasia (11.8%) and other reasons (4.9%). These themes are explored further in Table 1.

<insert table 1>

Materials: Barriers related to materials were most prominent and included a lack of appropriate assessments and resources for use in aphasia (especially severe aphasia), and lack of access to test materials that did exist. Respondents were concerned that materials were highly dependent on

language skills, and that there was a lack of appropriately normed, standardised, and functional assessments. Other reported barriers to assessing cognition in people with aphasia included: restrictions in the use of certain assessments (e.g., psychologists only), the prohibitive cost of assessments, a lack of assessments in multiple languages, the complexity of test instructions, an over reliance on formal scoring, and a lack of consensus on which tests to use.

Professional role: In a clinical setting, the lack of clarity of professional roles was a common theme. Respondents indicated that it was often unclear which professionals were responsible for cognitive assessment in aphasia. They also reported that the assessment of cognition was not considered within their scope of practice (five of the 12 responses were from SLPs, the remaining seven did not indicate their professional identity), and that it was considered the occupational therapist and/or neuropsychologist's role.

Time constraints were mentioned as a major barrier, due to the need to prioritise language assessments given the limited time often available to work with people with aphasia. A minority of responses included insurance constraints and staff availability as barriers. Three respondents indicated that they found it difficult to explain to other professionals the importance of assessing cognition in people with aphasia (two responses from SLPs, the remaining respondent did not indicate their professional identity).

Knowledge and skills: Some of the responses suggested that another barrier was their own lack of knowledge and experience with cognitive assessments.

Some professionals identified a further barrier as being that some (non-SLP) professionals had limited awareness that that non-linguistic cognitive deficits might also be present in aphasia.

Difficulties disentangling linguistic impairments from non-linguistic cognitive impairments using many of the available assessments was also mentioned, along with a limited understanding of aphasia amongst non-SLP professionals and limited understanding of cognition amongst SLPs.

Aphasia and comorbidities: This theme related to barriers inherent to aphasia and its associated co-morbidities. The negative impacts of both aphasia and motor deficits on the assessment of cognition were indicated by some respondents. These deficits impacted on the task demands of the assessment (i.e., understanding test instructions, writing test responses). Personal factors, such as self-awareness of aphasia, engagement or fatigue were reported as barriers.

Section 3: Tools for assessing cognition in aphasia.

The third part of the survey comprised open-ended questions about the tools respondents used to assess cognition in aphasia (Q9), the cognitive constructs they aimed to measure (Q9), the factors influencing the choice of tools (Q10), and perceived difficulties in choosing tools (Q11).

Question 9: The tools used to assess cognition in aphasia.

Table 2 provides a summary of the tools cited by 10 or more respondents (see Table 4 in Supplementary Materials for all responses and their classifications). The most frequently used tool reported (26.8% of the 164 respondents) was the Cognitive Linguistic Quick Test (Helm-Estabrooks, 2002), followed by the Trail Making Test (16.5%; Reitan, 1992), the Montreal Cognitive Assessment (15.8%; Nasreddine et al., 2005), and Raven's Colored Progressive Matrices (12.2%; Raven, 1962).

Responses varied as to how respondents named the tools available, both in the constructs that they thought the tools measured, and in the terminology that they used to refer to these constructs. For example, across respondents, the Rey Complex Figure test (Osterrieth, 1944) was reported to measure the following range of constructs: visual memory, memory, executive functions, visuospatial, visual memory, visual attention, visual perception, construction, planning, visuoconstructional abilities/memory, constructional praxis, short term and episodic memory, recognition and learning. To address the variation in terms used to describe constructs, we mapped the suggested constructs from responses onto cognitive domains using a widely used framework that adheres to the DSM-5 approach (i.e., Sachdev et al., 2014), these included: complex attention, executive function, learning and memory, language, social cognition and perceptual-motor function.

Question 10: Which factors are considered when choosing tools?

Question 10 asked respondents to outline, in an open-ended response, practical, clinical and/or theoretical factors considered when selecting tools to assess cognition in aphasia. Responses were classified into four main themes: tools, impairment, person with aphasia, and assessment (Figure 5).

<insert figure 5>

Assessment tool-related factors: The major factor related to tool selection was task administration, which included consideration of the suitability for people with aphasia in terms

of the language demands of the task itself, and of the task instructions. Another influential factor was reported to be the amount of time that the assessment took to perform (and how much time was available to the clinician and/or researcher) with some respondents indicating a preference for assessments that were less time consuming. For example:

‘Some tests (i.e., Wisconsin Card Sorting Test) are time-consuming and some aphasic persons have problems to understand the rules of the game. This is particularly true is [sic] persons with moderate-to-severe comprehension deficits.’

The availability of tests within the researcher’s and/or clinician’s context was the third most frequently reported factor concerning administration, followed by the complexity of administration, appropriateness given the person with aphasia’s background (language, culture, age, etc.), and the possibility for remote administration.

Respondents also mentioned factors that related to test construction and psychometrics such as validity (i.e., the extent to which a tool accurately measures a construct), reliability (i.e., how consistent a tool is if administered on different occasions) and the availability of normative data for different age groups, and educational and cultural backgrounds. Some respondents reported prioritising evidence-based tools in their practice.

A small proportion of respondents (5.3%) indicated a preference for using assessments of functional activities due to their ecological validity:

A few respondents (4.0%) indicated that, to better understand therapy outcomes, they prioritised tools that provided a comprehensive picture of an individual’s cognitive abilities (such as the Cognitive Linguistic Quick Test (Helm-Estabrooks., 2002), or the Montreal Cognitive Assessment (Nasreddine., 2005)), and primarily assessed domains that they believe most closely interact with language:

‘We choose the assessment tests considering those cognitive functions that may play a relevant role in language performance and its response to treatment. The selection also seeks to strike an appropriate balance between the need for a comprehensive assessment and not too time-consuming for people with aphasia.’

Factors related to the person with aphasia: Respondents reported factors related to both the clinical profile of the person with aphasia ('Impairment' in Figure 5) and also broader considerations ('Person with aphasia' in Figure 5).

There were four main ways in which respondents were influenced by the nature of the person with aphasia's impairment when choosing cognitive assessment tools: aphasia severity, comorbidities, lesion information and stroke severity, for example:

'I take into account the type of aphasia of the patient and the conserved language abilities, for example how is verbal comprehension, reading, image recognition ... I also take into account the severity of the lesion and the deficits that the patient and adjust the choice of tests to these abilities.'

The severity and type of language impairment was the main reported consideration, with nonverbal tasks often preferred. Other comorbidities such as vision, hearing and physical impairments, and/or apraxia of speech were also taken into consideration. A few responses also reported considering lesion information and stroke severity.

In terms of broader considerations, the level of engagement and the goals of the individual with aphasia were also reported as important when choosing a tool to measure cognition, these included the individual's stamina (i.e., extent and rapidity of fatigue), and emotional status (i.e., level of frustration) during the assessment. Age, languages spoken and cultural background, prior level of function, awareness and insight into their aphasia were also identified as factors that were taken into consideration.

Factors related to assessment: This theme focused on considerations influencing the choice of assessment tools more broadly. This included the need to gain a better understanding of a person's cognitive abilities while being mindful of their linguistic impairment. This was followed by a need for careful consideration of the main goals of the assessment process, the constructs that were required to be measured, the relevance to treatment planning, the availability of other members of a multidisciplinary team who may have access to assessment materials or experience with cognitive testing, and finally a consideration of which assessments the patient had already completed.

Question 11: What are the perceived difficulties in choosing tools?

In response to Question 11, most respondents (75%) reported difficulties choosing appropriate tools to assess cognition in aphasia. Three themes emerged from the responses: difficulties associated with the appropriateness of assessment tools, with professional roles, and with the differential diagnosis of language and cognitive impairments (Supplementary Table 4).

Difficulties associated with the appropriateness of assessment tools: the most frequently reported difficulty was finding assessment tools that were appropriate for people with aphasia, mainly due to available assessment tools being over-reliant on language ability. A particular concern was the availability of assessment tools with adequate psychometric properties, especially for individuals from diverse educational, linguistic and cultural backgrounds. The construction of assessment tools, in terms of stimuli, number of practice trials and validity was also indicated as a concern.

Furthermore, respondents reported finding it hard to find assessment tools that were suitable for people with aphasia with other comorbidities, such as motor deficits. A lack of available comprehensive cognitive assessment tools, as opposed to screening tools, was also mentioned.

Difficulties associated with professional roles: Issues concerning whose role it should be to assess cognition in aphasia and the lack of experience and knowledge with cognitive assessment were also evident in the responses. Some professionals also indicated they did not have enough time with people with aphasia to fit in a cognitive assessment.

Difficulties associated with the differential diagnosis of language and cognitive impairments: A final major difficulty reported, regarding assessment tools, was a lack of clear understanding of the interface between language and cognition, and the difficulty in separating them in order to understand the extent of cognitive and linguistic impairments.

Question 12: What are the changes made when assessing cognition in people with aphasia compared to people without aphasia?

In response to Question 12, the majority (88%) of respondents reported that they adapted their approach when assessing people with aphasia, compared to those without aphasia. For these respondents (n=155), a subsequent open-ended question probed the changes that they made. Three main themes emerged from these questions: modifications to the task, selection of assessment tools, and the assessment conditions (Table 3).

Modifications to the task: Modifications to the tasks included reducing the linguistic load of the task by simplifying language, slowing speech rate and providing repetitions. Providing visual support or using supported conversation techniques and allowing for multimodal input were also reported as ways of adapting tools for use with people with aphasia. Respondents indicated that they modified and adapted instructions so that they could be understood by people with aphasia prior to commencing the test. Some respondents indicated that to ensure that people with aphasia could be assessed, task standardisation procedures were not always followed (i.e., allowing participants more time to respond to a task that was permitted). Finally, some tasks were modified to accommodate comorbidities, (i.e., presenting the stimuli on the preferred side for people with hemianopia).

<insert table 3>

Selection of assessment tools: Differences in the selection of tools were reported, with respondents indicating that they tended to select tools that they believed were more appropriate for people with aphasia, that is, preferring tests that were not as reliant on language abilities. Most respondents also indicated a preference for using functional assessments, while some indicated a preference for informal assessments over formal assessments.

Respondents reported considering patients' needs, a preference for shortened versions of tasks, and that, during test administration, they tried to accommodate the requirements of an individual by adapting task demands to be appropriate for residual language ability and considering the language impairment when interpreting the results. Respondents also indicated they often worked with the multidisciplinary team to find the best assessment and the optimal way to administer it.

Assessment conditions: Finally, respondents indicated they tried to minimise distractions during testing, paced the administration depending on the person's energy levels, prioritised face to face testing, and requested support from caregivers.

Discussion

Despite the prevalence of non-linguistic cognitive deficits in people with aphasia (Fucetola et al., 2009; Gilmore et al., 2019; Mayer & Murray, 2012; Murray, 2012), the best approach to assessing cognition, beyond language, in this population is unclear. This study aimed to gauge professionals' understanding of cognition and identify the approaches currently being used to assess cognition in people with aphasia.

Our international, multidisciplinary survey of professionals who routinely worked with people with aphasia highlighted that 1) the assessment of cognition in this population is considered essential for effective clinical and research planning and management; but that 2) there is a dearth of research and guidance for determining how to reliably do this in both clinical and research environments, with most of the available assessment tools considered inappropriate for this population. In addition to identifying numerous barriers to assessing cognition in people with aphasia, this is the first study to probe the adaptations that are implemented in clinical and research settings to address some of these barriers.

Definition of cognition

In our data, cognition was defined as the skills, mental and brain processes responsible for thinking, processing and integrating information, problem solving and learning, with no major discrepancies between the definitions of cognition given. The aspects of cognition reported by our respondents aligned with the domains within current theoretical and neuropsychological assessment frameworks (Harvey, 2019; Helm-Estabrooks, 2002; Lezak et al., 2012; Sachdev et al., 2014b; Strauss et al., 2006; Sternberg & Sternberg, 2012) (i.e., attention, executive functions, memory, language, motor function and praxis, visuo-perceptual and visuospatial abilities). However, responses focused on the abilities that make us human and that cognition supports (e.g., to think, reason, remember, problem-solve, make decisions, do maths, plan, learn), rather than the cognitive function itself. The one point of divergence was whether language should be part of a definition of cognition.

A major difficulty endorsed by our respondents was the lack of clear understanding of the interface between language and cognition, and the challenges in separating them to disentangle cognitive and/ or language impairment. The inclusion of language as a construct of cognition has been debated for decades (e.g., Fodor, 1975, Nickels et al., 2023). There have been numerous theoretical approaches to understanding how linguistic impairment interacts with other cognitive domains in aphasia. For example, impaired attention (Hula et al., 2007; Hula & McNeil, 2008; McNeil et al., 2011), executive functioning (Thompson et al., 2018) and

short-term memory (Minkina et al., 2018) have been proposed as, at least partially, explaining language difficulties in aphasia. However, language has also been shown to impact cognitive functioning (see for example, Alderson-Day, B., & Fernyhough, 2015; Baldo et al., 2005; Granato et al., 2020).

Responses in our data highlight that defining cognition, and the contribution of language to cognition requires clarity in order to support professionals to provide optimal care for people with aphasia. Further empirical work is needed to understand the interaction between language and non-linguistic cognition, especially in people with aphasia.

Assessing cognition in aphasia is important

Our results illustrate that assessing cognition in aphasia is considered to serve different, equally important, purposes, mainly related to discharge and intervention planning.

Overall, respondents felt that it was important to assess all cognitive domains, but particularly attention, executive functioning, memory and learning. Previous literature reviews have also suggested memory and executive functioning to be a main focus of non-linguistic cognitive assessment in aphasia research (Fonseca et al., 2016; Tessaro et al., 2022). On the other hand, while our respondents indicated assessing attentional skills in aphasia was very important, such assessments were not frequently reported in published research (Fonseca et al., 2016; Tessaro et al., 2022). It seems unlikely that this is because attention tasks involve more language processing than assessments of other domains, as Tessaro et al.'s (2022) analysis suggested that 71.6% of tasks that assessed attention did not have language requirements. Given that attention deficits have been reported in people with aphasia, and both researchers and clinicians agree their assessment is desirable, further investigation and/or reporting in the literature seems important (Murray, 2002, 2012).

Although most respondents in the present study agreed that cognitive assessment is important in aphasia, several barriers were indicated that prevent the assessment of cognition in the clinical setting, particularly lack of time, staff, training, and appropriate testing materials. However, because professionals recognise the importance of this assessment, there were many indications that informal assessments were performed with people with aphasia.

Assessment Tools

Tool Appropriateness

Current tools and materials for assessing cognition were generally considered to be inappropriate to use with people with aphasia. Given this, professionals working with people

with aphasia face considerable challenges when assessing cognition beyond language given the reliance on language abilities in most of these cognitive assessments. For example, the Montreal Cognitive Assessment (Nasreddine et al., 2005), a commonly used cognitive screening tool that includes language tasks such as naming, repetition, and verbal fluency, uses tasks to measure memory skills that require unimpaired language abilities (e.g., retrieval of a list of words), which limits the feasibility of its use in aphasia (Barnay et al., 2014).

Another assessment tool reported to be frequently used was the Cognitive Linguistic Quick Test (Helm-Estabrooks, 2002), a screening assessment designed to screen the five cognitive domains in adults with acquired neurological disorders. This tool was designed specifically with people with aphasia in mind and five of the ten tasks aim to minimize language load (symbol cancellation, symbol trails, design memory, mazes and design generation). However, the scoring and reporting process in this test is predominantly completed by cognitive domain (attention, memory, executive functions, and of course, language) and, with the exception of visuospatial skills, the composite measure of each domain includes tasks reliant on language skills (e.g., generative and confrontation naming, story retelling, recalling personal facts). This is despite the fact that the CQLT also provides the normative data for individual non-linguistic tasks (albeit standardised with an English-speaking population in the USA).

Our results support the findings from a systematic review (Tessaro et al., 2022) that indicated that over half (54%) of the tests used to assess cognition in aphasia had some degree of language load (i.e., verbal stimuli, responses requiring language production or required mapping stimuli to its meaning). Given the challenges evident in modifying, or creating, cognitive assessments, the research and clinical communities are advised to systematically consider the language load of each assessment when selecting assessment tools to be used with people with aphasia. Tessaro et al. (2022) provide details of whether each cognitive test included verbal stimuli, required language production and/or required mapping verbal stimuli to meaning. This could be a useful resource for clinicians and researchers.

Tool adaptation

Our data illustrate that assessments require adaptations to make them more aphasiafriendly, with an overwhelming majority of respondents (88%) reporting that this was necessary. Whilst this will be no surprise to those working with aphasia, our survey has systematically identified that in clinical and research settings changes to assessments protocols are routinely implemented across all dimensions of the assessment process, from provision of

test instructions to the interpretation of results. Previous work by Guo et al. (2014) demonstrated that Singaporean SLPs adapted and modified cognitive assessment tools when working with people with aphasia. Our data provide evidence that this process of adaption occurs internationally and across clinical and research disciplines.

We found that administration procedures were modified to adapt for assessment of persons with aphasia (e.g., allowing more time to respond, and/or providing visual supports). Although typically performed by experienced clinicians and researchers, *ad hoc* adaptations and modifications can bring additional limitations to the assessment process. Deviating from standard administration procedures could impact on the accuracy and reliability of diagnosis when interpretation is based on normative data that have been established following standard administration procedures (Cicchetti, 1994). This presents as a major issue in clinical practice where the accuracy of tests scores assists in differential diagnosis, providing advice and possibly prognosis to affected individuals. Similarly, in a research setting, the accuracy of a test score can be used to determine suitability of inclusion in a study, or the impact and efficacy of an intervention. An unreliable score can lead to unethical inclusion, or exclusion in a study, or even false statistical results. Moreover, any deviations from standardization invalidates comparisons across studies, limiting the possibility of study replication, the ability to conduct reliable meta-analyses and the generalisability of results to people with aphasia beyond the experimental group. Finally, *ad hoc* adaptations (which may vary from person to person, clinical, research or person with aphasia) may appear to facilitate administration but do not ensure that they will fully clarify the test requirements for the testee. Consequently, errors in a modified test may reflect a deficit in the cognitive ability tested, a misinterpretation of test requirements, or both.

Informal and functional assessments

The use of informal and functional assessments to evaluate the cognitive status of people with aphasia was mentioned by respondents throughout the survey. Examples of informal and functional assessments included self-report by the person with aphasia or carer about daily activities such as washing up, dressing, and cooking. The reported success on these tasks was then used as an indicator of cognitive functioning.

Informal observations of the person with aphasia by the clinician were also part of cognitive assessment in clinical settings. Whilst the responses in our survey were not specific enough to allow us to explore informal and functional assessment practices in detail, an informal approach may be a useful way to gather information on the difficulties and strengths

of people with aphasia. However, Thomson et al., (2018) noted a lack of guidelines available for clinicians on conducting informal assessment, and that the success of an informal assessment is likely dependent on the clinicians' experience and skills. As such, although informative regarding a person's needs and strengths; informal assessments are limited by the idiosyncrasies of the tasks used and can be difficult to generalise across approaches.

Knowledge regarding the assessment of cognition in aphasia

One of the major barriers for assessing cognition in aphasia appears to stem from a lack of knowledge of the cognitive assessments available. SLP respondents reported a lack of knowledge of cognition and/or a lack of training in administering cognitive assessments. This also related to professional boundaries and scope of practice (see below). That SLPs felt they lacked training in the evaluation of cognition has been discussed in research regarding decision-making capacity assessment (Suleman & Hopper, 2016). Conversely, whilst the assessment of cognition is often the domain of neuropsychologists and/or occupational therapists, our respondents (who included these professions) indicated that these professionals may lack knowledge about, and experience working with aphasia. There were also reports that the wider multidisciplinary team may also have a limited understanding of the complexities of language impairment in aphasia.

Our results highlight the critical importance of inter-professional learning and continuing professional development, as well as working within meaningful multidisciplinary teams in which professionals bring their own discipline-specific expertise to the assessment process. This inter-professional learning should be available through in graduate training programs, national peak organizations, and best practice guidelines for aphasia and/or stroke. This would support SLPs in learning more about cognitive assessment, and non-SLPs in learning about aphasia and how it impacts cognitive assessment. Clarifying the scope of practice and ensuring any overlap of practice is rigorously taught across disciplines would support professionals to recognise their unique expertise and limitations when navigating the assessment of cognition in aphasia.

Whose role it is to assess cognition in aphasia?

The in-depth assessment of language is primarily conducted by SLPs. Our results showed that professional role becomes less clear when the assessment of non-linguistic cognition in aphasia is indicated. Despite the clear advantages to multidisciplinary work, with it may also come an ambiguity of responsibilities within a team; in our survey, professionals

reported a lack of clarity in their own role, but they were also unsure of the scope of practice of other professionals within their multidisciplinary team.

For example, the national stroke guidelines of the Royal College of Physicians (2023) recognise the role of SLPs in interpreting language assessments in people with aphasia. However, the guidelines also recommend there is close collaboration with SLPs during neuropsychological assessment beyond language, implicitly acknowledging that neuropsychological assessments of broader cognitive abilities, typically the role of clinical (neuro)psychologists, also involve language. These and other guidelines could be clearer about scope of practice so that the multidisciplinary team can provide mutual assistance to accurately interpret assessments of cognition bearing in mind their language content/bias.

As mentioned above, there was not only a lack of clarity regarding whose role it was to assess cognition, but also whose role it was to discuss the results of this assessment. Most participants (>80%) agreed that SLPs and neuropsychologists should both assess cognition and discuss the results of the assessments (see Figure 4, earlier). However, for other professionals (e.g., neurologist, physiotherapist, psychiatrist) respondents generally felt these professionals should have a role in discussing results rather than in performing assessments. For example, only a third of our respondents indicated that neurologists should assess cognition in people with aphasia, but half felt that neurologists should discuss the results with the person with aphasia and carers.

Echoing the broader findings by Rohde et al. (2013) that further improvement is needed within stroke multidisciplinary clinical guidelines, improved guidelines around professional roles are required to optimise the assessment of cognition and discussion of findings. This should ensure that the assessment of cognition is meaningful, appropriate and relevant, by empowering disciplines to develop their knowledge and practice, but also by providing accountability and therefore minimising the risk that cognition is either not assessed at all, or inappropriately assessed in people with aphasia.

Limitations

Our study collected and analysed a large number of responses, from a wide range of countries and professions. Whilst we deliberately sought participation from non-English speaking respondents, the majority of responses in our sample were from researchers and clinicians from developed and English-speaking countries. This limits the generalisation of our findings across developing and non-English speaking nations.

While it was a strength that we gathered responses from different countries, aphasia practices vary across these countries due to different healthcare and educational systems, and heterogeneity within the same country also exists, due to differences in training. Given the relatively small number of observations per country it was not possible to evaluate these differences and compare and contrast responses within- and across countries.

In addition, when considering differences across professionals relating to cognition and its assessment with aphasia, the inclusion of a range of professionals who work with aphasia allowed our results to have an important multidisciplinary perspective. Nevertheless, more than half of our sample was composed of SLPs (64%). Although this was anticipated given that they are the main professionals working with aphasia, the non-SLPS represented a large number of different professions, and this group was too heterogenous for cross-profession analysis to be possible. Future studies would benefit from a larger sample size in areas that were underrepresented in our data; this would allow further nuance in the analysis and interpretation of the data.

Apart from surveying clinical and research practice, clinical practice can be updated by drawing on findings from large-scale observational studies of different aspects of non-linguistic cognitive domains in aphasia rehabilitation¹ (e.g., El Hachiou et al., 2014; Fonseca et al., 2019; Marinelli et al., 2017), and studies that have used cognitive tests with relatively small linguistic demands. However, before embracing such tests in clinical practice, consideration should also be given to the psychometric properties of such tests, especially measurement error and other indices of reliability and validity (cf., Murray et al., 2018).

Conclusions

Although there have been many surveys on clinical practices in aphasia (for example, Aldous et al., 2014; Chang et al., 2018; Guo et al., 2014; Kiran et al., 2018; Rose et al., 2014; Suleman & Hopper, 2016; Verna et al., 2009), the few that relate to the assessment of cognition, have focused mostly on memory abilities alone (Greenspan et al., 2021; Salis et al., 2018). Therefore, our study provides novel insights into how aphasia researchers and clinicians define cognition, why and how they choose to assess cognition, what assessment tools they use, and the perceived barriers to assessing cognition in people with aphasia.

Our results have underscored the importance of assessing cognition in aphasia for a range of different purposes. Our data clearly highlights a lack of cognitive assessment tools

¹ We would like to thank Professor Matthew Lambon Ralph for suggesting this point.

that are appropriate for use with people with aphasia. Further research to develop appropriate assessment tools, and/or normative data specifically for this cohort is warranted (see table 4). Similarly, both researchers and clinicians would benefit from expert consensus recommendations to provide guidance on the most appropriate assessment tools available for assessing cognitive function in people with aphasia.

Whilst professionals in our study often reported relying on implementing pragmatic changes to the tasks in order to make them more suitable for people with aphasia, this lack of adherence to standardisation procedures can manifest as a barrier to accurate interpretation of results, reliable diagnostics and may impact generalisation of results across studies. To overcome this barrier, professionals expressed that they often use informal assessments to evaluate non-linguistic aspects of cognition in aphasia.

The lack of available materials and standardised administration appropriate for individuals with language impairment is also likely to contribute to the systematic exclusion of people with aphasia from stroke studies (Snaphaan et al., 2007; Srikanth et al., 2003; Shiggins et al., 2023). Given that a third of stroke survivors experience aphasia, systematically excluding this population will impact on the generalisability of stroke research more broadly.

Moreover, our data suggests that across disciplines, there is a perceived lack of appropriate training for conducting rigorous and informative cognitive assessments in people with aphasia. Thus, there is a need to ensure that appropriate training is provided to ensure all professions working with this population have a well-grounded understanding of cognition and an understanding of the consequences of aphasia when conducting cognitive assessment with those living with aphasia. Specifically, for the multidisciplinary team who work with people with aphasia, there is an urgent need to increase awareness about the importance of, and difficulties in, accurately assessing non-linguistic cognitive functions in people with aphasia.

Finally, we have demonstrated that to improve aphasia care, the relevant clinical guidelines for the many disciplines involved should clarify and specify the roles and contributions of the different professionals working with aphasia regarding the assessment of non-linguistic functions. To this end, in table 4, we provide actionable insights for the field, that we hope will support academics and clinicians to progress these important findings.

Declaration of interest statement

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Raw data were generated at Newcastle University. Derived data supporting the findings of this study are available from the first author [BT] on request.

References

Aldous, K., Tolmie, R., Worrall, L., & Ferguson, A. (2014). Speech-language pathologists' contribution to the assessment of decision-making capacity in aphasia: a survey of common practices. *International Journal of Speech-Language Pathology*, 16(3), 231–241. doi: 10.3109/17549507.2013.871751

American Speech-Language Hearing Association (2024). Retrieved 28 November 2024 <https://apps.asha.org/EvidenceMaps/Articles/ArticleSummary/c85526ef-3d44-ef11-8151-005056834e2b>

Australian Aphasia Rehabilitation Pathway (2024). Retrieved 25 November 2024 <https://www.aphasiapathway.com.au/?name=4-Assessing>

Barnay, J. L., Wauquiez, G., Bonnin-Koang, H. Y., Anquetil, C., Pérennou, D., Piscicelli, C., Lucas-Pineau, B., Muja, L., le Stunff, E., de Boissezon, X., Terracol, C., Rousseaux, M., Bejot, Y., Binquet, C., Antoine, D., Devilliers, H., & Benaim, C. (2014). Feasibility of the Cognitive Assessment scale for Stroke Patients (CASP) vs. MMSE and MoCA in aphasic left hemispheric stroke patients. *Annals of Physical and Rehabilitation Medicine*, 57(6), 422–435. doi: 10.1016/j.rehab.2014.12.001

Berg, E. A. (1948). A simple objective technique for measuring flexibility in thinking. *Journal of General Psychology*, 39, 15–22. <https://doi.org/10.1080/00221309.1948.9918159>

Block, C., Johnson-Greene, D., Pliskin, N., & Boake, C. (2017). Discriminating cognitive screening and cognitive testing from neuropsychological assessment: implications for professional practice. *Clinical Neuropsychologist*, 31(3), 487–500. doi: 10.1080/13854046.2016.1267803

Braun, V., Clarke, V., Hayfield, N., Terry, G. (2019). Thematic Analysis. In Liamputtong, P. (eds.) *Handbook of Research Methods in Health Social Sciences*. Springer. https://doi.org/10.1007/978-981-10-5251-4_103

Braun, V. V., & Clarke, V. (2006). Using thematic analysis in psychology. *Qualitative Research in Psychology*, 3(2), 77–101. doi: 10.1191/1478088706qp063oa

Brownsett, S. L. E., Warren, J., Geranmayeh, F., Woodhead, Z., Leech, R., & Wise, R. (2014). Cognitive control and its impact on recovery from aphasic stroke. *Brain*, 137(1), 242–254. doi: 10.1093/brain/awt289.

Butt, P. & Bucks, R. (2004). *The Butt Non-Verbal Reasoning Test*. Speechmark.

Canadian Stroke Best Practice. (2020). Recommendations. Retrieved 28 November 2024 <https://www.strokebestpractices.ca/recommendations>

Chang, H., Power, E., O'Halloran, R., & Foster, A. (2018). Stroke communication partner training: a national survey of 122 clinicians on current practice patterns and perceived implementation barriers and facilitators. *International Journal of Language & Communication Disorders*, 53(6), 1094–1109. doi: 10.1111/1460-6984.12421.

Cicchetti, D. (1994). Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychological Assessment*, 6(4), 284–290. <https://psycnet.apa.org/doi/10.1037/1040-3590.6.4.284>

Clarke, V., & Braun, V. (2017). Thematic analysis. *Journal of Positive Psychology*, 12(3), 297–298. <https://doi.org/10.1080/17439760.2016.1262613>

Demeyere, N., Riddoch, M. J., Slavkova, E. D., Bickerton, W.-L., & Humphreys, G. W. (2015). The Oxford Cognitive Screen (OCS): validation of a stroke-specific short cognitive screening tool. *Psychological Assessment*, 27(3), 883–894. doi: 10.1037/pas0000082

Diedrichs, V. A., Jewell, C. C., & Harnish, S. M. (2022). A scoping review of the relationship between nonlinguistic cognitive factors and aphasia treatment response. *Topics in Language Disorders*, 42(3), 212–235. doi: 10.1097/tld.0000000000000290.

El Hachoui, H., Visch-Brink, E. G., Lingsma, H. F., van de Sandt-Koenderman, M. W., Dippel, D. W., Koudstaal, P. J., Middelkoop, H. A. (2014). Nonlinguistic cognitive impairment in poststroke aphasia: a prospective study. *Neurorehabilitation and Neural Repair*, 28(3), 273–81. doi: 10.1177/1545968313508467.

Eysenbach, G. (2004). Improving the quality of web surveys: the Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *Journal of Medical Internet Research*, 6(3), 1–6. <https://doi.org/10.2196/jmir.6.3.e34>

Fodor, J. A. (1975). *The Language of Thought* (Vol. 5). Thomas Y. Crowell.

Fonseca, J., Ferreira, J., & Martins, I. (2016). Cognitive performance in aphasia due to stroke: a systematic review. *International Journal on Disability and Human Development*, 16(2), 127–139. <https://doi.org/10.1515/ijdhhd-2016-0011>

Fonseca, J., Raposo, A., & Martins, I. P. (2019). Cognitive functioning in chronic post-stroke aphasia. *Applied Neuropsychology: Adult*, 26(4), 355–364. doi: 10.1080/23279095.2018.1429442.

Folstein M.F., Folstein S. E., & McHugh, P. R. (1975). “Mini Mental State”: a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12, 189–98. doi: 10.1016/0022-3956(75)90026-6

Fucetola, R., Connor, L. T., Strube, M. J., & Corbetta, M. (2009). Unravelling nonverbal cognitive performance in acquired aphasia. *Aphasiology*, 23(12), 1418–1426. <https://doi.org/10.1080/02687030802514938>

Gilmore, N., Meier, E. L., Johnson, J. P., & Kiran, S. (2019). Nonlinguistic cognitive Factors predict treatment-induced recovery in chronic poststroke aphasia. *Archives of Physical Medicine and Rehabilitation*, 100(7), 1251–1258. doi: 10.1016/j.apmr.2018.12.024

Greenspan, W., Obermeyer, J., Tucker, C., Grunwald, H., Reinert, L., & Martin, N. (2021). Clinician perspectives on the assessment of short-term memory in aphasia. *Aphasiology*, 35(3), 334–356. <https://doi.org/10.1080/02687038.2020.1712584>

Guo, Y., Togher, L., & Power E. (2014). Speech pathology services for people with aphasia: what is the current practice in Singapore? *Disability and Rehabilitation*, 36(8), 691–704. doi: 10.3109/09638288.2013.804597.

Harvey, P. D. (2019). Domains of cognition and their assessment. *Dialogues in Clinical Neuroscience*, 21(3), 227–237. <https://doi.org/10.31887/DCNS.2019.21.3/pharvey>

Helm-Estabrooks, N. (2002). Cognition and aphasia: a discussion and a study. *Journal of Communication Disorders*, 35(2), 171–186. [https://doi.org/10.1016/S0021-9924\(02\)00063-1](https://doi.org/10.1016/S0021-9924(02)00063-1)

Hula, W. D., McNeil, M. R., & Sung, J. E. (2007). Is there an impairment of language-specific attentional processing in aphasia? *Brain and Language*, 103(1–2), 240–241. <https://doi.org/10.1016/j.bandl.2007.07.023>

Hula, W. D., & McNeil, M. R. (2008). Models of attention and dual-task performance as explanatory constructs in aphasia. *Seminars in Speech and Language*, 29(3), 169–187. doi: 10.1055/s-0028-1082882.

Kim, E. S., Suleman, S., & Hopper, T. (2020). Decision making by people with aphasia: a comparison of linguistic and nonlinguistic measures. *Journal of Speech, Language & Hearing Research*, 63(6), 1845–1860. https://psycnet.apa.org/doi/10.1044/2020_JSLHR-19-00182

Kiran, S., Cherney, L. R., Kagan, A., Haley, K. L., Antonucci, S. M., Schwartz, M., Holland, A. L., & Simmons-Mackie, N. (2018). Aphasia assessments: a survey of clinical and research settings. *Aphasiology*, 32(sup1), 47–49. <https://doi.org/10.1080/02687038.2018.1487923>

Knollman-Porter, K., Brown, J. A., Wallace, T., & Spitz, S. (2021). First-line health care providers' reported knowledge of and referrals to speech-language pathologists for clients with

mild traumatic brain injury. *American Journal of Speech-Language Pathology*, 30(5), 2214–2227. doi: 10.1044/2021_AJSLP-20-00373

Kohnert, K. (2004). Cognitive and cognate-based treatments for bilingual aphasia: a case study. *Brain and Language*, 91(3), 294–302. doi: 10.1016/j.bandl.2004.04.001

Kuzmina, E., & Weekes, B. S. (2017). Role of cognitive control in language deficits in different types of aphasia. *Aphasiology*, 31(7), 765–792. <https://doi.org/10.1080/02687038.2016.1263383>

LaCroix, A. N., McKayla, T., & Rogalsky, C. (2021). Assessment of alerting, orienting, and executive control in persons with aphasia using the Attention Network Test. *Aphasiology*, 35, 1318-1333. <https://doi.org/10.1080/02687038.2020.1795077>

Landis, J. R., & Koch, G. G. (1977). The measurement of observer agreement for categorical data. *Biometrics*, 33(1), 159. <https://doi.org/10.2307/2529310>

Lezak, M. D., Howieson, D. B., Bigler, E. D., Tranel, D. (2012). *Neuropsychological Assessment*. Oxford University Press.

Mayer, J. F., & Murray, L. L. (2012). Measuring working memory deficits in aphasia. *Journal of Communication Disorders*, 45(5), 325–339. doi: 10.1016/j.jcomdis.2012.06.002

Marinelli, C. V., Spaccavento, S., Craca, A., Marangolo, P., & Angelelli, P. (2017). Different cognitive profiles of patients with severe aphasia. *Behavioural Neurology*. 2017;2017:3875954. doi: 10.1155/2017/3875954.

Maxwell, R., O'Brien, M., O'Donnell, D., Christophers, L., & Kroll, T. (2021). Using a multidisciplinary approach to reveal decision-making capacity within acute care for an individual with aphasia. *SAGE Open Medical Case Reports*, 9. doi: 10.1177/2050313X211027098.

McNeil, M., Hula, W., & Sung, J. E. (2011). The role of memory and attention in aphasic language performance. In J. Guendouzi, F. Loncke, & M. J. Williams (Eds.), *The handbook of psycholinguistic and cognitive processes: Perspectives in communication disorders* (pp. 551–577). Psychology Press.

Minkina, I., Salis, C., & Martin, N. (2018). Short-term and working memory deficits in aphasia: current issues in theory, evidence, and treatment. *Journal of Neurolinguistics*, 48, 1–3. <https://doi.org/10.1016/j.jneuroling.2018.07.001>

Murray, L. L. (2002). Attention deficits in aphasia: presence, nature, assessment, and treatment. *Seminars in Speech and Language*, 23(2), 107–116. doi: 10.1055/s-2002-24987

Murray, L., Keeton, R., & Karcher, L. (2006). Treating attention in mild aphasia: evaluation of attention process training-II. *Journal of Communication Disorders*, 39(1), 37–61. doi: 10.1016/j.jcomdis.2005.06.001.

Murray, L. L. (2012). Attention and other cognitive deficits in aphasia: presence and relation to language and communication measures. *American Journal of Speech-Language Pathology*, 21(2), S51–64. doi: 10.1044/1058-0360(2012/11-0067)

Murray, L. L. (2017). Design fluency subsequent to onset of aphasia: a distinct pattern of executive function difficulties? *Aphasiology*, 31(7), 793–818. <https://doi.org/10.1080/02687038.2016.1261248>

Murray, L., Salis, C., Martin, N., & Dralle, J. (2018). The use of standardised short-term and working memory tests in aphasia research: a systematic review. *Neuropsychological Rehabilitation*, 28(3), 309–351. doi: 10.1080/09602011.2016.1174718.

Nasreddine, Z., Phillips, N., Bédirian, V., Charbonneau, S., Whitehead, V., Collin, I., Cummings, J., & Chertkow, H. (2005). The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society*, 53(4), 695–699. doi: 10.1111/j.1532-5415.2005.53221.x.

National Clinical Guideline for Stroke for the UK and Ireland (2023). [Rehabilitation and recovery - psychological effects of stroke - National Clinical Guideline for Stroke](https://www.strokeguideline.org/chapter/psychological-effects-of-stroke) [https://www.strokeguideline.org/chapter/psychological-effects-of stroke](https://www.strokeguideline.org/chapter/psychological-effects-of-stroke)

Nickels, L., Tessaro, B., Hameau, S., & Salis, C. (2024) The complex relationship between cognition & language: illustrations from acquired aphasia. In M. J. Ball, N. Muller & L. Spencer (Eds.), *Handbook of Clinical Linguistics*. (2nd ed.). Wiley.

Osterrieth, P. (1944). Le Test de copie d'une figure complexe. *Archives de Psychologie*, 30, 206–356.

Peñaloza, C., Martin, N., Laine, M., & Rodríguez-Fornells, A. (2022). Language learning in aphasia: a narrative review and critical analysis of the literature with implications for language therapy. *Neuroscience & Biobehavioral Reviews*, 104825. doi: 10.1016/j.neubiorev.2022.104825

R Core Team (2022). R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

Raven, J. C. (1962). *Coloured Progressive Matrices*. The Psychological Corporation.

Reitan, R. (1992). *Trail Making Test: manual for administration and scoring*. Reitan Neuropsychology Laboratory.

Rey, A. (1941). L'examen psychologique dans les cas d'encéphalopathie traumatique. *Archives de Psychologie*, 28, 286–340.

Robertson, I. H., Ward, T., Ridgeway, V., & Nimmo-Smith, I. (1994). *The Test of Everyday Attention*. Thames Valley Test Company.

Rohde, A., Worrall, L., & le Dorze, G. (2013). Systematic review of the quality of clinical guidelines for aphasia in stroke management. *Journal of Evaluation in Clinical Practice*, 19(6), 994–1003. <https://doi.org/10.1111/jep.12023>

Sachdev, P., Blacker, D., Blazer, D., Ganguli, M., Jeste, D., Paulsen, J., & Petersen, R. (2014). Classifying neurocognitive disorders: the DSM-5 approach. *Nature Reviews Neurology*, 10(11), 634–642. doi: 10.1038/nrneurol.2014.181

Salis, C., Murray, L., & Bakas, K. (2018). An international survey of assessment practices for short-term and working memory deficits in aphasia. *American Journal of Speech-Language Pathology*, 27(2), 574–591. doi: 10.1044/2017_AJSLP-17-0057

Schumacher, R., Halai, A. D., & Lambon Ralph, M. A. (2019). Assessing and mapping language, attention and executive multidimensional deficits in stroke aphasia. *Brain*, 142, 3202–3216. <https://doi.org/10.1093/brain/awz258>

Shiggins, C., Ryan, B., O'Halloran, R., Power, E., Bernhardt, J., Lindley, R. I., McGurk, G., Hankey, G. J., & Rose, M. L. (2022). Towards the consistent inclusion of people with aphasia in stroke research irrespective of discipline. *Archives of Physical Medicine and Rehabilitation*, 103(11), 2256–2263. doi: 10.1016/j.apmr.2022.07.004.

Simic, T., Bitan, T., Turner, G., Chambers, C., Goldberg, D., Leonard, C., & Rochon, E. (2020). The role of executive control in post-stroke aphasia treatment. *Neuropsychological Rehabilitation*, 30(10), 1853–1892. doi: 10.1080/09602011.2019.1611607

Simic, T., & Rochon, E. (2018). Executive control is a positive predictor of treatment maintenance following anomia therapy. *Aphasiology*, 32(sup1), 212–214. <http://dx.doi.org/10.1080/02687038.2018.1485861>

Simic, T., Rochon, E., Greco, E., & Martino, R. (2019). Baseline executive control ability and its relationship to language therapy improvements in post-stroke aphasia: a systematic review. *Neuropsychological Rehabilitation*, 29(3), 395–439. doi: 10.1080/09602011.2017.1307768

Snaphaan, L., & de Leeuw, F. E. (2007). Poststroke memory function in nondemented patients: a systematic review on frequency and neuroimaging correlates. *Stroke*, 38(1), 198–203. doi: 10.1161/01.STR.0000251842.34322.8f

Srikanth, V., Thrift, A., Saling, M., Anderson, J., Dewey, H., Macdonell, R., & Donnan, G. (2003). Increased risk of cognitive impairment 3 months after mild to moderate first-ever

stroke: community-based prospective study of nonaphasic English-speaking survivors. *Stroke*, 34(5), 1136–1143. doi: 10.1161/01.STR.0000069161.35736.39

Sternberg, R. J., & Sternberg, K. (2012). *Cognitive Psychology* (6th ed.). Wadsworth.

Strauss, E., Sherman, E. M. S., Spreen, O. (2006). *A Compendium of Neuropsychological Tests: administration, norms, and commentary*. Oxford University Press.

Suleman, S., & Hopper, T. (2016). Decision-making capacity and aphasia: speech-language pathologists' perspectives. *Aphasiology*, 30(4), 381–395. <https://psycnet.apa.org/doi/10.1080/02687038.2015.1065468>

Swinburn, K., Porter, G., & Howard, D. (2005). *The Comprehensive Aphasia Test*. Psychology Press.

Tessaro, B., Salis, C., Hameau, S. & Nickels, L. (2024). How cognition has been assessed in research with people with aphasia: a systematic literature review. *Language, Speech, and Hearing*, 27, 176-190. <https://doi.org/10.1080/2050571X.2024.2315801>

The Australian and New Zealand Clinical Guidelines for Stroke Management” (2024). Retrieved on 28 November 2024

<https://informme.org.au/guidelines/living-clinical-guidelines-for-stroke-management>

Thompson, H, Almaghyuli, A, Noonan, K., Barak, O, Lambon Ralph, M, & Jefferies, E. (2018). The contribution of executive control to semantic cognition: convergent evidence from semantic aphasia and executive dysfunction. *Journal of Neuropsychology*, 12(2), 312–340. doi: 10.1111/jnp.12142

Thomson, J., Gee, M., Sage, K., & Walker, T. (2018). What ‘form’ does informal assessment take? a scoping review of the informal assessment literature for aphasia. *International Journal of Language and Communication Disorders*, 53(4), 659–674. doi: 10.1111/1460-6984.12382.

van de Sandt-Koenderman, W. M. E., Wiegers, J., Wielaert, S. M., Duivenvoorden, H. J., & Ribbers, G. M. (2007). High-tech AAC and severe aphasia: candidacy for TouchSpeak (TS). *Aphasiology*, 21(5), 459–474. <https://doi.org/10.1080/02687030601146023>

Varkanitsa, M., Godecke, E., & Kiran, S. (2023). How much attention do we pay to attention deficits in poststroke aphasia? *Stroke*, 54(1), 55-66. <https://doi.org/10.1161/STROKEAHA.122.037936>

Verna, A., Davidson, B., & Rose, T. (2009). Speech-language pathology services for people with aphasia: a survey of current practice in Australia. *International Journal of Speech-Language Pathology*, 11(3), 191–205. <https://doi.org/10.1080/17549500902726059>

Wall, K. J., Cumming, T. B., Koenig, S. T., Pelecanos, A. M., & Copland, D. A. (2018). Using technology to overcome the language barrier: the Cognitive Assessment for Aphasia App. *Disability and Rehabilitation*, 40(11), 1333–1344. <https://psycnet.apa.org/doi/10.1080/09638288.2017.1294210>

Wechsler, D. (1997). *Wechsler Memory Scale–III*. The Psychological Corporation.

Wong, G. K. C., Ngai, K., Lam, S. W., Wong, A., Mok, V., & Poon, W. S. (2013). Validity of the Montreal Cognitive Assessment for traumatic brain injury patients with intracranial haemorrhage. *Brain Injury*, 27(4), 394–398. doi: 10.3109/02699052.2012.750746.

Zakariás, L., Kelly, H., Salis, & C., Code. (2019). The methodological quality of short-term/working memory treatments in poststroke aphasia: a systematic review. *Journal of Speech, Language & Hearing Research*, 62(6), 1979–2001. doi: 10.1044/2018_JSLHR-L-18-0057.

Zimmermann, P., & Fimm, B. (1994). *Test d'évaluation de l'attention (TEA)*. Wurselen, Germany: Psytes.

Tables

Table 1. Barriers to assessing cognition in aphasia reported (n of respondents = 158)

<i>Theme</i>	<i>Sub-themes</i>	<i>Illustrative quotes</i>
<i>(Number of responses)</i>	<i>(Number of responses)</i>	
Materials (169)	Appropriateness of assessment and resources (67)	<i>'Lack of appropriate tests/resources.'</i>
	Access to assessments and resources (29)	<i>'Sometimes, it is not accessible to me to get the means for a relevant evaluation.'</i>
	Materials are highly dependent on language (26)	<i>'Concerns which tests might be appropriate to assess cognitive functions in people with aphasia since most tests are heavily based on language.'</i>
	Lack of normed and standardised materials (16)	<i>'Having tools available that are normed and adapted for people with aphasia.'</i>
	Restrictions on use of assessments (7)	<i>'No access to psychology services which would be helpful to analyse assessment results.'</i>
	Lack of functional assessments (5)	<i>'Inability to assess patient functionally in a clinic setting (how they do pointing to safety hazards in a picture may not reflect their own judgement/safety out in the community).'</i>
	Cost of assessments (5)	<i>'Availability of measures and tools (and training in them). Barriers include 'we don't have that in our department and don't plan to get it' to too expensive to some publishers' restraint of trade, limiting use to neuropsychologists or others with at least a doctorate.'</i>

Professional role (116)	Especially for severe aphasia (4)	<i>'Standardised accessible cognitive and linguistic assessment for severe aphasia.'</i>
	Multilingual assessments (4)	<i>'Availability of test batteries in different Indian languages.'</i>
	Complexity of test instructions (3)	<i>'Using standardised assessment as outlined in manual can mean that the instructions are too complex for someone with aphasia. Modifying / simplifying the instruction can allow the person to answer the question / carry out the task but this impacts on the standardisation / validity of the result.'</i>
	Over reliance on scores of formal tests (2)	<i>'Overreliance on some cognitive tests which do not adequately look at aphasia.'</i>
	Lack of consensus (1)	<i>'Lack of a widely agreed upon test or tests to use for this purpose -- CLQT may come closest to filling this purpose.'</i>
	Clarity of MDT roles (43)	<i>'Lack of understanding of colleagues - misconceptions that you can't assess cognition effectively in people with aphasia. I find this in particular comes from neuropsychologists who leave these patients in the 'too hard basket' for the OTs and SLTs to assess.'</i>
	Time limitations (32)	<i>'Lack of time and have to prioritize between language and cognition for sake of evaluation report.'</i>
	Skills limitations (12)	<i>'Not having the knowledge and skills to assess as here it is the Occupation Therapists domain.'</i>
	Access to neuropsychology (10)	<i>'No access to psychology services which would be helpful to analyse assessment results.'</i>
	MDT communication (12)	<i>'Understanding the person with cognitive communication profile is the foundation to making progress, once you have achieved awareness you really start to make progress - getting the whole team on the same page really helps the person with cog communication.'</i>
	Insurance constraints (4)	<i>'Insurance does not pay that practice to a speech therapist.'</i>

	Occupational Therapists access cognition (4)	<i>'Cognitive assessment is the territory of Occupational Therapists in my workplace.'</i>
	Staff availability (4)	<i>'I am under resourced (staffing/competing workload priorities) to seek an increased role in cognitive assessment.'</i>
	Importance of cognitive assessment (3)	<i>'Difficulty justifying importance/role of cognitive assessment to MDT.'</i>
	Knowledge and experience with assessment (29)	<i>'Insufficient training/education in the multidisciplinary team to understand reports and evaluation results.'</i>
	Differential diagnosis in the presence of aphasia (20)	<i>'The presence of aphasia makes it difficult to determine whether any concerns are due to language or cognition.'</i>
Knowledge and skills (74)	Understanding of cognition (13)	<i>'A lack of agreement on the state of cognition in aphasia and the relationships that exist.'</i>
	Understanding of aphasia (7)	<i>'Healthcare providers not being sufficiently educated about aphasia.'</i>
	Awareness of comorbidities (5)	<i>'Reduced awareness and understanding by other professionals of cognitive difficulties for people with aphasia, particularly attention skills.'</i>
People with aphasia (51)	Impact of aphasia on assessment of cognition (33)	<i>'Severity of aphasia which may impair verbal responses for cognitive assessment instructions.'</i>
	Awareness of deficits (4)	<i>'Presence of poor insight into an aphasia in early stroke can make engagement in cognitive assessment challenging.'</i>

Engagement with assessment (3)	<i>'Severely depressed and/or apathetic subjects.'</i>
Fatigue (3)	<i>'Fatigue. This is a big issue. Often one cannot assess for longer than 30 min at a time in the early stages after stroke.'</i>
Impact of motor deficits on assessment of cognition (3)	<i>'Other stroke deficits affecting deficits e.g., if dominant hand is affected can't hold a pen, or if there is a limb apraxia may affect gestural use or assessment on a motor or functional task.'</i>
Knowledge of cognition (3)	<i>'Patient's and caregiver's knowledge and understanding of aphasia and cognitive impairments.'</i>
Unilateral spatial neglect (2)	<i>'Hemi attention / unilateral spatial neglect.'</i>
Estimating pre-morbid abilities (3)	<i>'Sometimes difficult to access families and therefore know about the patient's prior interests and capacities.'</i>
Location of assessment process (3)	<i>'Many cognitive functions are best assessed in the patients known environment (at home) and I meet them at the hospital'</i>

Notes. Numbers in brackets denote number of responses that were coded at each sub-theme. This question elicited 260 responses, but each response can be coded at more than one theme. Please note that the illustrative quotes may also have been coded at other themes and sub-themes.

¹ 17 responses were classified as unclear.

Table 2. Assessment tools used and constructs measured reported by respondents

Tool	Test frequency ¹	Complex attention	Executive functions	Learning and memory	Language	Perceptual-motor function	Social Cognition
Cognitive Linguistic Quick Test ²	44	24	22	23	9	16	0
Trail Making Test ³	27	13	18	0	0	2	0
Montreal Cognitive Assessment ⁴	26	4	2	4	3	2	0
Raven's Coloured Progressive Matrices ⁵	20	0	16	0	0	3	0
Mini Mental State Examination ⁶	14	3	1	4	2	2	0
Rey Complex Figure ⁷	14	1	5	10	0	10	0
Comprehensive Aphasia Test ⁸	13	2	4	6	1	3	0
Oxford Cognitive Screen ⁹	12	4	2	6	5	6	0
Test of Everyday Attention ¹⁰	12	10	1	1	0	0	0
The Butt Non-Verbal Reasoning Test ¹¹	12	0	8	0	0	0	0
Test of Attentional Performance ¹²	11	9	3	0	0	2	0
Weschler Adult Intelligence Scale ¹³	11	0	3	0	0	0	0
Wisconsin Card Sorting Test ¹⁴	11	1	10	1	0	0	0
Informal assessment	14	7	3	6	0	3	1

Notes. ¹The number of times each test was indicated by respondents. The numbers in each cognitive domain may not sum up to the numbers in ¹ as not all respondents indicated which construct they aim to measure with the cited tool. ²Helm-Estabrooks (2002). ³Reitan (1992). ⁴Nasreddine et al. (2005). ⁵Raven (1962). ⁶Folstein et al. (1975). ⁷Rey (1941). ⁸Swinburn et al. (2005). ⁹Demeyere et al. (2015). ¹⁰Robertson et al. (1994).

¹¹Butt and Bucks (2002). ¹²Zimmermann and Fimm (1994). ¹³Wechsler (1997). ¹⁴Berg, 1948.

Table 3. Adaptions made when assessing cognition in people with aphasia as compared to people without aphasia (n of respondents = 155)

<i>Theme (number of responses)</i>	<i>Sub-theme (number of responses)</i>	<i>Illustrative quotes</i>
Modifications to the task (172)	Reduce linguistic load (34)	<i>'Talk less and more slowly and use shorter sentences and'</i>
	Provide extra support (33)	<i>'Provide visual aids (e.g., number charts to point at/reference) to help them respond with a number).'</i>
	Modify instructions (28)	<i>'Aphasia-friendly instructions (reduce complexity, simplify instructions as needed).'</i>
	Adaptation to testing methodology (14)	<i>'Often break standardization of assessment to ensure ad'</i>
	Allow more time (12)	<i>'Repetition of commands, accept delayed responses if au'</i>
	Ensure understanding (6)	<i>'I take care ascertaining that they understand me well, I use shorter sentences, I use a communication board or a notepad if'</i>
	Slow speech rate (5)	<i>'Slow speed at which instructions are given.'</i>
	Pace the test (4)	<i>'Assessments divided into multiple sessions when attent'</i>
	Adaptations for other comorbidities (3)	<i>'Positioning visual stimuli on a preferred side, assuming'</i>
	Multimodal input (3)	<i>'I may pair visual, verbal, and imitation techniques to fa'</i>
Selection of tools (67)	Aphasia-friendly assessment tools (41)	<i>'Select assessments that are appropriate to the client's le'</i> <i>person has limited verbal output, then I select a nonverb'</i>
	Functional assessments (15)	<i>'I tend to focus towards assessing cognition within every'</i> <i>me to not highly depend upon an individual's language c'</i>
	Informal assessment (5)	<i>'Specific subtests may be omitted and/or informal assess'</i>
	Formal assessments (4)	<i>'My assessment would include more formalized test pro'</i>
	shortened task versions (1)	<i>'If I'm using an experimental task, I usually use a shorte'</i> <i>with aphasia.'</i>
Assessment conditions (35)	Patient needs (1)	<i>'Consider what the needs of the patient/caregivers are. I'</i> <i>and the next steps.'</i>
	Case specific requirements (15)	<i>'The protocol needs to be adapted to the language skills'</i> <i>will differ from person to person.'</i>
	Consider aphasia in assessments interpretation (9)	<i>'I take into consideration how their language impairment'</i> <i>performance on a test.'</i>

Work with MDT (5)	<i>'Executive function and many other deficits --> ask the p opinions on strategies and relevance of therapy.'</i>
Avoid distractions (2)	<i>'Choose a space that had least distraction, try to engena</i>
Consider fatigue (2)	<i>'Degree of fatigue noted may also impact tests selected.'</i>
Face to face (1)	<i>'Add visuals, change pacing, face-to-face, block informa distractions.'</i>
Support from caregiver (1)	<i>'Time, less planned, time of day, around meds, with fam</i>

Notes. Numbers in brackets denote number of answers that were coded at each theme.

MDT:Multidisciplinary team

Table 4: Actionable insights for the academic and clinical fields

<i>Finding</i>	<i>Insight/recommendation for the field</i>
Lack of guidance for reliable assessment of cognition in people with aphasia (PWA)	• Develop consensus guidance amongst international exp
Current assessment tools are inappropriate to use with people with aphasia	• Develop/update assessment tools that are suitable for pe
People with aphasia excluded from research that requires cognitive assessments.	• Account for exclusion of PWA when reviewing eviden • Researchers should include new assessments developed
Professionals reliant on adapting assessment tools for people with aphasia.	• Use caution when comparing performance to normative from standardization probably invalidates comparisons • Researchers must consider how this impacts study repli and generalisability of results. • Researchers and clinicians must consider the language l tool used. • Collect normative data for people living with aphasia.
Clarity around roles within the multidisciplinary team is needed.	• Update/create guidelines around scope of practice. • Inter-professional learning on this topic should be prior training programs, peak organizations, and best practice guidelines
And	
Perceived lack of appropriate training for conducting cognitive assessments in PWA.	

Figures

Figure 1. The cognitive domains that respondents indicated they included in their definition of cognition.

Figure 2. Considerations that influence respondents' decisions to assess cognition in aphasia.

Figure 3. Respondent's ratings of the importance of assessing cognition for specific purposes.

Figure 4. Responses to the multiple-choice questions 'Who should assess cognition in aphasia?' and 'Who should discuss the results?'

Figure 5. Responses to the question 'What factors do you consider when choosing tools to assess cognition in aphasia? (n of respondents = 150)

Figure 1. The cognitive domains that respondents indicated they included in their definition of cognition

Alt text for figure 1 [34 words]: A horizontal bar graph plotting the percentage of respondents that identified different domains (10 identified in total) as falling into a definition of cognition. Only praxis was reported by less than 75% of respondents.




Figure 2. Considerations that influence respondents' decisions to assess cognition in aphasia

Alt text for figure 2 [25 words]: A horizontal bar graph plotting the percentage of respondents that identified different considerations (13 were identified in total) that influence their decision to assess cognition.




Figure 3. Respondent's ratings of the importance of assessing cognition for specific purposes

Alt text for figure 3 [53 words]: A horizontal stacked bar graph plotting the respondents ratings of the importance of assessing cognition for seven different purposes identified. that identified ten different domains as falling into a definition of cognition. Most reported assessment thought assessment of cognition was important for providing strategies, returning to employment, determining dysfunction and for treatment planning.




Figure 4. Responses to multiple-choice questions: ‘Who should assess cognition in aphasia?’ and ‘Who should discuss the results?’

Alt text for figure 4 [53 words]: A horizontal stacked bar graph plotting the respondents responses to the multiple choice questions ‘Who should assess cognition in aphasia?’ and ‘Who should discuss the results?’.

Most respondents reported that the speech and language pathologist and the neuropsychologist should be responsible for both questions.




Figure 5. Summary of responses to the question ‘What factors do you consider when choosing tools to assess cognition in aphasia?’ (n of respondents = 150)

Alt text for figure 5 [24 words]: Four boxes titled, tools, impairment, person with aphasia and assessment. In each box are the associated factors respondents reported considering when choosing assessment tools.

<i>TOOLS</i> <i>(n = 143/95%)</i>	<i>IMPAIRMENT</i> <i>(n = 71/47%)</i>	<i>PERSON WITH APHASIA</i> <i>(n = 54/36%)</i>	<i>ASSESSMENT</i> <i>(n = 51/34%)</i>
⇒ Administration considerations (99): suitability for aphasia (40), time restrictions (29), availability of tools (20), complexity of administration (4), demographic appropriateness (4), possibility to administer remotely (2) ⇒ Test construction and psychometrics (21) ⇒ Ecological validity (8) ⇒ Comprehensiveness (6) ⇒ Evidence-based tools (4) ⇒ Tasks that assess cognitive functions relevant for language (4) ⇒ Screening (1)	⇒ Severity of aphasia (51) ⇒ Comorbidities (15) ⇒ Lesion information (2) ⇒ Stroke severity (2)	⇒ Engagement in assessment process (20) ⇒ Goals (17) ⇒ Linguistic and cultural appropriateness (5) ⇒ Awareness of aphasia (4) ⇒ Premorbid abilities (3) ⇒ Age (2)	⇒ Need for clarity regarding differential diagnosis (17) ⇒ Purpose of cognitive assessment (13) ⇒ Constructs to be tested (8) ⇒ Clinical management planning (7) ⇒ MDT availability (5) ⇒ Assessment history (4)

Notes. Numbers in brackets denote number of responses that were coded for each sub-theme.