



1 A meta-analysis of groundwater contamination by nitrates at the 2 African scale

3 Issoufou Ouedraogo¹ and Marnik Vanclooster¹

4 ¹Earth and Life Institute, Université catholique de Louvain, Croix du Sud 2, Box 2, B-1348 Louvain-la-Neuve, Belgium.

5 Correspondence to: I. Ouedraogo (ouediss6@yahoo.fr)

6 **Abstract.** Contamination of groundwater with nitrate poses a major health risk to millions of people around Africa.
7 Assessing the space-time distribution of this contamination, as well as understanding the factors that explain this
8 contamination is important to manage sustainable drinking water at the regional scale. This study aims assessing the
9 variables that contribute to nitrate pollution in groundwater at the pan African scale by statistical modeling. We compiled
10 a literature database of nitrate concentration in groundwater (around 250 studies) and combined it with digital maps of
11 physical attributes such as soil, geology, climate, hydrogeology and anthropogenic data for statistical model
12 development. The maximum, medium and minimum observed nitrate concentrations were analysed. In total, 13
13 explanatory variables were screened to explain observed nitrate pollution in groundwater. For the mean nitrate
14 concentration, 4 variables are retained in the statistical explanatory model: (1) Depth to groundwater (shallow
15 groundwater, typically <50m); (2) Recharge rate; (3) Aquifer type; and (4) Population density. The former three
16 variables represent intrinsic vulnerability of groundwater systems towards pollution, while the latter variable is a proxy
17 for anthropogenic pollution pressure. The model explains 65% of the variation of mean nitrate contamination in
18 groundwater at the pan Africa scale. Using the same proxy information, we could develop a statistical model for the
19 maximum nitrate concentrations that explains 42% of the nitrate variation. For the maximum concentrations, other
20 environmental attributes such as soil type, slope, rainfall, climate class and region type improves the prediction of
21 maximum nitrate concentrations at the pan African scale. As to minimal nitrate concentrations, in the absence of normal
22 distribution assumptions of the dataset, we do not develop a statistical model for these data. The data based statistical
23 model presented here represents an important step toward developing tools that will allow us to accurately predict nitrate
24 distribution at the African scale and thus may support groundwater monitoring and water management that aims
25 protecting groundwater systems. Yet they should be further refined and validated when more detailed and harmonized
26 data becomes available and/or combined with more conceptual descriptions of the fate of nutrients in the hydrosystem.

27

28



29 1 Introduction

30

31 Nitrate contamination of groundwater is a common problem in many parts of the world. Elevated nitrate concentrations
32 in drinking water can cause methemoglobinemia in infants and stomach cancer in adults (Yang et al., 1998; Knobeloch
33 et al., 2000; Hall et al., 2001). As such, the World Health Organization (WHO) has established a maximum contaminant
34 level (MCL) of 50 mg/L NO₃ (WHO, 2004). Nitrate in groundwater is generally from anthropogenic origin and
35 associated with leaching of nitrogen from agriculture plots or from waste and sewage sanitation systems. The heavy use
36 of nitrogenous fertilizers in cropping system is the largest contributor to anthropogenic nitrogen in groundwater
37 worldwide (Suthar et al., 2009). In particular, shallow aquifers in agricultural fields are highly vulnerable to nitrate
38 contamination (Böhlke, 2002; Kyoung-Ho et al, 2009). According to Spalding and Exner (1993), nitrate may be the
39 most widespread contaminant of groundwater.

40

41 In Africa, groundwater is recognized as playing a very important role in the development agenda. However, according
42 to Xu and Usher, (2006), degradation of groundwater is the most serious water resources problem in Africa. The two
43 main threats are overexploitation and contamination (MacDonald et al., 2013). Indeed, based on a review of 29 papers
44 from 16 countries, Xu and Usher (2006), have identified major groundwater pollution issues in Africa, considering the
45 following order of importance as follows: (1) nitrate pollution, (2) pathogenic agents, (3) organic pollution, (4)
46 salinization, and (5) acid mine drainage. These authors have identified that the major sources of groundwater
47 contamination are related to on-site sanitation, to the presence of solid waste dumpsites, including household waste pits,
48 to infiltration of surface water, to agricultural activities, to the presence of petrol service stations (underground storage
49 tanks), and to the mismanagement of wellfields. Nitrate contamination of groundwater is a problem that commonly
50 occurs in Africa, as illustrated in the studies for Algeria (Rouabhia et al., 2010; Messameh et al., 2014), Tunisia (Hamza
51 et al., 2007; Anane et al., 2014), Morocco (Bricha et al., 2007; Fetouani et al., 2008; Benabbou et al., 2014), Senegal
52 (Sall and Vanclooster, 2009; Diédhiou et al., 2012), Ivory Coast (Loko et al., 2013; Eblin et al., 2014), Ghana (Tay and
53 Kortatsi, 2008; Fianko et al., 2009), Nigeria (Wakida and Lerner, 2005; Akoteyon and Soladoye, 2011; Obinna et al.,
54 2014), South Africa (Maherry et al., 2009; Musekiwa and Majola, 2013), Ethiopia (BGS, 2001; Bonetto et al., 2005)
55 and Zambia (Wakida and Lerner, 2005). Several of these studies showed that pollution from anthropogenic activities
56 is the main source of high and variable nitrate levels. For example, Comte et al., (2012) illustrates that the groundwater
57 situated in the Quaternary sandy aquifer of the peninsula of Dakar is under strong anthropogenic pressure from the city
58 of Dakar, resulting in important nitrate loadings. Such contamination problems are often retrieved in many metropolises
59 in Africa. Notwithstanding the availability of all these studies at the local, regional or country level, no comprehensive
60 and synthetic study of nitrate contamination of groundwater at the scale of the African continent has been presented in
61 the literature. Assessing large scale groundwater contamination with nitrates is important for the planning of the large
62 scale groundwater exploitation programs and for designing transboundary water management policies. It yields also
63 important baseline information for monitoring progress in the implementation of the UN SDGs for water. It increase
64 awareness of citizens, international agencies and authorities (e.g., FAO, UNEP, and OECD, Water Sanitation for Africa
65 (WSA)) on the environmental factors likely to be significant to groundwater contamination. However, making an
66 appropriate pan-African synthesis of nitrate contamination of groundwater remains a scientific and technical challenge,
67 given the non-homogeneity of the nitrate monitoring programmes and the absence of administrative and institutional
68 capacity to collect and diffuse the data at the pan African scale. A concept that partially helps solving this urgent data
69 management problem is the concept of groundwater vulnerability. Groundwater vulnerability for nitrate contamination



70 is an expression of the likelihood that a given groundwater body will be negatively affected by nitrate contamination.
71 Given that the vulnerability is a likelihood, it is only an expression of the potential degradation of groundwater and
72 hence a proxy of groundwater contamination by nitrates. Groundwater vulnerability can be assessed based on available
73 generic data. It does therefore not depend on a strong and operational pan African groundwater quality monitoring
74 capacity. In this paper we propose and implement a methodology for assessing vulnerability of groundwater
75 contamination by nitrates at the pan African scale. We further consider nitrate in this study as a proxy for overall
76 groundwater pollution, which is consistent with the view of the US EPA (EPA, 1996).

77

78 In general, there are three categories of models for the assessment of groundwater vulnerability: (1) index methods or
79 subjective rating methods, (2) statistical methods and (3) process based modelling methods. Index-and-overlay methods
80 are one set of subjective rating methods that utilize the intersection of regional attributes with the qualitative
81 interpretation of data by indexing parameters and assigning a weighting scheme. The most widely used index method
82 is DRASTIC (Aller et al., 1985). Unfortunately, index methods are based on subjective rating methods (Focazio et al.,
83 2002) and should preferably be calibrated using measured proxies of vulnerability (Kihumba et al., 2015; Ouedraogo et
84 al., 2015). When a groundwater monitoring dataset is available, formal statistical methods can be used to integrate
85 groundwater contamination data directly in the vulnerability assessment. Finally, process-based methods refer to
86 approaches that explicitly simulate the physical, chemical and biological processes that affect contaminant behaviour in
87 the environment. They comprise the use of deterministic or stochastic process-simulation models, eventually linked to
88 physically based field observations (e.g., Coplen et al., 2000). Physically process-based methods are typically applied
89 at small scales, mostly to define well protection zones, rather than to assess groundwater vulnerability at broader scales
90 (Frind et al., 2006). A well-known example is the use of a physical based groundwater model (e.g. MODFLOW,
91 Harbaugh et al., 2000) that solve the governing equations of groundwater flow and solute transport. Such models have
92 explicit time steps and are often used to determine the time scales of contaminant transport to wells and streams, in
93 addition to the effects of pumping. However, they also have many parameters that require estimation. In this paper, we
94 use statistical models to assess vulnerability of groundwater systems towards nitrate pollution.

95

96 Formal statistical methods have often be deployed to assess vulnerability of groundwater at national and regional scales.
97 They are also often used to discriminate contaminant sources and to identify factors contributing to contamination
98 (Kolpin, 1997; Nolan and Hitt, 2006). Many authors used multiple linear regression (MLR) techniques. For example,
99 Bauder et al., (1993) investigated the major controlling factors for nitrate contamination of groundwater in agricultural
100 areas using MLR of land uses, climate, soil characteristics, and cultivations types. MLR was also used to relate pesticide
101 concentrations in groundwater to the age of the well, land use around the well, and the distance to the closest possible
102 source of pesticide contamination (Steichen, et al., 1988). Boy-Roura et al., (2013) used MLR to assess nitrate pollution
103 in the Osona region (NE Spain). Amini et al., (2008a) and Amini et al., (2008b) used MLR and Adaptive Neuro-Fuzzy
104 Inference System (ANFIS), a general non-linear regression technique, to study the global geogenic fluoride
105 contamination in groundwater and the global geogenic arsenic contamination in groundwater respectively. MLR has
106 the strong advantage that regression coefficients can directly be interpreted in terms of importance of explaining factors.
107 Many studies linking nitrate occurrence in groundwater to spatial variables have employed logistic regression (Hosmer
108 and Lameshow, 1989; Eckardt and Stackkelberg, 1995; Tesoriero and Voss, 1997; Gardner and Vogel, 2005; Winkel
109 et al., 2008; and Mair and El-kadi, 2013). According to Kleinbaum. (1994), MLR is conceptually similar to logistic
110 regression. Other authors have used more sophisticated approaches such as Bayesian methods (Worrall and Besien,



2005; Mattern et al., 2012) and, more recently, classification and regression tree modelling approaches (Burow et al., 2010; Mattern et al., 2012). Yet, according to our knowledge, a statistical model of groundwater nitrate contamination at the pan African scale does not exist yet.

In the present study, we used MLR techniques to assess the vulnerability of nitrate groundwater pollution at the pan African scale. To this end, we compiled a pan African groundwater pollution data base from the literature and combined it with environmental attributes inferred from a generic data basis. The generic data basis was developed in a former study to assess vulnerability using the DRASTIC index method (Ouedraogo et al., 2016). MLR models were subsequently identified to explain quantitatively the logtransformed observed nitrate contamination in terms of generic environmental attributes and finally, the regression models were interpreted in terms of characteristics of contaminants sources and hydrogeology of the African continent.

2 Study area

We studied the vulnerability of groundwater systems for nitrate contamination at the scale of the African continent. Groundwater is Africa's most precious natural resource, providing reliable water supplies to at least a third of the continent's population (MacDonald, 2010). However, the African continent is not blessed by a large quantity of groundwater resources, because it is the World's second-driest continent after Australia and water resources are limited. MacDonald et al., (2012) have estimated the volume of groundwater resource in Africa at 0.66 million km³.

Africa has a vast array of drainage networks, the most important ones are the Nile River, which drains northeast and empties into the Mediterranean Sea. The Congo River drains much of central Africa and empties into the Atlantic Ocean. The Niger River is the principal river of western Africa; it is the third-longest river after the Nile and the Congo River and empties into the Atlantic Ocean. Southern Africa is drained by the Zambezi River. Lake Chad constitutes one of the largest inland drainage areas of the continent. Other major lakes located in the east of Africa include Lake Tanganyika and Lake Victoria.

The elevation of Africa varies from below sea level to 5825 m above sea level. The average elevation is approximately 651 m (Ateawung, 2010). The geology of the African continent contains 13 lithological classes (Fig.1) with varying coverages: evaporites (0.6%), metamorphic rocks (27.6%), acid plutonic rocks (1.1%), basic plutonic rocks (0.2%), intermediate plutonic rocks (0.1%), carbonates sedimentary rocks (9.4%), mixed sedimentary rocks (6.4%), siliciclastic sedimentary rocks (16.4%), unconsolidated sediments (35.1%), acid volcanic rocks (0.1%), basic volcanic rocks (3.3%), intermediate volcanic rocks (0.6%) and water bodies (0.9%) (Hartmann and Moosdorf, 2012). The lithology describes the geochemical, mineralogical and physical properties of rocks.

3 Data and methods

3.1 Nitrate contamination data



For a large part of Africa there is very little, or no systematic monitoring of groundwater. In the absence of data systematic monitoring program, we compiled nitrate pollution data at the pan African scale from different literature sources. We considered approximately 250 published papers on nitrate contamination of groundwater in Africa. We consulted the web of sciences (Scopus™, Sciences Direct™, Google™, and Google Scholar™) and available books. Fig. 2 shows the spatial distribution of the considered field studies. Table 1 outlines criteria used in the web search.

153

3.2 Data quality evaluation

155

We used the following additional criteria to select the study:

- i. the publication should explicitly report on nitrate concentrations in groundwater; and
- ii. the publication should be published after 1999.

Also, when many articles have been published for the same field site, we used only the most recent study. We excluded older studies before 1999, since the intensity of human activities is expected to be significantly different after 1999. We eliminated thirty-seven articles because no quantitative data on nitrate concentration were reported. For the considered data set, 206 studies report on the maximum concentration of nitrate, 187 studies on minimum concentration of nitrate, and 94 studies on the mean concentration of nitrate. Out of the 94 datasets for which mean values were reported, 12 field sites have nitrate concentration smaller than 1 mg/L. We present the locations and references of the considered field studies in Table 2. In case spatial coordinates were not reported in the selected paper, we allocated the coordinates of the field study in Google Earth using the www.gps-coordinates.net and [www.mapcoordinates.net applications](http://www.mapcoordinates.net/applications). As an example, we present in Fig. 3 the identified locations and reported maximum nitrate values of the selected studies. The absence of exact spatial coordinates in many studies will therefore generate a positioning error in the analysis. However, given the extent of the study, i.e. the African continent, we consider that this positioning error will not have significant effects on the overall results.

171

3.3 Determination of spatial explanatory variables

173

Table 3, list the environmental attributes and data sources that we considered for explaining the observed nitrate contamination. These variables represent both anthropogenic and natural factors and were derived from multiple sources of information. The attributes are related to recharge, geology, hydrogeology, soil texture, land use, topography and pollution pressure and were partially inspired from the DRASTIC vulnerability mapping approach. We compiled all explanatory variables in a common GIS environment (ArcGIS 10.3™), using a common projection and resolution (15 km x 15 km) at the 1:60.000.000 scale. This spatial resolution was chosen because, we have considered that she was a reasonable compromise between different resolutions of the different datasets, computing constraints and regional extent. Indeed, this grid cell dimension has been used to map the vulnerability and risk pollution maps at the African scale (Ouedraogo et al., 2016). Generic variables at the grid scale were extracted to build our explanatory variables in this study. Most of these variables were categorical, but some were continuous.

184

Groundwater recharge is considered as primary explaining variable because recharge is the primary vehicle by which a contaminant is transported from the ground surface to groundwater. Groundwater recharge to an unconfined aquifer is



a function of precipitation, runoff, and evapotranspiration. The latter is related to vegetation and/or soil type. Groundwater recharge to a confined aquifer is generally more complex, as consideration must be given to the location of the recharge zone and the influence of any confining layers, vertical gradients, and groundwater pumping (Todd and Kennedy, 2010). In this study, we derived the African recharge map from the global-scale groundwater recharge model of Döll et al. (2008). We also considered independent climate data as alternative proxies of recharge. Hence, we considered the climate and region type data class as defined by Trambauer et al. (2014). We also considered the rainfall map as generated from the UNEP/FAO World and Africa GIS Data Base. The spatial resolution of this latter dataset is approximately 3.7 kilometers.

Subsequently we selected a set of environmental attributes related to aquifer type, groundwater position and the substrate that protects the aquifer. The depth to groundwater represents the distance that a contaminant must travel through the unsaturated zone before reaching the water table or to the first screen. We mapped the depth to water based on the data presented by Bonsor et al. (2011). The slope of the land surface is important with respect to groundwater vulnerability, because it determines the potential of a contaminant to infiltrate into the groundwater or be transported horizontally as runoff. We inferred the slope from the 90 meter Shuttle Radar Topography Mission (SRTM90) topographic map, using the Spatial Analyst software of ArcGIS10.2™. We derived the aquifer type and the impact of vadose zone material from the high resolution global lithological database (Glim) of Hartmann and Moosdorf (2012). We determined aquifer type and unsaturated lithological zone for each of the five hydro-lithological and lithological categories as defined by Gleeson et al., (2014). These categories are: unconsolidated sediments, siliciclastic sediments, carbonate rocks, crystalline rocks, and volcanic rocks (Gleeson et al., 2014). We constructed the soil type map from the 1 km resolution soil grid database developed by Hengl et al. (2012). We determined the hydraulic conductivity of aquifers from the Global Hydrogeology MaPS (GHYMPS) dataset (Gleeson et al., 2014). For the determination of the land use at the pan-African scale, we used the high resolution land cover/land use map from the GlobCoverdataset (Defourny et al., 2014). There are twenty two (22) classes of land cover that represents Africa in this dataset. We aggregated these 22 classes into 6 similar classes (water bodies, bare area, grassland/shrubland, forest, urban, croplands) as represented in the Fig. 4 and then regrouped them in 5 groups (water bodies, forest/bare area, grassland/shrubland, croplands, urban area).

Finally, we considered a set of variables related to possible pollution pressure. We considered the application of fertilizer in the agricultural sector as a possible explanatory variable. We generated the nitrogen fertilizer application map from the Potter and Ramankutty (2010) dataset. The values shown on this map represent an average application rate for all crops over a 0.5° resolution grid cell. Following this study, the highest N fertilizer application rate (i.e. 220 kg / ha) is found in Egypt's Nile Delta. We further considered population density as a proxy of pollution source. We considered the population density map for the year 2000, as produced by Nelson (2004).

3.4 Statistical model description

We used Multiple Linear Regression (MLR) as the statistical method for identifying the relationship between the observed nitrate concentrations in groundwater and the set of independent variables given in Table 3. MLR is based on least squares, which means that the model is fitted such that the sum of squares of differences of predicted and measured values is minimized (Koklu et al., 2009; Helsel and Hirsh, (1992)). The MLR model is denoted as by Eq. (1):



$$y_i = \beta_0 + \sum_{j=1}^n \beta_j x_{ij} + \varepsilon_i \quad i=1, m, \quad (1)$$

where y_i is the response variable at location i , β_0 is the intercept, β_j are the slope coefficients of the explanatory categorical or continuous variables x_{ij} , n the number of variables and m is the number of locations or wells (number of studies here). ε_i is the regression residual. In this study, the response variable is the logtransformed nitrate concentration in groundwater. The logtransformation was needed to stabilize the variance and to comply with the basic hypothesis of MLR. The logtransformed nitrate concentration is a continuous monotonic increasing function; it is therefore reasonable to accept that factors that contribute to the logtransformed nitrate load will also contribute to the nitrate load. The explanatory variables were defined using a stepwise procedure, using the Akaike Information Criterion (AIC) as test statistic (Helsel and Hirsch, 1992). We evaluated model performance based on the significance level of estimated coefficients, the coefficient of determination (R^2), the mean square error (MSE), the probability plots of model residuals (PRES), the plots of predicted versus observed values and the Akaike Information Criterion (AIC). High values of R^2 and low values of RMSE, PRES and AIC indicate a better performance of the model. To validate the model obtained by the stepwise procedure, the standard regression diagnostics were assessed. To test the heteroscedasticity in the model residuals, we use the Breusch-Pagan (BP) test by implementing with “lmtest” package. A Student statistic t test was finally used to check the statistical significance (with p -values <0.10) of variables in the final model. We assessed tolerance to examine if multicollinearity exists between variables. In this study, we performed the statistical analyses using the R version 3.1.1 (R Development Core team, 2015).

4 Results

4.1 Normality of the dependent variable

Prior to analysis, we carefully checked the data using descriptive statistics, such as boxplots and correlation analysis. The observed nitrate concentrations through meta-analysis ranges from 0 mg/L to 4625 mg/L for all categories, i.e. mean, maximum and minimum values of nitrate groundwater contamination. Descriptive statistics are summarized in Table 4. The average mean nitrate concentration is 27.85 mg/L. The positive skewness of the mean nitrate concentration data and the kurtosis suggest that the mean nitrate concentration is not normally distributed. In contrast, the lognormally transformed mean nitrate concentration obeys normality, as demonstrated by means of the non-parametric Shapiro-Wilk test (p -value=0.1432 >0.05). The histogram of mean and logtransformed concentration is shown in the Fig. 5. We also checked the minimum and maximum nitrate concentration for normality (results can be obtained from the authors upon request).

4.2 Correlation between nitrate in groundwater and explanatory variables

Land Cover/Land Use is a principle factor, controlling groundwater contamination. The box plot distribution of logtransformed mean nitrate concentration for different land use classes is presented in Fig. 6. Groundwater in agricultural and urban areas is clearly more susceptible to nitrate pollution as compared to forest/bare area land use. Also water bodies are susceptible to nitrate contamination but this result is likely spurious since only two studies support this category. We performed a similar analysis on the logtransformed maximum and minimum nitrate concentration. The corresponding boxplots results can be obtained from the authors upon request. High values for logtransformed maximum nitrate concentration are also found in urban and cropland areas. High values for logtransformed minimum



264 nitrate concentration are detected in croplands fields. All analyses confirm that the highest nitrate pollution is retrieved
265 in urban areas, immediately followed by agricultural areas.

266

267 In this study, the aquifer systems for Africa are divided into 5 categories based on the lithological formations. Fig. 7
268 shows the relation between mean logtransformed nitrate concentration and aquifer system type class. The carbonates
269 rocks, the unconsolidated sediments and the siliciclastic sedimentary rocks, represents respectively the first, the second
270 and the third class in terms of nitrate contamination. The crystalline rock and volcanic rock aquifer classes are less
271 contaminated. The high concentrations in the unconsolidated aquifer systems is a particular point of concern, since this
272 class is the most representative in terms of groundwater exploitation. The high concentrations in the carbonates rocks
273 and fractured basalt can be explained by their high vulnerability related to the presence of solution channels and
274 fractures.

275

276 The distribution of the logtransformed mean nitrate concentration data with depth is shown in Fig. 8. Apparently, no
277 clear relationship exist between depth to groundwater and nitrate contamination. The Pearson's correlation give a poor
278 correlation ($r=0.004$). However, the careful analysis of this figure shows clearly that shallower wells (7-25 m bgl and
279 25-50 m bgl) are associated with higher values of logtransformed mean nitrate concentration, in contrast to the low
280 values of logtransformed nitrate concentrations found in the deeper groundwater systems (>250 m bgl).

281

282 The relationship between the logtransformed mean nitrate concentration and groundwater recharge can also be observed
283 in Fig. 8. This figure shows that nitrate concentration in the groundwater of shallow aquifers generally increases with
284 recharge, except for the very low recharge class (0-45 mm/year). The high nitrate observations for this latter low
285 recharge class may be due to irrigation water return that feed the groundwater and that is not integrated in the recharge
286 calculations. The analysis of Pearson's correlation between recharge and logtransformed mean nitrate give an $r=-0.292$.

287

288 The relation between the logtransformed mean nitrate concentration and the population density is given in Fig. 8. We
289 observe an increasing nitrate in groundwater related to increasing population. This explicit relationship between
290 population density and nitrate concentration has a Pearson's correlation of 0.632. This obviously confirms the
291 importance of studying the population as a potential polluting parameter and its relevant correlation to nitrate occurrence
292 in the groundwater at the pan African scale.

293

294 Nitrogen fertilizer contributes significantly to an increase in crop yields, but excess nitrogen fertilizer generally pollutes
295 groundwater (Green et al., 2005; Nolan et al., 2002). In the case of Africa, the impact of the nitrogen fertilizer application
296 rate on logtransformed mean nitrate concentration is illustrated in Fig. 8. Pearson's correlation give a low relation
297 ($r=0.09$). The analysis in this figure confirms that no clear relationship exist between fertilizer load and groundwater
298 nitrate contamination. This can be linked to the relatively low fertilizer use in Africa, as compared to other continents.
299 Indeed, most studies have nitrogen fertilizer dressings that are below 50 kg/ha. According the FAO (2012), Africa
300 accounts only for about 2.9 percent of the world fertilizer consumption in 2011.

301

302



We performed similar correlation analysis on the logtransformed maximum concentration and logtransformed minimum concentration respectively. Details can be obtained from the authors upon request. Results of these analyses are coherent with the results for logtransformed mean nitrate concentration.

4.3 Development of the multi-variate statistical model

We developed a set of multiple variable regression models for the logtransformed mean and maximum nitrate concentration in terms of above mentioned explanatory variables. A positive regression coefficient indicates a positive correlation between a significant explanatory variable and a target contaminant, while a negative coefficient suggests an inverse or negative correlation. We retained only explanatory variables with p-values ≤ 0.1 .

The best final model that explains the log transformed mean nitrate concentration includes only 4 explanatory variables: (1) Depth to groundwater, (2) Recharge, (3) Aquifer type, and (4) Population density. Table 5 summarizes the results of this linear regression model. This model can explain 65 percent of the logtransformed mean nitrate concentration observations. The sign of the parameter coefficient indicates the direction of the relationship between independent and dependent variable (Boy-Roura et al., 2013). The p-value expresses the «attained significance level» for that slope coefficient, which is the significance level attained by the data (Helsel and Hirsch, 1992). The lower the p-value, the more significant is the model parameter.

The regression analysis confirms the strong relationship between population density and logtransformed mean nitrate concentration. As the p-value is far below 0.05, we are more than 95 % confident that the population density strongly affects the nitrate occurrences in groundwater.

The aquifer medium is another important explanatory variable for logtransformed mean nitrate concentration. Three categories of aquifer media are significantly explaining the dependent variables: carbonates rocks, crystalline and unconsolidated sediments rocks. Indeed, the analyse of regression coefficients shows that the likelihood of nitrate contamination decreases with the presence of unconsolidated sediments and crystalline rocks. Other aquifer types tested include siliciclastic sedimentary rocks and volcanic rocks aquifers were found not statistically significant in the model. However, the aquifer media type is an important variable to assess groundwater vulnerability and to bring information about the hydrogeological system in the assessment. It allows differentiating the vulnerability in terms of aquifer lithology. Variables such as hydraulic conductivity could be surrogates for aquifer media, because hydraulic conductivity data were developed based on the lithological formation. Nevertheless they were not statistically significant in the final model.

The third variable represents the depth to groundwater. The three first classes (0-7; 7-25 and 25-50 m bgl) of groundwater depth are all statistically significant. The water table corresponding to the 0-7 m class has the strongest statistical significance. The positive parameter coefficient indicates large contamination for shallow groundwater depths. By analysing the table of coefficient, we observe that the largest groundwater depth class (100-250 m bgl) is not statistically significant (p-value >0.05). We can conclude that the shallow groundwater systems in African scale are most vulnerable to nitrate pollution.



344 The fourth variable included in the final model is the recharge. The recharge rate in the 45-123 mm/year and 123-224
345 mm/year class are statistically significant. In general, these rates correspond to semi-arid and dry sub-humid regions.
346 The high concentrations in these areas can be due to intensive agricultural activities.

347
348 Other explanatory variables such as rainfall or land cover/land use were not considered in the final model. Indeed,
349 notwithstanding a variable such as land cover/land use strongly influences observed logtransformed mean nitrate
350 concentration (Fig. 6), it is related to other variables such as population density. Hence, to avoid multicollinearity in
351 the final model, the land cover/land use variable is no longer included in the final model.

352
353 The final multiple linear regression (MLR) model using the four variables yields an R^2 of 0.65, indicating that 65% of
354 the variation in observed logtransformed mean nitrate concentration at the pan African scale is explained by the model.
355 The result of the model is globally significant because the p-value = $2.422e-10$ at 95% of the significant level. The
356 observed versus predicted log transformed mean nitrate concentration is shown in Fig. 9 and indicates that the MLR fits
357 the data well. A probability plot of model residuals indicates that they follow a normal distribution (Fig. 10). We
358 performed the Shapiro-Wilk test as an additional check on the distribution of nitrate residuals. Because the probability
359 associated with the test statistic is larger than 0.05, we accept the null hypothesis that the residuals follow a normal
360 distribution. Despite the fact that a few points have higher Cook's D values compared to the rest of the observation, they
361 were kept in the MLR to represent the whole range of nitrate concentration data. In order to check the regressions
362 assumptions of homoscedasticity, a plot of the residuals of logtransformed mean nitrate versus the predicted
363 logtransformed mean values is illustrated in Fig.11. We observe that the majority of observations are in the range of -2
364 to 2 except for two outliers observed in the bottom left part of the graph. The residual standard error of the
365 logtransformed mean nitrate is 0.91116 (ln (mg/L)). We observe that the residuals decreases with increasing predicted
366 nitrate concentrations. The Breusch-Pagan test was used to assess heteroscedasticity in the model residuals (BP=24.2773
367 and p-value= 0.042). With a p-value of 0.042, we reject the null hypothesis that the variance of the residuals is constant
368 and infer that heteroscedasticity is indeed present. As a results, we may expect some bias in the MLR model.

369
370 Similarly to the logtransformed mean nitrate concentration modelling, we developed another model corresponding to
371 the logtransformed maximum nitrate concentration. This model yielded only an R^2 = 0.42 for the maximum values. The
372 explanatory variables which influence the logtransformed maximum nitrate concentration in groundwater are: depth to
373 groundwater, soil media, topography, rainfall, climate class and type of region. For the logtransformed minimum
374 concentration, the absence of normal distribution assumptions did not allow to develop a MLR model.

375 5 Discussion

376 We present in this study a data based method to assess the vulnerability of groundwater systems for water quality
377 degradation. We used the logtransform of reported nitrate concentration as a proxy for groundwater vulnerability. We
378 present a statistical model to explain this proxy in terms of generic data at the pan-African scale. In a previous study we
379 evaluated the groundwater vulnerability for pollution at the pan-African scale using the generic DRASTIC approach
380 (Ouedraogo et al., 2016). Yet, the uncalibrated DRASTIC model predictions are subjected to quite some uncertainty, in
381 particularly due to the subjectivity in assigning the generic DRASTIC model parameters. In contrast to this previous
382 study we focus in this paper on nitrate pollution which is a parameter that is strongly related to vulnerability and that



383 often is measured in on-going monitoring programmes. We integrate published nitrate in groundwater data explicitly in
384 the assessment, thereby eliminating completely the subjectivity of the DRASTIC approach. The study also targets the
385 optimal use of the available data for the prediction purposes. Certain data might be redundant or biased. We addressed
386 in this study the quality of data (Sect. 3.2). In this study, we used multiple linear regression (MLR) for explaining nitrate
387 groundwater in terms of other generic spatially distributed environmental parameters. MLR is an approach to model the
388 relationship between a response variable and multiple set of explanatory variables (Rawlings et al., 1998). MLR analysis
389 is capable of both predicting and explaining a response variable using explanatory variables without compromise
390 (Kleinbaum et al., 1988). Previous studies of MLR using spatial variables for nitrate concentration in groundwater
391 showed R^2 values of 0.52 and 0.64 in shallow alluvial aquifers (Gardner and Vogel, 2005; Kaown et al., 2007) and R^2
392 of 0.82 in deep sandy tertiary aquifers (Mattern et al., 2009). For the application in this study, we selected the parameters
393 using stepwise MLR regression, allowing to select only those parameters which have significant impact on the
394 logtransformed concentration values of nitrate.

395

396 The explanatory variables with the strongest influence on the mean log transformed nitrate concentration at the pan
397 African scale are the population density and groundwater depth, which is in agreement with results from other studies
398 such as Nolan, (2001), Nolan et al., (2002), Nolan and Hitt (2006), Liu et al., (2013), Bonsor et al., (2011) and
399 Sorichetta et al. (2013). Both explanatory variables are directly related to the probability of having high nitrate
400 concentrations in groundwater. The strong influence of the population density variable can be explained by the serious
401 problem of sanitation in Africa townships. This is consistent with the conclusions of the UNEP/UNESCO project
402 ‘Assessment of Pollution Status and Vulnerability of Water Supply Aquifers Cities’, stating that the major pollution
403 pressure on African water bodies are related to poor on-site sanitation, solid waste dumpsites including household waste
404 pits and surface water influences (Xu and Usher, 2006). This is also consistent with other studies stating that that
405 leaking septic tanks and sewer systems are considerably causing nitrate contamination of groundwater in urban areas
406 (Bohlke, 2002; Showers et al., 2008). The magnitude of contamination is not only affected by the population density,
407 but also by the socio-economic setting (UNEP/DEWA, 2014). A high population density is therefore often associated
408 with the lack of adequate sanitation in many slums/shanty towns in Africa. The strong influence of population density
409 in our model suggests that high concentrations in groundwater are mainly from subsurface leakage of municipal sewage
410 systems, petrol service station (underground storage tanks), and agricultural chemicals in small scale farming. Hence,
411 sanitation programmes in Africa must not be delinked from groundwater protection and controlling the use of fertilizer
412 products in agriculture.

413

414 Nitrate concentrations were generally higher values for shallower wells than for deeper groundwater systems. The
415 inverse relation between depth and nitrate is consistent with previous groundwater studies that considered well depth or
416 depth of the screened interval as explanatory variables (Nolan and Hitt, 2006; Nolan et al., 2014; Wheeler et al., 2015;
417 Ouedraogo and Vanclooster, 2016). Deeper groundwater typically is older and may predate periods of intensive fertilizer
418 application (1950–present). Also, given the larger travel times associated with the recharge of deep groundwater
419 systems, there is an enhanced opportunity for denitrification (Wheeler et al., 2015). The same author showed that the
420 lagged increase in groundwater nitrate concentration relative to the increase of animal feeding operations suggests
421 groundwater recharge times of years or decades. Similar large recharge times for deep sandy aquifers were also
422 identified by Mattern and Vanclooster (2009).



423

424 The strong relation between nitrate contamination and both, groundwater depth and population density is a particular
425 point of concern given the fact that the majority (85 %) of Africa's population lives in regions where depth to
426 groundwater is shallow (0-50 m bgl) and where humps may be used to abstract water. Eight percent of these people (i.e.
427 nearly 66 million people) are likely to live in areas where depth to groundwater is 0-7 m bgl. A significant minority (8
428 %) of Africa's population lives in regions where the depth to groundwater is between 50 and 100 m bgl and common
429 hand pump technologies (e.g. India Mark) are inoperable in these cases. These areas are mainly within southern Africa
430 and to a lesser extent situated in the Sahel.

431

432 A third important explanatory variable that was included in the model was the groundwater recharge rate. This is
433 consistent with studies like Hanson (2002) and Saffigna and Keeney (1997). According to UNEP/DEWA (2014),
434 recharge from multiple sources influences groundwater microbial and chemical water quality. Groundwater recharge
435 rate is interlinked with many other environmental variables including, but not limiting, soil type, aquifer type, antecedent
436 soil water content, land use / land cover type and rainfall (Sophocleous, 2004; Ladekarl et al., 2005; Anuraga et al.,
437 2006). Hence, to avoid multi-collinearity, variables like land use/land cover type, rainfall, and soil type were not
438 considered in the final model.

439

440 Despite land cover/land use type is not explicitly included in the final model, the exploratory analysis clearly shows a
441 strong relationship between nitrate concentration and land use/land cover type. Indeed, nitrate concentrations are
442 generally higher in urban areas. This is consistent with many other studies such as Showers et al., (2008). The high
443 contamination in urban areas jeopardises groundwater exploitation in urban areas. Urbanization is a pervasive
444 phenomenon around the world, and groundwater demands in urban areas are increasingly growing. The degradation of
445 groundwater bodies in urban areas is therefore a particular point of concern. Also agricultural land exhibit an impact on
446 groundwater nitrate concentrations compared to the grassland/shrubland, water bodies and forest/bare area, but this
447 effect is less important as compared to agricultural land effects in other parts of the world (e.g. Europe).

448

449 The influence of aquifer type to the nitrate contamination was demonstrated by Boy-Roura et al.,(2013) and the
450 influence of soil type by Liu et al.,(2013). As with land cover/ land use type, these variables were not retained in the
451 final model to avoid collinearity with recharge.

452

453 The advantage of the MLR technique is that it can be easily implemented, and that model parameters can be easily
454 interpreted if the possible interaction between variables is ignored. However, MLR cannot represent well the many non-
455 linear dynamics that are associated to the contamination of groundwater systems. The violation of the homoscedasticity
456 hypothesis for instance, indicate that some bias will be present in our MLR model. Standard statistical models employed
457 in distribution modelling, such as MLR, work under the assumption of independence in the residuals and
458 homoscedasticity. When heteroscedasticity is present, residuals may be autocorrelated. This will lead to inflated
459 estimates in degrees of freedom, an underestimation of the residual variances and an overestimation of the significance
460 of effects (Legendre and Fortin, 1989; Legendre, 1993; Dale and Fortin, 2002; Keitt et al., 2002). This may show that
461 others variables should be included in the model or that the system may be highly non-linear.

462

463



464

465

466

467 We could avoid heteroscedasticity and improve the modelling performance by introducing non-linear regression
468 techniques (Prasad et al.,2006) or by introducing additional variables in the model. . Indeed, many studies showed that
469 non-linear statistical models of groundwater contamination outperform as compared to linear models (e.g. Pineros-
470 Garcet et al.,2006; Mattern et al., 2009; Oliveira et al.,2012 and Wheeler et al.,2015). To uncover nonlinear relationships
471 non-parametric data mining approaches provide obvious advantages (Olden et al., 2008; Wiens, 1989; Dungan et al.,
472 2002). Machine learning provides a framework for identifying other explanatory variables, building accurate
473 predictions, and exploring other nonlinear mechanistic relationships in the system. We may therefore expect that non-
474 linear statistical models will improve the explanatory capacity of the model and remove heteroscedasticity from the
475 model.

476

477 However, we believe that this theoretical constraint of heteroscedasticity does not questions the overall results. The
478 observed heteroscedasticity can be considered modest in view of large extent of the study, and the violation of statistical
479 design criteria when collecting data through a meta-analysis. Also, the interpretation of the factors and coefficients
480 associated with non-linear regression techniques become more complicated. We therefore prefer to maintain in this
481 paper the MLR techniques as a first approach to screen the factors that contribute to logtransformed mean nitrate
482 concentration risk. We suggest however that future studies should address the added value that can be generated with
483 non-linear modelling techniques. Such non-linear modelling techniques are particularly needed for the maximum
484 concentration for which the R^2 of simple MLR remains currently too poor and also for the minimum concentration who
485 shows the absence of normal distribution assumptions.

486

487 Also, in this study, we only identified a MLR model based on a meta-analysis spanning the pan-African continent.
488 Since, the data collected through the meta-analysis are very heterogeneous, the quality of the data set remains rather
489 poor. Therefore, future studies should critically address the validity of the identified model and explore how the model
490 can be improved and be used in a predictive model. It is however suggested that such model improvement and validation
491 step should be based on a more homogeneous data set. We therefore suggest to perform this future model validation
492 and model improvement step using data collected at the regional scale using more homogeneous data collection
493 protocols.

494

495 6 Conclusion

496 Contamination of groundwater by nitrate is an indicator of groundwater quality degradation and remains a point of
497 concern for groundwater development programmes all over the world. It is also a good proxy of overall groundwater
498 vulnerability for quality degradation. We address in this paper the issue of nitrate contamination of groundwater at the
499 African scale. We inferred the spatial distribution of nitrate contamination of groundwater from a meta-analysis of
500 published field studies of groundwater contamination. We analysed the literature for reported mean, maximum and
501 minimum concentration of nitrate contamination. We subsequently analysed, using box-plots, the reported



contamination in terms of spatially distributed environmental attributes related to pollution pressure and attenuation capacity. We extracted the explanatory variables from a geographic information system with the ArcGIS 10.3TM tool. We finally developed a MLR statistical model allowing to explain quantitatively the logtransformed observed contamination that is a proxy of vulnerability in terms of spatially distributed attributes. We selected the explanatory variables using a stepwise regression method.

We show that groundwater contamination by nitrates is reported throughout the continent, except for a large part of the Sahara desert. The observed nitrate concentrations through meta-analysis ranges from 0 mg/L to 4625 mg/L for all categories, i.e. mean, maximum and minimum values of nitrate groundwater contamination. The average mean nitrate concentration is 27 mg/L. The distribution of the reported nitrate contamination data is strongly skewed. We therefore build the statistical models for the logtransformed reported nitrate mean and maximum concentrations.

The graphical box plot analysis shows that nitrate contamination is important in shallow groundwater systems and strongly influenced by population density and recharge rate. Nitrate contamination is therefore a particular point of concern for groundwater systems in urban sectors.

The MLR model for the log transformed mean nitrate concentration uses the depth to groundwater, groundwater recharge rate, and aquifer type and population density as explanatory variable. The total variability explained of the log transformed reported mean nitrate concentration by this analysis was 65 %, suggesting that other variables not accounted for in the available ancillary data sets (such as climate zones) or better representations of the variable we do considered may be needed to improve understanding of nitrate concentrations. These findings highlight the challenges in developing appropriate regional variables to predict the conditions most vulnerable to high nitrate concentrations. The MLR shows that the population density parameter is the most statistically significant variable. This authenticates that leaking cesspits and sewer systems are considerably causing nitrate contamination of groundwater predominantly in urban areas. We identified similar MLR models for the log transformed maximum reported nitrate concentrations. Yet, for this latter attribute, the explained variation using the simple MLR techniques (42 %) remains small.

One of the main strengths of our study is that it is based on a large database of groundwater contamination reports from different countries, spanning the African continent and linked to environmental attributes that are available in a spatially distributed high resolution format. In addition, the developing a continental-scale model of nitrate contamination in groundwater of Africa, with its highly variable climate zones (hyperarid, arid, semi-arid, dry sub-humid, humid, tropical, and Mediterranean) allowed to determine which explanatory variables mainly influence the presence of nitrate. This represents an important step in managing and protecting both water resources and human health, particularly in semi-arid and arid regions. The main weakness or the major constraints of the modelling lies in the lack of detailed information available at the pan African scale, particularly the lack and uneven distribution of measured nitrate points. In spite of weaknesses and uncertainties caused by a moderate heteroscedasticity from residuals model, the modelling approach presented here has great potential. Although the meta-analysis should not replace nitrate testing, it gives a first indication of possible contamination; it can be also applied to preliminary assessment of nitrate using spatial variables and thus may support the planning process and guidelines for transboundary aquifers managers and regional basin organizations. This is particularly important as the demand for drinking water is increasing rapidly due to climate change and population growth, which will undoubtedly increase the pressure on groundwater resources.



543 Finally, further development may include the use of non-linear modelling techniques such as Random Forest techniques
544 to identify the causal mechanism behind autocorrelation and heteroscedasticity in nitrate distributions over large extents
545 such as Africa. Such techniques have the potential to improve the quality of explanation and eventually prediction by
546 incorporate spatial autocorrelation, but complicate the physical explanation of observed trends. In addition, the model
547 should further be validated using more homogeneous data sets. There is a need for a process-based continental scale
548 nitrate estimate that uses a consistent approach and data, as the basis for studying potential environmental factors
549 impacts on groundwater resources in Africa. In a predictive mode, the model could be used for exposure estimate in
550 epidemiological studies on the effect of polluted groundwater on human health. Also, an application of the statistical
551 model to others contaminants could be explored.

552



References

- Abate, E. Y.: Anthropogenic Impacts on Groundwater Resources in the urban Environment of Dire Dawa, Ethiopi. Master Thesis in Geosciences, Department of Geosciences, Faculty of Mathematics and Natural Sciences, University of Oslo: 64pp., 2010.
- Abd El-Salam, M. M. and Abu-Zuid, G. I.: Impact of landfill leachate on the groundwater quality: A case study in Egypt. *Journal of Advanced Research*, 2014.
- Abdalla, F. A. and Scheytt, T.: Hydrochemistry of surface water and groundwater from a fractured carbonate aquifer in the Helwan area, Egypt. *J. Earth Syst. Sci.* 121(1): 109–124., 2012.
- Abdel-Lah, A. K. and Shamruk, M.: Impact of septic system on ground water quality in a Nile valley village, Egypt. Sixth International Water Technology Conference, IWTC 2001, Alexandria, Egypt: pp.237-245., 2001.
- Abdelbaki, C., Asnoui, F., Assoud, I., Cherif, Z. E. A. and Yahiaoui, I.: Contribution to the cartography of the groundwater quality of the urban group of Tlemcen (Algeria). *Larhyss Journal* n°16: pp. 7-19., 2013.
- Abdellah, A. M., Abdel-Magid, H. M. and Shommo, E. I.: Assessment of Groundwater Quality in Southern Suburb of the Omdurman City of Sudan. *Greener Journal of Environmental Management and Public Safety* Vol. 2 (2): pp. 083-090., 2013.
- Abdesselam, S., Halitim, A., Jan, A., Trolard, F., and Bourrié, G.: Anthropogenic contamination of groundwater with nitrate in arid region: case study of southern Hodna (Algeria). *Environmental Earth Sciences* 70(5): 2129-2141., 2012.
- Abenan, A. T., Jean, K. K., Bachir, S. M., Armand, A. K., Serge, D. K., Tanoh, K. J. J., Adiow, E. T., Kouassi, S. H., Jean-Patrice, J. and Jean, B.: Contribution of multicriteria analysis and GIS for the specific vulnerability mapping to agricultural inputs of groundwaters in Bonoua area (Southeast of Côte d'Ivoire), 2012.
- Abou, M. M.: Pollution des eaux souterraines en milieu urbain, cas de la nappe phréatique de la ville de Niamey. *Institut Supérieur des Techniques de l'Eau*, No.112, ID cas: 462, Maticule: NIG002: 8pp., 2000.
- Ackah, M., Agyemang, O., Anim, A. K., Osei, J., Bentil, N. O., Kpattah, L., Gyamfi, E. T. and Hanson, J. E. K.: Assessment of groundwater quality for drinking and irrigation: the case study of Teiman-Oyarifa Community, Ga East Municipality, Ghana. *Proceedings of the International Academy of Ecology and Environmental Sciences* 1(3-4): 186-194., 2011.
- Adelekan, B. and Ogunde, O.: Quality of water from dug wells and the lagoon in Lagos Nigeria and associated health risks. *Scientific Research and Essays* 7(11): 1195-1211., 2012.
- Affum, A. O., Osae, S. D., Nyarko, B. J., Afful, S., Fianko, J. R., Akiti, T. T., Adomako, D., Acquah, S. O., Dorleku, M., Antoh, E., Barnes, F. and Affum, E. A.: Total coliforms, arsenic and cadmium exposure through drinking water in the Western Region of Ghana: application of multivariate statistical technique to groundwater quality. *Environ Monit Assess* 187(2): 4167., 2015.
- Aghzar, N., Berdai, H., Bellouti, A. and Soudi, B.: Pollution nitrique des eaux souterraines au Tadla (Maroc). *Revue des sciences de l'eau / Journal of Water Science* 15(2): p. 459-492., 2002.
- Ahmed, A. L. M., Sulaiman, S., Osman, M. M., Saeed, E. M. and Mohamed, Y. A.: Groundwater quality in Khartoum state, Sudan. *Journal of Environmental Hydrology* 8(Paper 12): 7pp., 2000.
- Ahoussi, K. E., Youan, T. M., Loko, S., Adja, M. G., Lasm, T. and Jourda, J. P.: Étude hydrogéochimique des eaux des aquifères de fractures du socle Paléo-protérozoïque du Nord-Est de la Côte d'Ivoire : Cas de la région de Bondoukou. *Afrique SCIENCE* 08(3): 51 - 68., 2012.
- Ake, G. E., Kouadio, H. B., Dongo, K., Dibi, B., Kouame, F. K. and Biemi, J.: Application des méthodes DRASTIC et SI pour l'étude de la vulnérabilité à la pollution par les nitrates (NO₃⁻) de la nappe de Bonoua (Sud-Est de la Côte d'Ivoire). *Int. J. Biol. Chem. Sci* 4(5): 1676-1692., 2010.
- Akinbile, C. O.: Environmental Impact of Landfill on Groundwater Quality and Agricultural Soils in Nigeria." *Soil & Water Res.* 7(1): 18–26., 2012.
- Akoteyon, I. S. and Soladoye, O.: Groundwater Quality Assessment in Eti-Osa, Lagos-Nigeria using Multivariate Analysis. *J. Appl. Sci. Environ. Manage.* 15(1): 121 - 125., 2011.
- Aladejana, J. A. and Talabi, A. O.: Assessment of Groundwater Quality in Abeokuta Southwestern, Nigeria." *International Journal Of Engineering And Science* 2(6): pp.21-31., 2013.
- Alaoui, H. L., Oufdou, K. and Mezrioui, N.: Environmental pollutions impacts on the bacteriological and physicochemical quality of suburban and rural groundwater supplies in Marrakesh area (Morocco). *Environ Monit Assess* 145(1-3): 195-207., 2008.
- Aller, L., Bennet, T., Lehr, J. H., and Petty, R. J.: DRASTIC: A standardized system for evaluating groundwater pollution using hydrological settings. Ada, OK, USA: Prepared by the National water Well Association for the US EPA Office of Research and Development. EPA/600/2-85-018, 163pp, 1985.



- 611 Amini, M., Mueller, K., Abbaspour, K. C., Rosenberg, T., Afyuni, M., Møller, K. N., Sarr, M. and Johnson, C. A.:
612 Statistical modeling of global geogenic fluoride contamination in groundwaters. *Environmental science*
613 & technology 42(10): 3662-3668., 2008a.
- 614 Amini, M., Abbaspour, K. C., Berg, M., L. Winkel, L., Hug, S. J., Hoehn, E., Yang, H. and C. A. Johnson.:
615 Statistical modeling of global geogenic arsenic contamination in groundwater. *Environmental science &*
616 *technology* 42(10): 3669-3675., 2008b.
- 617 Anane, M., Selmi, Y., Limam, A., Jedidi, N. and Jellali, S.: Does irrigation with reclaimed water significantly
618 pollute shallow aquifer with nitrate and salinity? An assay in a perurban area in North Tunisia." *Environ*
619 *Monit Assess* 186(7): 4367-4390., 2014.
- 620 Angola Water Works.: Annual Drinking Water Quality Report.. IN5276001.
621 <http://www.angolain.org/Water/documents/Angola2013AnnualWaterReport.pdf>: 2pp., 2013 (last acces:
622 13 August, 2014).
- 623 Ansa- Asare, O. D., Darko, H. F. and Asante, K. A.: Groundwater quality assessment of Akatsi, Adidome and Ho
624 districts in the Volta Region of Ghana. *Desalination* 248(1-3): 446-452., 2009.
- 625 Anthony, K.-O.: Impact of municipal solid wastes on underground water sources in Nigeria. *European Scientific*
626 *Journal* Vol.8(No.11): 19.pp., 2012.
- 627 Anuraga, T. S., Ruiz, L., Kumar, M. S. M., Sekhar, M. and Leijnse, A.: Estimating groundwater recharge using
628 land use and soil data: A case study in South India. *Agricultural Water Management* 84(1-2): 65-76.,
629 2006.
- 630 Armah, Y.: Hydrochemical Analysis of Groundwater In The Lower Pra Basin of Ghana. *Journal of Water Resource*
631 *and Protection* 2(10): pp. 864-871., 2010.
- 632 Armah, F. A., Luginaah, I. and Benjamin, A.: Water Quality Index in the Tarkwa Gold Mining Area in Ghana."
633 *The Journal of Transdisciplinary Environmental Studies* 11(2): 15pp., 2012.
- 634 Assloul, J. E., Kholtei, S., Amrani-Paaza, N. E. and Hilali, A.: Impact des activités anthropiques sur la qualité
635 des eaux souterraines de la communauté Mzamza (Chaouia, Maroc). *Revue des sciences de l'eau* 20(3):
636 309-321, 2007.
- 637 Ateawung, J. N.: A GIS Based Water Balance Study of Africa.. Master of Physical Land Resources, Universiteit
638 Gent Vrije Universiteit Brussel Belgium., 55pp., 2010.
- 639 Aza-Gnandji, C. D. R., Xu, Y., Raitt, L. and Levy, J.: Salinity of irrigation water in the Philippi farming area of
640 the Cape Flats, Cape Town, South Africa. *Water SA* 39(2), 199-210, 2013.
- 641 Bagalwa, M., Majaliwa, M., Mushagalusa, N. and Karume, K.: Estimation of Transported Pollutant Load from
642 Small Urban Kahuwu River Micro-catchment in Lake Kivu, Democratic Republic of Congo. *Journal of*
643 *Environmental Science and Engineering B* 2: 460-472., 2013.
- 644 Bahri, F. and Saibi, H.: Characterization, classification, bacteriological, and evaluation of groundwater from 24
645 wells in six departments of Algeria. *Arabian Journal of Geosciences* 5(6): 1449-1458., 2012.
- 646 Balogun, I. I., Akoteyon, I. S. and Adeaga, O.: Evaluating Land Use Effects on Groundwater Quality in Lagos-
647 Nigeria Using Water Quality Index. *Journal of Scientific Research* 4(2), 397-409, 2012.
- 648 Barhe, T. A. and Bouaka, F.: Physicochemical Characterization and Chlorination of Drilling Water Consumed In
649 Brazzaville-Congo. *Journal of Chemical, Biological and Physical Sciences, An International Peer Review*
650 *E-3 Journal of Sciences* 3(3): 2328-2336., 2013.
- 651 Batisani, N. : Groundwater hydrochemistry evaluation in rural Botswana: A contribution to integrated water
652 resources management. *Ethiopian Journal of Environmental Studies and Management* 5(4)., 2012.
- 653 Bauder, J.W, Sinclair, K.N. and Lund, R.E.: Physiographic and land use characteristics associated with nitrate
654 nitrogen-nitrogen in montana groundwater. *Journal of Environmental Quality* 22: 255-262., 1993.
- 655 Belghiti, M. L., Chahlaoui, A., Bengoumi, D. and El Moustaine, R.: Etude de la qualite physico-chimique et
656 bacteriologique des eaux souterraines de la nappe plio-quaternaire dans la région de Meknès (Maroc).
657 *Larhyss Journal* n°14: pp. 21-36., 2013.
- 658 Belkhir, L. and Mouni, L.: Hydrochemical analysis and evaluation of groundwater quality in El Eulma area,
659 Algeria." *Appl Water Sci* 2: 127-133., 2012.
- 660 Ben Abbou, M., Fadil, F., El Haji, M. and Zemzami, M.: Impact anthropique sur la qualite des eaux souterraines
661 du bassin versant de l'Oued Taza (Maroc). *European Scientific Journal*.,10(5), 117-137., 2014.
- 662 Benabbou, A., Benhoussa, A., Fekhaoui, M., El blidi, S., El Abidi, A. and Bounagua, M.: Assessment the risk
663 of impact of waste treatment center on groundwater quality in Oum Azza (Rabat, Morocco)." *J. Mater.*
664 *Environ. Sci.* 5(1): 143-152., 2014.
- 665 Böhlke, J.-K.: Groundwater recharge and agricultural contamination." *Hydrogeology Journal* 10(1): 153-179.,
666 2002.
- 667 Bonetto, S., De Luca Domenico, A., Fornaro, M. and Lasagna, M.: Drinking and irrigation groundwater quality
668 in Bulbule and Zway area (Ethiopia). *Proceedings of "EWRA 2005 – 6th International Conference*: 16pp.,
669 2005.
- 670 Bonsor, H. C. and MacDonald, A. M.: An initial estimate of depth to groundwater across Africa. *British Geological*
671 *Survey Open Report OR/11/067*: 26pp., 2011.



- 672 Bordalo, A. A. and Savva-Bordalo, J.: The quest for safe drinking water: an example from Guinea-Bissau (West
673 Africa). *Water Res* 41(13): 2978-2986., 2007.
- 674 Bossa, A. Y., Diekkrüger, B., Giertz, S., Steup, G., Sintondji, L. O., Agbossou, E. K. and Hiepe, C.: Modeling
675 the effects of crop patterns and management scenarios on N and P loads to surface water and groundwater
676 in a semi-humid catchment (West Africa). *Agricultural Water Management* 115: 20-37., 2012.
- 677 Boy-Roura, M., Nolan, B. T., Menció, A. and Mas-Pla, J.: Regression model for aquifer vulnerability assessment
678 of nitrate pollution in the Osona region (NE Spain). *Journal of Hydrology* 505: 150-162., 2013
- 679 Brandvold, V.: Nitrate Pollution of the Thiaroye Aquifer in the Suburban Parts of the Dakar Region, Senegal.
680 Using Chemical and Stable Isotope Analyses to Identify the Sources of Nitrate and the Processes at the
681 Groundwater/Surface Water Interface. Thesis of Norwegian University of Science and
682 Technology/Department of Geology and Mineral Resources Engineering: 133pp., 2013.
- 683 Bricha, S., Ounine, K., Oulkheir, S., El Haloui, N. and Attarassi, B.: Etude de la qualité physicochimique et
684 bactériologique de la nappe phréatique M'nasra (Maroc). *Afrique SCIENCE* 03(3): 391 - 404., 2007.
- 685 British Geological Survey, BGS.: Groundwater Quality: Ghana, 4pp., 2000.
- 686 British Geological Survey, BGS.: Groundwater Quality: Tanzania, 4pp., 2000.
- 687 British Geological Survey, BGS.: Groundwater Quality: Ethiopia, 6pp., 2001
- 688 British Geological Survey, BGS.: Groundwater Quality: Uganda, 4pp., 2001.
- 689 British Geological Survey, BGS.: Groundwater Quality: Zambia, 4pp., 2001.
- 690 British Geological Survey, BGS.: Groundwater Quality: Burkina Faso, 4pp., 2002.
- 691 British Geological Survey, BGS.: Groundwater Quality: Nigeria, 9pp., 2003.
- 692 British Geological Survey, BGS.: Groundwater Quality: Groundwater Quality: Tanzania. 4pp., 2000.
- 693 British Geological Survey, BGS.: Groundwater Quality: Water Quality Fact Sheet: nitrate (Benin and Senegal).
694 Produced for WaterAid. London: 6pp., 2003.
- 695 Burow, K. R., Nolan, B. T., Rupert, M. G. and Dubrovsky, N. M.: Nitrate in groundwater of the United States,
696 1991– 2003." *Environmental science & technology* 44(13): 4988-4997., 2010.
- 697 Bushra, A. H.: Quantitative status, vulnerability and pollution of groundwater resources in different environmental
698 and climatic contexts in Sardinia and in Ethiopia. PhD Thesis in Earth Science. 205pp., 2011.
- 699 Charfi, S., Zouari, K., Feki, S. and Mami, E.: Study of variation in groundwater quality in a coastal aquifer in
700 north-eastern Tunisia using multivariate factor analysis. *Quaternary International* 302: 199-209., 2013.
- 701 Chippaux, J.-P., Houssier, S., Gross, P., Bouvier, C. and Brissaud, F.: Étude de la pollution de l'eau souterraine
702 de la ville de Niamey, Niger. *Bull Soc Pathol Exot* 94(2): 119-123., 2002.
- 703 Cobbina, S. J., Nyame, F. K. and Obiri, S.: Groundwater Quality in the Sahelian Region of Northern Ghana, West
704 Africa. *Research Journal of Environmental and Earth Sciences* 4(4): 482-491., 2012.
- 705 Comte, J., Banton, O., Sambou, S., Travi, Y. and Ouedraogo, I.: L'aquifère des sables de la presqu'île de Dakar
706 (Sénégal) : état de la ressource et impacts anthropiques. Dix-huitièmes journées techniques du Comité
707 Français d'Hydrogéologie de l'Association International des Hydrogéologues. «Ressources et gestion des
708 aquifères littoraux. Cassis2012»: 276-274., 2012.
- 709 Coplen, T. B., Herczeg, A. L., and Barnes, C.: Isotope engineering—using stable isotopes of the water molecule
710 to solve practical problems. *Environmental tracers in subsurface hydrology*, Springer: 79-110., 2000.
- 711 Cronin, A. A., Pedley, S., Hoadley, A. W., Komou, F. K., Haldin, L., Gibson, J. and Breslin, N.: Urbanisation
712 effects on groundwater chemical quality: findings focusing on the nitrate problem from 2 African cities
713 reliant on on-site sanitation. *J Water Health* 5(3): 441-454., 2007.
- 714 Dale, M. R. T. and Fortin, M.-J.: Spatial autocorrelation and statistical tests in ecology. *Ecoscience* 9: 162-167.,
715 2002.
- 716 Dammo, M. N., Deborah, J. M., Yusuf, I. A. and Sangodoyin, A. Y.: Evaluation of ground water quality of
717 konduga town, Nigeria. *European International Journal of Science and Technology* Vol.2(5): pp.13-20.,
718 2013.
- 719 Dan-Hassan, M., Olasehinde, P., Amadi, A., Yisa, J. and Jacob, J.: Spatial and temporal distribution of nitrate
720 pollution in groundwater of Abuja, Nigeria. *International Journal of Chemistry* 4(3): p104., 2012.
- 721 Defourny, P., Kirches, G., Brockmann, C., Boettcher, M., Peters, M., Bontemps, S., et al.: Land cover CCI product
722 user guide version 2. 2014.
- 723 De Witte, I.: Characterization of the status of Dar es Salaam aquifer in view of salt-water intrusion and nitrate
724 contamination. Master in de Geologie, Faculteit Wetenschappen Vakgroep Geologie en Bodemkunde,
725 Universiteit Gent: pp.141., 2012.
- 726 Dibi, B., et al.,: Characterization of the Origin of Pollutants in Groundwater from Biostatistical Tests: The Case
727 of Catchment Ehania, South-Eastern Cote d'Ivoire. *Journal of Water Resource and Protection*
728 05(12): 1178-1185., 2013.
- 729 Diédihiou, M., et al.: Tracing groundwater nitrate sources in the Dakar suburban area: an isotopic multi-tracer
730 approach. *Hydrological Processes* 26(5): 760-770., 2012.



- 731 Djebebe-Ndjiguim, C. L. et al.: Characterization of the aquifers of the Bangui urban area, Central African
732 Republic, as an alternative drinking water supply resource. *Hydrological Sciences Journal* 58(8): 1760-
733 1778., 2013.
- 734 Döll, P. and Fiedler, K.: Global-scale modeling of groundwater recharge. *Hydrology and Earth System*
735 *Sciences*, 12, 863-885., 2008.
- 736 Douagui, A. G. et al.: Assessment of the bacteriological quality and nitrate pollution risk of Quaternary
737 groundwater in the southern part of Abidjan District (Côte d'Ivoire). *Journal of Hydro-environment*
738 *Research* 6(3): 227-238., 2012.
- 739 Drissa, S. T., Ernest, A. K., Marie-Solange, O. Y., Gbombélé, S. and Nagnin, S.: A Multivariate Statistical
740 Analysis of Groundwater Chemistry Data in the Highest Bandama Basin at Tortiya (Northern Côte
741 D'Ivoire). *Earth Resources* 1(3): 72., 2013.
- 742 Dungan, J.L., Perry, J.N., Dale, M.R.T., Legendre, P., Citron-Pousty, S., Fortin, M.J., Jakomulska, A., Miriti, M.,
743 Rosenberg, M.S.: A balanced view of scale in spatial statistical analysis. *Ecography* 25:626–240., 2002.
- 744 Dzwaïro, B., Hoko, Z., Love, D. and Guzha, E.: Assessment of the impacts of pit latrines on groundwater quality
745 in rural areas: A case study from Marondera district, Zimbabwe. *Physics and Chemistry of the Earth,*
746 *Parts A/B/C* 31(15-16): 779-788., 2006.
- 747 Easa, A. and Abou-Rayhan, A.: Domestic wastewater effect on the pollution of the groundwater in rural areas in
748 Egypt. Fourteenth International Water Technology Conference, IWTC 14 2010, Cairo, Egypt: pp.909-
749 923., 2010.
- 750 Eblin, S. G., Soro, G. M., Sombo, A. P., Aka, N., Kambiré, O. and Soro, N.: Hydrochimie des eaux souterraines
751 de la région d'Adiaké (Sud-Est côtier de la Côte d'Ivoire). *Larhyss Journal* n°17: pp.193-214., 2014.
- 752 Eckhardt, D. A. V. and Stackelberg, P. E.: Relation of groundwater quality to land use on Long Island, New York." *Ground Water* 33(6): 1019-1033., 1995.
- 753 Edet, A., Nganje, T., Ukpogon, A. and Ekwere, A.: Groundwater chemistry and quality of Nigeria: A status
754 review." *African Journal of Environmental Science and Technology* 5(13): 1152-1169., 2011.
- 755 Ekere, N. R.: Nitrates in rural communities water sources in Uzouwani, South eastern Nigeria, west Africa. *Journal*
756 *of Environmental Research And Development* Vol 7(2), 2012.
- 757 EL Bouqdaoui, K., Aachib, M., Blaghen, M. and Kholtei, S.: Modélisation de la pollution par les nitrates de la
758 nappe de Berrechid, au Maroc. *Afrique SCIENCE* 05(1): 99 - 113., 2009.
- 759 EL Hammoumi, N., Naima, M., Lekhlif, B. and Lakhdar, M.: Use of multivariate statistical and geographic
760 information system (GIS)-based approach to evaluate ground water quality in the irrigated plain of Tadla
761 (Morocco). *International Journal of Water Resources and Environmental Engineering* 5(2): pp. 77-93.,
762 2013.
- 763 El Hammoumi, N., Sinan, M., Lekhlif, B. and El Mahjoub, L.: Évaluation de la qualité des eaux souterraines pour
764 l'utilisation dans l'eau potable et l'agriculture : plaine de Tadla, Maroc. *Afrique SCIENCE* 08(1): 54 -
765 66., 2012.
- 766 Engida, Z. A.: Groundwater study of Addis Ababa area. Report No. 01/93: pp.28., 2001.
- 767
768
- 769 FAO-SWALIM.: Hydrogeological Survey and Assessment of Selected Areas in Somaliland and Puntland.
770 Technical Report No. W-20, FAO-SWALIM (GCP/SOM/049/EC) Project, Nairobi, Kenya., 2012.
- 771 Fedrigoni, L., M. Krimsa, M., Zouari, K., Maliki, A. and Zuppi, G. M.: Origine de la minéralisation et
772 comportement hydrogéochimique d'une nappe phréatique soumise à des contraintes naturelles et
773 anthropiques sévères : exemple de la nappe de Djebeniana (Tunisie). *C. R. Acad. Sci. Paris, Sciences de*
774 *la Terre et des planètes / Earth and Planetary Sciences* 332: 665–671., 2001.
- 775 Fekkoul, A., Zarhlou, Y., Boughriba, M. Machmachi, I. and Chafi, A.: Contamination par les pesticides
776 organochlores et les nitrates des eaux souterraines du système aquifère de la plaine des Triffa (Maroc Nord
777 Oriental). *ScienceLib Editions Mersenne* Vol.3(No.110801), 2011.
- 778 Fetouani, S., Sbaa, M., Vanclooster, M., and Bendra, B.: Assessing ground water quality in the irrigated plain of
779 Triffa (north-east Morocco). *Agricultural Water Management* 95(2): 133-142., 2008.
- 780 Fianko, J. R., Osae, S., Adomako, D. and Achel, D. G.: Relationship between land use and groundwater quality
781 in six districts in the eastern region of Ghana. *Environ Monit Assess* 153(1-4): 139-146., 2009.
- 782 Focazio, M., J., Reilly, T. E., Rupert, M. G. and Helsel, D. R.: Assessing Ground-Water Vulnerability to
783 Contamination: Providing Scientifically Defensible Information for Decision Makers. *USGS Circular,*
784 1224, 33pp., 2002.
- 785 Friedel, M. J.: Inventory and Review of Existing PRISM Hydrogeologic Data for the Islamic Republic of
786 Mauritania, Africa: U.S. Geological Survey. Open-File Report 2008-1138. 69 p., 2008.
- 787 Frind, E. O., Molson, J. W. and Rudolph, D. L.: Well vulnerability: a quantitative approach for source water
788 protection. *Ground Water* 44(5): 732-742., 2006.
- 789 Gardner, K. K. and Vogel, R. M.: Predicting Ground Water Nitrate Concentration from Land Use. *Ground Water*
790 43(3): pages 343–352., 2005.



- 791 Gleeson, T., Moosdorf, N., Hartmann, J. and van Beek, L. P. H.: A glimpse beneath earth's surface: GLobal
792 HYdrogeology MaPS (GLHYMPS) of permeability and porosity. *Geophysical Research Letters* 41(11):
793 3891-3898., 2014.
- 794 Greene, E. A., LaMotte, A. E. and Cullinan, K. A.: Ground-Water Vulnerability to Nitrate Contamination at
795 Multiple Thresholds in the Mid-Atlantic Region Using Spatial Probability Models." *Scientific*
796 *Investigations Reports* 2004-5118., 2004.
- 797 Grimason, A. M., Beattie, T. K., Masangwi, S. J., Jabu, G. C., Taulo, S. C. and Lungu, K. K.: Classification and
798 quality of groundwater supplies in the Lower Shire Valley, Malawi – Part 2: Classification of borehole
799 water supplies in Chikhwawa, Malawi. *Water SA* 39(4)., 2013.
- 800 Guideal, R., Bala, A. E. and Ikpokonte, E. A.: Physical and Chemical Quality of Groundwater in the Quaternary
801 Aquifer in N'djamena Region, Chad Republic. *Journal of Applied Sciences Research* 6(11): 1885-1890.,
802 2010.
- 803 Hamza, M. H., Added, A., Francés, A. and Rodríguez, R.: Validité de l'application des méthodes de vulnérabilité
804 DRASTIC, SINTACS et SI à l'étude de la pollution par les nitrates dans la nappe phréatique de Metline–
805 Ras Jebel–Raf Raf (Nord-Est tunisien). *Comptes Rendus Geoscience* 339(7): 493-505., 2007.
- 806 Hanson, C. R.: Nitrate concentrations in Canterbury ground water – A review of existing data. Report no. R02/17.
807 Environment Canterbury, 2002.
- 808 Harbaugh, A. W., Banta, E. R., Hill, M. C. and McDonald, M. G.: MODFLOW-2000, The U.S. Geological Survey
809 Modular Groundwater Model-User Guide to Modularization Concepts and the Groundwater and Flow
810 process. Open-File Report 00-92. 121 pp., 2000.
- 811 Hartmann, J. and Moosdorf, N.: The new global lithological map database GLiM: A representation of rock
812 properties at the Earth surface. *Geochemistry, Geophysics, Geosystems*, 13(12),
813 Q12004,doi:10.29/2012GC004370., 2012.
- 814 Hassane, A. B.: Aquifères superficiels et profonds et pollution urbaine en Afrique : cas de la communauté urbaine
815 de Niamey (Niger). These de Doctorat, Université Abdou Moumouni de Niamey: pp.198+annexes., 2010.
- 816 Haylamicheal, D. and Moges, A.: Assessing water quality of rural water supply schemes as a measure of service
817 delivery sustainability: A case study of WondoGenet district, Southern Ethiopia." *African Journal of*
818 *Environmental Science and Technology* 6(5): 229-236.: 2012.
- 819 Helsel, D.R., Hirsch, R.M.: Statistical Methods in Water Resources: Techniques of Water-Resources
820 Investigations, Book 4. US Geological Survey, p. 522 (Chapter A3), 1992.
- 821 Hengl, T., et al.: SoilGrids1km - Global Soil Information Based on Automated Mapping. *PLoS One*, 9(8),
822 e105992., 2014.
- 823 Hentati, I., Zaïri, M. and Ben Dhia, H. : A statistical and geographical information system analysis for groundwater
824 intrinsic vulnerability: a validated case study from Sfax-Agareb, Tunisia. *Water and Environment Journal*
825 25(3): 400-411., 2011.
- 826 Hosmer D.W., and Lemeshow S.: Applied Logistic Regression. John Wiley and Sons, New York, NY, pp. 307.,
827 1989.
- 828 Idriss, H., Salih, I. and Sam, A. K.: Study of radon in ground water and physicochemical parameters in Khartoum
829 state. *Journal of Radioanalytical and Nuclear Chemistry* 290(2): 333-338., 2011.
- 830 Imoisi, O., Ayesanmi, A. and Uwumarongie-Ilori, E.: Assessment of groundwater quality in a typical urban
831 settlement of resident close to three dumpsites in South-south, Nigeria." *Journal of Environmental*
832 *Science and Water Resources* 1(1): 12-17., 2012.
- 833 Ishaku, J. M.: Assessment of groundwater quality index for Jimeta- Yola area, Northeastern Nigeria. *Journal of*
834 *Geology and Mining Research* 3(9): pp. 219-231., 2011.
- 835 Kahssay, G., Olivier, J., Ayenew, T. and Ramose, M.: Low Cost Sanitation and its impact on Quality of
836 Groundwater in Addis Ababa. 19pp., 2010.
- 837 Kanyerere, T., Levy, J., Xu, Y. and Saka, J.: Assessment of microbial contamination of groundwater in upper
838 Limphasa River catchment, located in a rural area of northern Malawi. *Water SA* 38(4)., 2012.
- 839 Kaown, D., Hyun, H., Bae, G.-O. and Lee, K.-K.: Factors affecting the spatial pattern of nitrate contamination
840 in shallow groundwater. *Journal of environmental quality* 36(5): 1479-1487., 2007.
- 841 Kashaigili, J. J.: Assessment of groundwater availability and its current and potential use and impacts in Tanzania.
842 Final Report. International Water Management Institute (IWMI): 58pp., 2010.
- 843 Keitt, T. H., Bjørnstad, O.N., Dixon, P.M., and Citron-Pousty, S.: Accounting for spatial pattern when modeling
844 organism-environment interactions. *Echography* 25: 616-625., 2002.
- 845 Ketata, M., Gueddari, M. and Bouhlila, R.: Use of geographical information system and water quality index to
846 assess groundwater quality in El Khairat deep aquifer (Enfidha, Central East Tunisia). *Arabian Journal*
847 *of Geosciences* 5(6): 1379-1390., 2011.
- 848 Kissao, G. and Housséni, A.: Flooding and groundwater quality in a peripheral urban zone: the case of the district
849 of Agoè-Zongo in Lomé, Togo. 4th AMMA Interntional Conference, 2-6 July 2012, Toulouse, France:
850 23pp., 2012.



- 851 Kleinbaum, D.G., Kupper, L. L, Muller, K. E.: Applied regression analysis and other multivariate methods (2nd
852 ed.). PWS Publishing Co., Boston; 366p., 1988.
- 853 Kleinbaum, D.G.: Logistic Regression, A Self-Learning Text. Springer-Verlag, Berlin, p. 282., 1994.
- 854
- 855 Knobeloch, L., Salna, B., Hogan, A., Postle, J. and Anderson, H.: Blue babies and nitrate-contaminated well
856 water. *Environmental Health Perspectives* 108(7): 675-678., 2000.
- 857 Koklu, R., Sengorur, B. and Topal, B.: Water Quality Assessment Using Multivariate Statistical Methods—A
858 Case Study: Melen River System (Turkey). *Water Resources Management* 24(5): 959-978., 2009.
- 859 Kolpin, D.W.: Agricultural chemicals in groundwater of the Midwestern United States: relations to land use.
860 *Journal of Environmental Quality* 26: 1025-1037., 1997.
- 861 Kouame, K. J., Jourda, J. P., Saley, M. B., Deh, S. K., Anani, A. T. and Biémi, J.: Mapping of groundwater
862 vulnerability zones to pollution in various hydrogeological environments of Côte d'Ivoire by DRASTIC
863 method. *International Journal of Scientific & Engineering Research* 4(5): 915-923., 2013.
- 864 Kouassi, M. A., Yao, A. K., Ahoussi, E. K., Seki, C. L., Yao, A. N., Kouassi, I. K. and Biemi, J.: Apports des
865 méthodes statistiques et hydrochimiques à la caractérisation des eaux des aquifères fissurés de la région
866 du N'zi-Comoé (Centre-Est de la Côte d'Ivoire). *Int. J. Biol. Chem. Sci* 4(5): 1816-1838., 2010.
- 867 Kouassi, M. A., Ahoussi, E. K., Koffi, B. Y., Ake, Y. A. and Biemi, J.: Caractérisation hydrogéochimique des
868 eaux des aquifères fissurés de la zone Guiglo-Duekoué (Ouest de la Côte d'Ivoire). *International Journal*
869 *of Biological and Chemical Sciences* 6(1), 2012.
- 870 Kuitcha, D., Takounjou, A. L. F. and Ndjama, J.: Apport de l'hydrochimie et de l'isotope de l'environnement à
871 la connaissance des ressources en eaux souterraines de Yaoundé, Cameroun. *Journal of Applied*
872 *Biosciences* 67: 5194-5208., 2013.
- 873 Kulabako, N., Nalubega, M. and Thunvik, R.: Study of the impact of land use and hydrogeological settings on
874 the shallow groundwater quality in a peri-urban area of Kampala, Uganda." *Science of the Total*
875 *Environment* 381(1): 180-199., 2007.
- 876 Kyoung-Ho, K., Yun, S. T., Choi, B. Y., Chae, G. T., Joo, Y., Kim, K. and Kim, H.-S. : Hydrochemical and
877 multivariate statistical interpretations of spatial controls of nitrate concentrations in a shallow alluvial
878 aquifer around oxbow lakes (Osong area, central Korea)." *Journal of contaminant hydrology* 107(3): 114-
879 127., 2009.
- 880 Labar, S., A. Hani and L. Djabri.: Assessing groundwater contamination by hydrocarbons and heavy metals
881 (Northeast of Algeria). *Journal of Environmental Research and Management* 3(2): pp. 017-021., 2012a.
- 882 Labar, S., Hani, A. and Djabri, L.: Biochemical Approach to Assess Groundwater Pollution by Petroleum
883 Hydrocarbons (Case Skikda Algeria). *Journal of Water Resource and Protection* 04(07): 493-496., 2012b
- 884 Ladekarl, U. L., Rasmussen, K. R., Christensen, S., Jensen, K. H. and Hansen, B.: Groundwater recharge and
885 evapotranspiration for two natural ecosystems covered with oak and heather. *Journal of Hydrology* 300(1-
886 4): 76-99., 2005.
- 887 Laftouhi, N.-E., Vanclooster, M., Jalal, M., O. Witam, O., Aboufirassi, M., Bahir, M. and Persoons, É.:
888 Groundwater nitrate pollution in the Essaouira Basin (Morocco). *Comptes Rendus Geoscience* 335(3):
889 307-317., 2003.
- 890 Lagnika, M., Ibikounle, M., Montcho, J. C., Wotto, V. D. and Sakiti, N. G.: Caractéristiques physico-chimiques
891 de l'eau des puits dans la commune de Pobè (Bénin, Afrique de l'ouest). *Journal of Applied Biosciences*
892 79(1): 6887., 2014.
- 893 Lamribah, A., M. H. Benajiba, Y. Saoud, Ahrikat, M. and Benzakour, M.: Impact de la pollution urbaine sur la
894 contamination par les nitrates et les nitrites de la nappe phréatique de martil (Maroc). *Larhyss Journal*
895 n°14: pp. 79-91., 2013.
- 896 Laurent, M., François, A. and Marie, M. J.: Assessment of groundwater quality during dry season in southeastern
897 Brazzaville, Congo. *International Journal of Applied Biology and Pharmaceutical Technology Volume*
898 *I:(3): pp.762-769., 2010.*
- 899 Laurent, M. and Marie, M. J.: Groundwater Quality of Southeastern Brazzaville, Congo. *E-Journal of Chemistry*
900 *7(3): pp.861-869., 2010.*
- 901 Legendre, P.: Spatial autocorrelation: trouble or new paradigm? *Ecology* 74: 1659-1673., 1993.
- 902
- 903 Legendre, P. and Fortin, M.-J.: Spatial pattern and ecological analysis. *Vegetation* 80: 107-138., 1989.
- 904
- 905 Liu, C. W., Wang, Y. B. and Jang, C. S.: Probability-based nitrate contamination map of groundwater in Kinmen.
906 *Environ Monit Assess* 185(12): 10147-10156., 2013.
- 907 Loko, S., KE, A. and Biemi, J.: Anthropogenic Activities and the Degradation of the Environmental Quality in
908 Poor Neighborhoods of Abidjan, Côte d'Ivoire: Abia Koumassi Village. *Journal of Environmental*
909 *Protection, (4), 1099-1107., 2013a.*
- 910 Loko, S., Ahoussi, K. E., Koffi, Y. B., Kakou, N. F., Kouassi, A. M. and Biemi, J.: Microbiological and
911 physico-chemical quality of groundwater from artisanal sites of mining exploitation in the South-West of



- 912 Côte d'Ivoire: case of the area of Hiré. International Journal of Scientific & Engineering Research 4(9):
913 567-574., 2013b.
- 914 Longo, J. N.: Apport des outils hydrogéochimiques et isotopiques à la gestion de l'aquifère du mont Amba,
915 Kinshasa / République Démocratique du Congo). Thèse en cotutelle internationale Académie d' Aix
916 Marseille Université d'Avignon et des Pays de Vaucluse, Université de Kinshasa Faculté des Sciences
917 Agronomiques. 203pp., 2009.
- 918 MacDonald, A.: Groundwater, health and livelihoods in Africa. British Geological Survey © NERC 2010
919 Earthwise 26, 2pp, 2010.
- 920 MacDonald, A. M., Bonsor, H. C., Dochartaigh, B. É. Ó. and Taylor, R. G.: Quantitative maps of groundwater
921 resources in Africa. Environmental Research Letters 7(2): 024009., 2012.
- 922 MacDonald, A. M., Taylor, R. G. and Bonsor, C. H.: (eds.) Groundwater in Africa – is there sufficient water to
923 support the intensification of agriculture from “Land Grabs”. Hand book of land and water grabs in
924 Africa, pp 376-383., 2013.
- 925 Madioune, D. H., Faye, S. C. and Faye, S.: Etude de la vulnérabilité intrinsèque à la pollution de la nappe libre
926 des sables quaternaires de Thiarye par la méthode DRASTIC. Journal des Sciences et Technologies 9(2),
927 pp.1-11., 2011.
- 928 Maherry, A., Clarke, S., Tredoux, G. and Engelbrecht, P.: Spatial and temporal analysis of the nitrate
929 concentrations in groundwater for South Africa. 12pp., 2009.
- 930 Mair, A. and El-Kadi, A. I.: Logistic regression modeling to assess groundwater vulnerability to contamination in
931 Hawaii, USA. J Contam Hydrol 153: 1-23., 2013.
- 932 Mande, S. A. S., Liu, M., Djaneye-Boundjou, G., Liu, F., Bawa, M. L. and Chen, H.: Nitrate in drinking water:
933 A major polluting component of groundwater in gulf region aquifers, south of Togo. International Journal
934 of the Physical Sciences 7(1), pp. 144 - 152., 2012.
- 935 Masoud, A. A.: Spatio-temporal evaluation of the groundwater quality in Kafr Al-Zayat District, Egypt.”
936 Hydrological Processes. Hydrol. Process. 27, 2987–3002., 2013.
- 937 Matini, L., Moutou, J. M. and Kongo-Mantono, M. S.: Evaluation hydro-chimique des eaux souterraines en milieu
938 urbain au Sud-Ouest de Brazzaville, Congo. Afrique SCIENCE 05(1): 82 - 98., 2009.
- 939 Matini, L., Tathy, C. and Moutou, J. M.: Seasonal Groundwater Quality Variation in Brazzaville, Congo.”
940 Research Journal of Chemical Sciences 2(1): 7-14., 2012.
- 941 Mattern S., Fasbender, D. and Vanclooster, M.: Discriminating sources of nitrate pollution in a sandy aquifer.
942 Journal of hydrology 376:275-284., 2009.
- 943 Mattern, S. and Vanclooster, M.: Estimating travel time of recharge water through the unsaturated zone using
944 transfer function model Environmental Fluid Mechanics. DOI 10.1007/s10652-009-9148-1., 2009.
- 945 Mattern, S., Raouafi, W., Bogaert, P., Fasbender, D. and Vanclooster, M.: Bayesian Data Fusion (BDF) of
946 monitoring data with a statistical groundwater contamination model to map groundwater quality at the
947 regional scale. Journal of Water Resource and Protection Vol.4 (No.11), 2012.
- 948 Mbewe, S.: Characterisation and temporal variability assessment of groundwater quality in petauke town, Eastern
949 province, Zambia. Master of Science in Integrated Water Resources Management. Department of
950 Geology, School of Mines. University of Zambia. 134pp., 2013.
- 951 McKenzie, J. M., Mark, B. G., Thompson, L. G., Schotterer, U. and Lin, P.-N.: A hydrogeochemical survey of
952 Kilimanjaro (Tanzania): implications for water sources and ages. Hydrogeology Journal 18(4): 985-995.,
953 2010.
- 954 Messameh, A., Benkhaled, A. and Achour, B.: Numerical Modelling of Groundwater Anthropic Pollution in
955 Urban Area: Biskra City, Algeria. 13th International Conference on Urban Drainage, Sarawak, Malaysia,
956 7–12 September 2014: 6pp., 2014.
- 957 Kihumba, A. M., Longo, J. N., and Vanclooster, M.: Modelling nitrate pollution pressure using a multivariate
958 statistical approach: the case of Kinshasa groundwater body, Democratic Republic of Congo.
959 Hydrogeology Journal: 1-13, 2015.
- 960 Mjemah, I. C.: Saltwater Intrusion and Nitrate Pollution in the Coastal Aquifer of Dar es Salaam.” International
961 Workshop, Tanzania, Rome, 22 April 2013: 31pp.: 2013.
- 962 Mkandawire, T.: Quality of groundwater from shallow wells of selected villages in Blantyre District, Malawi.
963 Physics and Chemistry of the Earth, Parts A/B/C 33(8-13): 807-811., 2008.
- 964 Mpakam, H. G., Kenmogne, G. R. K., Tatiétsé, T. T., Maire, E., Boeglin, J.-L., Ekodeck, G. E. and Dupré, B.:
965 Étude des facteurs de pollution des ressources en eau en milieu urbain : cas de Bafoussam (Ouest-
966 Cameroun). 27pp., 2009.
- 967 Mpenyana-Monyatsi, L. and Momba, M.: Assessment of groundwater quality in the rural areas of the North West
968 Province, South Africa. Scientific Research and Essays 7(8): 903-914., 2012.
- 969 Mtoni, Y., Mjemah, I. C., Bakundukize, C., Van Camp, M., Martens, K. and Walraevens, K.: Saltwater intrusion
970 and nitrate pollution in the coastal aquifer of Dar es Salaam, Tanzania. Environmental earth sciences
971 70(3): 1091-1111., 2013.



- Muiuane, E.: The Quality of Groundwater in and around Maputo city, Mozambique. Department of Geology, Eduardo Mondlane University
<http://www.waternetonline.ihe.nl/downloads/uploads/symposium/zambia2007/Water%20and%20Society/Muiuane.pdf>: 11pp., 2007. (last access: 13 August 2014).
- Musekiwa, C. and Majola, K.: Groundwater Vulnerability Map for South Africa." South African Journal of Geomatics 2(2): pp.152-163., 2013.
- Nachiyunde, K., Ikeda, H., Tanaka, K. and Kozaki, D.: Evaluation of portable water in five provinces of Zambia using a water pollution index. African Journal of Environmental Science and Technology 7(1): pp. 14-29., 2013.
- Nair, G. A., Bohjuari, J. A., Al-Mariami, M. A., Attia, F. A. and El-Touml, F.: Groundwater quality of north-east Libya. Journal of Environmental Biology 27(4): 695-700., 2006.
- Napacho, Z. A. and Manyele, S. V.: Quality assessment of drinking water in Temeke District (part II): Characterization of chemical parameters. African Journal of Environmental Science and Technology 4(11): pp. 775-789., 2010.
- Nedaw, D.: Water Balance and Groundwater Quality of Koraro Area, Tigray, Northern Ethiopia. (MEJS) 2(2): 110-127., 2010.
- Nelson, A.: African Population Database, UNEP GRID Sioux Falls. Retrieved "date of download" from Please cite documentation as follows: Nelson, Andy, 2004. African Population Database Documentation, UNEP GRID Sioux Falls. Retrieved 1/27/2011 from UNEP/GRID Sioux Falls., 2004.
- Ngatcha, B. N. and Daira, D.: Nitrate pollution in groundwater in two selected areas from Cameroon and Chad in the Lake Chad basin. Water Policy 12(5): 722-733., 2010.
- Nkansah, M.A., Ofosuah, J. and Boakye, S.: Quality of groundwater in the Kwahu West district of Ghana. American Journal of Scientific and Industrial Research 1(3): 578-584., 2010.
- Nolan, B. T.: Relating Nitrogen Sources and Aquifer Susceptibility to Nitrate in Shallow GroundWaters of the United States. U.S. Geological Survey, 23pp., 2001.
- Nolan, B. T., Hitt, K. J. and Ruddy, B. C.: Probability of Nitrate Contamination of Recently Recharged Groundwaters in the Conterminous. Environmental science & technology 36(10), 2138-2145., 2002.
- Nolan, B. T. and Hitt, K. J.: Vulnerability of shallow groundwater and drinking-water wells to nitrate in the United States. Environmental science & technology 40(24): 7834-7840., 2006.
- Nolan, B.T., Gronberg, J.M., Faunt, C.C., Eberts, S.M., Belitz, K.: Modeling nitrate at domestic and public-supply well depths in the Central Valley, California. Environ. Sci. Technol. 48 (10), 5643–5651, 2014.
- Nougang, M. E., Nola, M., Djuikom, E., Euphrasie, O. V., Moungang, L. M. and Bessa, H. A.: Abundance of Faecal Coliforms and Pathogenic E. coli Strains in Groundwater in the Coastal Zone of Cameroon (Central Africa), and Relationships with Some Abiotic Parameters. Current Research Journal of Biological Sciences 3(6), pp.622-632., 2011.
- Nwankwoala, H. O. and Udom, G. J.: Hydrogeochemical evaluation of groundwater in parts of Eastern Niger Delta, Nigeria. Journal of Academic and Applied Studies 1(1): pp. 33-58., 2011.
- Nyarko, A. A.: Assessment of groundwater quality and urban water provision: A case of Taifa township in the Ga-east district of the greater Accra region, Ghana. MSc Environmental Science, Kwame Nkrumah University of Science and Technology(KNUST): 72pp., 2008.
- Obi, C. N. and George, P.: The Microbiological and Physico-chemical Analysis of Borehole waters used by Off-Campus Students of Michael Okpara University of Agriculture Umudite (MOUUAU), Abia-State, Nigeria. Research Journal of Biological Sciences 6(11): 602-607., 2011.
- Obinna, N. C., Ezeabasili, A. C. C. and Okoro, B. U.: Assessment of ground water quality in Awka, Anambra state, Nigeria. Journal of Environmental Science and Water Resources 3(4): pp. 080 - 085., 2014.
- Obuobie, E. and Barry, B.: Groundwater in sub-Saharan Africa: Implications for food security and livelihoods, Ghana Country Status on Groundwater. Final Report: 48pp., 2010.
- Ojuri, O. O. and Bankole, O. T. : Groundwater Vulnerability Assessment and Validation for a Fast Growing City in Africa: A Case Study of Lagos, Nigeria. Journal of Environmental Protection 04(05): 454-465., 2013.
- Olden, J. D., Lawler, J. T., and Poff, N.L.: Machine learning methods without tears: a primer for ecologists. The Quarterly review of biology 83(2): 171-193., 2008.
- Oliveira, S., Oehler, F., San-Miguel-Ayanz, J., Camia, A. and Pereira, J. M. C.: Modeling spatial patterns of fire occurrence in Mediterranean Europe using Multiple Regression and Random Forest. Forest Ecology and Management 275: 117-129., 2012.
- Omoboriwo, A. O., Chiaghanam, O. I., Soronnadi-Ononiwu, G. C., Acra, E. J., Okengwu, K. O., Ugwueze, C. U., Yikarebogha, Y. and Momta, P. S.: Appraisal of the Groundwater Quality in Arochukwu Area, Afikpo Basin, Nigeria. International Journal of Science and Technology 2(11): pp.788-793., 2012.
- Ornguga, T. T.: Groundwater Quality Assessment and Sanitary Surveillance of Boreholes in Rural Areas in Benue State Of Nigeria. Academic Journal of Interdisciplinary Studies 3(5), pp153-158., 2014.
- Osemwegie, I., Oga, Y. M.-S., Ahoussi, K. E., Koffi, Y. B., Kouassi, A. M. and Biémi, J.: Influence of Anthropogenic Activities of Groundwater from Hand Dug Wells within the Precarious Settlements of



- 1033 Southern Abidjan, Côte d'Ivoire: Case of the Slums of Anoumabo (Marcory) and Adjouffou (Port-
1034 Bouët). *Journal of Water Resource and Protection* 05(04): 427-439., 2013.
- 1035 Ouedraogo, I; Defourny, P; Vanclooster, M.: Mapping the groundwater vulnerability for pollution at the pan
1036 African scale. *Science of the Total Environment*, 544, 939–953,(December 2015),2016.
- 1037 Ouedraogo, I ; Vanclooster, M.: Shallow groundwater poses pollution problem for Africa. In: *SciDev.Net*, 2016.
1038
- 1039 Pavelic, P., Giordano, M., Keraita, B., Ramesh, V. and Rao, T.: Eds. *Groundwater availability and use in sub-*
1040 *saharan africa: a review of 15 countries*. Colombo, Sri Lanka: International Water Management Institute
1041 (IWMI). 274 p., 2012.
- 1042 Pineros-Garcet J.D., Ordonnez, A., Roosen, J. and Vanclooster, M.: Metamodelling: theory, concepts and
1043 application to nitrate leaching modelling. *Ecological modelling* 193: 629-644., 2006.
- 1044 Potter, P., Ramankutty, N., Bennett, E. M., and Donner, S.D.: Characterizing the Spatial Patterns of Global
1045 Fertilizer Application and Manure Production. *Earth Interactions* 14: 1-22., 2010
- 1046 Prasad, A. M., Iverson, L.R. and Liaw, A.: Newer classification and regression tree techniques: bagging and
1047 random forests for ecological prediction. *Ecosystems* 9(2): 181-199., 2006.
- 1048 Rawlings, J.O, Pantula, S.G, Dickey, D.A.: *Applied regression analysis, a research tool*, Springer; 658p., 1998
1049
- 1050 R Development Core Team.: *A language and environment for statistical computing*. Vienna: R Fundation
1051 Statistical Computing. (Accessed free. Available at <http://www.r-project.org/>),2015.
- 1052 Rim-Rukeh, A., Ikhifa, G. O. and Okokoyo, P. A.: Physico-chemical characteristics of some waters used for
1053 drinking and domestic purposes in the Niger Delta, Nigeria. *Environ Monit Assess* 128(1-3): 475-482.,
1054 2007.
- 1055 Rosillon, F., Savadogo, B., Kabore, A., Bado-Sama, H. and Dianou, D.: Attempts to Answer on the Origin of the
1056 High Nitrates Concentrations in Groundwaters of the Sourou Valley in Burkina Faso. *Journal of Water*
1057 *Resource and Protection* 04(08): 663-673., 2012.
- 1058 Rouabhia, A., Baali, F., Hani, A. and Djabri, L.: Impact des activités anthropiques sur la qualité des eaux
1059 souterraines d'un aquifère en zone semi-aride: Cas de la plaine de la Merdja, Nord Est de l'Algérie.
1060 *Sécheresse* 20(3), 2009.
- 1061 Rouabhia, A., Baali, F. and Fehdi, C.: Impact of agricultural activity and lithology on groundwater quality in the
1062 Merdja area, Tebessa, Algeria. *Arabian Journal of Geosciences* 3(3): 307-318.,2010.
- 1063 Sadek, M. and El-Samie, A. S.: Pollution vulnerability of the Quaternary aquifer near Cairo, Egypt, as indicated
1064 by isotopes and hydrochemistry. *Hydrogeology Journal* 9(3): 273-281.,2001.
- 1065 Saffigna, P. G. and Keeney, D. R.: Nitrate and Chloride in Groundwater under irrigated agriculture in Central
1066 Wisconsin. *Groundwater* 15(2): 170-177., 1997.
- 1067 Salem, M. A. and Alshergawi, M. I.: Physico-chemical Evaluation of Drinking Water Quality in Alshati District
1068 of Libya. *IOSR Journal Of Environmental Science, Toxicology And Food Technology* 4(1): pp.41-51.,
1069 2013.
- 1070 Salim, L. H. M., Elhassan, B. M. and Elawad, M. A.: Physicochemical Analysis of Surface and Ground Water in
1071 Karrary Locality, Sudan. *International Journal of Science, Engineering and Technology Research*
1072 *(IJSETR)* 3(3): pp.436-438., 2014.
- 1073 Sall, M. and Vanclooster, M.: Assessing the well water pollution problem by nitrates in the small scale farming
1074 systems of the Niayes region, Senegal. *Agricultural Water Management* 96(9): 1360-1368., 2009.
- 1075 Sanok, F. M., Toriman, M. E. b. and Mokhtar, M.: Evaluation of Spatial and Temporal Variation in North East
1076 Jabal Al Hasawna wellfields Water quality, Libya. *International Journal of Advances in Agricultural &*
1077 *Environmental Engineering* 1(1).pp23-28., 2014.
- 1078 Sbargoud, S.: Validité de l'application de la méthode de vulnérabilité DRASTIC à l'étude de la pollution par les
1079 nitrates dans la Mitidja (Nord d'Algérie). *Geotunis* 2013: 22pp., 2013.
- 1080 Seeber, K., Daira, D., Bala, A. M. and Vassolo, S.: Lake Chad Basin: Sustainable Water Management,
1081 Groundwater Quality Investigations in the Lower Logone Floodplain in April – May 2013. 47pp., 2014.
- 1082 Semar, A., Saïbi, H. and Medjerab, A.: Contribution of multivariate statistical techniques in the hydrochemical
1083 evaluation of groundwater from the Ouargla phreatic aquifer in Algeria. *Arabian Journal of Geosciences*
1084 6(9), 3427-3436., 2013.
- 1085 Sharaky, A. M., Atta, S. A., El Hassanein, A. S. and Khallaf, K. M. A.: Hydrogeochemistry of groundwater in
1086 the western Nile delta aquifers, Egypt. 2nd International Conference on the Geology of Tethys, 19-21
1087 March, 2007, Cairo University: 23pp., 2007.
- 1088 Showers, W. J., Genna, B., McDade, T., Bolich, R. and Fountain, J. C.: Nitrate contamination in groundwater on
1089 an urbanized dairy farm. *Environmental science & technology* 42(13): 4683-4688., 2008.
- 1090 Smahi, D.: Environmental Impact of Casablanca Landfill on Groundwater Quality, Morocco. *International Journal*
1091 *of Geosciences* 04(01): 202-211., 2013.



- Smida, H., Abdellaoui, C., Zairi, M. and Dhia, H. B.: Cartographie des zones vulnérables à la pollution agricole par la méthode DRASTIC couplée à un Système d'information géographique (SIG) : cas de la nappe phréatique de Chaffar(sud de Sfax, Tunisie). *Sécheresse* 21(2), 131-146., 2010.
- Sophocleous, M.: GROUNDWATER RECHARGE, in Groundwater, [Eds. Luis Silveira, Stefan Wohnlich, and Eduardo J. Usunoff], in Encyclopedia of Life Support Systems (EOLSS), Developed under the Auspices of the UNESCO, Eolss Publishers, Oxford, UK, [http://www.eolss.net]., 2004.
- Sorichetta, A., Ballabio, C., Masetti, M., Robinson, G. R. Jr. and Sterlacchini, S.: A comparison of data-driven groundwater vulnerability assessment methods. *Ground Water* 51(6): 866-879., 2013.
- Sorlini, S., Palazzini, D., Sieliechi, J. and Ngassoum, M.: Assessment of Physical-Chemical Drinking Water Quality in the Logone Valley (Chad-Cameroon). *Sustainability* 5(7): 3060-3076., 2013.
- Spalding, R. F. and Exner, M. E.: Occurrence of nitrate in groundwater- a review. *J. Environ. Qual.* 22(392-402)., 1993.
- Stadler, S., Osenbrück, K., Knöller, K., Himmelsbach, T., Suckow, A. and Weise, S. M.: Origin of nitrate in groundwaters of Kalahari, Botswana. *European Society for Isotope Research, eggauberg Austria* 27.6 – 1.7.2004: 3pp., 2004.
- Stadler, S., Osenbrück, K., Knöller, K., Suckow, A., Sültenfuß, J., Oster, H., Himmelsbach, T. and Hötzel, H.: Understanding the origin and fate of nitrate in groundwater of semi-arid environments. *Journal of Arid Environments* 72(10): 1830-1842., 2008.
- Steichen, J., Koelliker, J., Grosh, D., Heiman, A., Yearout, R. and Robbins, V.: Contamination of farmstead wells by pesticides, volatile organics, and inorganic chemicals in Kansas. *Groundwater Monitoring & Remediation* 8(3), 153-160., 1988.
- Suthar, S., Bishnoi, P., Singh, S., Mutiyar, P. K., Nema, A. K. and Patil, N. S.: Nitrate contamination in groundwater of some rural areas of Rajasthan, India. *J Hazard Mater* 171(1-3): 189-199., 2009.
- Tabue, Y. J. G., Feumba, R., Wethe, J., Ekodeck, G. E. and De, M. G.: Evaluation of Groundwater Suitability for Domestic and Irrigational Purposes: A Case Study from Mingoa River Basin, Yaounde, Cameroon." *Journal of Water Resource and Protection*, 2012.
- Tabue Y. J. G., Ntamack, D., Feumba, R., Ngnikam, E., Wéthé, J. and Tanawa, É.: Vulnérabilité des eaux souterraines et périmètres de protection dans le bassin versant de la Mingoa (Yaoundé, Cameroun). *Revue de l'Université de Moncton* 40(2): 71., 2009.
- Tadesse, N., Bheemalingeswara, K. and Abdulaziz, M.: Hydrogeological Investigation and Groundwater Potential Assessment in Haromaya Watershed, Eastern Ethiopia. *Mekelle University* 2(1): 26-48., 2010.
- Tagma, T., Hsissou, Y., Bouchaou, L., Bouragba, L. and Boutaleb, S.: Groundwater nitrate pollution in Souss-Massa basin (south-west Morocco). *African Journal of Environmental Science and Technology* Vol. 3(10): pp. 301-309., 2009.
- Taha, G. M. E. E.: Determination, Source Identification and GIS Mapping for Nitrate Concentration in Ground water from Bara Aquifer. B.Sc. Honor (Chemistry), Faculty of Science, University of Khartoum: 95pp., 2010.
- Talabi, A. O.: Geostatistical Assessment of Groundwater Quality from coastal aquifers of Igbokoda, Southwestern Nigeria. *Journal Of Environmental Science, Toxicology And Food Technology (IOSR-JESTFT)* 2(2): PP 18-24., 2012.
- Tay, C. and Kortatsi, B.: Groundwater Quality Studies: A Case Study of the Densu Basin, Ghana. *West African Journal of Applied Ecology* Vol.12: 18pp., 2008.
- Tegegn, F. E.: Physico-chemical pollution pattern in Akaki River basin, Addis Ababa, Ethiopia. Master's thesis Physical Geography and Quaternary Geology, Stockholm University, 38pp., 2012.
- Temgoua, E.: Chemical and Bacteriological Analysis of Drinking Water from Alternative Sources in the Dschang Municipality, Cameroon. *Journal of Environmental Protection* 02(05), 620-628., 2011.
- Tesoriero, A. J. and Voss, F. D.: Predicting the probability of Elevated Nitrate Concentrations in the Puget Sound-Basin, Implications for Aquifer Susceptibility and Vulnerability. *Ground Water* 35(6): 1029-1039., 1997.
- Tilahun, K. and Merkel, B. J.: Assessment of groundwater vulnerability to pollution in Dire Dawa, Ethiopia using DRASTIC. *Environmental Earth Sciences* 59(7): 1485-1496., 2010.
- Todd Engineers and Kennedy/Jenks Consultants. Revised Final Groundwater Vulnerability Study Santa Clara County, California. Project Report, 92 pp. October 2010.
- Totin, H. S. V., Boukari, M., Faye, S., Alassane, A., Orékan, V. and Boko, M.: Climate and Land Use Change Impacts on Groundwater Quality in the Beninese Coastal Basin of the Transboundary Aquifer System Benin-Nigeria-Togo. International Conference "Transboundary Aquifers: Challenges and New Directions"(ISARM2010), 6-8 December 2010: 4pp., 2010.
- Totin, H. S. V., Amoussou, E., Odoulami, L., Edorh, P. A., Boukari, M. and Boko, M.: Groundwater pollution and the safe water supply challenge in Cotonou town, Benin (West Africa). *Proceedings of H04, IAHS-IAPSO-IASPEI Assembly, Gothenburg, Sweden, July 2013 IAHS Publ.* 361., 2013.



- 1151 Trambauer, P., E. Dutra, S. Maskey, M. Werner, F. Pappenberger, L. P. H. van Beek and S. Uhlenbrook.:
1152 Comparison of different evaporation estimates over the African continent. *Hydrol. Earth Syst. Sci.* 18(1):
1153 193-212., 2014.
- 1154 UNEP/DEWA.: Sanitation and Groundwater Protection –a UNEP Perspective UNEP/DEWA,
1155 [http://www.bgr.bund.de/EN/Themen/Wasser/Veranstaltungen/symp_sanitatgwprotect/present_mmayi_](http://www.bgr.bund.de/EN/Themen/Wasser/Veranstaltungen/symp_sanitatgwprotect/present_mmayi_.pdf.pdf?__blob=publicationFile&v=2)
1156 [pdf.pdf?__blob=publicationFile&v=2](http://www.bgr.bund.de/EN/Themen/Wasser/Veranstaltungen/symp_sanitatgwprotect/present_mmayi_.pdf.pdf?__blob=publicationFile&v=2): 18pp.,(last access: 13 August 2014).
- 1157 U.S. Environmental Protection Agency: Environmental indicators of water quality in the United States:
1158 Washington, D.C., U.S. Environmental Protection Agency, Office of Water, EPA 841- R-96-002, 25p.,
1159 1996.
- 1160 Wakida, F. T. and Lerner, D. N.: Non-agricultural sources of groundwater nitrate: a review and case study." *Water*
1161 *research* 39(1): 3-16., 2005.
- 1162 Wheeler, D. C., Nolan, B. T., Flory, A. R., DellaValle, C. T. and Ward, M. H.: Modeling groundwater nitrate
1163 concentrations in private wells in Iowa. *Sci Total Environ* 536: 481-488., 2015.
- 1164 Wiens, J. A.: Spatial scaling in ecology. *Functional ecology* 3(4): 385-397., 1989.
- 1165
- 1166 Winkel, L., Berg, M., Amini, M., Hug, S. J., and Annette Johnson, C.: Predicting groundwater arsenic
1167 contamination in Southeast Asia from surface parameters. *Nature Geosci* 1(8): 536-542., 2008.
- 1168 World Health Organization (WHO):. Rolling Revision of the WHO Guidelines for Drinking-Water Quality. Draft
1169 for review and comments, Nitrates and nitrites in drinking-water. WHO/SDE/WSH/04.08/56: 133pp.,
1170 2004 (last access, 15 September 2015).
- 1171 Worrall, F. and Besien, T.: The vulnerability of groundwater to pesticide contamination estimated directly from
1172 observations of presence or absence in wells. *Journal of Hydrology* 303(1-4): 92-107., 2005.
- 1173 Wotany, E. R., Ayonghe, S. N., Fantong, W. Y., Wirmvem, M. J. and Ohba, T.: Hydrogeochemical and
1174 anthropogenic influence on the quality of water sources in the Rio del Rey Basin, South Western,
1175 Cameroon, Gulf of Guinea. *African Journal of Environmental Science and Technology* 7(12): pp.1053-
1176 1069., 2013.
- 1177 Xu, Y. and Usher, B.: Groundwater pollution in Africa, Taylor&Francis/Balkema, The Netherlands. 353pp., 2006.
- 1178 Yang, C. Y., Cheng, M. F., Tsai, S.S., and Hsieh, Y. L.: Calcium, magnesium, and nitrate in drinking water and
1179 gastric cancer mortality. *Cancer Science* 89(2): 124-130., 1998.
- 1180 Yao, T. K., Oga, M.S., Fouche, O., Baka, D., Pernelle, C. and Biemi, J.: Évaluation de la potabilité chimique des
1181 eaux souterraines dans un bassin versant tropical :Cas du Sud-Ouest de la Côte d'Ivoire." *International*
1182 *Journal of Biological and Chemical Sciences* 6(6)., 2013.
- 1183 Zghibi, A., Tarhouni, J. and Zouhri, L.: Assessment of seawater intrusion and nitrate contamination on the
1184 groundwater quality in the Korba coastal plain of Cap-Bon (North-east of Tunisia). *Journal of African*
1185 *Earth Sciences* 87: 1-12., 2013.
- 1186 Zingoni, E., Love, D., Magadza,C., Moyce, W. and Musiwa, K.: Effects of a semi-formal urban settlement on
1187 groundwater quality." *Physics and Chemistry of the Earth, Parts A/B/C* 30(11-16): 680-688., 2005.
- 1188
- 1189
- 1190
- 1191



1192

Table 1. Criteria used to identify nitrate data studies within web data bases.

Search engine	Search criteria
Google, Google Scholar, and Google Books	Groundwater pollution + Africa Nitrate in groundwater + “African country name” or “African capital city name” Groundwater quality + Africa Nitrate and agricultural practices in Africa Groundwater vulnerability + “African country name” Pollution des eaux souterraines par les nitrates+ ‘nom du pays Africain’ (in French) Pollution des eaux souterraines + "nom du pays Africain" (in French) Nitrate concentrations under irrigated agriculture + “African country name”
Web of Sciences, Scopus and Sciences Direct	Groundwater pollution by nitrate + “African country name” Nitrate in groundwater + “Africa capital city name” Pollution des eaux souterraines par les nitrates + "nom du capital des pays"(in French) Groundwater contamination by nitrate + “Africa countries” or “African capital city name” Africa irrigated agriculture + nitrate Groundwater contamination by nitrate + “Africa country name” or “African capital city name” Nitrate concentrations under irrigated agriculture + “African country name” Groundwater vulnerability to nitrate contamination + “Africa country name” or “African capital city name”
Books	Groundwater pollution in Africa (Xu and Usher, 2006)

1193

1194

1195

1196



Table 2. Localisation of study sites considered in the meta-analysis

Country	Localisation	Number of studies per country	References
Algeria	North east of Algeria	11	Labar et al., 2012a
	Ouargla phreatic aquifer in Algeria: Valley of OuedM'y a		Semar et al., 2013
	Nord-Algerian aquifer (Mitidja)		Sbargoud, 2013
	Medja area		Rouabhia et al., 2010
	Biskra		Messameh et al., 2014
	Case Skikda		Labar et al., 2012b
	El Eulma		Belkhiri and Mouni, 2012
	Mostaganem, Mecheria, Naama, Tiaret, Bechar, and Adrar		Bahri and Saibi, 2012
	Southern Hodna		Abdesselam et al., 2012
	Tlemcen		Abdelbaki et al., 2013
	Merdja plain		Rouabhia et al., 2009
Angola	Angola	1	Angola Water Works, 2013
Benin	Cotonou	6	Totin et al., 2013
	Beninese coastal basin		Totin et al., 2010
	Municipality of Pobè		Lagnika et al., 2014
	Dongo-pont		Bossa et al., 2012
	Cotonou		BGS, 2003
	Cotonou		Xu et al. 2006
Bostwana	Rural Bostwana	3	Batisani, 2012
	Kalahari		Stadler et al., 2004
	Eastern fringe of the Kalahari near Serowe		Stadler et al., 2008
Burkina Faso	Burkina Faso	4	BGS, 2002
	Burkina Faso		Pavelic et al., 2012
	Sourou Valley		Rosillon et al., 2012
	Ouagadougou		Xu et al. 2006
	Mingoa River basin/Yaounde		Tabue et al., 2012
Cameroon	Bafoussam	10	Mpakam et al., 2009
	Coastal zone of Cameroon/Douala		Nougang et al., 2011
	Logon Valley/Chad-Cameroon		Sorlini et al., 2013
	Anga's river		Kuitcha et al., 2013
	Rio del Rey Basin/South Western Coast		Wotany et al., 2013
	Mingoa/Yaounde catchment		Tabue et al., 2009



Central African Republic	2 areas of Cameroon and Chad in the Lake Chad basin Dschang Municipality Cameroon			Ngatcha and Daira, 2010 Temgoua, 2011 Xu et al.2006
	Bangui area	1		Djebebe-Ndjiuim et al., 2013
Congo-Brazzaville	Brazzaville	5		Matini et al., 2012
	Brazzaville			Barhe and Bouaka, 2013
	Soult East Brazzaville			Laurent et al., 2010
	Soult East Brazzaville			Laurent and Marie, 2010
Egypt	South West Brazzaville	7		Matini et al., 2009
	Alexandria			Abd El-Salam and Abu-Zuid, 2014
	Helwan			Abdalla and Scheytt, 2012
	Nile Valley			Abdel-Lah and Shamukh, 2001
	Tahta			Easa and Abou-Rayhan, 2010
	Kafr Al-Zayet District			Masoud, 2013
	Nile Delta aquifers / Western Nile Delta			Sharaky et al., 2007
	Cairo, Egypt / province of Giza			Sadek and El-Samie, 2001
	Dire Dawa			Abate, 2010
	Ethiopia			BGS, 2001
Ethiopia	Raya valley	13		Bushra, 2011
	Adis Ababa			Engida, 2001
	Addis Ababa			Kahsay et al.
	Koraro/Tigray			Nedaw, 2010
	Ethiopia			Pavelic et al., 2012
	Bulbule and Zway			Bonetto et al., 2005
	Haromaya Watershed, Eastern Ethiopia			Tadesse et al., 2010
	Akaki			Tegegn, 2012
	Adis Ababa			Xu et al.2006
	Dire Dawa of Sabian area			Tilahun and Merkel, 2010
	Wondo Genet District, Southern Ethiopia			Haylamicheal and Moges, 2012
	Ga East			Ackah et al., 2011
	Sawla-Tuna-Kalba District			Cobbina et al., 2012
Ghana	Akatsi, Adidome and Ho Districts	14		Ansa- Asare et al., 2009
	Ghana			BGS, 2000
	Six districts in the eastern region of Ghana			Fianko et al., 2009
	Kwahu West District			Nkansah et al., 2010
	Ga-East District of Accra(Taifa)			Nyarko, 2008
	Ghana			Obuobie and Barry, 2010
	Ghana			Pavelic et al., 2012
	Ghana			



Densu basin Contamination in Ghana Western Region of Ghana Gold Mining area in Ghana/Tarkwa Lower Pra Basin of Ghana			Tay and Kortatsi, 2008 Xu et al.2006 Affum et al., 2015 Armah et al., 2012 Armah, 2010
Guinea-Biseau	1		Bordalo and Savva-Bordalo, 2007
Boloma			Abenan et al., 2012
Bonoua			Ahoussi et al., 2012
Bondoukou region			Ake et al., 2010
Bonoua aquifer (South-East Ivory Coast)			Douagui et al., 2012
Abidjan District			Eblin et al., 2014
Adiaké Region			Kouame et al., 2013
Abidjan and Korhogo			Xu et al.2006
Abidjan aquifer			Yao et al., 2013
South-West Ivory Coast			Kouassi et al., 2010
N'zi-Comoé (Centre East Ivory Coast)			Kouassi et al., 2012
Guiglo-Douekoué (West Ivory Coast)			Ahoussi et al., 2012
N'zi, N'Zianouan municipality(South Ivory Coast)			Drissa et al., 2013
Bandama basin at Tortiya(Nothern Ivory Coast)	16		Loko et al., 2013a
Abia Koumassi village/Abidjan			Osemwegie et al., 2013
Slums of Anoumabo (Marcory) and AdioufouPort-Bouet			Dibi et al., 2013
Catchment Ehania, South-Eastern Ivory Coast			Loko et al., 2013b
Hiré , South-West of Ivory Coast			Xu et al.2006
Kisaumi, Mombasa	1		Nair et al., 2006
North-East Libya			Salem and Alshergawi, 2013
Alshati	3		Sanok et al., 2014
North East Jabal Al Hasawnah			Xu et al.2006
Lake Chilwa basin			Grimason et al., 2013
Chikhwawa	4		Kanyere et al., 2012
Upper Limphasa River/Nkhata-Bay district			Mkandawire, 2008
Blantyre			Xu et al.2006
Bamako city			Pavelic et al., 2012
Mali	3		Cronin et al., 2007
Timbuktu			Friedel, 2008
Mauritania	1		Ben Abbou et al., 2014
Oued Taza			Aghzar et al., 2002
Taldla plain			Alaoui et al., 2008
Marrakesh	16		Belghiti et al., 2013
Meknès region			Benabbou et al., 2014
Oum Azza of Rabat			



Phreatic aquifer of M'nasra		Bricha et al., 2007
Taldla plain		EL Hammoumi et al., 2013
Mzamza-Chouia		Asslouj et al., 2007
Berrechid plain		EL Bouqdaoui et al., 2009
Taldla plain		EL Hammoumi et al., 2012
Triffa plain		Fekkoul et al., 2011
Triffa plain		Fetouani et al., 2008
Essaouira Basin		Laftouhi et al., 2003
Phreatic aquifer of Martil		Lamribah et al., 2013
Casablanca		Smahi, 2013
Souss-Massa basin (South-west Morocco)		Tagma et al., 2009
Mozambique		
Lichinga		Cronin et al., 2007
Maputo city	2	Muiuane, 2007
Niamey		Chippaux et al., 2002
Niamey	3	Hassane, 2010
Niamey		Abou, 2000
Uzouwani (South Eastern Nigeria)		Ekere, 2012
Lagos		Adelekan and Ogunde, 2012
Ondo State		Akinbile, 2012
Southwestern Abeokuta		Aladejana and Talabi, 2013
Lagos		Anthony, 2012
Lagos-State		Balogun et al., 2012
Nigeria		BGS, 2003
Konduga town		Dammo et al., 2013
Abuja		Dan-Hassan et al., 2012
Nigeria		Edet et al., 2011
Edo State/ South-South		Imoisi et al., 2012
Jimeta-Yola (Northeastern of Nigeria)		Ishaku, 2011
Eastern Niger Delta		Nwankwoala and Udom, 2011
Anambra State		Obinna et al., 2014
Lagos		Ojuri and Bankole, 2013
Afikpo basin		Omaboriwo et al., 2012
Benue State	24	Ornguga, 2014
Nigeria		Palevlic et al., 2012
Niger Delta		Rim-Rukeh et al., 2007
Igbokoda, Southwestern Nigeria		Talabi, 2012
Lagos		Wakida and Lerner, 2005
Nigeria		Xu et al., 2006
Abia state		Obi and George, 2011



	Eti-Osa, Lagos			Akotey and Soladoye, 2011
Republic Democratic of Congo	Kahuzi-Biega National Parks Kinshasa		2	Bagalwa et al., 2013 Longo, 2009
Senegal	Dakar Region Thiaroye Niayes region Dakar Dakar Dakar Yeumbeul/Dakar		7	Brandvold, 2013 Madioune et al., 2011 Sall and Vanclooster, 2009 Wakida and Lerner, 2005 Xu et al., 2006 Diédhiou et al., 2012 BGS, 2003
Somalia	Somaliland and Puntland		1	FAO-SWALIM, 2012
South Africa	South Africa Philippi/Western Cape Mpumalanga Province South Africa South Africa Hex River Valley; Sandveld; Hertzogville		6	Maherry et al., 2009 Aza-Gnandji et al., 2013 Mpenyana-Monyatsi and Momba, 2012 Musekiwa and Majola, 2013 Pavelic et al., 2012 Xu et al., 2006
Sudan	Southern Suburb of the Ondurman Khartoun Karray Karray Khartoun		5	Abdellah et al., 2013 Ahmed et al., 2000 Salim et al., 2014 Taha, 2010 Idriss et al., 2011
Tanzania	Tanzania Dar es Salam Dodoma Kilimandjaro region Dar es Salam Dar es Salam Tanzania Temekedistic/Dar es Salam		8	BGS, 2000 De Witte, 2012 Kashaigili, 2010 McKenzie et al., 2010 Mjemah, 2013 Mtoni et al., 2013 Palevic et al., 2012 Napacho and Manyele, 2010
Tchad	N'djamena Lake Chad basin Chad basin		3	Guideal et al., 2010 Seeber et al., 2014 Ngarcha and Daura, 2010
Togo	Agodè-Zongo Gulf/South of Togo		2	Kissao and Housséni, 2012 Mande et al., 2012
Tunisia	North-east of Tunisia (Korba aquifer) Cap Bon Cap Bon Djebeniana		8	Zghibi et al., 2013 Anane et al., 2014 Charfi et al., 2013 Fedrigoni et al., 2001



Uganda	Metline-Ras Jebel-Raf Raf/North-East Sfax-Agareb El Khairat aquifer Chaffar/ South of Sfax		Hamza et al., 2007 Hentati et al., 2011 Ketata et al., 2011 Smida et al., 2010
	Uganda Kampala/Bwaise III	2	BGS, 2001 Kulabako et al., 2007
	Petauke Town John Laing and Misisi de Lusaka Copperbelt Province/(North Western Province; Lusaka Province ; Central Province ; Southern Province) Lusaka	4	Mbewe, 2013 Xu et al.2006 Nachiyunde et al., 2013 Wakida and Lerner, 2005
Zambia			
Zimbabwe	Kamangara Epworth at Harare	2	Dzwairo et al., 2006 Zingoni et al., 2005



Table 3. Explanatory variables used in the MLR analysis.

Explanatory variables	Type	Units or Categories	Spatial resolution/Scale	Date	Data source(s)
Land Cover/Land Use	Categorical data	-	300 m	2014	¹ UCL/ELIe-Geomatics (Belgium)
Population density	Continuous point data	people/km ²	2.5 km	2004	ESRI : www.arcgis.com/home
Nitrogen application	Continuous point data	kg/ha	0.5° x 0.5°	2009	² SEDAC : www.sedac.ciesin.columbia.edu
Climate class data	Categorical data	-	0.5°	1997	Global-Aridity values (UNEP, 1987)/ (UNESCO-IHE, Delft, The Netherlands)
Type of regions	Categorical data	-	0.5°	2014	Global-Aridity values (UNEP, 1987)/ (UNESCO-IHE, Delft, The Netherlands)
Rainfall class	Categorical data	mm/year	3.7 km	1986	UNEP : http://www.grid.unep.ch
Depth to groundwater	Categorical data	m	0.5° x 0.5°	2012	British Geological Survey: www.bgs.ac.uk/
Aquifer type	Categorical data	-	1:3750 000	2012	³ GLiM data (Hamburg University)
Soil type	Categorical data	-	1 km x 1 km	2014	ISRIC, World Soil Information: www.isric.org/content/soilgrids
Unsaturated zone (impact of vadose zone)	Categorical data	-	1:3750 000	2012	GLiM data (Hamburg University)
Topography/Slope	Continuous point data	Percentage (%)	90 m	2000	⁴ UCL/ELIe-Geomatics (Belgium) and CGIAR/CSI
Recharge	Continuous point data	mm/year	5 km	2008	Global-scale modelling of groundwater recharge (University of Frankfurt)
Hydraulic conductivity	Continuous point data	m/day	Average size of polygon ~100km ²	2014	⁵ GLHYMPS data (McGill University)

¹Université Catholique de Louvain/Earth and Life Institute/Environnemental sciences;

²Socioeconomic Data and Applications Center (SEDAC);

⁴Consultative Group for International Agricultural Research (CGIAR)/ Consortium for Spatial Information (CSI);

³The new global lithological map database GLiM: A representative of rock properties at the Earth surface;

⁵A glimpse beneath earth's surface: Global Hydrogeology MaPS (GLHYMPS) of permeability and porosity.



Table 4. Summary statistics of original and log (ln) transformed nitrate data.

Statistic	Maximum NO ₃ ⁻ concentration	Maximum ln(NO ₃ ⁻) concentration	Mean NO ₃ ⁻ concentration	Mean ln(NO ₃ ⁻) concentration	Minimum NO ₃ ⁻ concentration	Minimum ln(NO ₃ ⁻) concentration
Number of data (-)	206	206	82	82	185	185
Minimum (mg/l or ln(mg/l))	0.08	-2.52	1.26	0.231	0	0
Maximum (mg/l or ln(mg/l))	4625	8.43	648	6.473	180	5.19
Median (mg/l or ln(mg/l))	73.64	4.29	27.58	3.317	0.55	0.43
Mean (mg/l or ln(mg/l))	190.05	3.99	54.85	3.169	8.91	1.08
Variance ((mg/l) ² or ln(mg/l) ²)	183778.94	3.39	163.92	43.901	537.07	1.78
CV (-)	225.56	46.18	8085.08	1.935	260.08	123.04
Standard Deviation (mg/l or ln(mg/l))	428.69	1.84	89.91	1.391	23.17	1.33
Kurtosis	60.24	0.90	23.99	-0.167	25.57	0.37
Skewness	6.75	-0.74	4.31	-0.294	4.56	1.2



Table 5. Optimal linear regression model for explaining the logtransformed mean nitrate concentration

Coefficients:				
	Estimate	Std. Error	t value	Pr (> t)
(Intercept)	3.348e+00	6.624e-01	5.055	3.56e-06 ***
Depth [0-7]	1.160e+00	3.895e-01	2.977	0.00404 **
Depth [7-25]	6.563e-01	3.693e-01	1.778	0.08002*
Depth [25-50]	1.114e+00	4.755e-01	2.342	0.02216 **
Depth [50-100]	6.536e-01	4.005e-01	1.632	0.10744
Depth [100-250]	4.258e-01	6.766e-01	0.629	0.53129
Recharge [0-45]	-2.506e-01	6.089e-01	-0.412	0.68200
Recharge [45-123]	-1.187e+00	6.055e-01	-1.961	0.05407*
Recharge [123-224]	-1.112e+00	6.134e-01	-1.812	0.07440*
Recharge [224-355]	-8.856e-01	6.089e-01	-1.455	0.15047
Aquifer media [Crystalline rocks]	-9.851e-01	3.374e-01	-2.920	0.00477 **
Aquifer media [Siliciclastic sedimentary rocks]	1.893e-02	3.916e-01	0.048	0.96158
Aquifer media [Unconsolidated sediments rocks]	-7.632e-01	3.384e-01	-2.255	0.02740 **
Aquifer media [Volcanic rocks]	-5.245e-01	6.123e-01	-0.857	0.39469
Population density (people/km ²)	5.611e-04	6.887e-05	8.147	1.30e-11 ***
Residual standard error: 0.9116 on 67 degrees of freedom				
Multiple R-squared: 0.65				
F-statistic: 8.693 on 14 and 67 DF, p-value=2.422e-10 < 0.001				

Note: *** significant at $p < 0.001$; ** significant at $p < 0.05$ and * significant at $p < 0.1$




Fig. 1. Hydrogeological setting of the African continent (from Hartmann and Moosdorf, 2012).




Fig. 2. Distribution of studies identified across Africa.




Fig. 3. The locations and the maximum values of nitrate in Africa superimposed on risk pollution map as generated in the previous generic vulnerability study of Ouedraogo et al., 2016.




Fig. 4. Land Cover/Land Use map of Africa (modified from Defourny et al., 2014).




Fig. 5. Histograms of observed mean nitrate concentration (mg/l) and logtransformed mean nitrate concentration (ln (mg/l)).




Fig. 6. Log transformed mean nitrate concentration for different land use classes.




Fig. 7. Log transformed mean nitrate concentration for different aquifer system classes.




Fig. 8. Log transformed mean nitrate concentration for different groundwater depth classes, recharge classes, population density classes and nitrogen application rate classes.




Fig. 9. Predicted versus observed mean logtransformed nitrate concentration ($R^2=0.65$).




Fig. 10. Normal probability distribution of model residuals for the predicted logtransformed mean nitrate concentration.




Fig. 11. Relation between residuals and predicted logtransformed mean nitrate concentration.