

Chapter 3

Dyle Catchment: Description of the Study Area

In order to avoid redundancy in the subsequent chapters, the study area is described in this chapter.

The study area is the Walloon region part of the Dyle catchment (580 km²) of central Belgium (Figure 3.1). This is a typical ‘arbitrary’ scale in the sense of Baveye and Boast (1999), i.e. not related to the physical processes occurring in the system under study; but it is entirely embedded in the hydrogeological catchment of a sandy aquifer, thus making this choice consistent. The climate is temperate maritime, with a mean annual temperature of 9.7°C and mean summer and winter temperatures of 16.4°C and 3.0°C, respectively. Average annual precipitation is 804 mm.

Figure 3.2 shows the main land uses in the Dyle catchment. Typical land uses are urban (generally located in the valleys; about 20% of land use in the study area), pasture and forests (found on valley slopes; about 16 and 18% respectively), and arable (silt and silty loam soils on the plateau). The simulations of atrazine leaching presented in the following chapters were restricted to arable land, which covers approximately 46% of the study area (*Système Intégré de Gestion Et de Contrôle* (SIGEC), 1999).

In chapter 4, the study area is extended to the whole Brusselian aquifer. This aquifer is located in tertiary sands overlain by a quaternary loess layer of variable thickness (0 to 15 m). This groundwater body is of primary importance for the regional supply of drinking water and has been classified




Figure 3.1: Location of the study area in Belgium.

as a ‘vulnerable area’, resulting from the implementation of the European Union Nitrate Directive (EEC, 1991). Figure 3.3 shows a hydrogeological section between Louvain-la-Neuve and Wavre (cf. section line in Figure 3.2). Figure 3.3 shows that the loess layer is mainly present on the plateau, while it is eroded in the valleys where the Brusselian sands appear. From this section it can be noticed that groundwater table depth varies from 0 to more than 30 metres¹.

Figure 3.4 displays a map of the main soil types Dyle catchment. This map results from a reclassification of the 1:20,000 soil map, using texture and drainage properties, profile development and the presence/absence of a change of substrate.

Silt is clearly the dominant soil type in the area, especially on the plateau in the east, south-east and south-west of the catchment. Silt with no profile development corresponds to alluvial deposits in the small depressions. In

¹The hydrogeological mapping project from which this section was obtained is still under development, and maps were not available for the whole study area at the time this thesis was undertaken




Figure 3.2: Land use in the Dyle catchment. Sources: SIGEC (1999) and Landsat TM image (1999). The red line shows the location of the hydrogeological section displayed in Figure 3.3.

the valleys, the loess layer has been partially to completely eroded, leaving shallow soils on a sandy substrate or even the Brusselian sands uncovered.

A mask over non-arable areas was added to Figure 3.5. This shows that arable land use is almost exclusively located on silty soils, thus reducing the variability of soil properties for the simulations presented in chapters 6 to 8.




Figure 3.3: Hydrogeological section between Louvain-la-Neuve and Wavre.
Source: DGRNE (2002).




Figure 3.4: Soil types in the Dyle catchment (scale 1:20,000).




Figure 3.5: Soil map with a white mask on non-arable areas.