

Review Article on Modern Technology in Liver Surgery and Transplantation

Three-dimensional modeling in complex liver surgery and liver transplantation

Jian-Peng Liu^a, Jan Lerut^b, Zhe Yang^a, Ze-Kuan Li^c, Shu-Sen Zheng^{a,c,d,*}^a Department of Hepatobiliary and Pancreatic Surgery, Department of Liver Transplantation, Shulan (Hangzhou) Hospital, Zhejiang Shuren University School of Medicine, Hangzhou 310000, China^b Starzl Unit of Abdominal Transplantation, University Hospitals Saint Luc, Université catholique Louvain, Brussels, Belgium^c Division of Hepatobiliary Pancreatic Surgery, the First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China^d National Clinical Research Center of Infectious Diseases, Hangzhou 310003, China

ARTICLE INFO

Article history:

Received 22 September 2021

Accepted 24 May 2022

Available online 1 June 2022

Keywords:

3D printing models

Liver surgery

Liver transplantation

ABSTRACT

Liver resection and transplantation are the most effective therapies for many hepatobiliary tumors and diseases. However, these surgical procedures are challenging due to the anatomic complexity and many anatomical variations of the vascular and biliary structures. Three-dimensional (3D) printing models can clearly locate and describe blood vessels, bile ducts and tumors, calculate both liver and residual liver volumes, and finally predict the functional status of the liver after resection surgery. The 3D printing models may be particularly helpful in the preoperative evaluation and surgical planning of especially complex liver resection and transplantation, allowing to possibly increase resectability rates and reduce postoperative complications. With the continuous developments of imaging techniques, such models are expected to become widely applied in clinical practice.

© 2022 First Affiliated Hospital, Zhejiang University School of Medicine in China. Published by Elsevier B.V. All rights reserved.

Introduction

Hepatobiliary cancer and chronic liver failure are diagnosed more and more frequently. For these diseases, liver resection and liver transplantation (LT) are the most effective treatments [1,2]. Although well standardized, postoperative complications and deaths still occur after such procedures [3,4]. Especially advanced hepatobiliary surgery is challenging because surgeons have to deal with the complexity and variability of the Glissonean pedicles, the hepatic venous anatomy and the relationship between these intrahepatic structures and the lesions [5]. Accurate preoperative assessment and preparation are of utmost importance both to obtain an R0 resection and to reduce postoperative complications.

Presently computed tomography (CT) and magnetic resonance imaging (MRI) are the main tools used to evaluate the extent, spread, resectability and transplantability of hepatobiliary tumors [6]. These two-dimensional (2D) imaging techniques allow to visualize the hepatic arterial (HA) and portal vein (PV) blood supply, the accompanying biliary structures, the draining hep-

atic veins and their frequently present anatomic variations [7–10]. Moreover, angio-CT and MRI can determine, through identification of the “feeding and draining” vascular structures rather precisely, both anatomic and functional, total and residual liver volumes (RLV) [11–14]. Some shortcomings in relation to a detailed view of the liver anatomy and lesion can be resolved by 3D modeling and printing [15]. 3D printing is the process in which an excavated model is transformed into a 3D morphology. Such models have already been widely used in the fields of stomatology, neurosurgery and nephrology. More recently their use has been introduced in complex liver resection and (living donor) LT procedures [16–21]. These models help in the preoperative surgical planning by providing precise anatomical details and spatial intrahepatic relationships. This information is crucial especially in the accurate planning of complex surgical procedures including vascular reconstructions [22–24].

Methods

A systematic literature published in English was searched in PubMed and EMBASE using the search terms: “3D print”, OR “3D model”, OR “Three-dimensional print”, OR “Three-dimensional model”, AND “liver surgery”, OR “liver transplantation”, OR “liver”, and the search date was defined 1990/01/01–2021/12/31. Finally, 18 (3.2%) of 566 papers were retained for the analysis

* Corresponding author at: Department of Hepatobiliary and Pancreatic Surgery, Department of Liver Transplantation, Shulan (Hangzhou) Hospital, 848 Dongxin Road, Hangzhou 310000, China.

E-mail address: shusenzheng@zju.edu.cn (S.-S. Zheng).




Fig. 1. Flow diagram related to the systematic review of the English literature related to 3D printing models and liver resection and transplantation.

(Table 1, Fig. 1) [25–42]. The specific process of the 3D printing models in liver surgery is shown in Fig. 2.

3D printing models in advanced liver surgery

The complexity and variability of the hepatic vessels and their corresponding biliary tracts represent a main technical difficulty in complex liver resections. To enhance safety and efficacy, one must fully understand the location and the spatial relationship between the lesion(s) and the biliary and vascular structures. 3D printing models can precisely visualize the shape and location of the HA, the PV and the biliary tract, the size and location of tumors and finally the inter-relationship between all those structures. A detailed cartography helps the liver surgeon preserve or resect these structures and calculate the functional RLV, both of which enhance the safety of the planned procedure [25].

3D printing models and primary hepatobiliary tumors

In children, hepatoblastoma and undifferentiated embryonal sarcoma are the most frequent liver tumors [43–45]. As these tumors are often extended and therefore intermingled with the blood vessels, resection may compromise the residual liver function. Moreover, in these patients the fragility of the vasculature increases the risk for intra- and postoperative complications [46–48]. The exact preoperative evaluation of tumor size, the relationship with the adjacent blood vessels and the determination of RLV are all of particular importance in these pediatric patients.

Souzaki et al. reported a liver resection in a 3-year-old girl with hepatoblastoma after chemotherapy located between the left and right PV branches. Intraoperative findings correlated well with the 3D printing liver models, allowing to perform an extended left hepatectomy. The pathology of the resected specimen confirmed a negative surgical margin [28].

In adults, large HCC are common and are often accompanied by invasion of the portal and/or hepatic veins and impairment of liver function because they mostly develop in a fibrotic/cirrhotic liver [49,50]. This frequently present underlying liver disease im-

plies the necessity for a detailed preoperative evaluation of both the liver function and the functional reserve of the RLV [51–53]. Xiang et al. [54] reported the application of 3D printing technology in the resection of a complex large HCC in the presence of a PV variation that a segment 4 vein branch coming from the right anterior PV. 3D printing models can clearly visualize this anomaly. A right hepatectomy would have reduced RLV to only 21.37%; adaptation of the surgical technique based on the 3D printing models can preserve the S4 PV branch and thereby increase the RLV from 21.37% to 57.25%.

The anatomic resection of an HCC, including its draining PV branch, can reduce intraoperative bleeding and improve tumor-free survival [55,56]. Such approach requires a thorough understanding of the segmental liver anatomy [57–59]. Kuroda et al. [26] performed 3D printing models guided anatomic R0 resections of segment 7, segment 4 and right anterior ventral segment. Based on their experience the authors concluded that 3D printing models guided surgery make these procedures safer.

When liver tumors invade large blood vessels, vascular reconstructions are often required to ensure an R0 resection [60]. Huber et al. [61] summarized a series of 10 complex liver resections guided by 3D printing models. Seven required a vascular reconstruction. Again, the authors believed that 3D printing models are of great help not only assess the relationship between a tumor and the blood vessels but also give useful information in relation to vascular reconstruction. Based on this experience they proposed to set up prospective studies to evaluate the real clinical impact of 3D imaging and printing in advanced liver surgery. Similarly, a multi-center study [41] involving 35 patients from 8 centers confirmed the strong correlation between 3D printing models and the analysis of the hepatectomy specimen. The obtained information, however, did not affect the results of surgery. More studies are needed to demonstrate the impact of 3D printing models on complications and prognosis of liver resection.

Perihilar cholangiocarcinoma (PHCCA) is featured by an insidious onset, a difficult early diagnosis, a rapid disease progression and, consequently, a poor prognosis [62–65]. Due to the complex relationship with the accompanying HA and PV as well as the high anatomical variability of these Glissonean structures, R0 surgical resection is difficult to achieve. Enhancing the accuracy of preoperative evaluation in relation to local and environmental tumor extension is very important to optimize the surgical planning [66]. 3D printing models may be an effective tool [29].

Larghi Laureiro et al. [30] reported a 29-year-old woman with PHCCA invading the right PV in which an R0 right trisectionectomy with removal of the entire englobed thrombosed PV followed by a complex vascular reconstruction was successfully performed based on 3D printing models. 3D printing models were highly accurate to identify the anatomical relationship between the tumorous bile duct and the invaded accompanying vessels. Zeng et al. [67] combined 3D visualization and 3D printing models in the individualized precision surgical treatment of 10 patients presenting Bismuth-Corlette III and IV PHCCA. Despite several vascular variations, all complex surgical procedures could be performed safely based on the preoperative 3D visualization and printing models which fit the different encountered intraoperative findings very closely.

3D printing models and secondary liver tumors

Colorectal cancer is one of the most common cancers and the liver is the most common site of metastasis [68,69]. Unlike primary liver cancer, metastatic liver cancer is usually diagnosed in an advanced stage and therefore usually requires neoadjuvant therapy to improve resectability and outcome [70]. 3D printing models can help assess the efficacy of this complementary medical ther-

Table 1
Selected review of the literature in relation to 3D printing models and advanced liver surgery and transplantation.

Studies	Year	n	Disease	Aim	Imaging method	Significant findings made by the authors in relation to 3D printing
Liver resection						
Madurska et al. [25]	2017	1	HCC	Preoperative planning	CT/MRI	Providing good anatomical details of the relationship between liver tumors and adjacent structures.
Kuroda et al. [26]	2017	2	HCC	Preoperative planning	CT	The simplified 3D models of intrahepatic vessels without liver parenchyma are sufficient to effectively guide intraoperative operations and are suitable for all HCC surgeries.
Perica et al. [27]	2017	1	HCC	Preoperative planning	CT	3D printing models are of limited value in diagnostic radiology. However, they still have guiding significance in the surgical planning and intraoperative guidance of hepatocellular carcinoma treatment.
Souzaki et al. [28]	2015	1	Hepatoblastoma	Preoperative planning	CT	3D printed models help in surgical planning of pediatric liver malignancies.
Takagi et al. [29]	2014	1	PHCCA	Preoperative planning	CT	The reproducibility of 3D modeling by the latest imaging has been clarified and future preoperative simulation should be dramatically changed.
Larghi Laureiro et al. [30]	2020	1	PHCCA	Preoperative planning	CT	The application of 3D models in liver surgery helps clarify some useful preoperative questions and the accuracy of the model is high.
Igami et al. [31]	2014	2	CRLM	Preoperative planning	CT	It is a simple and feasible method to use 3D printing liver to perform hepatectomy for small tumors that cannot be seen by intraoperative ultrasound examination.
Oshiro et al. [32]	2017	1	CRLM	Preoperative planning	CT	This model can clearly simulate the resection line and help safely perform the surgery; and it is cost effective, and has a shorter production time.
Witowski et al. [33]	2017	1	CRLM	Preoperative planning	CT	A cost-effective technique for the preparation of 3D-printed liver models is proposed that can reduce operative time and improve surgical outcomes.
LDLT						
Zein et al. [34]	2013	6	Liver cirrhosis	Preoperative planning LDLT	CT/MRI	These models are proved to be very accurate, and this was the first time that a human liver has been replicated and validated at the time of surgery with an actual local liver.
Baimakhanov et al. [35]	2015	1	HCC	Preoperative planning LDLT	CT	3D printing models are recommended for LDLT with complex vascular or biliary tract anatomy that require multiple or complex anastomoses.
Soejima et al. [36]	2016	1	Secondary biliary cirrhosis post Kasai	Preoperative planning LDLT	CT	Preoperative simulation using 3D-printed liver models is a technological innovation in pediatric LDLT, especially for small infants.
Ishii et al. [37]	2020	1	Secondary biliary cirrhosis post Kasai	Preoperative planning LDLT	CT	Patient-specific 3D liver model facilitated the identification of vessels during living donor liver transplantation in situs inversus totalis.
Wang et al. [38]	2019	10	Biliary atresia (6), congenital cholangiectasis (1), biliary cirrhosis (1), Wilson's disease (1), autoimmune hepatitis (1)	Preoperative simulation LDLT	CT/MRI	In small infants and complex cases of pediatric LDLT, 3D printing models can help minimize the risk of large-size syndrome and graft reduction.
Rhu et al. [39]	2021	90	HCC (59); others (31)	Preoperative simulation LDLT	CT/MRI	3D printing models can improve clinical outcomes, especially in bile-duct division.
Raichurkar et al. [40]	2021	1	Wilson disease	Preoperative simulation LDLT	CT	Three-dimensional printing of liver models can help predict the actual size of the graft after donor hepatectomy, in patients undergoing LDLT.
Others						
Lopez-Lopez et al. [41]	2021	35	CRLM (12), ICC (7), Klatskin tumor (5), hemangioma (1), adenoma (1), focal nodular hyperplasia (1), HCC (3), gallbladder (1), primary sarcoma (2), sarcoma metastases (1), adrenal metastases (1).	Preoperative planning	CT/MRI	The 3D printing liver models have a good correlation with CT/MRI, which is helpful for preoperative simulation, but do not affect postoperative outcomes.
Chen et al. [42]	2021	291	N/A	Preoperative simulation	CT/MRI	It is possible to take medial segment graft as a donor liver with the help of 3D printing technology.

CT: computed tomography; MRI: magnetic resonance imaging; LDLT: living donor liver transplantation; HCC: hepatocellular carcinoma; ped tumors: pediatric tumors; CRLM: colorectal liver metastases; PHCCA: perihilar cholangiocarcinoma; ICC: intrahepatic cholangiocarcinoma; N/A: not available.




Fig. 2. Operating processes using 3D printing models in liver surgery and transplantation. CT: computed tomography; MRI: magnetic resonance imaging; 3D: three-dimensional; MHV: middle hepatic vein.

apy and reevaluate the initial preoperative imaging following their shrinking process [71]. Igami et al. [31] performed hepatectomy in two patients evaluated for multiple colorectal metastases using 3D printing models. Following chemotherapy 3D printing models can identify small “missing” tumors by preoperative ultrasound and adapt the resection lines, facilitating an appropriate hepatectomy.

3D printing models in LT

LT is the most effective therapy for end-stage liver disease [72,73]. Living donor liver transplantation (LDLT) and split-LT are two means to enlarge the liver allograft pool [74–76]. Comprehensive accurate imaging of all vascular structures of these allografts is vital in the application of the precision LDLT and split-LT surgery allowing to increase the safety both in donor and recipient [77–79]. 3D printing models are means to identify better the anatomical structures and to accurately determine both donor and recipient standard liver volumes (SLVs) and graft volumes. By simulating the perfusion of all segmental territories the corresponding LV can be measured to precisely calculate the respective graft body weight ratios, which are necessary for an optimal function of the liver graft and for the safety of the donor [34].

3D printing models and LDLT in adults

Adult to adult LDLT is a complex and risky undertaking, the surgical difficulty of adult being related not only to the various vascular and biliary constellations but also to the risk for liver insufficiency both in donor and recipient [80–82]. In 2013 Zein et al. [34] first printed translucent 3D liver models of three donors and three recipients. Their models were highly accurate in relation to the diameters of the vascular structures (with an average size error of less than 4 mm for the main portal and hepatic veins and of less than 1.3 mm for the HA). Table 2 shows that the errors of the 3D printing models group in cases are 3.3%, 3.3% and 16.6%, while the errors of the CT group are 4.3%, 28.8% and 30.1%. In relation to the volume measurements 3D printing models outperformed 3D imaging model [83]. Based on this small experience, the authors concluded that physical 3D printing models give very precise information and that they may also allow to improve the tactile and spatial relationships of blood vessels compared to those

Table 2

Differences between liver volume measured using 3D printing models and 3D imaging compared to CT evaluation (Zein et al. [34])

Patient	Native LV (mL)	3D printing LV (mL)	CT LV (mL)
Case 1	1215	1175 (3.3%*)	1267 (4.3%*)
Case 2	1195	1235 (3.3%*)	851 (28.8%*)
Case 3	1250	1458 (16.6%*)	1626 (30.1%*)

*: errors presented in percentage. CT: computed tomography; LV: liver volume.

generated from computerized 3D graphic models. Baimakhanov et al. [35] transplanted a right liver including the middle hepatic vein guided by 3D printing models. This donor had a middle hepatic vein (MHV) larger than the right hepatic vein (RHV), and this MHV partially drained a vast area, including segment 6. 3D printing models measured a liver volume corresponding to 47.5% of the recipient SLV. Hence, they decided to use an extended right lobe graft and an autologous portal vein Y-graft interposition for the hepatic vein anastomosis. The solid 3D printing models made it easier to imagine the reconstructed shape and angulation of the anastomosis based on a better spatial perception. Finally, the Y-type shaped hepatic vein reconstruction was performed by connecting the right portal branch to the MHV and the left portal branch to the RHV.

Rhu et al. [39] analyzed 200 adult LDLTs. Compared with the non-image guided group, 90 LDLTs completed with image guidance did not reveal differences in relation to the type of bile duct anatomy ($P = 0.144$), but 3D printing models obtained a significantly higher number of single bile duct orifices (80.0% vs. 52.7%, $P = 0.001$). The 3D printing models may therefore be of great value to obtain a more accurate and safer transection of the biliary tract in the donor.

3D printing models and LDLT in pediatric patients

In pediatric LDLTs, the left lateral lobe (segments 2 and 3) grafting is a very well standardized and safe procedure [84]. However, large-for-size grafts [a graft to recipient weight ratio (GRWR) > 4] and inappropriate hepatic venous outflow still remain common reasons of surgical failure. Large grafts placed in a small ab-

dominal cavity can lead to an impaired allograft oxygenation and a compression or torsion of the portal and/or hepatic veins causing PV thrombosis and/or hepatic outflow obstruction [85–87]. Therefore, a very precise surgical planning and implantation procedure are needed in such constellation.

Soejima et al. [36] evaluated an 11-month-old girl post-Kasai surgery using 3D printing models. The estimated left lateral segment graft volume was 295 mL, corresponding to a GRWR of 4.9%. After reassessment, including a partial S2 liver resection, the weight of the allograft could be reduced to 245 g corresponding to a GRWR of 3.8%, avoiding a large-for-size grafting. Ishii et al. [37] used 3D printing technology to evaluate vascular variations in a case of biliary atresia in situs in versus totalis. The sterilized 3D printing models were brought into the operative field and used as an intraoperative navigation tool. This recipient presented an unconventional arterial anomaly, the right hepatic artery originating from the celiac axis and reaching dorsally in the hepatoduodenal ligament at the right edge of the caudate lobe. Hepatectomy was challenging based on the documented anatomical variations, and the liver could be removed without damaging the right hepatic artery. Moreover, the industrial CT scan of the patient-specific 3D liver model revealed that the gaps between the liver model and the original data were < 0.4 mm in the 90% area, < 0.8 mm in the 98% area, and 1.53 mm at the maximum.

Wang et al. [38] looked at the impact of 3D printing models comparing ten 3D printing guided LDLTs to 20 non-3D printing guided LDLTs. The 3D printing group had a significantly shorter operative time (2.3 ± 0.4 vs. 3.0 ± 0.4 h, $P < 0.001$) and less cost (17.1% lower) ($34\,600 \pm 6600$ vs. $41\,700 \pm 10\,400$ RMB, $P = 0.03$) compared to the non-3D printing group. They also concluded that 3D printing models can help minimize the risk of large-for-size syndrome and graft reduction.

Conclusions

3D printing technology is used more and more frequently in complex liver surgery and transplantation. 3D printing models can visualize and calculate very precisely the anatomy and diameter of blood vessels, bile ducts, their relationship with liver tumors and finally to give a precise idea about total and RLVs. Although 3D modeling and printing may be superior to “traditional” state-of-the-art 2D imaging in relation to the planning of advanced liver resection and LDLT procedures and to improve the tactile evaluation of the operative field, more research will be needed to confirm its real superiority. Future technical development is expected to overcome the inconvenience of increased cost and time-consumption.

Acknowledgements

None.

CRedit authorship contribution statement

Jian-Peng Liu: Conceptualization, Writing – original draft. **Jan Lerut:** Writing – review & editing. **Zhe Yang:** Conceptualization, Funding acquisition. **Ze-Kuan Li:** Writing – original draft. **Shu-Sen Zheng:** Conceptualization, Validation, Funding acquisition, Writing – review & editing.

Funding

This study was supported by grants from the National S&T Major Project (2017ZX10203205) and the Natural Science Foundation of Zhejiang Province (Y21H160259).

Ethical approval

Not needed.

Competing interest

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article.

References

- [1] Piano S, Tonon M, Vettore E, Stanco M, Pilutti C, Romano A, et al. Incidence, predictors and outcomes of acute-on-chronic liver failure in outpatients with cirrhosis. *J Hepatol* 2017;67:1177–1184.
- [2] Wang JZ, Xiong NY, Zhao LZ, Hu JT, Kong DC, Yuan JY. Review fantastic medical implications of 3D-printing in liver surgeries, liver regeneration, liver transplantation and drug hepatotoxicity testing: a review. *Int J Surg* 2018;56:1–6.
- [3] Serenari M, Han KH, Ravaioli F, Kim SU, Cucchetti A, Han DH, et al. A nomogram based on liver stiffness predicts postoperative complications in patients with hepatocellular carcinoma. *J Hepatol* 2020;73:855–862.
- [4] Chan ACY, Chok KSH, Dai J, Tsang SHY, Cheung TT, Poon R, et al. Transferability of liver transplantation experience to complex liver resection for locally advanced hepatobiliary malignancy – lessons learnt from 3 decades of single center experience. *Ann Surg* 2020.
- [5] Zhao D, Lau WY, Zhou W, Yang J, Xiang N, Zeng N, et al. Impact of three-dimensional visualization technology on surgical strategies in complex hepatic cancer. *Biosci Trends* 2018;12:476–483.
- [6] Bajpai S, Kambadakone A, Guimaraes AR, Arellano RS, Gervais DA, Sahani D. Image-guided treatment in the hepatobiliary system: role of imaging in treatment planning and posttreatment evaluation. *Radiographics* 2015;35:1393–1418.
- [7] Alonso-Torres A, Fernández-Cuadrado J, Pinilla I, Parrón M, de Vicente E, López-Santamaría M. Multidetector CT in the evaluation of potential living donors for liver transplantation. *Radiographics* 2005;25:1017–1030.
- [8] Nakamura T, Tanaka K, Kiuchi T, Kasahara M, Oike F, Ueda M, et al. Anatomical variations and surgical strategies in right lobe living donor liver transplantation: lessons from 120 cases. *Transplantation* 2002;73:1896–1903.
- [9] Sureka B, Patidar Y, Bansal K, Rajesh S, Agrawal N, Arora A. Portal vein variations in 1000 patients: surgical and radiological importance. *Br J Radiol* 2015;88:20150326.
- [10] Gruttadauria S, Foglieni CS, Doria C, Luca A, Lauro A, Marino IR. The hepatic artery in liver transplantation and surgery: vascular anomalies in 701 cases. *Clin Transplant* 2001;15:359–363.
- [11] Sonnemans LJ, Hol JC, Monshouwer R, Prokop M, Klein WM. Correlation between liver volumetric computed tomography results and measured liver weight: a tool for preoperative planning of liver transplant. *Exp Clin Transplant* 2016;14:72–78.
- [12] Radtke A, Sotiropoulos GC, Molmenti EP, Schroeder T, Peitgen HO, Frilling A, et al. Computer-assisted surgery planning for complex liver resections: when is it helpful? A single-center experience over an 8-year period. *Ann Surg* 2010;252:876–883.
- [13] Vauthey JN, Chaoui A, Do KA, Bilimoria MM, Fenstermacher MJ, Charnsangavej C, et al. Standardized measurement of the future liver remnant prior to extended liver resection: methodology and clinical associations. *Surgery* 2000;127:512–519.
- [14] Urata K, Kawasaki S, Matsunami H, Hashikura Y, Ikegami T, Ishizone S, et al. Calculation of child and adult standard liver volume for liver transplantation. *Hepatology* 1995;21:1317–1321.
- [15] Bangeas P, Tsioukas V, Papadopoulos VN, Tsoulfas G. Role of innovative 3D printing models in the management of hepatobiliary malignancies. *World J Hepatol* 2019;11:574–585.
- [16] Liaw CY, Guvendiren M. Current and emerging applications of 3D printing in medicine. *Biofabrication* 2017;9:024102.
- [17] Pugliese L, Marconi S, Negrello E, Mauri V, Peri A, Gallo V, et al. The clinical use of 3D printing in surgery. *Updates Surg* 2018;70:381–388.
- [18] Dawood A, Marti Marti B, Sauret-Jackson V, Darwood A. 3D printing in dentistry. *Br Dent J* 2015;219:521–529.
- [19] Liu K, Yan L, Li R, Song Z, Ding J, Liu B, et al. 3D printed personalized nerve guide conduits for precision repair of peripheral nerve defects. *Adv Sci (Weinh)* 2022;9:e2103875.
- [20] García-Sevilla M, Moreta-Martínez R, García-Mato D, Arenas de Frutos G, Ochandiano S, Navarro-Cuellar C, et al. Surgical navigation, augmented reality, and 3D printing for hard palate adenoid cystic carcinoma en-bloc resection: case report and literature review. *Front Oncol* 2022;11:741191.
- [21] Chaudhary A, Chopra S, Sinha VD. Role of three-dimensional printing in neurosurgery: an institutional experience. *Asian J Neurosurg* 2021;16:531–538.
- [22] Perica ER, Sun Z. A systematic review of three-dimensional printing in liver disease. *J Digit Imaging* 2018;31:692–701.
- [23] Tack P, Victor J, Gemmel P, Annemans L. 3D-printing techniques in a medical setting: a systematic literature review. *Biomed Eng Online* 2016;15:115.
- [24] Marro A, Bandukwala T, Mak W. Three-dimensional printing and medical imaging: a review of the methods and applications. *Curr Probl Diagn Radiol* 2016;45:2–9.

- [25] Madurska MJ, Poyade M, Eason D, Rea P, Watson AJ. Development of a patient-specific 3D-printed liver model for preoperative planning. *Surg Innov* 2017;24:145–150.
- [26] Kuroda S, Kobayashi T, Ohdan H. 3D printing model of the intrahepatic vessels for navigation during anatomical resection of hepatocellular carcinoma. *Int J Surg Case Rep* 2017;41:219–222.
- [27] Perica E, Sun Z. Patient-specific three-dimensional printing for pre-surgical planning in hepatocellular carcinoma treatment. *Quant Imaging Med Surg* 2017;7:668–677.
- [28] Souzaki R, Kinoshita Y, Ieiri S, Hayashida M, Koga Y, Shirabe K, et al. Three-dimensional liver model based on preoperative CT images as a tool to assist in surgical planning for hepatoblastoma in a child. *Pediatr Surg Int* 2015;31:593–596.
- [29] Takagi K, Nanashima A, Abo T, Arai J, Matsuo N, Fukuda T, et al. Three-dimensional printing model of liver for operative simulation in perihilar cholangiocarcinoma. *Hepatogastroenterology* 2014;61:2315–2316.
- [30] Larghi Laureiro Z, Novelli S, Lai Q, Mennini G, D'andrea V, Gaudenzi P, et al. There is a great future in plastics: personalized approach to the management of hilar cholangiocarcinoma using a 3-D-printed liver model. *Dig Dis Sci* 2020;65:2210–2215.
- [31] Igami T, Nakamura Y, Hirose T, Ebata T, Yokoyama Y, Sugawara G, et al. Application of a three-dimensional print of a liver in hepatectomy for small tumors invisible by intraoperative ultrasonography: preliminary experience. *World J Surg* 2014;38:3163–3166.
- [32] Oshiro Y, Mitani J, Okada T, Ohkohchi N. A novel three-dimensional print of liver vessels and tumors in hepatectomy. *Surg Today* 2017;47:521–524.
- [33] Witowski JS, Pędziwiatr M, Major P, Budzyński A. Cost-effective, personalized, 3D-printed liver model for preoperative planning before laparoscopic liver hemihepatectomy for colorectal cancer metastases. *Int J Comput Assist Radiol Surg* 2017;12:2047–2054.
- [34] Zein NN, Hanounieh IA, Bishop PD, Samaan M, Eghtesad B, Quintini C, et al. Three-dimensional print of a liver for preoperative planning in living donor liver transplantation. *Liver Transpl* 2013;19:1304–1310.
- [35] Baimakhanov Z, Soyama A, Takatsuki M, Hidaka M, Hirayama T, Kinoshita A, et al. Preoperative simulation with a 3-dimensional printed solid model for one-step reconstruction of multiple hepatic veins during living donor liver transplantation. *Liver Transpl* 2015;21:266–268.
- [36] Soejima Y, Taguchi T, Sugimoto M, Hayashida M, Yoshizumi T, Ikegami T, et al. Three-dimensional printing and biotexture modeling for preoperative simulation in living donor liver transplantation for small infants. *Liver Transpl* 2016;22:1610–1614.
- [37] Ishii T, Fukumitsu K, Ogawa E, Okamoto T, Uemoto S. Living donor liver transplantation in situs inversus totalis with a patient-specific three-dimensional printed liver model. *Pediatr Transplant* 2020;24:e13675.
- [38] Wang P, Que W, Zhang M, Dai X, Yu K, Wang C, et al. Application of 3-dimensional printing in pediatric living donor liver transplantation: a single-center experience. *Liver Transpl* 2019;25:831–840.
- [39] Rhu J, Choi GS, Kim MS, Kim JM, Joh JW. Image guidance using two-dimensional illustrations and three-dimensional modeling of donor anatomy during living donor hepatectomy. *Clin Transplant* 2021;35:e14164.
- [40] Raichurkar KK, Lochan R, Jacob M, Asthana S. The use of a 3D printing model in planning a donor hepatectomy for living donor liver transplantation: first in India. *J Clin Exp Hepatol* 2021;11:515–517.
- [41] Lopez-Lopez V, Robles-Campos R, García-Calderon D, Lang H, Cugat E, Jiménez-Galanes S, et al. Applicability of 3D-printed models in hepatobiliary surgery: results from "LIV3DPRINT" multicenter study. *HPB (Oxford)* 2021;23:675–684.
- [42] Chen CY, Yeh YT, Liu C, Lee RC, Huang SS, Loong CC. The feasibility of medial segment graft in pediatric liver transplantation revisited by three-dimensional printing. *J Pediatr Surg* 2021;56:1162–1168.
- [43] Tsoulfas G, Mekras A, Agorastou P, Kiskinis D. Surgical treatment for large hepatocellular carcinoma: does size matter? *ANZ J Surg* 2012;82:510–517.
- [44] Sharma D, Subbarao G, Hepatoblastoma Saxena R. *Semin Diagn Pathol* 2017;34:192–200.
- [45] Iacob ER, Popoiu CM, Nyiredi A, Mogoantă L, Badea O, Boia ES. Mesenchymal hamartoma of the left liver lobe in an 18-month-old female patient. *Rom J Morphol Embryol* 2016;57:841–847.
- [46] Lim IIP, Bondoc AJ, Geller JI, Tiao GM. Hepatoblastoma-the evolution of biology, surgery, and transplantation. *Children (Basel)* 2018;6:1.
- [47] Murawski M, Dakowicz L, Losin M, Krawczuk-Rybak M, Czaderna P. Isolated caudate lobe (Spiegel lobe) resection for hepatoblastoma. Is it enough to achieve a sufficient resection margin? A case report. *J Pediatr Surg* 2013;48:E25–E27.
- [48] Yang T, Tan T, Yang J, Pan J, Hu C, Li J, et al. The impact of using three-dimensional printed liver models for patient education. *J Int Med Res* 2018;46:1570–1578.
- [49] Chen ZL, Zhang CW, Liang L, Wu H, Zhang WG, Zeng YY, et al. Major hepatectomy in elderly patients with large hepatocellular carcinoma: a multicenter retrospective observational study. *Cancer Manag Res* 2020;12:5607–5618.
- [50] Fang Q, Xie QS, Chen JM, Shan SL, Xie K, Geng XP, et al. Long-term outcomes after hepatectomy of huge hepatocellular carcinoma: A single-center experience in China. *Hepatobiliary Pancreat Dis Int* 2019;18:532–537.
- [51] Belghiti J, Kianmanesh R. Surgical treatment of hepatocellular carcinoma. *HPB (Oxford)* 2005;7:42–49.
- [52] Yang LY, Chang RM, Lau WY, Ou DP, Wu W, Zeng ZJ. Mesohepatectomy for centrally located large hepatocellular carcinoma: Indications, techniques, and outcomes. *Surgery* 2014;156:1177–1187.
- [53] Ruiz E, Pineau P, Flores C, Fernández R, Cano L, Cerapio JP, et al. A preoperative nomogram for predicting long-term survival after resection of large hepatocellular carcinoma (>10 cm). *HPB (Oxford)* 2022;24:192–201.
- [54] Xiang N, Fang C, Fan Y, Yang J, Zeng N, Liu J, et al. Application of liver three-dimensional printing in hepatectomy for complex massive hepatocarcinoma with rare variations of portal vein: preliminary experience. *Int J Clin Exp Med* 2015;8:18873–18878.
- [55] Chien SC, Chen CY, Cheng PN, Liu YS, Cheng HC, Chuang CH, et al. Combined transarterial embolization/chemoembolization-based locoregional treatment with sorafenib prolongs the survival in patients with advanced hepatocellular carcinoma and preserved liver function: a propensity score matching study. *Liver Cancer* 2019;8:186–202.
- [56] Subbotin VM. A hypothesis on paradoxical privileged portal vein metastasis of hepatocellular carcinoma. Can organ evolution shed light on patterns of human pathology, and vice versa? *Med Hypotheses* 2019;126:109–128.
- [57] Jarufe N, Figueroa E, Muñoz C, Moisan F, Varas J, Valbuena JR, et al. Anatomic hepatectomy as a definitive treatment for hepatolithiasis: a cohort study. *HPB (Oxford)* 2012;14:604–610.
- [58] Nishino H, Hatano E, Seo S, Nitta T, Saito T, Nakamura M, et al. Real-time navigation for liver surgery using projection mapping with indocyanine green fluorescence: development of the novel medical imaging projection system. *Ann Surg* 2018;267:1134–1140.
- [59] Wang S, Yue Y, Zhang W, Liu Q, Sun B, Sun X, et al. Dorsal approach with Glissonian approach for laparoscopic right anatomic liver resections. *BMC Gastroenterol* 2021;21:138.
- [60] Radulova-Mauersberger O, Weitz J, Riediger C. Vascular surgery in liver resection. *Langenbecks Arch Surg* 2021;406:2217–2248.
- [61] Huber T, Huettl F, Trippel V, Baumgart J, Lang H. Experiences with three-dimensional printing in complex liver surgery. *Ann Surg* 2021;273:e26–e27.
- [62] Lauterio A, De Carlis R, Centonze L, Buscemi V, Incarboni N, Vella I, et al. Current surgical management of peri-hilar and intra-hepatic cholangiocarcinoma. *Cancers (Basel)* 2021;13:3657.
- [63] Serrablo A, Serrablo L, Alikhanov R, Tejedor L. Vascular resection in perihilar cholangiocarcinoma. *Cancers (Basel)* 2021;13:5278.
- [64] Gunasekaran G, Bekki Y, Lourdasamy V, Schwartz M. Surgical treatments of hepatobiliary cancers. *Hepatology* 2021;73:128–136.
- [65] Rizvi S, Gores GJ. Pathogenesis, diagnosis, and management of cholangiocarcinoma. *Gastroenterology* 2013;145:1215–1229.
- [66] Sapisochin G, Ivanics T, Subramanian V, Doyle M, Heimbach JK, Hong JC. Multidisciplinary treatment for hilar and intrahepatic cholangiocarcinoma: A review of the general principles. *Int J Surg* 2020;82S:77–81.
- [67] Zeng N, Yang J, Xiang N, Wen S, Zeng S, Qi S, et al. Application of 3D visualization and 3D printing in individualized precision surgery for Bismuth-Corlette type III and IV hilar cholangiocarcinoma. *Nan Fang Yi Ke Da Xue Xue Bao* 2020;40:1172–1177.
- [68] Datta J, Narayan RR, Kemeny NE, D'Angelica MI. Role of hepatic artery infusion chemotherapy in treatment of initially unresectable colorectal liver metastases: a review. *JAMA Surg* 2019;154:768–776.
- [69] Malafosse R, Penna C, Sa Cunha A, Nordlinger B. Surgical management of hepatic metastases from colorectal malignancies. *Ann Oncol* 2001;12:887–894.
- [70] Wang LJ, Wang HW, Jin KM, Li J, Xing BC. Comparison of sequential, delayed and simultaneous resection strategies for synchronous colorectal liver metastases. *BMC Surg* 2020;20:16.
- [71] Choi YR, Kim JH, Park SJ, Hur BY, Han JK. Therapeutic response assessment using 3D ultrasound for hepatic metastasis from colorectal cancer: Application of a personalized, 3D-printed tumor model using CT images. *PLoS One* 2017;12:e0182596.
- [72] Chan KM, Hung HC, Lee JC, Wu TH, Wang YC, Cheng CH, et al. A review of split liver transplantation with full right/left hemi-liver grafts for 2 adult recipients. *Medicine (Baltimore)* 2021;100:e27369.
- [73] Silveira F, Silveira FP, Macri MM, Nicoluzzi JE. Analysis of liver waiting list mortality in Paraná, Brazil: what shall we do to face organ shortage? *Arq Bras Cir Dig* 2012;25:110–113.
- [74] Rela M, Rammohan A. Why are there so many liver transplants from living donors in Asia and so few in Europe and the US? *J Hepatol* 2021;75:975–980.
- [75] Linecker M, Krones T, Berg T, Niemann CU, Steadman RH, Dutkowski P, et al. Potentially inappropriate liver transplantation in the era of the "sickest first" policy - A search for the upper limits. *J Hepatol* 2018;68:798–813.
- [76] Chan KM, Wang YC, Wu TH, Cheng CH, Lee CF, Wu TJ, et al. Encouraging split liver transplantation for two adult recipients to mitigate the high incidence of wait-list mortality in the setting of extreme shortage of deceased donors. *J Clin Med* 2019;8:2095.
- [77] Togashi J, Akamatsu N, Kokudo N. Living donor liver transplantation for hepatocellular carcinoma at the University of Tokyo Hospital. *Hepatobiliary Surg Nutr* 2016;5:399–407.
- [78] Matsushima H, Fujiki M, Sasaki K, Rotroff DM, Sands M, Bayona Molano MDP, et al. Predictive value of hepatic venous pressure gradient for graft hemodynamics in living donor liver transplantation. *Liver Transpl* 2019;25:1034–1042.
- [79] Cai L, Yeh BM, Westphalen AC, Roberts JP, Wang ZJ. Adult living donor liver imaging. *Diagn Interv Radiol* 2016;22:207–214.
- [80] Hara T, Soyama A, Hidaka M, Kitasato A, Ono S, Natsuda K, et al. Analysis of early relaparotomy following living donor liver transplantation. *Liver Transpl* 2016;22:1519–1525.

- [81] Barbetta A, Aljehani M, Kim M, Tien C, Ahearn A, Schilperoort H, et al. Meta-analysis and meta-regression of outcomes for adult living donor liver transplantation versus deceased donor liver transplantation. *Am J Transplant* 2021;21:2399–2412.
- [82] Durairaj MS, Shaji Mathew J, Mallick S, Nair K, Manikandan K, Titus Varghese C, et al. Middle hepatic vein reconstruction in adult living donor liver transplantation: a randomized clinical trial. *Br J Surg* 2021;108:1426–1432.
- [83] Ikegami T, Maehara Y. Transplantation: 3D printing of the liver in living donor liver transplantation. *Nat Rev Gastroenterol Hepatol* 2013;10:697–698.
- [84] Troisi RI, Elsheikh Y, Alnemary Y, Zidan A, Sturdevant M, Alabbad S, et al. Safety and feasibility report of robotic-assisted left lateral sectionectomy for pediatric living donor liver transplantation: a comparative analysis of learning curves and mastery achieved with the laparoscopic approach. *Transplantation* 2021;105:1044–1051.
- [85] Raghu VK, Carr-Boyd PD, Squires JE, Vockley J, Goldaracena N, Mazariegos GV. Domino transplantation for pediatric liver recipients: Obstacles, challenges, and successes. *Pediatr Transplant* 2021;25:e14114.
- [86] Lu YG, Pan ZY, Zhang S, Lu YF, Zhang W, Wang L, et al. Living donor liver transplantation in children: perioperative risk factors and a nomogram for prediction of survival. *Transplantation* 2020;104:1619–1626.
- [87] Barbetta A, Butler C, Barhouma S, Hogen R, Rocque B, Goldbeck C, et al. Living donor versus deceased donor pediatric liver transplantation: a systematic review and meta-analysis. *Transplant Direct* 2021;7:e767.