

More than one third of the organic carbon exposed by the world's largest thaw slump (Batagay, Siberia) is not directly available for mineralization but geochemically stabilized

Maxime Thomas^{1*}, Loeka L. Jongejans², Jens Strauss², Chloé Vermeylen¹, Sacha Calcutt¹, Arthur Monhonval¹, Thomas Opel³, Guido Grosse², Sophie Opfergelt¹

¹ Earth and Life Institute, Université catholique de Louvain, Louvain-la-Neuve, Belgium

² Permafrost Research Section, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Potsdam, Germany

³ Polar Terrestrial Environments Section, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Potsdam, Germany

*Corresponding author: Maxime Thomas: maxime.thomas@uclouvain.be

Croix du Sud, 1 bte L7.05.10, B-1348 Louvain-la-Neuve, Belgium | Phone: +32 (0)10 47 36 29

Abstract submission for EUCOP 2023 in session 19: Carbon stocks, soil properties, greenhouse gas fluxes and atmospheric feedbacks of permafrost regions.

Mineral-organic carbon (OC) interactions account for 30 – 80 % of the total permafrost OC pool. Quantifying the nature and controls of mineral-OC interactions is necessary to better assess permafrost-carbon-climate feedbacks. This is particularly true for ice-rich environments that are impacted by rapid thaw and the development of thermokarst landforms. Retrogressive thaw slumps are amongst the most dynamic forms of slope thermokarst and they expand through the years due to the ablation of an ice-rich headwall. These phenomena are important to consider in the permafrost carbon budget since they expose deep OC sometimes tens of thousands of years old that would not have re-entered the modern carbon cycle if these disturbances had not occurred. Here, we analyzed sediment samples collected from the headwall of the Batagay megaslump, East Siberia, locally reaching 55 m high. The series of discontinuous deposits comprises also older sediment up to ~650 ka old. We present total element concentrations, mineralogy, and mineral-OC interactions in the different stratigraphic units. The mineralogy in the deposits is very similar across the sedimentary series. Our data show that up to $34 \pm 8\%$ of the total OC pool is stabilized by mineral-OC interactions. For most of the analyzed samples, associations to poorly crystalline iron oxides do not have a significant role in OC stabilization. Hypothesizing a retreat rate of 26000 m²/yr and constant thickness of stratigraphic units within the headwall, we provide a first order estimate of $\sim 2 \times 10^7$ kg of OC is exported annually downslope of the headwall, with $\sim 38\%$ being geochemically stabilized by complexation with metals or associations to poorly crystalline iron oxides. These data support that more than one third of the organic carbon exposed by this massive thaw slump is not directly available for mineralization, but rather stabilized geochemically.

Keywords: mineral-organic carbon interactions, retrogressive thaw slump, mass wasting, thermokarst landform, Batagay

This abstract participates in the Outstanding PYRN Oral Communication / Poster Award.