

Histamine Challenge Test in Children Using Forced Oscillation to Measure Total Respiratory Resistance*

Patrick Lebecque, M.D.; Sheldon Spier, M.D.C.M.;
Jean-Guy Lapierre, M.D.; Andre Lamarre, M.D.;
Raezelle Zinman, M.D.C.M., and Allan L. Coates, M.D.C.M.

This study analyzed the relationship between total respiratory resistance (Rrs) measured by forced oscillation technique and FEV₁ during histamine provocation test in 31 children between seven and 17 years of age. Rrs was measured at frequencies between 6 (R₆) and 26 Hz (R₂₆). (R₆ - R₂₆)/R₂₆ was used as an index of frequency dependency of Rrs. A positive histamine test was defined as PC₂₀ less than 8 mg/ml. Seventeen subjects had a positive test, and all of these had increases from baseline of R₆ greater than 50 percent and (R₆ - R₂₆)/R₂₆ greater than 0.45. Of the 14

subjects whose PC₂₀ was greater than 8 mg/ml, only two had changes in R₆ and (R₆ - R₂₆)/R₂₆ of this magnitude. These two subjects had changes in FEV₁ of 16 and 18 percent. There was a strong linear relationship between the changes in FEV₁ and both R₆ and (R₆ - R₂₆)/R₂₆ from baseline to the final value at the end of the test ($r = 0.87$ and 0.91 respectively). In conclusion, this study demonstrated that the evaluation of airway reactivity by histamine challenge may be done by forced oscillation technique. It is easy to administer and may allow testing of children unable to perform spirometry.

Asthma in children is frequently underdiagnosed,¹ in part because of the lack of specific tests. The histamine challenge test to demonstrate suspected airway lability or to find an explanation for a cough in the absence of apparent pathology has recently gained favor for use in both adults and children. The degree of positivity is usually assessed by the concentration of inhaled histamine necessary to provoke a 20 percent fall in initial FEV₁ (PC₂₀). In Europe, a 15 percent fall in FEV₁ (PC₁₅) is sometimes used as the marker of a positive test. Because this test depends on the ability of the subject to perform a reproducible forced expiratory maneuver, it is usually limited to children older than six or seven years. For those children too young to cooperate with the usual spirometry of pulmonary function testing, one frequently must rely on history alone. In 1956, Dubois et al² described the use of forced oscillation as a measure of total respiratory system impedance which was noninvasive and required minimal cooperation from the subject. Such a technique might lend itself to testing children too young to cooperate with the usual histamine chal-

lenge. The purpose of this study was to compare the changes in FEV₁ to the changes in respiratory system resistance as measured by forced oscillation during a histamine challenge test in children.

Since the original description, the technique of forced oscillation has become much easier to perform and can be applied to infants and older children.³⁻¹⁴ Fourier analysis of the frequency spectrum in the flow signal (resulting from the application of a pressure signal applied to the respiratory system) allows separation of the components of the impedance of the system at various frequencies.¹⁵

MATERIALS AND METHODS

The subjects for this study consisted of 31 Caucasian children between seven and 17 years of age who were referred for a histamine challenge test. Twelve were boys and 19 girls. No one was receiving medication at the time of the study. All had baseline values for total respiratory resistance (Rrs), measured by forced oscillation, that were within normal limits,¹⁶ as were their baseline spirometry values.¹⁷ Two-thirds had been referred because of symptoms suggestive of exercise-induced asthma, and one third for investigation of unexplained cough. Informed consent was obtained and the study had ethics committee approval.

Spirometry was done on an Eagle One (Collins, Braintree, MA) automated water seal spirometer. Values derived were forced vital capacity (FVC), forced expiratory volume in one second (FEV₁) and forced expiratory flow at 25 and 75 percent of FEV (FEF_{25-75%}). Baseline measures were repeated at least three times and were reproducible within 5 percent for FEV₁. The effort associated with the greatest sum of FVC and FEV₁ was used to derive FEV₁ and FEF_{25-75%}. For this study, lower normal limits were defined as 80 percent of predicted for FVC and FEV₁, 60 percent for FEF_{25-75%} and an FEV₁/FVC ratio of 80 percent. Predicted values were taken from Knudson et al.¹⁸

Measurement of Rrs by forced random noise used a commercially available unit (Oscillaire, Jones Medical Instrument Co, Oak Brook,

*From the Department of Pediatrics, Respiratory Medicine Service, McGill University Montreal Children's Hospital Research Institute, and the Département de Pédiatrie, Service Cardio-Pulmonaire, Université de Montréal, Institut de Recherche, Hôpital Ste-Justine, Montreal, Canada.

Supported by the Canadian Cystic Fibrosis Foundation and the McGill University Montreal Children's Hospital Research Institute, Montreal, Canada.

This is publication no. 87003 of the McGill University Montreal Children's Hospital Research Institute.

Manuscript received October 1; revision accepted January 27.

Reprint requests: Dr. Coates, D-380 Montreal Children's Hospital, 2300 Tupper Street, Montreal, Quebec, Canada H3H 1P3

IL) based on the work of Lándsér and colleagues.¹⁹⁻²¹ The device has been described elsewhere in detail.²²⁻²³ With this equipment the subject breathes quietly across a pneumotachograph. A varying pressure signal consisting of pseudorandom noise with a peak never more than 2 cm H₂O and containing energy at all frequencies is superimposed on the regular respirations. The microprocessor in the instrument uses Fourier analysis of the pressure and flow signals to separate the components of impedance into resistive and reactive elements and calculate the frequency spectrum. The values are averaged over 16-sec intervals of quiet breathing, and the resistance and reactance components of the impedance are calculated for even frequencies between 2 and 26 Hz. Reactance (X) is the summation of the elastic and inertial components of the respiratory system which are opposite in sign to each other. For each even frequency the coherence function of the values for the breaths used to derive the results is calculated. It has been empirically determined that when this is greater than 0.95 the relative error in either resistance or reactance will not exceed 10 percent.¹⁹ It is to be noted that a value of 1 for the coherence function indicates the total absence of noise or a linearities. The components of the impedances of interest were the resistances at the relatively low frequency of 6 Hz (R_6) and the higher frequency of 26 Hz (R_{26}), the arithmetic mean resistance between these frequencies (R_m), and the reactance at 26 Hz (X_{26}). Change in resistance as a function of frequency was expressed as $(R_6 - R_{26})/R_{26}$. Only those values where the calculated coherence function was 0.95 or greater were used in the results. Values for measurements at 2 and 4 Hz were not calculated because of the length of time required for adequate data acquisition at these low frequencies.

Prior to the onset of testing, during a period of 20 to 30 minutes, baseline parameters for spirometry and Rrs were measured to acquaint the subject with the procedure. Rrs was measured 21 times in each subject. Each measurement consisted of a 16-sec period during which the subject breathed into a mouthpiece connected to the machine. Each subject wore nose clips and supported his cheeks with his hands, and was encouraged not to swallow or move his head. These maneuvers allowed each child to adopt a comfortable position on the apparatus and to be able to generate reproducible results.

Histamine testing was performed according to the recommendations of Cockcroft et al;²⁴ the subject inhaled increasing concentrations of histamine during 2 min of tidal breathing at each concentration. Two milliliters of the solution were nebulized by a Hudson nebulizer model 1720 supplied with a constant air flow of 5 L/min²⁵ and delivered by a mask covering the nose and mouth. The subject did not wear nose clips but was encouraged to breathe through the mouth. The first inhalation consisted of phosphate-buffered saline solution (PBS) and was followed 5 min later by an initial concentration of histamine of 0.5 mg/ml, which was doubled at each 5-min interval until a maximum of 8 mg/ml was reached. Spirometry was done 3 min after the end of each inhalation. At each stage of the histamine test, three or four measurements of Rrs were made over a period of 150 sec prior to the measurement of FEV₁, and the mean of each parameter was used. The test was terminated when FEV₁ fell by at least 20 percent of baseline, or the maximum concentration was reached. Following administration of the concentration of histamine which led to a 20 percent fall in FEV₁, the subject was given an inhalation of 0.5 ml of salbutamol (5 mg/ml in 1.5 ml of normal saline solution) and 5 min later Rrs was determined using the mean of six to eight measurements. FEV₁ was also measured to ensure that it had returned to normal. The concentration of histamine that provoked a 20 percent fall (PC₂₀) was determined graphically²⁴ from plotting FEV₁ on the Y-axis against the logarithm of the concentration of histamine on the X-axis. The value of the parameters of Rrs at PC₂₀ can be determined when plotted against the log dose of histamine. Similarly, PC₁₅ and the Rrs values at PC₁₅ can be derived.

All data were analyzed by comparing the values measured following the inhalation of histamine to the baseline values. From the baseline measurements, intrasubject and intragroup mean values

Table 1—Anthropometric, Spirometric and Rrs Data

	Group A (n = 17)		Group B (n = 14)	
	Mean	SD	Mean	SD
Age (years)	11.8	3.1	11.6	1.29
Height (cm)	148.1	15.7	147.6	16.2
FEV ₁ (L)	2.43	0.76	2.57	0.86
FEF25-75% (L/sec)	2.56	0.83	3.04	1.21
R ₆ (cm H ₂ O/L/sec)	3.78	1.34	3.53	1.29
R _m (cm H ₂ O/L/sec)	3.80	1.09	3.59	1.19
R ₂₆ (cm H ₂ O/L/sec)	3.68	0.90	3.52	1.01
X ₂₆ (cm H ₂ O/L/sec)	1.16	0.47	1.21	0.46
(R ₆ - R ₂₆)/R ₂₆	0.01	0.21	0.00	0.16

and standard deviation (SD) for Rrs and Xrs and coefficients of variations for Rrs were calculated. Rrs values obtained following histamine inhalation that were more than ± 3 SD of baseline were considered significantly different from baseline.

RESULTS

There were 17 patients with a positive response to histamine as indicated by a PC₂₀ of 8 mg/ml or less. These constituted group A, and the 14 subjects with negative test results made up group B. Allergic rhinitis was present in five members of group A and one of group B. Anthropometric data and mean baseline values for R₆, R₂₆, R_m, X₂₆ and $(R_6 - R_{26})/R_{26}$ for both groups are given in Table 1. There was no significant difference in any of these parameters between the two groups. Range of baseline values for R₆ was 1.99 to 6.99; R₂₆, 2.03 to 5.75; for R_m 2.07 to 6.28; and for X₂₆ 0.29 to 2.29, all in cm H₂O/L/sec. The index of frequency dependence of resistance, $(R_6 - R_{26})/R_{26}$, was between -0.2 and 0.4. The mean of individual coefficients of variation for R₆ was 9.3 percent, for R₂₆, was 10.2 percent, and for R_m was 8.4 percent. These latter values were not influenced by the age of the subject or whether the subject had a positive histamine test reaction. Table 2 gives the coefficients of correlation between the resistance values for R₆, R_m, and R₂₆, and the spirometric values for the baseline test values. All of these were significant and were better for R₆ than for R₂₆. For group A, geometric mean PC₂₀ was 3.6 mg/ml of histamine. Following inhalation of PBS solution, FEV₁ fell a mean of 1.6 percent for all subjects.

Figure 1 demonstrates the typical changes in Rrs during a positive test with increasing concentrations of inhaled histamine. In this case, PC₂₀ was 4 mg/ml. It

Table 2—Correlation Coefficients for Parameters Derived from Forced Oscillation and Spirometry*

	FEV ₁	FEF25-75%
R ₆	.84	.81
R _m	.84	.81
R ₂₆	.76	.72

*p<0.001 in all cases




FIGURE 1. Typical response for a positive histamine challenge. Baseline values (dot solid line), the value after 2 mg/ml histamine where the FEV₁ had fallen 9 percent (dot long dash), after 4 mg/ml histamine where the FEV₁ had dropped 20 percent (dot short dash) and after inhalation of salbutamol where the FEV₁ was back to 105 percent of initial (triangle dash) are given for both resistance and reactance plotted against frequency.

can be seen that there is a considerable increase in R_6 but little change in R_{26} ; in other words, a marked increase in the frequency dependence of resistance and a progressive diminution of reactance. The mean linear correlation coefficient between R_{rs} and frequency between 6 and 26 Hz at the end of positive challenges was 0.95 ± 0.02 .

Table 3 compares the change in FEV₁, R_6 , R_m , X_{26} , and $(R_6 - R_{26})/R_{26}$ from baseline values to those at the final concentration of histamine (paired t -test). For group A, all changes were significant ($p < 0.005$). At PC₂₀ derived from interpolation, the mean increases in R_6 , R_m and R_{26} were 78, 42 and 12 percent respectively. There is a striking relation between the change from baseline to that after the highest concentration of histamine administered in R_6 (ΔR_6) and FEV₁ (ΔFEV_1), with $r = 0.87$, $p < 0.001$ for the 31 tests (Fig 2). Change in frequency dependence of resistance is also strikingly related to ΔFEV_1 (Fig 3); $r = 0.91$, $p < 0.001$. There are similar relationships with lower r -values between

ΔFEV_1 and ΔX_{26} ($r = 0.71$, $p < 0.001$) and R_m ($r = 0.75$, $p < 0.001$) but no significant relation with ΔR_{26} . Using a threshold increase of 50 percent from baseline in R_6 or 0.45 for $(R_6 - R_{26})/R_{26}$, all patients with positive histamine test results (*ie*, PC₂₀ less than 8 mg/ml) would be included, giving a test sensitivity for the technique of forced oscillation of unity. In the 14 patients with negative histamine test results, these thresholds were exceeded in only two, giving a test specificity of 0.86. For group A, values for R_6 were 3 SD different from the mean of baseline in all 17 cases; for R_m , X_{26} and $(R_6 - R_{26})/R_{26}$ this difference pertained in 16 of 17 cases. The difference of 3 SD had been reached for these three parameters at a mean of one dose before the final dose of histamine. In group B, the limit of 3 SD had been reached at the end of the test six of 14 times for R_6 , five times for X_{26} , four times for $(R_6 - R_{26})/R_{26}$ and three times for R_m .

Finally, Table 4 indicates that the administration of salbutamol at the end of a positive test results in the

Table 3—Changes in Parameters from Baseline to Final Histamine Concentration*

	Group A (n = 17)		Group B (n = 14)	
	Mean	Range	Mean	Range
ΔFEV_1	-26.9†	-20 to -48	-8	1 to -18
ΔR_6	104†	60 to 194	31.6†	4 to 139
ΔR_m	47.8†	28 to 88	17.1‡	-2 to 81
ΔR_{26}	14.6†	-4 to 45	4.6§	-14 to 28
ΔX_{26}	-1.73†	-0.33 to -2.9	-0.70‡	0.1 to -2.74
$\Delta(R_6 - R_{26})/R_{26}$	0.80†	0.50 to 1.56	0.23‡	-0.02 to 0.70

* ΔFEV_1 , ΔR_6 , ΔR_m and ΔR_{26} expressed as percent change; ΔX_{26} and $\Delta(R_6 - R_{26})/R_{26}$ expressed as absolute change.

† $p < 0.0005$

‡ $p < 0.01$

§ $p < 0.05$




FIGURE 2. Change in R_6 from baseline to the final concentration of histamine, plotted against a similar change in FEV_1 . Both are expressed as a percentage of baseline for all subjects. The relationship is highly significant ($r = 0.87$, $p < 0.001$.)




FIGURE 3. Change in absolute values for $(R_6 - R_{26})/R_{26}$ before and at the end of the test plotted against the similar change in FEV_1 , as in Figure 2. The relationship is highly significant ($r = 0.91$, $p < 0.001$.)

Table 4— FEV_1 and Rrs in Group A after Final Dose of Histamine and After Salbutamol Inhalation

	Before Salbutamol	After Salbutamol
FEV_1^*	73.1 (6.5)†	102.5 (6.9)†
R_6^*	204 (35.9)†	86.5 (13.2)†
R_{26}^*	114.6 (12.7)†	96.4 (15.8)†
Rm^*	147.8 (18.1)†	91.5 (12.3)†
$X_{26}‡$	-6.9	+1.1
$(R_6 - R_{26})/R_{26}‡$	+8.9	-1.2

*Expressed as a percentage of the initial value.

†Standard deviation.

‡Expressed as standard deviations from the initial value.

return to baseline of not only FEV_1 but also parameters measured by the oscillatory technique. This can also be seen in Figure 1. The change in FEV_1 with salbutamol administration correlated with the changes in R_6 and in $(R_6 - R_{26})/R_{26}$ ($r = 0.64$, $p < 0.01$ for both).

DISCUSSION

Using the definition of a positive histamine challenge test as a PC_{20} not greater than 8 mg/ml, the method of forced oscillation to measure Rrs using a threshold of an increase in R_6 of 50 percent or greater or an increase in $(R_6 - R_{26})/R_{26}$ of 0.45 or greater will include all positive tests; in other words, sensitivity of 100 percent. Those two patients whose PC_{20} was greater than 8 mg/ml but who were positive by the criteria of forced oscillation had changes in FEV_1 of 16 and 18 percent. If, as in many places in Europe, PC_{15} not greater than 8 mg/ml defined a positive test, forced oscillation test results would have resulted in total separation of the groups. In addition, three patients in group A showed changes in R_6 and $(R_6 - R_{26})/R_{26}$ exceeding 6 SD from baseline values one dose prior to the provocative dose causing a fall in FEV_1 of 20 percent or more. This increased sensitivity of the forced oscillation technique may be due in part to elimination of changes in bronchomotor tone accompanying maximal inspiration when measuring FEV_1 ,²⁶⁻³¹ a maneuver that is not necessary when measuring Rrs . The correlations between baseline values of R_6 , Rm , R_{26} and FEV_1 were also very acceptable and similar to the observations of König et al.²³ In terms of reproducibility, coefficients of variation were similar and on the order of 10 percent; in other words, comparable to that normally seen for FEV_1 . These results are in agreement with those of others.^{26,27}

There are two theoretic limitations to using Rrs for histamine challenge testing. The first is that Rrs represents the resistance of the thoracic cage and lung tissue, as well as airway resistance (Raw). However, Raw is the predominant factor and there are good cor-

relations between Raw and Rrs in health and disease in adults and children.^{16,22,33} Moreover, in determining Rrs by forced oscillation and measuring the combination of lung tissue resistance plus Raw with esophageal balloon techniques, Nagels et al²¹ measured the resistance attributable to the thoracic cage and demonstrated that the way this varies with frequency was similar in normal adults and those with COPD. In other words, when comparing normal subjects to those with COPD, this implies that the differences observed in Rrs are due to changes at the level of the airways and pulmonary parenchyma. The other potential handicap is that lung volumes are not determined during the test. Nagels et al²¹ have shown in both normal adults and those with COPD that the measurement of Rrs at a volume lower than FRC increases the Rrs and leads to or accentuates the frequency dependence of Rrs. The inverse was also true. In fact, the inhalation of histamine may be accompanied by hyperinflation, which may be due to changes in inspiratory muscle activity³³ or changes in glottic aperture.³⁴ The latter would lead to an increase in Rrs, while the former would have the opposite effect, rendering physiologic interpretation of the isolated effects difficult if lung volumes were not measured.

A possible advantage of this technique is the help it could provide in locating the site of action of histamine. According to the model of Mead³⁵ refined by Pimmel et al,³⁶ the respiratory system is best represented by central and peripheral airway resistive compartments with a capacitative shunt between them. They demonstrated that an increase in the contribution of the peripheral compartment increased the frequency dependency of resistance. In light of these models, it would be tempting to consider accentuation of the frequency dependency of resistance in this study, as well as in studies in adults with challenges by methacholine³⁷ and cold air³⁸ inhalation to indicate a peripheral site of action of these stimuli. However, this study does not take into account the effects of shunting of pressure at the level of the upper airways with increasing intrathoracic airways resistance upon the frequency dependency of resistance. Recent works³⁹⁻⁴⁰ caution against an interpretation using this technique of peripheral airway changes. This needs to be explored in more detail.

In conclusion, in the population of children studied we found that airway reactivity to histamine can be assessed as reliably by the method of forced oscillation as by the forced expiratory technique. The results of this study suggest that there may be a role for the evaluation of bronchial reactivity using the forced oscillation technique in those children who are capable of quiet breathing into the mouthpiece of the machine, even though they may be incapable of performing a reproducible forced expiratory maneuver.

REFERENCES

- 1 Speight ANP, Lee DA, Hey EN. Underdiagnosis and undertreatment of asthma in childhood. *Br Med J* 1983; 286:1253-56
- 2 DuBois AB, Brody AW, Lewis DH, Burgess BF. Oscillation mechanics of lungs and chest in man. *J Appl Physiol* 1956; 8:587-94
- 3 Wohl MEB, Stigol LC, Mead J. Resistance of the respiratory system in healthy infants and infants with bronchiolitis. *Pediatrics* 1969; 43:495-508
- 4 Mansell A, Levison H, Kruger K, Tripp TL. Measurement of respiratory resistance in children by forced oscillations. *Am Rev Respir Dis* 1972; 106:710-14
- 5 Cogswell JJ. Forced oscillation technique for determination of resistance to breathing in children. *Arch Dis Child* 1973; 48:259-66
- 6 Landau LI, Phelan PD. Evaluation of two techniques for measurement of respiratory resistance by forced oscillation. *Thorax* 1973; 28:136-41
- 7 Rutter N, Milner AD, Hiller EJ. Effect of bronchodilators on respiratory resistance in infants and young children with bronchiolitis and wheezy bronchiolitis. *Arch Dis Child* 1975; 50:719-22
- 8 Wanner A, Zarzecki S, Marks MB. Continuous measurement of respiratory resistance in asthmatic children. *Respiration* 1977; 34:61-68
- 9 Aronson H, Solymar L, Dempsey J, Bjure J, Olsson T, Bake B. A modified forced oscillation technique for measurements of respiratory resistance. *J Appl Physiol* 1977; 42:650-55
- 10 Lenney W, Milner AD. At what age do bronchodilator drugs work? *Arch Dis Child* 1978; 53:532-35
- 11 Lenney W, Milner AD. Alpha and beta adrenergic stimulants in bronchiolitis and wheezy bronchitis in children under 18 months of age. *Arch Dis Child* 1978; 53:707-09
- 12 Stanescu D, Moavero NE, Veriter C, Brasseur L. Frequency dependence of respiratory resistance in healthy children. *J Appl Physiol* 1979; 47:268-72
- 13 Williams SP, Fullton JM, Tsai MJ, Pimmel RL, Collier AM. Respiratory impedance and derived parameters in young children by forced random noise. *J Appl Physiol* 1979; 47:169-74
- 14 Berdel D, Magnussen H, Bockler M. Sollwerte für den oszillatorischen Atemwiderstand im Kindesalter. *Prax Pneumol* 1979; 33:35-37
- 15 Michaelson ED, Grassman ED, Peters WR. Pulmonary mechanics by spectral analysis of forced random noise. *J Clin Invest* 1975; 56:1210-30
- 16 Duiverman EJ, Bak A, Van den Bergh, et al. Forced oscillation technique (Ládsér type). Reference values for resistance and reactance of the pulmonary system in healthy children aged 2.3 to 12.6 years. *Bull Eur Physiopathol Respir* 1983; 19:818-19
- 17 Dickman ML, Schmidt CD, Gardner RM. Spirometric standards for normal children and adolescents (ages 5 years through 18 years). *Am Rev Respir Dis* 1971; 104:680-87
- 18 Knudson RJ, Slatin RC, Lebowitz MD, Burrows B. The maximal expiratory flow-volume curve. Normal standards, variability and effects of age. *Am Rev Respir Dis* 1976; 113:587-600
- 19 Ládsér FJ, Nagels J, Demedts M, Billet L, Van de Woestijne KP. A new method to determine frequency characteristics of the respiratory system. *J Appl Physiol* 1976; 41:101-06
- 20 Ládsér FJ, Nagels J, Van de Woestijne KP. Implementation by means of microprocessor techniques for the measurement of total respiratory impedance during spontaneous breathing. *Prog Respir Res* 1979; 11:135-43
- 21 Nagels J, Ládsér FJ, Van der Linden L, Clément J, Van de Woestijne KP. Mechanical properties of lungs and chest wall during spontaneous breathing. *J Appl Physiol* 1980; 49:408-16
- 22 Nussbaum E, Galant SP. Measurements of total respiratory resistance in children by a modified forced oscillation method.

- Pediatr Res 1984; 18:139-45
- 23 Konig P, Hordvik NL, Pimmel RL. Forced random noise resistance determination in childhood asthma. *Chest* 1984; 86:884-90
 - 24 Cockcroft DW, Killan DN, Mellon JA, Hargreave FE. Bronchial reactivity to inhaled histamine: a method and clinical survey. *Clin Allergy* 1977; 7:235-43
 - 25 Cropp GJA, Bernstein IL, Boushey HA, et al. Guidelines for bronchial inhalation challenges with pharmacologic and antigenic agents. *ATS News* 1980; 2:11-19
 - 26 Duiverman EJ, Neijens HJ, Affourtit M, de Jongste JS, Kerrebijn KF. Bronchial hyperreactivity by the forced oscillation technique. *Eur J Respir Dis* 1983; 64(suppl 128):415-16
 - 27 Solymar L, Aronsson PH, Engstrom I, Bake B, Bjure J. Forced oscillation technique and maximum expiratory flows in bronchial provocation tests in children. *Eur J Respir Dis* 1984; 65:486-95
 - 28 Nadel JA, Tierney DF. Effect of a previous deep inspiration on airway resistance in man. *J Appl Physiol* 1961; 16:717-19
 - 29 Bouhuys A, Hunt VK, Kim BM, Zapletal A. Maximum expiratory flow rates in induced bronchoconstriction in man. *J Clin Invest* 1969; 48:1159-68
 - 30 Higgenbottom T, Clark TJH. Practical importance of a preceding full inhalation or exhalation upon the measurement of airway resistance. *Clin Sci* 1980; 58:249-53
 - 31 Orehek J, Nicoli MM, Delpierre S, Beaupré A. Influence of the previous deep inspiration on the spirometric measurement of provoked bronchoconstriction in asthma. *Am Rev Respir Dis* 1981; 123:269-72
 - 32 Matthys H, Klein G, Kohler D. Standardization of bronchial responsiveness in children. *Prog Respir Res* 1981; 17:155-66
 - 33 Martin J, Powell E, Shore S, Emrich J, Engel L. The role of respiratory muscles in the hyperinflation of bronchial asthma. *Am Rev Respir Dis* 1980; 121:441-47
 - 34 Collett PW, Brancatisano T, Engel L. Changes in the glottic aperture during bronchial asthma. *Am Rev Respir Dis* 1983; 128:719-23
 - 35 Mead J. Contribution of compliance of airways to frequency-dependent behavior of lungs. *J Appl Physiol* 1969; 26:670-73
 - 36 Pimmel RL, Tsai MJ, Winter DC, Bromberg PA. Estimating central and peripheral respiratory resistance. *J Appl Physiol* 1978; 45:375-80
 - 37 Pimmel RL, Fullton JM, Ginsberg JF, Haak ED, McDonnell WE, Bromberg PA. Correlation of airway resistance with forced random noise resistance parameters. *J Appl Physiol* 1981; 51:33-39
 - 38 Decramer M, Demedts M, Van de Woestijne KP. Isocapnic hyperventilation with cold air in healthy non-smokers, smokers and asthmatic subjects. *Bull Eur Physiopathol Respir* 1984; 20:237-43
 - 39 Peslin R, Duvivier C, Gallina C, Cervantes P. Upper airways artifact in respiratory impedance measurements. *Am Rev Respir Dis* 1985; 132:712-14
 - 40 Peslin R, Duvivier C, Didelon J, Cervantes P. Respiratory impedance measured with head generator to minimize airway shunt. *J Appl Physiol* 1985; 59:1790-95

Who are the pulmonary specialists in Tucson?

I need a cardiology consultant in Cleveland.

How can I find qualified consultants in cardiothoracic surgery in the southwest suburbs of Boston?

THE NEW, COMPREHENSIVE 1987-88 ACCP MEMBERSHIP DIRECTORY

The new Directory can answer your questions about referrals better than ever before. It will be an authoritative guide to current interests, residency and specialty training, education and principal specialty of qualified consultants in the US, Canada and worldwide.

The Directory, an invaluable reference to members and colleagues, will now feature alphabetic, geographic and specialty listings for easier referral.

FCCP: Cite these letters with pride. They represent peer recognition of your unique status as a consultant in the disciplines of the American College of Chest Physicians.

