



Repetitive Thinking, Executive Functioning, and Depressive Mood in the Elderly

Journal:	<i>Aging and Mental Health</i>
Manuscript ID	CAMH-2016-0097.R1
Manuscript Type:	Original Article
Keywords:	executive functions, depression,, repetitive thinking, mental rumination

SCHOLARONE™
Manuscripts

Abstract

Repetitive Thinking, Executive Functioning, and Depressive Mood in the Elderly

Aging & Mental Health

Objectives: Previous findings and the depressive-executive dysfunction hypothesis (Alexopoulos, 2003) suggest that the established association between executive functioning and depression is accounted for by repetitive thinking. Investigating the association between executive functioning, repetitive thinking, and depressive mood, the present study empirically tested this mediational model in a sample of older adults, while focusing on both concrete and abstract repetitive thinking. This latter distinction is important given the potential protective role of concrete repetitive thinking, in contrast to the depletive effect of abstract repetitive thinking.

Method: A sample of 43 elderly volunteers, between 75 and 95 years of age, completed tests of executive functioning (the Stroop test, the Trail Making test, and the Fluency test), and questionnaires of repetitive thinking and depression.

Results: Positive correlations were observed between abstract repetitive thinking and depressive mood, and between concrete repetitive thinking and executive functioning; a negative correlation was observed between depressive mood and executive functioning. Further, mediational analysis evidenced that the relation between executive functioning and depressive mood was mediated by abstract repetitive thinking.

Conclusion: The present data provide, for the first time, empirical support to Alexopoulos' (2003) depressive-executive dysfunction hypothesis: the lack of executive resources would favor a mode of abstract repetitive thinking, which in turn would deplete mood. It suggests that clinical intervention targeting depression in the elderly should take

into consideration repetitive thinking modes and the executive resources needed to disengage from rumination.

Keywords: executive functions, rumination, repetitive thinking, depression, elderly

For Peer Review Only

Repetitive Thinking, Executive Functions, and Depressive Mood in the Elderly

Since the seminal work of Susan Nolen-Hoeksema (Nolen-Hoeksema, 1991, 2000), it is well established that mental rumination, a negative and evaluative mode of repetitive thinking (RT), is the main psychological process that precipitates and maintains depressive mood. In an extensive literature review, Watkins (2008) has distinguished two modes of RT: abstract/evaluative RT versus concrete/experiential RT. The former implies thinking repetitively in an abstract mode with a focus on self-evaluation and on the cause and consequence of one's mood state, for example "what will happen to me if I continue to have such a bad sleep? Why can't I relax?". This mode is characteristic of depressive rumination. Indeed, a wealth of research has demonstrated that abstract-evaluative RT is causally related to mood depletion, predicts later depression, poor response to psychological intervention, and is related to procrastination and poor interpersonal problem solving (Nolen-Hoeksema, Wisco, & Lyubomirsky, 2008; Watkins et al., 2011). In contrast, concrete/experiential RT focuses on how the present experience unfolds and how it relates to the individual's needs and goals, for example "what sensation do I feel in my body now? What is the issue at stake, now?". This RT mode has been shown to restore positive mood, to activate goal-directed behaviors, and to promote better interpersonal problem solving (Watkins, 2008; Watkins et al., 2011). Based on these findings, a new psychological treatment of depression has been suggested, the Rumination-Focused Cognitive Behaviour Therapy (Watkins, 2015) that aims at preventing abstract/evaluative RT and at developing concrete/experiential thinking.

Developing concrete/experiential thinking necessitates systematic training. Indeed, concrete/experiential thinking requires important executive control resources (Dalgleish et al., 2007). In particular, concrete/experiential thinking implies the capacity to focus on specific, precise information defining the present experience and the capacity to disengage from other information, such as judgments or general thoughts that are automatically activated. In other

words, concrete/experiential thinking requires extensive attention control resources in order for the individual to remain focused on the immediate data of the present experience and goal, and to prevent automatic capture by judgments or other distracting thoughts. Conversely, experimental evidence has shown that abstract/evaluative RT hampers executive resources in people with depleted mood (Philippot & Brutoux, 2008; Watkins & Brown, 2002).

Interestingly, ample empirical evidence has demonstrated that depression is characterized by a depletion of executive resources (Castaneda, Tuulio-Henriksson, Marttunen, Suvisaari, & Lönnqvist, 2008; Fossati, Ergis, & Allilaire, 2002; Hasselbalch, Knorr, & Kessing, 2011). A model relating abstract/evaluative RT, executive capacities, and depression has been suggested (Belzung, Willner, & Philippot, 2015). Specifically, according to this model, exposure to stressors, such as life events or daily hassles, is likely to trigger abstract/evaluative RT, especially in people who lack the executive resources needed to engage in concrete experiential thinking. An abstract/evaluative RT would in turn precipitate and maintain depressive mood. A further feedback loop would be instantiated by the fact that depressive mood would hamper executive resources, heightening the vulnerability to abstract/evaluative RT that in turn maintains depression.

This cognitive model of depression has a particular relevance for geriatric depression. Indeed, old age is characterized by a decline of cognitive capacities, especially the more complex and elaborated or more fluid cognitive functions (Baltes, 1987). Executive functions (EF) seem particularly vulnerable in old age (Johnson, Lui, & Yaffe, 2007). It follows that the older adults present a higher risk of developing abstract/evaluative RT, given their potentially challenged EF. This might account for the positive association between age and depression (Knight, McMahon, Green, & Skeaff, 2004) or the high prevalence of depression in the elderly (Beekman et al., 1995). In line with this hypothesis, Ricarte, Ros, Serrano, Martínez-Lorca, and Latorre (2015) reported a relation between brooding (a form of abstract

RT) and depressive mood in an elderly sample. Also congruent with this view, Opdebeeck, Nelis, Quinn, and Clare (2014) have observed that mood and rumination explain a significant amount of variance in cognitive test scores in elderly people with lower levels of cognitive reserve.

Synthetizing these ideas, Alexopoulos (2003) has proposed the depressive-executive dysfunction hypothesis to account for geriatric depression. According to that hypothesis, aging would impact negatively on the executive control of attention. The poorer capacity of executive control would favor the development of abstract/evaluative RT, which in turn would deplete mood. The depressive state so induced would further handicap the executive control of attention, hence locking a feedback loop maintaining depression.

Current Study

The association between EF, abstract and concrete RT, and depressive mood has not been systematically investigated yet. Previous findings and the depressive-executive dysfunction hypothesis suggest a mediational model where the association between EF and depression is accounted for by RT. Our goal was to empirically test this mediational model in a sample of older adults, while focusing on both concrete and abstract RT. This study constitutes, to our knowledge, the first empirical test of Alexopoulos' (2003) depressive-executive dysfunction hypothesis. Further, the distinction between both forms of RT is important given the potential protective role of concrete RT, in contrast to the depletive effect of abstract RT.

Method

Participants

A final sample of 43 volunteers (39 women), aged 75 to 95 ($M = 83.21$; $SD = 5.88$), was used in the study. They were recruited from elderly homes (77%), the geriatric outpatient consultation of a general hospital (14%), and acquaintances (9%). To be included in the

study, participants had to be 75 years old or more, to be physically able to carry out the testing, (i.e. functional motor control, hearing, and vision after correction when needed), and to present no signs of dementia or poor memory performance (as diagnosed by a neurologist and indexed by scores below the 10th percentile for free recall and cued recall at the Buschke 16 test; Grober & Buschke, 1987). From the original sample ($N = 53$), ten participants were excluded based on these criteria: 6 because of their poor performance at the Buschke 16, and 4 because of physical problems (e.g., inability to hold a pen, poor sight).

Procedure

Potential candidates for the study were first given a written presentation of the study. Those who manifested interest to participate were met by the experimenter who gave full explanation of the study and invited them to give a written informed consent to participate. Testing was then carried individually in a quiet room, alone with the experimenter, in one or two sessions according to the alertness capacity of the participants. The experimenter was a last year master student in psychology, with clinical experience in geriatrics, and a proper training in neuropsychological testing. The study protocol was approved by the ethical committee of Mont-Godinne university hospital.

Measures

The measures were presented in the same order to the participants, as follows:

Socio-demographic variables. Participants indicated their age, sex, and level of educational attainment.

Neuropsychological tests

The Stroop test (Stroop, 1935) proceeds in three parts. First, participants have to name colors of printed colored rectangles (denomination). Second, they have to read the names of colors printed in black (lecture). Third, they have to name the color of the ink of color words

that do not match the color of their ink (interference). The final score is the number of errors during the interference part.

For the Trail Making Test (TMT, Reitan, 1958) the subjects had to link in ascending order with a pen, without holding it from the sheet, a set of letters and numbers printed in random spaces. In a first section, the participants had to link only numbers (1-2-3...) or letters (a-b-c...). In a second section, participants must alternate numbers and letters (1-a-2-b-3-c...). If a mistake was observed by the experimenter, a correction was requested on line. The final score was the number of errors while performing section 2.

The Fluency test (Cardebat, Doyon, Puel, Goulet, & Joanne, 1990) requires the participants to produce as many words (a) beginning by the same letter (phonetic part) and (b) belonging to the same semantic category (semantic part). The total number of correct words produced was used to compute the two scores of phonetic and of semantic flexibility, respectively. Performing this test implies the activation of effective search strategies in semantic memory.

These neuropsychological tests allowed us computing an EF composite. First, each cognitive indicator was transformed to z-scores. Then, the final EF score was computed as the standardized average of the four indicators (Cronbach's alpha = .82).

Questionnaires

Depressive symptomatology. As a measure of depressive mood, the participants filled out the short version of the Geriatric Depression Scale (GDS, Sheikh & Yesavage, 1986). This scale contains 15 items requiring a "yes" or "no" answer. It presents good psychometric properties (Cronbach's alpha = .75) and demonstrated concurrent and discriminant validity (Freidman, Heisel, & Delavan, 2005).

Repetitive thoughts. The Mini Cambridge-Exeter Repetitive Thought Scale (Douilliez et al., 2014) is a 15-item self-report inventory that assesses two dimensions of ruminative

thinking: concrete and abstract RT. Items were rated on a 4-point scale from 1 (*almost never*) to 4 (*almost always*). The internal consistency, as measured by Cronbach's alpha, is **weak** to good for the subscales concrete RT (Cronbach's alpha=.61) and unconstructive repetitive abstract RT (Cronbach's alpha=.80; Douilliez et al., 2014).

Data Analysis

We examined the association between EF and **depressive mood** and the mediational role of concrete and abstract repetitive thinking (RT) using a multiple mediation model (**Figure 1**) estimated by the SPSS PROCESS macro (Hayes, 2013). This macro tests the significance of the association between EF and the two indicators of RT (paths a_1 [concrete RT] and a_2 [abstract RT]), between the mental rumination and **depressive mood** (paths b_1 and b_2), and between EF and **depressive mood** (paths c and c'). The difference between path c (total effect) and path c' (direct effect) is that c' is estimated while adjusting for the two mediator variables (i.e., concrete and abstract RT). The indirect effects of EF on **depressive mood** via concrete and abstract RT (path $a_1 * \text{path } b_1$ and path $a_2 * \text{path } b_2$) were estimated using a bootstrapping approach ($N = 1,000$).¹ Given the cross-sectional nature of the data we also examined the same multiple mediation model with EF as an outcome and depressive mood as a predictor.

Results

The means, standard deviations, and the inter-correlations between all the study variables are presented in Table 1. It can be seen that depressive mood is strongly related to abstract/evaluative RT, but not to concrete experiential thinking. Further, depressive mood is strongly and negatively related to **EF**.

The results of the mediation analysis are presented in Table 2. It includes the bootstrapped estimates and 95% bias-corrected confidence intervals (CI) for the total, direct,

¹ The same pattern of results was obtained when age, sex, and education were included as covariates.

and indirect effects of EF on **depressive mood**. As expected, EF was positively associated with concrete RT, $a_1 = 1.25$, $SE = .431$, $p = .006$, and negatively associated with abstract RT, $a_2 = -1.56$, $SE = .599$, $p = .013$. In turn, abstract RT was significantly associated with **depressive mood**, $b_2 = .35$, $SE = .088$, $p < .001$. Higher levels of abstract RT correspond to higher levels of **depressive mood**. The indirect effect of EF on **depressive mood** via abstract RT was significant, estimate = $-.54$, $SE = .254$, $p = .034$, **effect size (percent mediation) = .48**. The direct effect was not significant, estimate = $-.60$, $SE = .391$, $p = .131$, which according to the traditional statistical approach to test mediational models based on causal steps (Baron & Kenny, 1986) would imply the existence of a full mediation (Figure 1). When the same mediational model was estimated with EF as an outcome, the indirect effects of depressive mood via abstract and concrete RT were not significant estimates : $-.03$, $SE = .039$, $p = .411$; $-.02$, $SE = .022$, $p = .300$, respectively.

Discussion

In the present data, depressive mood was clearly related to abstract/evaluative RT, while the relation with concrete/experiential RT did not reach significance ($r = -.19$, p (one tail) $< .11$). Previous studies, conducted in student and adult populations, have systematically evidenced a positive correlation between depressive mood and abstract/evaluative RT (ranging from .44 to .53, according to the study) while the reverse correlation for concrete/experiential RT was not systematically reaching statistical significance, albeit always negative, ranging from $-.12$ to $-.39$ (Di Schiena, Luminet, Philippot, & Douilliez, 2012; DiSchiena, Luminet, & Philippot, 2011; Douilliez et al., 2014; Douilliez & Philippot, 2012; Heeren & Philippot, 2011). The present study thus replicates in a sample of adults over 75 year of age what has been observed in samples of younger adults: depressive mood is strongly associated with abstract/evaluative RT, even more so in the present sample. **In contrast**, the reverse relation with concrete experiential thinking is much weaker. As suggested by many

1
2
3 authors (Nolen-Hoeksema, 1991; Teasdale, 1993; Watkins, 2008, 2015), depression is
4
5 specifically associated with a mode of abstract, overgeneralizing and judgmental thinking,
6
7 mostly about the self. This is true along the whole lifespan, and the present study is the first
8
9 to observe this specific relation in people over 75 year of age. In the present data,
10
11 concrete/experiential RT does not appear to be a significant protective factor against
12
13 depression. This latter mode of RT might be more protective against anxiety than depression
14
15 (Douilliez et al., 2014).
16
17

18
19 As reported by many previous studies in the adult population (Hasselbalch et al.,
20
21 2011), depressive mood is also clearly related to executive dysfunction in the present elderly
22
23 sample. In the present data, it should be noted that executive capacities are equally related to
24
25 both modes of RT, positively so with concrete/experiential RT, and negatively so for
26
27 abstract/evaluative RT. This pattern of results is congruent with the notion that
28
29 concrete/experiential RT is an effortful process, requiring the capacity to disengage from
30
31 automatically activated information and to willfully elaborate information relevant for the
32
33 target task (Dagleish et al., 2007). In contrast, abstract/evaluative RT would be favored by
34
35 the lack of such executive control in RT and would represent a more automatic mode of RT.
36
37

38
39 Importantly, the mediational hypothesis revealed a pattern of results that is consistent
40
41 with the notion that the relationship between executive dysfunction and depressive mood is
42
43 partially mediated by abstract/evaluative RT. People with less executive resources would be
44
45 more vulnerable to abstract/evaluative RT, which in turn concurs to depressive mood. This
46
47 pattern of results corroborates Alexopoulos' (2003) depressive-executive dysfunction
48
49 hypothesis. It suggests that training elderly people to disengage from abstract/evaluative RT
50
51 might be a fruitful strategy to alleviate depressive mood, as already established in younger
52
53 adults (Watkins, 2015). In elderly people this training might more particularly entail a
54
55
56
57
58
59
60

specific cognitive training of the executive resources needed to disengage from automatically activated verbal information.

The correlational nature of the present data prevents firm causal interpretations. However, the fact that the reverse mediational model (EF as a mediator of the effect of RT on depressive mood) was not significant strengthens the support to our directional hypothesis. Future studies might investigate this issue by examining how training EF in the elderly might impact on their RT and, consecutively on their depleted mood.

The present study did not include a comparative sample of young adults. Hence, it cannot be determined whether abstract/evaluative RT is more present in elderly people than in young adults, and whether differences in EF might account for such a difference. Still, a comparison can be made with the data reported by Douilliez et al. (2014) for a normative sample of 243 adults aged between 18 and 75 years. In that sample, abstract/evaluative thinking reached a mean of 18.93, very similar to the mean of 18.30 observed in the present sample. For concrete/experiential thinking, the normative adults' sample ($M = 17.06$, $SD = 3.71$) is slightly higher than the one observed in the present elderly sample ($M = 15.74$, $SD = 3.03$). Yet, a recent study comparing young and old adults suggests less brooding (a form of RT) in the latter (Ricarte et al., 2015). This question requires further attention by future research.

A limit of the present study is that the sample is mostly constituted by women. While this reflects the demography of this age group, future studies should ascertain that the same patterns of covariations are observed in men. Another limit is that all the data have been collected at the same time. A better test of the mediational hypothesis would be to conduct a longitudinal study, examining how the causal cascade unfolds in time.

In sum, the present study is the first to examine the relationship between executive capacities, two modes of RT, and depressive mood in a sample over 75 year of age. Like in

1
2
3 their younger counter-parts, elderly people show a strong relation between abstract evaluative
4
5 RT and depressive mood. Further, the present data provide, for the first time, empirical
6
7 support to Alexopoulos' (2003) depressive-executive dysfunction hypothesis. It suggests that
8
9 clinical intervention targeting depressive mood in the elderly should take into consideration
10
11 abstract/evaluative RT and the executive resources needed to disengage from it.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1.
Means, Standard Deviations, and Inter-correlations among the Study Variables (n=43)

	<i>M (SD)</i> <i>or %</i>	1	2	3	4	5	6	7	8	9	10
1. Age	83.21 (5.88)										
2. Sex (Women)	90.7%	.002									
3. Education											
Primary school	37.2%										
Secondary School	51.2%	.002	.003								
Higher Education	11.6%										
4. Semantic Fluency	18.44 (7.08)	-.16	-.01	.34*							
5. Phonemic Fluency	10.30 (4.94)	-.06	-.05	.58***	.70***						
6. TMT	3.21 (2.43)	-.22	.16	.13	.54***	.56***					
7. Stroop	5.43 (5.67)	-.06	.10	.13	.50**	.49**	.52***				
8. Executive Functioning	0 (1.00)	-.14	.06	.37*	.84***	.84***	.80***	.78***			
9. Abstract RT	18.30 (4.14)	.11	-.26	-.07	-.21	-.34*	-.34*	-.39*	-.38*		
10. Concrete RT	15.74 (3.03)	-.14	-.05	-.02	.35*	.24	.33*	.42**	.41**	-.23	
11. Depressive Mood	4.60 (2.73)	.20	-.19	.21	-.23	-.34*	-.42**	-.46**	-.41**	.60***	-.19

M = mean; *SD* = standard deviation; ****p*<.001 ***p*<.01. **p*<.05

Table 2.

Total, Direct, and Indirect Effects of Executive Functioning on Depressive Mood

	Bootstrap Estimate	SE	95% BC CI Lower	95% BC CI Upper
Total Effect	-1.12	.388	-1.904	-.336
Direct Effect	-.60	.391	-1.392	.188
Indirect Effect Concrete RT	.02	.161	-.415	.259
Indirect Effect Abstract RT	-.54	.230	-1.063	-.155

Note: Based on 1,000 bootstrap samples; SE = standard error; BC CI = bias-corrected confidence intervals;

RT = Repetitive Thinking.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Captions

Figure 1.

The Multiple Mediation Model of the Association between Executive Functioning and Depressive Mood via Concrete and Abstract Repetitive Thinking

Unstandardized Regression Coefficients from a Bootstrap Procedure are Presented

**** $p < .001$, ** $p < .01$, * $p < .05$; $R^2 = .40$ *** (Direct Effect Model)*

References

- Alexopoulos, G. S. (2003). Role of executive function in late-life depression. *Journal of Clinical Psychiatry*, 64, 18-23.
- Baltes, P. B. (1987). Theoretical propositions of life-span developmental psychology: On the dynamics between growth and decline. *Developmental Psychology*, 23, 611-626.
- Baron, R. M., & Kenny, D. A. (1986). The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*, 51, 1173-1182.
- Beekman, A. T., Deeg, D. J., van Tilburg, T., Smit, J. H., Hooijer, C., & van Tilburg, W. (1995). Major and minor depression in later life: A study of prevalence and risk factors. *Journal of Affective Disorders*, 36, 65-75. doi:10.1016/0165-0327(95)00061-5
- Belzung, C., Willner, P., & Philippot, P. (2015). Depression: From psychopathology to pathophysiology. *Current Opinion in Neurobiology*, 30, 24-30. doi:10.1016/j.conb.2014.08.013
- Cardebat, D., Doyon, B., Puel, M., Goulet, P., & Joanette, Y. (1990). Formal and semantic lexical evocation in normal subjects. Performance and dynamics of production as a function of sex, age and educational level. *Acta Neurologica Belgica*, 90, 207-217.
- Castaneda, A. E., Tuulio-Henriksson, A., Marttunen, M., Suvisaari, J., & Lönnqvist, J. (2008). A review on cognitive impairments in depressive and anxiety disorders with a focus on young adults. *Journal of Affective Disorders*, 106, 1-27. doi:10.1016/j.jad.2007.06.006
- Dalglish, T., Williams, J. M. G., Golden, A. M., Perkins, N., Barrett, L. F., Barnard, P. J., . . . Watkins, E. (2007). Reduced specificity of autobiographical memory and depression:

The role of executive control. *Journal of Experimental Psychology: General*, 136, 23-42. doi:10.1037/0096-3445.136.1.23

Di Schiena, R., Luminet, O., Philippot, P., & Douilliez, C. (2012). Adaptive and maladaptive perfectionism in depression: Preliminary evidence on the role of adaptive and maladaptive rumination. *Personality and Individual Differences*, 53, 774-778. doi:10.1016/j.paid.2012.05.017

DiSchiena, R., Luminet, O., & Philippot, P. (2011). Alexithymia and ruminative thinking styles in depression: A multidimensional investigation. *Personality and Individual Differences*, 50, 10-14.

Douilliez, C., Heeren, A., Lefèvre, N., Watkins, E., Barnard, P., & Philippot, P. (2014). Validation of the French version of a questionnaire that evaluates constructive and non-constructive repetitive thoughts. *Canadian Journal of Behavioural Science*, 46, 185-192. doi:10.1037/a0033185

Douilliez, C., & Philippot, P. (2012). Constructive and unconstructive repetitive thinking in maladaptive perfectionists. *Journal de Thérapie Comportementale et Cognitive*, 22, 68-74.

Fossati, P., Ergis, A. M., & Allilaire, J. F. (2002). Executive functioning in unipolar depression: A review. *Encephale*, 28, 97-107.

Freidman, B., Heisel, M. J., & Delavan, R. L. (2005). Psychometric properties of the 15-item Geriatric Depression Scale in a functionally impaired, cognitively intact, community-dwelling elderly primary care patients. *Journal of the American Geriatrics Society*, 53, 1570-1576. doi:10.1111/j.1532-5415.2005.53461.x

Grober, E., & Buschke, H. (1987). Genuine memory deficits in dementia. *Developmental Neuropsychology*, 3, 13-36.

- Hasselbalch, B. J., Knorr, U., & Kessing, L. V. (2011). Cognitive impairment in the remitted state of unipolar depressive disorder: A systematic review. *Journal of Affective Disorders, 134*, 20-31. doi:10.1016/j.jad.2010.11.011
- Hayes, A. F. (2013). *Introduction to Mediation, Moderation, and Conditional Process Analysis: A Regression-Based Approach*. New York: The Guilford Press.
- Heeren, A., & Philippot, P. (2011). Changes in ruminative thinking mediate the clinical benefits of mindfulness: Preliminary findings. *Mindfulness, 2*, 8-13. doi:10.1007/s12671-010-0037-y
- Johnson, J. K., Lui, L. Y., & Yaffe, K. (2007). Executive function, more than global cognition, predicts functional decline and mortality in elderly women. *Journals of Gerontology. Series A: Biological Sciences and Medical Sciences, 62A*, 1134-3341.
- Knight, R. G., McMahon, J., Green, T. J., & Skeaff, C. M. (2004). Some normative and psychometric data for the Geriatric Depression Scale and the Cognitive Failures Questionnaire from a sample of health older persons. *New Zealand journal of psychology, 33*, 163-170.
- Nolen-Hoeksema, S. (1991). Responses to depression and their effects on the duration of depressive episodes. *Journal of Abnormal Psychology, 100*, 569-582.
- Nolen-Hoeksema, S. (2000). The role of rumination in depressive disorders and mixed anxiety/depressive symptoms. *Journal of Abnormal Psychology, 109*, 504-511. doi:101037/10021-843X.109.3.504
- Nolen-Hoeksema, S., Wisco, B. E., & Lyubomirsky, S. (2008). Rethinking rumination. *Perspectives on Psychological Science, 3*, 400-424. doi:10.1111/j.1745-6924.2008.00088.x

Opdebeeck, C., Nelis, S. M., Quinn, C., & Clare, L. (2014). How does cognitive reserve impact on the relationships between mood, rumination, and cognitive function in later life? *Aging & Mental Health*, 19, 705-712. doi:10.1080/13607863.2014.962005

Philippot, P., & Brutoux, F. (2008). Induced rumination dampens executive processes in dysphoric young adults. *Journal of Behavior Therapy and Experimental Psychiatry*, 39, 219-227. doi:10.1016/j.jbtep.2007.07.001

Reitan, R. M. (1958). Validity of the Trail Making Test as an indicator of organic brain damage. *Perceptual and Motor Skills*, 8, 271-276. doi:10.2466/PMS.8.7.271-276

Ricarte, J., Ros, L., Serrano, J. P., Martínez-Lorca, M., & Latorre, J. M. (2015). Age differences in rumination and autobiographical retrieval. *Aging & Mental Health*, 2, 1-7. doi:10.1080/13607863.2015.1060944

Sheikh, J. I., & Yesavage, J. A. (1986). Geriatric Depression Scale (GDS): Recent evidence and development of a shorter version. *Clinical Gerontology*, 5, 165-173.

Stroop, J. R. (1935). Studies of interference in serial verbal reactions. *Journal of Experimental Psychology*, 18, 643-662.

Teasdale, J. D. (1993). Emotion and two kinds of meaning: Cognitive therapy and applied cognitive science. *Behaviour Research and Therapy*, 31, 339-354.


Watkins, E. R. (2008). Constructive and unconstructive repetitive thought. *Psychological Bulletin*, 134, 163-206. doi:10.1037/0033-2909.134.2.163

Watkins, E. R. (2015). Psychological treatment of depressive rumination. *Current Opinion in Psychology*, 4, 32-36.

Watkins, E. R., & Brown, R. G. (2002). Rumination and executive function in depression: An experimental study. *Journal of Neurology, Neurosurgery, and Psychiatry*, 72, 400-402. doi:10.1136/jnnp.72.3.400

Watkins, E. R., Mullan, E., Wingrove, J., Rimes, K., Steiner, H., Bathurst, N., . . . Scott, J.
(2011). Rumination-focused cognitive-behavioural therapy for residual depression:
Phase II randomised controlled trial. *The British Journal of Psychiatry*, 199, 317-322.
doi:10.1192/bjp.bp.110.090282

For Peer Review Only



254x190mm (96 x 96 DPI)