

EUR Research Information Portal

Human Leukocyte Antigen Mismatching and Survival in Contemporary Hematopoietic Cell Transplantation for Hematologic Malignancies

Published in:

Journal of Clinical Oncology

Publication status and date:

Published: 21/08/2024

DOI (link to publisher):

[10.1200/JCO.24.00582](https://doi.org/10.1200/JCO.24.00582)

Document Version

Publisher's PDF, also known as Version of record

Document License/Available under:

Article 25fa Dutch Copyright Act

Citation for the published version (APA):

Arrieta-Bolaños, E., Bonneville, E. F., on behalf of the Cellular Therapy and Immunobiology Working Party of the EBMT, Crivello, P., Robin, M., Gedde-Dahl, T., Salmenniemi, U., Kröger, N., Yakoub-Agha, I., Crawley, C., Choi, G., Broers, A. E. C., Forcade, E., Carre, M., Poiré, X., Huynh, A., Lenhoff, S., Ciceri, F., Tholouli, E., ... Fleischhauer, K. (2024). Human Leukocyte Antigen Mismatching and Survival in Contemporary Hematopoietic Cell Transplantation for Hematologic Malignancies. *Journal of Clinical Oncology*, 42(28), 3287-3299. <https://doi.org/10.1200/JCO.24.00582>

[Link to publication on the EUR Research Information Portal](#)

Terms and Conditions of Use

Except as permitted by the applicable copyright law, you may not reproduce or make this material available to any third party without the prior written permission from the copyright holder(s). Copyright law allows the following uses of this material without prior permission:

- you may download, save and print a copy of this material for your personal use only;
- you may share the EUR portal link to this material.

In case the material is published with an open access license (e.g. a Creative Commons (CC) license), other uses may be allowed. Please check the terms and conditions of the specific license.

Take-down policy

If you believe that this material infringes your copyright and/or any other intellectual property rights, you may request its removal by contacting us at the following email address: openaccess.library@eur.nl. Please provide us with all the relevant information, including the reasons why you believe any of your rights have been infringed. In case of a legitimate complaint, we will make the material inaccessible and/or remove it from the website.

Human Leukocyte Antigen Mismatching and Survival in Contemporary Hematopoietic Cell Transplantation for Hematologic Malignancies

Esteban Arrieta-Bolaños, PhD^{1,2} ; Edouard F. Bonneville, PhD^{3,4} ; Pietro Crivello, PhD¹ ; Marie Robin, MD⁵ ; Tobias Gedde-Dahl, MD⁶ ; Urp Salmenniemi, MD⁷ ; Nicolaus Kröger, MD⁸ ; Ibrahim Yakoub-Agha, MD⁹ ; Charles Crawley, MD¹⁰; Goda Choi, MD¹¹; Annoek E.C. Broers, MD¹² ; Edouard Forcade, MD¹³; Martin Carre, MD¹⁴; Xavier Poiré, MD¹⁵; Anne Huynh, MD¹⁶; Stig Lenhoff, MD¹⁷; Fabio Ciceri, MD^{18,19}; Eleni Tholouli, MD²⁰; Thomas Schroeder, MD²¹ ; Eric Deconinck, MD²² ; Kristina Carlson, MD²³ ; Liesbeth C. de Wreede, PhD³ ; Jorinde D. Hoogenboom, MSc⁴; Florent Malard, MD²⁴ ; Annalisa Ruggeri, MD¹⁸; and Katharina Fleischhauer, MD^{1,2} ; on behalf of the Cellular Therapy and Immunobiology Working Party of the EBMT

DOI <https://doi.org/10.1200/JCO.24.00582>

ABSTRACT

ACCOMPANYING CONTENT

 Editorial, p. 3263
 Data Supplement

Accepted June 14, 2024
Published August 21, 2024

J Clin Oncol 42:3287-3299
© 2024 by American Society of
Clinical Oncology



View Online
Article

PURPOSE Human leukocyte antigen (HLA) mismatching can reduce survival of patients with blood cancer after hematopoietic cell transplantation (HCT). How recent advances in HCT practice, in particular graft-versus-host disease (GVHD) prophylaxis by post-transplantation cyclophosphamide (PTCy), influence HLA risk associations is unknown.

PATIENTS AND METHODS The study included 17,292 unrelated HCTs with 6-locus high-resolution HLA typing, performed mainly for acute leukemia or related myeloid neoplasms between 2016 and 2020, including 1,523 transplants with PTCy. HLA risk associations were evaluated by multivariable Cox regression models, with overall survival (OS) as primary end point.

RESULTS OS was lower in HLA mismatched compared with fully matched transplants (hazard ratio [HR], 1.23 [99% CI, 1.14 to 1.33]; $P < .001$). This was driven by class I HLA-A, HLA-B, HLA-C (HR, 1.29 [99% CI, 1.19 to 1.41]; $P < .001$) but not class II HLA-DRB1 and HLA-DQB1 (HR, 1.07 [99% CI, 0.93 to 1.23]; $P = .19$). Class I antigen-level mismatches were associated with worse OS than allele-level mismatches (HR, 1.36 [99% CI, 1.24 to 1.49]; $P < .001$), as were class I graft-versus-host peptide-binding motif (PBM) mismatches compared with matches (HR, 1.42 [99% CI, 1.28 to 1.59]; $P < .001$). The use of PTCy improved GVHD, relapse-free survival compared with conventional prophylaxis in HLA-matched transplants (HR, 0.77 [0.66 to 0.9]; $P < .001$). HLA mismatching increased mortality in PTCy transplants (HR, 1.32 [1.04 to 1.68]; $P = .003$) similarly as in non-PTCy transplants (interaction $P = .43$).

CONCLUSION Class I but not class II HLA mismatches, especially at the antigen and PBM level, are associated with inferior survival in contemporary unrelated HCT. These effects are not significantly different between non-PTCy compared with PTCy transplants. Optimized HLA matching should still be considered in modern HCT.

INTRODUCTION

Allogeneic hematopoietic cell transplantation (HCT) is the only potentially curative treatment option for many patients with high-risk blood cancers.¹⁻³ Owing to important advances in transplant procedures, including high-accuracy human leukocyte antigen (HLA) tissue typing, pharmacological approaches to combat infection, new conditioning regimens, and new protocols for graft-versus-host disease

(GVHD) prophylaxis and treatment, HCT outcomes have improved consistently over the past decades.⁴⁻⁸ However, donor-recipient HLA mismatching can significantly impair patient survival.^{9,10} Hence, clinically tolerable HLA mismatches are sought and may include class- or locus-specific risks,^{9,11-13} B-leader disparity,^{14,15} or the divergence between the repertoire of antigenic peptides, that is, the immunopeptidome, of mismatched HLA-DPB1 or HLA class I molecules.¹⁶⁻¹⁹ These models have been developed in the

CONTEXT

Key Objective

This study investigates the role of human leukocyte antigen (HLA) mismatching in contemporary unrelated hematopoietic cell transplantation (HCT), in particular under post-transplant cyclophosphamide (PTCy)–based graft-versus-host disease (GVHD) prophylaxis.

Knowledge Generated

Class I but not class II HLA mismatches, especially at the antigen and peptide-binding motif level, are associated with inferior survival after contemporary unrelated HCT. This effect is not significantly different between transplants under conventional GVHD prophylaxis compared with PTCy. On the basis of this, optimized HLA matching should still be considered in modern HCT.

Relevance (C.F. Craddock)

This large registry study confirms the adverse impact of HLA mismatch on outcome in patients transplanted from an unrelated donor which, in this study, is not mitigated by the use of PTCy as a GVHD prophylaxis regimen. Ongoing prospective randomized trials will be of importance in definitively addressing the potential of PTCy to improve outcomes after mismatched HCT.*

*Relevance section written by JCO Associate Editor Charles F. Craddock, MD.

setting of GVHD prophylaxis mainly based on calcineurin inhibitors, with or without in vivo T-cell depletion (TCD) by anti-thymocyte globulin (ATG) or alemtuzumab. The introduction of post-transplantation cyclophosphamide (PTCy) has markedly improved outcomes of allogeneic HCT across HLA mismatches, with similar survival rates after HLA-matched HCT under conventional GVHD prophylaxis and full-haplotype mismatched HCT under PTCy.^{6,20–22} Recently, the use of PTCy has been reported to improve GVHD, relapse-free survival (GRFS) also in unrelated HLA-matched HCT.^{23,24} If and how the HLA risk associations observed under conventional GVHD prophylaxis apply to PTCy-based transplant platforms is currently unknown.

Here, we investigated the impact of HLA in contemporary unrelated HCT, leveraging a cohort of over 17,000 recent transplants with 6-locus, high-resolution HLA typing collected through the European Society for Blood and Marrow Transplantation (EBMT) and performed under conventional or PTCy-based GVHD prophylaxis. This allowed us to study the potential interaction between HLA matching and the type of GVHD prophylaxis to comprehensively address the role of HLA in recent HCT for hematologic malignancies.

PATIENTS AND METHODS

Study Population and Design

The study included 17,292 adult patients with hematologic malignancies who underwent a first unrelated HCT from bone marrow or peripheral blood stem cells between 2005 and 2020 and had available 6-locus, high-resolution HLA typing for both donors and recipients (Table 1). Patients transplanted with cord blood units and those receiving grafts

with ex-vivo TCD were excluded. Clinical data were collected and stored by the EBMT Registry. Transplant centers reporting to the Registry commit to obtain informed consent in agreement with the principles of the Declaration of Helsinki and according to the local regulations applicable at the time to report pseudonymized data. The study was approved by the Cellular Therapy and Immunobiology Working Party of the EBMT.

HLA Matching

The number of allele (ie, second field) mismatches at HLA-A, HLA-B, HLA-C, HLA-DRB1, and/or HLA-DQB1 (henceforth referred to as the five main HLA loci) was used to classify transplants as fully matched (ie, 10 of 10) or 9 of 10 and eight of 10 matched, respectively (Table 1). Single HLA mismatches in the 9 of 10 pairs were further classified according to (1) their class (ie, HLA class I v HLA class II), (2) their locus (ie, any of the five main HLA loci), and (3) their resolution (ie, second-field allelic or first-field antigenic).²⁶ Single mismatches at HLA-B were also classified as B-leader¹⁵ matched or mismatched. Moreover, single mismatches at HLA-A, HLA-B, or HLA-C were classified as peptide-binding motif (PBM)–matched or mismatched in the GVH vector, as previously described.¹⁹ Finally, mismatches at HLA-DPB1 were divided into permissive or nonpermissive in both 9 of 10 and 10 of 10 pairs according to the functional distance-based, bidirectional T-cell epitope (TCE) model.^{16,18,27–30}

Clinical Outcomes and Statistical Analysis

Clinical outcome was studied in the full cohort and in subgroups according to the aforementioned models. The

TABLE 1. Patient-, Disease-, and Transplant-Related Characteristics of the Cohort

Variable	Groups	PTCy No (n = 15,630)	PTCy Yes (n = 1,523)
Age, years, median (IQR)	Patients	56 (44-63)	55 (43-63)
	Donors	29 (24-37)	28 (24-36)
	Missing, ^a No. (%)	1,182 (7)	51 (3)
Disease at transplant, No. (%)	AML	5,706 (37)	522 (34)
	MDS	2,137 (14)	204 (13)
	MPN	1,096 (7)	78 (5)
	ALL	1,656 (11)	178 (12)
	Other ^b	5,035 (31)	541 (36)
Disease risk index, ^c No. (%)	Low	861 (5)	77 (5)
	Intermediate	11,405 (74)	1,109 (73)
	High	2,888 (18)	290 (19)
	Very high	476 (3)	47 (3)
Recipient-donor sex match, No. (%)	Male-male	7,154 (46)	713 (47)
	Male-female	1,974 (13)	210 (14)
	Female-male	3,971 (26)	371 (24)
	Female-female	2,396 (15)	222 (15)
	Missing	135 (<1)	7 (<1)
Recipient-donor CMV match, No. (%)	-/-	4,769 (31)	390 (26)
	-/+	1,364 (9)	105 (7)
	+/-	3,985 (25)	436 (28)
	+/+	5,332 (34)	577 (38)
	Missing	180 (1)	15 (1)
KPS, No. (%)	≥90	10,579 (68)	997 (65)
	80	3,315 (21)	407 (27)
	<80	736 (5)	74 (5)
	Missing	1,000 (6)	45 (3)
HCT-CI, No. (%)	Low risk (0)	6,230 (40)	728 (48)
	Intermediate risk (1-2)	2,815 (18)	299 (19)
	High risk (≥3)	3,010 (19)	360 (24)
	Missing	3,575 (23)	136 (9)
Graft source, No. (%)	BM	1,411 (9)	75 (5)
	PBSC	14,219 (91)	1,448 (95)
Conditioning regimen, No. (%)	MAC	4,600 (30)	593 (39)
	MAC + TBI	2,024 (13)	243 (16)
	RIC	7,377 (47)	492 (32)
	RIC + TBI	1,537 (9.9)	178 (12)
	Missing	92 (<1)	17 (1)
T-cell depletion, ^d No. (%)	No	3,026 (19)	1,159 (76)
	Yes	12,604 (81)	364 (24)
Conventional GVHD prophylaxis, No. (%)	CsA + MTX	5,495 (35)	0 (0)
	CsA + MMF	3,928 (25)	0 (0)
	CsA	1,491 (10)	0 (0)
	TAC-based	1,639 (10)	495 (33)
	Other	3,077 (20)	1,028 (67)
Transplant year, No. (%)	2005-2010	1,631 (10)	3 (<1)
	2011-2015	4,000 (26)	171 (11)
	2016-2020	9,999 (64)	1,349 (89)

(continued on following page)

TABLE 1. Patient-, Disease-, and Transplant-Related Characteristics of the Cohort (continued)

Variable	Groups	PTCy No (n = 15,630)	PTCy Yes (n = 1,523)
HLA matching status, No. (%)	10 of 10	12,210 (78)	924 (61)
	9 of 10	3,004 (19)	522 (34)
	8 of 10	416 (3)	77 (5)
HLA class mM, ^e No. (%)	Class I	2,117 (72)	418 (80)
	Class II	827 (28)	104 (20)
HLA class I PBM status, ^f No. (%)	PBM-GVH match	672 (31)	132 (32)
	PBM-GVH mM	1,182 (54)	219 (52)
	Not informative	323 (15)	67 (16)
HLA class I mM resolution, ^g No. (%)	Low-res match	530 (24)	68 (16)
	Low-res mM	1,647 (76)	350 (84)
HLA-DPB1 matching status, No. (%)	Allele-matched	3,998 (26)	360 (24)
	Permissive	6,462 (41)	619 (41)
	Nonpermissive	5,170 (33)	544 (36)

NOTE. 139 pairs with missing data for PTCy use are not included in the table.

Abbreviations: AL, acute leukemia; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BM, bone marrow; CMV, cytomegalovirus; CsA, cyclosporin A; GVH, graft-versus-host; GVHD, graft-versus-host disease; HCT-CI, hematopoietic cell transplantation comorbidity index; HLA, human leukocyte antigen; KPS, Karnofsky Performance Score; Low-res, low resolution; MDS, myelodysplastic neoplasm; mM, mismatch; MMF, mycophenolate mofetil; MPN, myeloproliferative neoplasm; MTX, methotrexate; PBM, peptide binding motif group; PBSC, peripheral blood stem cells; PTCy, post-transplantation cyclophosphamide; RIC, reduced intensity conditioning; TAC, tacrolimus; TBI, total body irradiation.

^aMissing data refer to donor age only.

^bIncludes non-Hodgkin lymphoma (n = 1,509), chronic lymphocytic leukemia (n = 446), chronic myeloid leukemia (n = 481), Hodgkin lymphoma (n = 371), multiple myeloma (n = 643), MDS/MPN including chronic myelomonocytic leukemia (n = 1,377), and other acute leukemia (n = 793).

^cDisease risk index calculated as described.²⁵

^dIn vivo T-cell depletion by antithymocyte globulin, anti-T-lymphocyte globulin, or alemtuzumab.

^eReferring to the 9 of 10 group.

^fReferring to the HLA class I mismatched 9 of 10 group.

^gLow-res match, allelic mismatch at the second field. Low-res mismatch, antigenic mismatch at the first field. Numbers and percentages refer to the HLA class I mismatched 9 of 10 group.

primary end point was overall survival (OS). GRFS, RFS, nonrelapse mortality (NRM), disease relapse or progression, and grades 2 to 4 and 3 to 4 aGVHD and chronic (c)GVHD were secondary end points. End point definitions were as follows: OS, time from transplant to death from any cause; GRFS, time to aGVHD grades 3 to 4, extensive cGVHD, relapse/progression (henceforth referred to jointly as relapse), or death, whichever occurred first³¹; RFS, time to relapse or death; NRM, time to death without relapse.

Survival probabilities and cumulative incidence estimates were calculated using the Kaplan-Meier and the Aalen-Johansen method, respectively. Competing risks models were used to calculate the cumulative incidences of NRM and aGVHD/cGVHD, for which relapse and death or second allogeneic HCT were considered as competing risks, respectively. Univariable comparisons were performed with the log-rank and Gray tests. Regression analyses were performed using multivariable Cox proportional hazards models, including a gamma-distributed frailty term for transplant center.³² For outcomes with competing risks, cause-specific hazard models were fitted. To analyze the differential effects of HLA mismatch models in the subgroups with conventional

or PTCy-based GVHD prophylaxis, interaction analyses were implemented. Complete-case analysis was used to deal with missing data in baseline covariates other than HCT-CI, Karnofsky Performance Score, and donor age, where the missing indicator method was used instead.^{33,34} To account for multiple testing, a predefined threshold of $P < .01$ was used to determine statistical significance. All statistical analyses were performed using R Statistical Software (version 4.3.1).³⁵

Further details on the clinical outcomes and statistical methods can be found in the Data Supplement (online only).

RESULTS

Overall Outcomes

The median follow-up time in the overall cohort was 39.7 (95% CI, 38.8 to 40.8) months. At 60 months post-transplant, the OS was 50.8% (95% CI, 49.9 to 51.7) while GRFS and RFS were 21.4% (95% CI, 20.7 to 22.2) and 44.2% (95% CI, 43.3 to 45.1), respectively. The overall cumulative incidences of relapse, NRM, and cGVHD at 60 months were




FIG 1. Survival after contemporary HCT according to HLA mismatching. Kaplan-Meier and forest plots show overall survival (OS) probabilities (shaded areas) and adjusted HR estimates and their 99% CI for the entire cohort stratified according to (A, B) the number of mismatches (mM) at HLA-A, HLA-B, HLA-C, HLA-DRB1, or HLA-DQB1; (C, D) the specific mismatched HLA locus and class among the 9 of 10 pairs; (E, F) the typing resolution for HLA class I mM in the 9 of 10 pairs; and (G, H) PBM GVH (continued on following page)

FIG 1. (Continued). matching for HLA class I mM in the 9 of 10 pairs. The fully matched 10 of 10 pairs are used as reference in the HR estimates (vertical lines). The numbers below the Kaplan-Meier plots correspond to the number of individuals at risk at the different time points according to the groups indicated in the legend from top to bottom. HR and *P* values in the forest plots are derived from multivariable analyses shown in Table 2. GVH, graft-versus-host; HCT, hematopoietic cell transplantation; HLA, human leukocyte antigen; HR, hazard ratio; Low-res, low resolution (ie, antigen level); OS, overall survival; PBM, peptide-binding motif.

32.5% (95% CI, 31.7 to 33.3), 23.3% (95% CI, 22.6 to 24), and 40.7% (95% CI, 39.9 to 41.5) while those of grades 2 to 4 and grades 3 to 4 aGVHD at 4 months were 30.5% (95% CI, 29.8 to 31.2) and 11.5% (95% CI, 11 to 12), respectively.

TABLE 2. Multivariable Cox Proportional Hazards Models of HLA Associations in the Full Cohort

End Point	Subgroups	No. (events)	99% CI	<i>P</i>
OS	10 of 10	12,841 (5,143)	1	
	9 of 10	3,435 (1,603)	1.23 (1.14 to 1.33)	<.001
	8 of 10	473 (234)	1.29 (1.08 to 1.54)	<.001
	10 of 10	12,841 (5,143)	1	
	Class I mM	2,525 (1,218)	1.29 (1.19 to 1.41)	<.001
	Class II mM	910 (385)	1.07 (0.93 to 1.23)	.19
	10 of 10	12,841 (5,143)	1	
	Low-res ^a match	579 (257)	1.12 (0.95 to 1.33)	.08
	Low-res ^a mM	1,946 (961)	1.36 (1.24 to 1.49)	<.001
	10 of 10	12,841 (5,143)	1	
	PBM-GvH ^b match	776 (362)	1.16 (1 to 1.34)	.008
	PBM-GvH ^b mM	1,367 (679)	1.42 (1.28-1.59)	<.001
GRFS	10 of 10	11,992 (6,879)	1	
	9 of 10	3,131 (1,942)	1.17 (1.09 to 1.25)	<.001
	8 of 10	425 (261)	1.19 (1.01 to 1.4)	.007
	10 of 10	11,992 (6,879)	1	
	Class I mM	2,301 (1,466)	1.23 (1.14 to 1.33)	<.001
	Class II mM	830 (476)	1 (0.88 to 1.13)	.98
	10 of 10	11,992 (6,879)	1	
	Low-res ^a match	528 (315)	1.06 (0.91 to 1.24)	.30
	Low-res ^a mM	1,773 (1,151)	1.3 (1.19 to 1.41)	<.001
	10 of 10	11,992 (6,879)	1	
	PBM-GvH ^b match	711 (450)	1.15 (1.01 to 1.31)	.004
	PBM-GvH ^b mM	1,234 (799)	1.33 (1.2 to 1.46)	<.001

NOTE. Multivariable models were adjusted for patient and donor age, sex (mis)matching, CMV serostatus matching, comorbidity score (HCT-CI),³⁶ Karnofsky Performance Score, disease at transplant, disease risk index,²⁵ year of HCT, stem cell source, conditioning regimen and use of total-body irradiation, use of in vivo T-cell depletion, PTCy use, and HLA-DPB1 TCE matching status, and include a gamma-distributed frailty term for transplant center.

Abbreviations: GRFS, graft-versus-host disease, relapse-free survival; HCT, hematopoietic cell transplantation; HLA, human leukocyte antigen; HR, hazard ratio; mM, mismatch; OS, overall survival; PBM-GvH, peptide-binding motif graft-versus-host; TCE, T-cell epitope.

^aLow-resolution (Low-res).

^bPBM-GvH matching for HLA class I mM.

HLA Mismatching and HCT Outcome

We first examined the effect of the number of mismatches at any of the five main HLA loci on HCT outcome in the overall cohort. The presence of one or two mismatches significantly reduced the 5-year probability of OS from 52.2% (95% CI, 51.1 to 53.2) in the 10 of 10 pairs to 46.7% (95% CI, 44.8 to 48.6) and 45.3% (95% CI, 40.4 to 50.1) in the 9 of 10 and 8 of 10 pairs, respectively (*P* < .001; Fig 1). In multivariable analysis, one or two mismatches were associated with higher risk of mortality (both OS and GRFS) compared with full matches (Fig 1, Table 2). Similar significant associations were observed for RFS, NRM, and aGVHD II-IV and III-IV but not for relapse or cGVHD (Data Supplement, Table S1 and Fig S1). Additional covariates also significantly associated with OS included patient and donor age and disease- and performance status-related factors (Data Supplement, Table S2).

Next, we performed a more granular risk evaluation in 9 of 10 transplants according to HLA mismatch class, locus, or resolution. Five-year OS was significantly worse for single HLA class I but not HLA class II mismatches, both in univariable and multivariable analysis (Fig 1, Table 2). Of note, there was evidence for time dependency of these associations, apparently driven by strong nonproportionality in the HLA class II effect (Data Supplement, Fig S2). Among mismatched class I loci, HLA-A (hazard ratio [HR], 1.33 [95% CI, 1.18 to 1.50], *P* < .001) and HLA-B (HR, 1.43 [95% CI, 1.22 to 1.68], *P* < .001) conferred higher adjusted risks than HLA-C (HR, 1.17 [95% CI, 1.02 to 1.34], *P* = .003) compared with the 10 of 10 reference (Fig 1). Instead, the adjusted risks for HLA-DRB1 (HR, 1.03 [95% CI, 0.82 to 1.29], *P* = .71) and DQB1 (HR, 1.09 [95% CI, 0.92 to 1.29], *P* = .17) mismatches were similar and not significantly different from the 10 of 10 reference (Fig 1). HLA class I low-resolution mismatches (ie, at the first field or antigen level) were associated with worse OS compared with high-resolution mismatches (ie, at the second field or allele level), both in univariable and multivariable analysis (HR, 1.21 [95% CI, 1.01 to 1.46]; *P* = .007; Fig 1, Table 2). Class- and locus-specific associations with lower OS were mirrored by lower GRFS (Table 2), RFS, and higher NRM and aGVHD and cGVHD, whereas no significant associations were observed with relapse (Data Supplement, Table S1).

Models for Tolerable HLA Mismatches

In the 3,526 patients transplanted from 9 of 10 donors, we tested three models of tolerable HLA mismatches, that is,




FIG 2. HCT outcome in the full cohort according to the use of PTCy. Kaplan-Meier/cumulative incidence and forest plots show probabilities and adjusted (continued on following page)

FIG 2. (Continued). HR estimates and their 99% CI stratified according to the use of PTCy-based or conventional GVHD prophylaxis. (A, B) OS; (C, D) GRFS; (E, F) aGVHD III-IV; and (G, H) cGVHD. *P* values in the Kaplan-Meier/cumulative incidence plots correspond to the results of univariable comparisons in the overall cohort using log-rank or Gray tests, as appropriate. Individuals at risk at different time points post-transplant are shown underneath each plot (top row: transplants without PTCy; bottom row: transplants with PTCy). HR and *P* values in the forest plots are derived from multivariable analyses for the fully 10 of 10 matched pairs shown in Data Supplement, Table S5 with conventional GVHD prophylaxis as reference (vertical lines). GRFS, graft-versus-host disease, relapse-free survival; (a/c)GVHD, (acute/chronic) graft-versus-host disease. GVHD, graft-versus-host disease; HCT, hematopoietic cell transplantation; HR, hazard ratio; OS, overall survival; PBM, peptide-binding motif; PTCy, post-transplant cyclophosphamide.

immunopeptidome divergence between HLA-A, HLA-B, HLA-C or HLA-DPB1 mismatches as classified by PBM or TCE groups, respectively,^{16,19} and B-Leader mismatches.¹⁵ The 5-year probabilities of OS dropped from 52.2% (95% CI, 51.1 to 53.2) in the 10 of 10 reference to 46.7% (95% CI, 42.7 to 50.6) and 43.5% (95% CI, 40.5 to 46.4) in the HLA class I PBM-GVH matched and mismatched groups, respectively ($P < .001$, Fig 1). In multivariable analysis, the hazards of OS, NRM, and aGVHD II-IV were significantly worse both in HLA class I PBM-GVH matched and mismatched pairs compared with the 10 of 10 reference; however, this difference was more marked for HLA class I PBM-GVH mismatches (Fig 1, Table 2 and Data Supplement, Table S1). Of note, HLA class I PBM-GVH mismatched pairs had significantly worse OS than HLA class I PBM-GVH matched pairs in direct comparison (HR, 1.23 [95% CI, 1.04 to 1.46], $P = .002$). This was mirrored by significantly worse GRFS, RFS, aGVHD III-IV, and cGVHD compared with the 10 of 10 reference for HLA class I PBM-GVH mismatches but not for matches (Table 2 and Data Supplement, Table S1). Significant associations with nonpermissive but not permissive HLA-DPB1 TCE mismatches were observed in the 9 of 10 cohort only for aGVHD II-IV, whereas in the 10 of 10 group nonpermissive mismatches were associated with worse NRM and aGVHD III-IV (Data Supplement, Table S3). In the 10 of 10, the hazards of relapse were significantly reduced while those of aGVHD II-IV were significantly increased for both permissive and nonpermissive HLA-DPB1 mismatches. Single B-Leader matches were associated with significantly worse OS, GRFS, RFS and NRM compared with the 10 of 10 reference, whereas B-leader mismatches were significantly associated only with NRM (Data Supplement, Table S4). However, we did not observe significant differences between B-leader matched and mismatched pairs for any of the outcomes tested (HR, 0.89 [95% CI, 0.60 to 1.34], $P = .47$ for OS; not shown for the other end points).

Effects of PTCy on the Role of HLA Mismatching in HCT Outcome

To investigate the effects of PTCy on transplant outcome, we studied the 1,523 patients who received PTCy-based GVHD prophylaxis. The median follow-up time in this group of patients was 28.9 (95% CI, 27.1 to 31.1) months. Overall, the

use of PTCy improved the 5-year probability of OS from 50.5% (95% CI, 49.6 to 51.5) under conventional prophylaxis to 54.5% (95% CI, 50.6 to 58.2%; $P = .003$). This was mirrored by significantly better GRFS and reduced aGVHD and cGVHD compared with conventional prophylaxis (Fig 2). Multivariable models for the 10 of 10 matched transplants confirmed these findings for GRFS and GVHD, whereas no significant differences were observed for other end points (Fig 2 and Data Supplement, Table S5).

To better understand the impact of PTCy on the role of a mismatch at any of the five main HLA loci, we comparatively analyzed outcomes in patients that had received a 9 of 10 or a 10 of 10 matched transplant under conventional or PTCy-based GVHD prophylaxis. The effects of a single HLA mismatch were similar regardless of PTCy use (nonsignificant interaction; $P = .22$), conferring significantly increased risks of mortality (both OS and GRFS) in the presence and absence of PTCy (Fig 3, Table 3 and Data Supplement, Fig S3). Similar findings were obtained when excluding the 364 patients who had received *in vivo* TCD in addition to PTCy (Table 1) from the analysis (not shown). We did not find significant differences in the effects of HLA class I mismatches, low-resolution mismatches, and PBM-GVH mismatches between the groups that did or did not receive PTCy (nonsignificant interactions; $P \geq .42$; Fig 3, Table 3, and Data Supplement, Fig S3). Similar data were also found for the other end points studied (Data Supplement, Table S6). Risk associations between permissive or nonpermissive HLA-DPB1 TCE mismatches observed in 10 of 10 transplants under conventional GVHD prophylaxis appeared to be less pronounced when PTCy was used, although the relevant interaction tests were not significant (Data Supplement, Table S7).

DISCUSSION

HLA matching between donor and recipient has been considered a mainstay for the clinical success of unrelated HCT^{1,2,37} on the basis of compelling evidence from numerous independent retrospective studies.^{9,12,13,15} Consequently, the ideal unrelated donor should be fully matched at the allelic level for these loci and, if possible, carry a tolerable HLA-DPB1 mismatch according to the TCE model.^{16,18,27,28,38-40}




FIG 3. Survival after HCT according to HLA mismatching and GVHD prophylaxis. Kaplan-Meier and forest plots show overall survival (OS) probabilities and adjusted HR estimates and their 99% CI stratified according to (A, B) the presence (9 of 10) or absence (10 of 10) of a mismatch (mM) at HLA-A, HLA-B, HLA-C, HLA-DRB1, or HLA-DQB1, (C, D) the specific mismatched HLA class, (E, F) the typing (continued on following page)

FIG 3. (Continued). resolution for HLA class I mM, and (G, H) PBM GVH matching for HLA class I mM. The Kaplan-Meier plots and relevant *P* values correspond to the results of univariable comparisons in the subgroup of patients who received PTCy-based GVHD prophylaxis, using log-rank tests. Individuals at risk at different time points post-transplant are shown underneath each plot, according to the groups indicated in the legend from top to bottom. HR and 99% CI in the forest plots are derived from multivariable analyses shown in Table 3, separately for the conventional and PTCy-based GVHD prophylaxis, using the 10 of 10 pairs from each PTCy subgroup as reference. Interaction tests were nonsignificant in all cases ($P \geq .15$). The thick vertical lines correspond to the HR estimates for single (ie, 9 of 10), class I/class II, Low-res, and PBM-GVH matched/mismatched pairs in the full cohort. Low-res, low resolution (ie, antigen level). GVHD, graft-versus-host disease; HCT, hematopoietic cell transplantation; HLA, human leukocyte antigen; HR, hazard ratio; PBM, peptide-binding motif; PTCy, post-transplant cyclophosphamide.

Nevertheless, the studies at the basis for the current practice in donor selection originate mainly from the beginning of the century, and it is unknown if and how they have been affected by the developments in transplantation and histocompatibility since that time. In particular, high-accuracy next-generation sequencing–based HLA typing has been established in most tissue typing laboratories around the world within the past decade.^{5,41,42} This has led to a dramatic increase in the number of unrelated donors with high-quality HLA data in the worldwide registries, allowing for granular selection of well-matched donors for increasing numbers of patients.^{43–45} Perhaps the most important change in transplantation protocols over the past decade, regards the introduction of PTCy-based regimens for GVHD prophylaxis.^{6,22,46–48} Similar survival probabilities after HLA-matched HCT under conventional immune prophylaxis and after HLA-mismatched HCT under PTCy have suggested that its use can overcome the HLA barrier.^{21,49} However, this notion is at least partly challenged by observations that HLA mismatches are still relevant when their role is evaluated within the PTCy platforms^{50,51} and that tolerable mismatches can be identified also in the setting of haploidentical HCT under PTCy.^{52,53}

In this study, we reassessed the role of HLA mismatches in contemporary unrelated HCT, leveraging a data set of more than 17,000 transplants, including over 1,500 under PTCy-based GVHD prophylaxis, collected through the EBMT Registry mainly within the past decade. Our results show that HLA disparity remains an important factor determining mortality risks also to date, but that this role is driven by HLA-A, HLA-B, HLA-C and not by HLA-DRB1, HLA-DQB1 mismatches, a finding in line with a recent independent international study.¹⁵ Of note, the HLA class II effects were time-dependent, that is, an impact on OS could be appreciated in the first months after transplantation but not at longer follow-up, possibly mirroring a protective effect on cGVHD, although this was not accompanied by a lower incidence of relapse.

Toxicity by HLA class I mismatches resulted in increased severe aGVHD and NRM, without a concomitant decrease in relapse incidence. This underlines the importance of continuing endeavors to prevent and treat GVHD^{7,8} for improved

transplant outcomes. We also find that high-resolution HLA class I mismatches are less deleterious than low-resolution mismatches. This might at least partly mirror the advantageous effect we demonstrated for HLA class I PBM-GVH matches compared with mismatches, the former being more likely between alleles that are matched at the low-resolution level.²⁶ Again, toxicity from PBM-GVH mismatches resulted in higher risks of severe aGVHD and NRM, validating our previous observations made in an independent, mainly North American cohort.¹⁹ Together, these findings provide further compelling evidence for the role of immunopeptide divergence between mismatched HLA molecules as biomarker for risk stratification in allogeneic HCT and provide a means for intelligent mismatched unrelated donor selection to improve transplant outcome. This notion is also at the basis of the risk associations between nonpermissive HLA-DPB1 TCE mismatches and clinical outcome that have been repeatedly validated.^{17,54} These findings were confirmed in the 10 of 10 and extended to the 9 of 10 setting in this study.

Our data set provided the important opportunity to investigate the impact of PTCy on HCT outcome overall and on the role of HLA mismatches. PTCy improved GFRS compared with conventional GVHD prophylaxis as previously described^{23,24} and reduced severe acute and chronic GVHD without a significant increase in the hazard of relapse. There were no significant differences in the detrimental effects of mismatches at the five main HLA loci on outcome between conventional and PTCy-based regimens. In particular, although the numbers in each subgroup are still limited, our findings suggest that the negative effect of HLA class I disparity at the low-resolution and PBM-GVH level may apply also to transplantation under PTCy-based immune prophylaxis. Of note, PTCy effects seemed to be more pronounced for HLA-DPB1 mismatches. If confirmed, this might suggest different impacts of PTCy on alloreactive CD4+ compared with CD8+ T-cell responses, as previously suggested.^{55,56} Further investigations are warranted to test this intriguing hypothesis.

Our study has limitations, in particular regarding its retrospective nature and potential underlying sources of heterogeneity related to the use of registry data. It should be

TABLE 3. Multivariable Cox Proportional Hazards Models of HLA Associations by PTCy Use

End Point	Subgroups	PTCy No			PTCy Yes			Interaction <i>P</i> ^a
		No. (events)	HR (99% CI)	<i>P</i>	No. (events)	HR (99% CI)	<i>P</i>	
OS	10 of 10	11,943 (4,867)	1		898 (276)	1		.43
	9 of 10	2,924 (1,393)	1.22 (1.13 to 1.33)	<.001	511 (210)	1.32 (1.04 to 1.68)	.003	
	10 of 10	11,943 (4,867)	1		898 (276)	1		.42
	Class I mM	2,117 (1,050)	1.29 (1.18 to 1.41)	<.001	408 (168)	1.32 (1.02 to 1.71)	.006	
	Class II mM	807 (343)	1.05 (0.91 to 1.21)	.39	103 (42)	1.32 (0.86 to 2.04)	.09	
	10 of 10	11,943 (4,867)	1		898 (276)	1		.69
	Low-res ^b match	515 (230)	1.1 (0.92 to 1.32)	.15	64 (27)	1.31 (0.77 to 2.2)	.19	
	Low-res ^b mM	1,602 (820)	1.36 (1.23 to 1.51)	<.001	344 (141)	1.33 (1.01 to 1.75)	.007	
	10 of 10	11,943 (4,867)	1		898 (276)	1		.75
	PBM-GVH ^c match	648 (310)	1.14 (0.98 to 1.33)	.03	128 (52)	1.28 (0.86 to 1.9)	.11	
	PBM-GVH ^c mM	1,153 (586)	1.42 (1.26 to 1.59)	<.001	214 (93)	1.5 (1.09 to 2.06)	<.001	
GRFS	10 of 10	11,147 (6,472)	1		845 (407)	1		.72
	9 of 10	2,645 (1,662)	1.16 (1.08 to 1.25)	<.001	486 (280)	1.2 (0.97 to 1.47)	.02	
	10 of 10	11,147 (6,472)	1		845 (407)	1		.74
	Class I mM	1,912 (1,240)	1.24 (1.14 to 1.34)	<.001	389 (226)	1.22 (0.98 to 1.52)	.02	
	Class II mM	733 (422)	0.99 (0.87 to 1.13)	.79	97 (54)	1.1 (0.76 to 1.61)	.50	
	10 of 10	11,147 (6,472)	1		845 (407)	1		.36
	Low-res ^b match	469 (287)	1.09 (0.93 to 1.28)	.14	59 (28)	0.82 (0.5 to 1.36)	.32	
	Low-res ^b mM	1,443 (953)	1.29 (1.18 to 1.42)	<.001	330 (198)	1.31 (1.04 to 1.65)	.002	
	10 of 10	11,147 (6,472)	1		845 (407)	1		.90
	PBM-GVH ^c match	589 (381)	1.15 (1 to 1.32)	.008	122 (69)	1.16 (0.83 to 1.63)	.26	
	PBM-GVH ^c mM	1,032 (676)	1.32 (1.18 to 1.47)	<.001	202 (123)	1.39 (1.06 to 1.82)	.002	

NOTE. Multivariable models were adjusted for patient and donor age, sex (mis)matching, CMV serostatus matching, comorbidity score (HCT-CI),³⁶ Karnofsky Performance Score, disease at transplant, disease risk index,²⁵ year of HCT, stem cell source, conditioning regimen and use of total-body irradiation, use of in vivo T-cell depletion, and HLA-DPB1 TCE matching status and include a gamma-distributed frailty term for transplant center. Abbreviations: GRFS, graft-versus-host disease, relapse-free survival; HCT, hematopoietic cell transplantation; HLA, human leukocyte antigen; HR, hazard ratio; mM, mismatch; N, number; PBM, peptide-binding motif; OS, overall survival; PTCy, post-transplantation cyclophosphamide; TCE, T-cell epitope.

^aInteraction analyses between the respective matching models and the use of PTCy.

^bLow-resolution (Low-res).

^cPBM matching for class I mM.

emphasized that, although, to our knowledge, our data set contains one of the largest numbers of unrelated transplants under PTCy-based GVHD prophylaxis with 6-locus HLA typing for patients and donors pairs analyzed so far, its size is still limited to draw any definitive conclusions on the role of HLA disparity in this setting. Compared with other recent studies from international registries,^{57,58} ours has the distinct feature of including both PTCy and non-PTCy transplants, thereby enabling us to investigate the statistical interaction between HLA matching status and GVHD prophylaxis. The lack of evidence for such interaction in our data set suggests that optimal HLA matching should still be considered in the selection of stem cell donors for HCT also under PTCy-based GVHD prophylaxis. Nevertheless, the relative role of HLA compared with other relevant clinical variables such as donor

age could not be addressed because of the limited size of the PTCy cohort and are hence warranted in future studies. Moreover, these investigations should also leverage the results from ongoing and future prospective clinical trials on HLA-matched and mismatched PTCy-based unrelated HCT.⁵⁹⁻⁶¹

Taken together, our findings suggest that HLA class I, in particular at the antigen and PBM level, should be prioritized over class II matching in contemporary unrelated donor selection and that these recommendations still hold also for transplants performed under PTCy-based GVHD prophylaxis. However, compiling larger, preferentially multiregistry data and evidence from prospective clinical trials is highly needed to better understand the role of HLA in the era of PTCy.

AFFILIATIONS

¹Institute for Experimental Cellular Therapy, University Hospital Essen, Essen, Germany

²German Cancer Consortium (DKTK), Partner Site Essen/Düsseldorf, Essen, Germany

³Department of Biomedical Data Sciences, LUMC, Leiden, the Netherlands

⁴EBMT Leiden Study Unit, Leiden, the Netherlands

⁵Saint-Louis Hospital, BMT Unit, Paris, France

⁶Oslo University Hospital, Rikshospitalet, Oslo, Norway

⁷HUCH Comprehensive Cancer Center, Helsinki, Finland

⁸University Medical Center Hamburg, Hamburg, Germany

⁹CHU de Lille, Université de Lille, INSERM U1286, Lille, France

¹⁰Addenbrookes Hospital, Cambridge, United Kingdom

¹¹University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

¹²Erasmus MC Cancer Institute, Rotterdam, the Netherlands

¹³CHU Bordeaux, Hôpital Haut-Leveque, Pessac, France

¹⁴CHU Grenoble Alpes - Université Grenoble Alpes, Grenoble, France

¹⁵Cliniques Universitaires St Luc, Brussels, Belgium

¹⁶CHU - Institut Universitaire du Cancer Toulouse, Toulouse, France

¹⁷Skane University Hospital, Lund, Sweden

¹⁸IRCCS Ospedale San Raffaele, Milan, Italy

¹⁹University Vita-Salute San Raffaele, Milan, Italy

²⁰Manchester Royal Infirmary, Manchester, United Kingdom

²¹Department of Hematology and Stem Cell Transplantation, University Hospital Essen, Essen, Germany

²²Hôpital Jean Minjot, Besançon, France

²³University Hospital Uppsala, Uppsala, Sweden

²⁴Sorbonne Université, Hôpital Saint-Antoine, AP-HP, Paris, France

CORRESPONDING AUTHOR

Katharina Fleischhauer, MD; Twitter: @IztE; e-mail: katharina.fleischhauer@uk-essen.de.

EQUAL CONTRIBUTION

A.R. and K.F. shared senior authorship.

REFERENCES

- Copelan EA: Hematopoietic stem-cell transplantation. *N Engl J Med* 354:1813-1826, 2006
- Singh AK, McGuirk JP: Allogeneic stem cell transplantation: A historical and scientific overview. *Cancer Res* 76:6445-6451, 2016
- Cieri N, Maurer K, Wu CJ: 60 years young: The evolving role of allogeneic hematopoietic stem cell transplantation in cancer immunotherapy. *Cancer Res* 81:4373-4384, 2021
- Gooley TA, Chien JW, Pergam SA, et al: Reduced mortality after allogeneic hematopoietic-cell transplantation. *N Engl J Med* 363:2091-2101, 2010
- Lange V, Bohme I, Hofmann J, et al: Cost-efficient high-throughput HLA typing by MiSeq amplicon sequencing. *BMC Genomics* 15:63, 2014
- Rimando J, McCurdy SR, Luznik L: How I prevent GVHD in high-risk patients: Posttransplant cyclophosphamide and beyond. *Blood* 141:49-59, 2023
- Jamy O, Zeiser R, Chen YB: Novel developments in the prophylaxis and treatment of acute GVHD. *Blood* 142:1037-1046, 2023
- Zeiser R: Novel approaches to the treatment of chronic graft-versus-host disease. *J Clin Oncol* 41:1820-1824, 2023
- Lee SJ, Klein J, Haagenson M, et al: High-resolution donor-recipient HLA matching contributes to the success of unrelated donor marrow transplantation. *Blood* 110:4576-4583, 2007
- Bolon YTAR, Allbee-Johnson M, Estrada-Merly N, et al: Current use and outcome of hematopoietic stem cell transplantation: CIBMTR summary slides, 2023.
- Flomenberg N, Baxter-Lowe LA, Confer D, et al: Impact of HLA class I and class II high-resolution matching on outcomes of unrelated donor bone marrow transplantation: HLA-C mismatching is associated with a strong adverse effect on transplantation outcome. *Blood* 104:1923-1930, 2004
- Ayuk F, Beelen DW, Bornhauser M, et al: Relative impact of HLA matching and non-HLA donor characteristics on outcomes of allogeneic stem cell transplantation for acute myeloid leukemia and myelodysplastic syndrome. *Biol Blood Marrow Transplant* 24:2558-2567, 2018
- Furst D, Muller C, Vucinic V, et al: High-resolution HLA matching in hematopoietic stem cell transplantation: A retrospective collaborative analysis. *Blood* 122:3220-3229, 2013
- Petersdorf EW, Stevenson P, Bengtsson M, et al: HLA-B leader and survivorship after HLA-mismatched unrelated donor transplantation. *Blood* 136:362-369, 2020
- Petersdorf EW, Carrington M, O'Huigin C, et al: Role of HLA-B exon 1 in graft-versus-host disease after unrelated haemopoietic cell transplantation: A retrospective cohort study. *Lancet Haematol* 7:e50-e60, 2020
- Fleischhauer K, Shaw BE, Gooley T, et al: Effect of T-cell-epitope matching at HLA-DPB1 in recipients of unrelated-donor haemopoietic-cell transplantation: A retrospective study. *Lancet Oncol* 13:366-374, 2012
- Meurer T, Crivello P, Metzger M, et al: Permissive HLA-DPB1 mismatches in HCT depend on immunopeptidome divergence and editing by HLA-DM. *Blood* 137:923-928, 2021
- Arrieta-Bolaños E, Crivello P, He M, et al: A core group of structurally similar HLA-DPB1 alleles drives permissiveness after hematopoietic cell transplantation. *Blood* 140:659-663, 2022
- Crivello P, Arrieta-Bolaños E, He M, et al: Impact of the HLA immunopeptidome on survival of leukemia patients after unrelated donor transplantation. *J Clin Oncol* 41:2416-2427, 2023
- Kanaky CG, Coffey DG, Towler AMH, et al: Origin and evolution of the T cell repertoire after posttransplantation cyclophosphamide. *JCI Insight* 1:e86252, 2016
- Bashy A, Zhang X, Sizemore CA, et al: T-cell-replete HLA-haploidentical hematopoietic transplantation for hematologic malignancies using post-transplantation cyclophosphamide results in outcomes equivalent to those of contemporaneous HLA-matched related and unrelated donor transplantation. *J Clin Oncol* 31:1310-1316, 2013
- O'Donnell PV, Jones RJ: The development of post-transplant cyclophosphamide: Half a century of translational team science. *Blood Rev* 62:101034, 2023

SUPPORT

Supported by grants from the Deutsche Knochenmarkspenderdatei to EAB (DKMS-SLS-JHRG-2021-02), and from the Deutsche José Carreras Leukämie Stiftung (DJCLS 17R/2023), the Dr Werner Jackstädt Stiftung and the Joseph Senker Stiftung to KF. The authors would like to thank the staff at the participating transplant centers collecting and contributing clinical data for this study.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO.24.00582>.

AUTHOR CONTRIBUTIONS

Conception and design: Esteban Arrieta-Bolaños, Edouard F. Bonneville, Liesbeth C. de Wreede, Florent Malard, Annalisa Ruggeri

Administrative support: Jorinde D. Hoogenboom

Provision of study materials or patients: Marie Robin, Tobias Gedde-Dahl, Urvu Salmenniemi, Nicolaus Kröger, Ibrahim Yakoub-Agha, Charles Crawley, Goda Choi, Annoek E.C. Broers, Edouard Forcade, Martin Carre, Xavier Poiré, Anne Huynh, Stig Lenhoff, Fabio Ciceri, Eleni Tholouli, Thomas Schroeder, Eric Deconinck, Krizstina Carlson

Collection and assembly of data: Esteban Arrieta-Bolaños, Marie Robin, Tobias Gedde-Dahl, Urvu Salmenniemi, Nicolaus Kröger, Ibrahim Yakoub-Agha, Charles Crawley, Goda Choi, Annoek E.C. Broers, Edouard Forcade, Martin Carre, Xavier Poiré, Anne Huynh, Stig Lenhoff, Fabio Ciceri, Eleni Tholouli, Thomas Schroeder, Eric Deconinck, Kristina Carlson, Liesbeth C. de Wreede, Jorinde D. Hoogenboom, Annalisa Ruggeri, Katharina Fleischhauer

Data analysis and interpretation: Esteban Arrieta-Bolaños, Edouard F. Bonneville, Pietro Crivello, Liesbeth C. de Wreede, Annalisa Ruggeri, Katharina Fleischhauer

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

23. Bolanos-Meade J, Reshef R, Fraser R, et al: Three prophylaxis regimens (tacrolimus, mycophenolate mofetil, and cyclophosphamide; tacrolimus, methotrexate, and bortezomib; or tacrolimus, methotrexate, and maraviroc) versus tacrolimus and methotrexate for prevention of graft-versus-host disease with haematopoietic cell transplantation with reduced-intensity conditioning: A randomised phase 2 trial with a non-randomised contemporaneous control group (BMT CTN 1203). *Lancet Haematol* 6:e132-e143, 2019
24. Bolanos-Meade J, Hamadani M, Wu J, et al: Post-transplantation cyclophosphamide-based graft-versus-host disease prophylaxis. *N Engl J Med* 388:2338-2348, 2023
25. Saccardi R, Putter H, Eikema DJ, et al: Benchmarking of survival outcomes following haematopoietic stem cell transplantation (HSCT): An update of the ongoing project of the European Society for Blood and Marrow Transplantation (EBMT) and Joint Accreditation Committee of ISCT and EBMT (JACIE). *Bone Marrow Transplant* 58:659-666, 2023
26. Nunes E, Heslop H, Fernandez-Vina M, et al: Definitions of histocompatibility typing terms. *Blood* 118:e180-e183, 2011
27. Zino E, Frumento G, Marktel S, et al: A T-cell epitope encoded by a subset of HLA-DPB1 alleles determines nonpermissive mismatches for hematologic stem cell transplantation. *Blood* 103:1417-1424, 2004
28. Arrieta-Bolanos E, Crivello P, Shaw BE, et al: In silico prediction of nonpermissive HLA-DPB1 mismatches in unrelated HCT by functional distance. *Blood Adv* 2:1773-1783, 2018
29. Fleischhauer K, Ahn KW, Wang HL, et al: Directionality of non-permissive HLA-DPB1 T-cell epitope group mismatches does not improve clinical risk stratification in 8/8 matched unrelated donor hematopoietic cell transplantation. *Bone Marrow Transplant* 52:1280-1287, 2017
30. Crivello P, Zito L, Sizzano F, et al: The impact of amino acid variability on alloreactivity defines a functional distance predictive of permissive HLA-DPB1 mismatches in hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant* 21:233-241, 2015
31. Ruggeri A, Labopin M, Ciceri F, et al: Definition of GVHD-free, relapse-free survival for registry-based studies: An ALWP-EBMT analysis on patients with AML in remission. *Bone Marrow Transplant* 51:610-611, 2016
32. Balan TA, Putter H: A tutorial on frailty models. *Stat Methods Med Res* 29:3424-3454, 2020
33. Bonneville EF, Schetelig J, Putter H, et al: Handling missing covariate data in clinical studies in haematology. *Best Pract Res Clin Haematol* 36:101477, 2023
34. Blake HA, Leyrat C, Mansfield KE, et al: Estimating treatment effects with partially observed covariates using outcome regression with missing indicators. *Biom J* 62:428-443, 2020
35. R_Core_Team: R: A Language and Environment for Statistical Computing. Vienna, Austria, R Foundation for Statistical Computing, 2023. <https://www.R-project.org/>
36. Sorror M, Storer B, Sandmaier BM, et al: Hematopoietic cell transplantation-comorbidity index and Karnofsky performance status are independent predictors of morbidity and mortality after allogeneic nonmyeloablative hematopoietic cell transplantation. *Cancer* 112:1992-2001, 2008
37. Petersdorf EW: Optimal HLA matching in hematopoietic cell transplantation. *Curr Opin Immunol* 20:588-593, 2008
38. Crocchiolo R, Zino E, Vago L, et al: Nonpermissive HLA-DPB1 disparity is a significant independent risk factor for mortality after unrelated hematopoietic stem cell transplantation. *Blood* 114:1437-1444, 2009
39. Pidala J, Lee SJ, Ahn KW, et al: Nonpermissive HLA-DPB1 mismatch increases mortality after myeloablative unrelated allogeneic hematopoietic cell transplantation. *Blood* 124:2596-2606, 2014
40. Oran B, Saliba RM, Carmazzi Y, et al: Effect of nonpermissive HLA-DPB1 mismatches after unrelated allogeneic transplantation with in vivo T-cell depletion. *Blood* 131:1248-1257, 2018
41. Gabriel C, Furst D, Fae I, et al: HLA typing by next-generation sequencing - getting closer to reality. *Tissue Antigens* 83:65-75, 2014
42. Mayor NP, Wang T, Lee SJ, et al: Impact of previously unrecognized HLA mismatches using ultrahigh resolution typing in unrelated donor hematopoietic cell transplantation. *J Clin Oncol* 39:2397-2409, 2021
43. Gragert L, Spellman SR, Shaw BE, et al: Unrelated stem cell donor HLA match likelihood in the US registry incorporating HLA-DPB1 permissive mismatching. *Transplant Cell Ther* 29:244-252, 2023
44. Auletta JJ, Kou J, Chen M, et al: Real-world data showing trends and outcomes by race and ethnicity in allogeneic hematopoietic cell transplantation: A report from the center for international blood and marrow transplant Research. *Transplant Cell Ther* 29:346.e1-346.e10, 2023
45. Fleischhauer K: Improved accessibility of optimally matched stem cell donors. *Transplant Cell Ther* 29:213-214, 2023
46. Kanakry CG, Fuchs EJ, Luznik L: Modern approaches to HLA-haploidentical blood or marrow transplantation. *Nat Rev Clin Oncol* 13:10-24, 2016
47. Nunes NS, Kanakry CG: Mechanisms of graft-versus-host disease prevention by post-transplantation cyclophosphamide: An evolving understanding. *Front Immunol* 10:2668, 2019
48. Luznik L, O'Donnell PV, Symons HJ, et al: HLA-haploidentical bone marrow transplantation for hematologic malignancies using nonmyeloablative conditioning and high-dose, posttransplantation cyclophosphamide. *Biol Blood Marrow Transplant* 14:641-650, 2008
49. Rashidi A, Hamadani M, Zhang MJ, et al: Outcomes of haploidentical vs matched sibling transplantation for acute myeloid leukemia in first complete remission. *Blood Adv* 3:1826-1836, 2019
50. Gooptu M, Romee R, St Martin A, et al: HLA-haploidentical vs matched unrelated donor transplants with posttransplant cyclophosphamide-based prophylaxis. *Blood* 138:273-282, 2021
51. Bacigalupo A: HLA still matters in allogeneic transplants. *Blood* 138:209-211, 2021
52. Fuchs EJ, McCurdy SR, Solomon SR, et al: HLA informs risk predictions after haploidentical stem cell transplantation with posttransplantation cyclophosphamide. *Blood* 139:1452-1468, 2022
53. Fleischhauer K: Haplo-PtCy: Adjusting the HLA barrier. *Blood* 139:1431-1433, 2022
54. Fleischhauer K, Shaw BE: HLA-DP in unrelated hematopoietic cell transplantation revisited: Challenges and opportunities. *Blood* 130:1089-1096, 2017
55. Hadjis AD, Nunes NS, Khan SM, et al: Post-transplantation cyclophosphamide uniquely restrains alloreactive CD4(+) T-cell proliferation and differentiation after murine MHC-haploidentical hematopoietic cell transplantation. *Front Immunol* 13:796349, 2022
56. McCurdy SR, Radojicic V, Tsai HL, et al: Signatures of GVHD and relapse after posttransplant cyclophosphamide revealed by immune profiling and machine learning. *Blood* 139:608-623, 2022
57. Sanz J, Labopin M, Choi G, et al: Younger unrelated donors may be preferable over HLA match in the PTCy era: A study from the ALWP of the EBMT. *Blood* 143:2534-2543, 2024
58. Auletta JJ, Al Malki MM, DeFor T, et al: Post-transplant cyclophosphamide eliminates disparity in GVHD-free, relapse-free survival and overall survival between 8/8 matched and 7/8 mismatched unrelated donor hematopoietic cell transplantation in adults with hematologic malignancies. *Blood* 142:780, 2023
59. Shaw BE, Jimenez-Jimenez AM, Burns LJ, et al: National marrow donor program-sponsored multicenter, phase II trial of HLA-mismatched unrelated donor bone marrow transplantation using post-transplant cyclophosphamide. *J Clin Oncol* 39:1971-1982, 2021
60. Shaw BE, Jimenez-Jimenez AM, Burns LJ, et al: Three-year outcomes in recipients of mismatched unrelated bone marrow donor transplants using post-transplantation cyclophosphamide: Follow-up from a National marrow donor Program-Sponsored prospective clinical trial. *Transplant Cell Ther* 29:208.e1-e208.e6, 2023
61. National Institutes of Health: HLA-mismatched unrelated donor hematopoietic cell transplantation with post-transplantation cyclophosphamide (ACCESS). <https://classic.clinicaltrials.gov/ct2/show/NCT04904588>

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Human Leukocyte Antigen Mismatching and Survival in Contemporary Hematopoietic Cell Transplantation for Hematologic Malignancies

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/authors/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](http://OpenPayments)).

Marie Robin

Research Funding: AbbVie (Inst), Medac (Inst)

Travel, Accommodations, Expenses: Medac

Urp Salmenniemi

Consulting or Advisory Role: Mylan

Nicolaus Kröger

Honoraria: Novartis, Celgene (Inst), Sanofi, Jazz Pharmaceuticals (Inst), Kite/Gilead, RIEMSER (Inst), AOP Orphan Pharmaceuticals, BMS GmbH & Co. KG, Neovii, Alexion Pharmaceuticals, Takeda, Pierre Fabre

Consulting or Advisory Role: Neovii, Sanofi, Jazz Pharmaceuticals, Novartis, Celgene, RIEMSER, Gilead Sciences

Speakers' Bureau: AOP Orphan Pharmaceuticals

Research Funding: Neovii (Inst), Novartis (Inst), Celgene (Inst), RIEMSER (Inst)

Travel, Accommodations, Expenses: Neovii, Novartis, Gilead Sciences, Jazz Pharmaceuticals, Sanofi, Celgene

Ibrahim Yakoub-Agha

Honoraria: Celgene, Novartis, Gilead Sciences, Biotest, Janssen, Jazz Pharmaceuticals

Travel, Accommodations, Expenses: Gilead Sciences

Charles Crawley

Travel, Accommodations, Expenses: Funding for attending ASH meeting in Dec 2019

Edouard Forcade

Consulting or Advisory Role: MaaT Pharma, Novartis

Speakers' Bureau: Jazz Pharmaceuticals, Kite/Gilead, Alexion Pharmaceuticals, Novartis, SOBI, GlaxoSmithKline

Travel, Accommodations, Expenses: Alexion Pharmaceuticals, Novartis, Kite/Gilead

Xavier Poiré

Consulting or Advisory Role: Takeda (Inst), Pfizer (Inst)

Travel, Accommodations, Expenses: Gilead Sciences (Inst), Takeda (Inst), Servier (Inst)

Eleni Tholouli

Consulting or Advisory Role: Gilead Sciences, Jazz Pharmaceuticals, Autolus Therapeutics, Vertex, Bristol Myers Squibb/Celgene, Janssen, Takeda

Speakers' Bureau: Kite/Gilead, Jazz Pharmaceuticals, Pfizer, Vertex, Takeda

Travel, Accommodations, Expenses: Jazz Pharmaceuticals, Menarini, Autolus, Kite/Gilead

Eric Deconinck

Consulting or Advisory Role: Stemline Menarini

Research Funding: Immunogen (Inst), Roche (Inst), Novartis/Pfizer (Inst)

Travel, Accommodations, Expenses: Stemline Therapeutics, Immunogen, Novartis/Pfizer, Janssen-Cilag, Gilead Sciences

Kristina Carlson

Stock and Other Ownership Interests: PFM Medical, AstraZeneca

Florent Malard

Honoraria: Mallinckrodt/Therakos, Janssen, Sanofi, Jazz Pharmaceuticals, Astellas Pharma, Biocodex, Gilead Sciences, Novartis, Celgene/Bristol Myers Squibb, AstraZeneca, MSD

Travel, Accommodations, Expenses: Pfizer, Takeda, Sanofi

Katharina Fleischhauer

Patents, Royalties, Other Intellectual Property: Royalties for a commonly developed commercial product from Genome Diagnostics. Royalties are paid to my Institution and used for Research Funding purposes (Inst)

No other potential conflicts of interest were reported.