



Abstract #14916

Abstract Information

Abstract ID:

14916

Abstract Title:**Evolution of Fe oxides crystallinity in permafrost deposits from mid-Pleistocene to Holocene: implications for mineral organic carbon interactions****Authors (presenting author in bold):****Maxime Thomas**¹, Loeka L Jongejans^{2,3}, Jens Strauss², Chloé Vermeylen¹, Sacha Calcus¹, Thomas Opel² and Sophie Opfergelt⁴, (1)UCLouvain, (2)Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, (3)University of Potsdam, (4)UCLouvain, Earth and Life Institute**Abstract Text:**

Mineral-organic carbon (OC) interactions are involved in the geochemical stability of OC and thus in the susceptibility of permafrost to release greenhouse gases. Those so-called stabilizing or protecting interactions - accounting for ~30 % to ~80 % of permafrost soil OC - includes organo-mineral associations, such as OC sorbed onto Fe-oxides, or organo-metallic complexes. Over timescales of soil development (millennia), the capacity of soils to stabilize OC is linked to soil development through changes in soil mineralogy. Specifically, weathering products such as short-range order minerals (e.g., poorly crystalline Fe-oxides) have an extensive surface area to bind OC. However, over time, these minerals evolve towards more crystalline phases with a lower surface area available to bind OC. Freezing conditions are considered to minimize changes in Fe oxides crystallinity at short time scale, but can we consider the mineralogy of Fe oxides in a frozen deposit as stable over millennial timescale? We investigate this question along a sequence of permafrost deposits from the headwall of the Batagay megaslump, Siberia, comprising sediment up to ~650 ka old. We analyzed the proportion of Fe as poorly crystalline and crystalline Fe oxides and organo-metallic complexes, and the proportion of total OC pool forming mineral-OC interactions in the different stratigraphic units. Our data show that: (i) the proportion of iron as poorly crystalline iron oxides significantly drops with increasing age of the deposit, from $28 \pm 14\%$ for Holocene deposits to $6 \pm 2\%$ for mid-Pleistocene deposits; (ii) the proportion of iron as crystalline oxides increases from $15 \pm 20\%$ to $34 \pm 2\%$ for the same deposits, respectively; (iii) the proportion mineral-bound OC relative to the total decreases over time from $45 \pm 13\%$ to $32 \pm 6\%$. These findings highlight that the mineral surfaces available for OC stabilization can evolve over millennial timescales in a frozen deposit. This raises the need to better constrain mineral OC interactions for older OC exposed by abrupt thawing of permafrost such as in megaslumps.

Final Paper##:

7

Keywords:

Fe-oxides, mineral organic carbon interactions and permafrost

Submitter's E-mail Address:

maxime.thomas@uclouvain.be

Submitter Full Name:

Maxime Thomas

Preferred Presentation Style:

Oral

Conference Attendance:

In person

Registration:

I understand that the presenting author named in this submission must register for the conference by 31 May 2023.

Submission Permission and Copyright:

I grant non-exclusive copyright for my abstract and any related uploaded content to the Conference Exchange and the conference owners. I give permission for my details to be stored in the conference database. I, and the presenting author (if different than the submitter), agree to be contacted by the conference organizers about this submission. I agree to the recording and online streaming of any presentations scheduled as a result of this submission. I confirm all listed authors have agreed to these terms.

First Author

Presenting Author

Maxime Thomas

Email: maxime.thomas@uclouvain.be

UCLouvain

Earth and Life Institute

PhD student & teaching assistant

Croix du Sud 1, bte L7.05.10

Boltzmann building floor 0

Ottignies-Louvain-la-Neuve 1348

Belgium

Second Author

Loeka L Jongejans

Email: loeka.jongejans@live.nl

Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research

Permafrost Research Section

Potsdam

Germany

University of Potsdam

Institute of Geosciences

Potsdam

Germany

Third Author

Jens Strauss

Email: jens.strauss@awi.de

Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research

Permafrost Research Section

Potsdam

Germany

Fourth Author

Chloé Vermynen

Email: chloe.vermylen@student.uclouvain.be

UCLouvain

Earth and Life Institute

Ottignies-Louvain-la-Neuve 1348

Belgium

Fifth Author

Sacha Calcus

Email: sacha.calcus@student.uclouvain.be

UCLouvain

Earth and Life Institute

Ottignies-Louvain-la-Neuve

Belgium

Sixth Author

Thomas Opel

Email: thomas.opel@awi.de

Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research

Polar Terrestrial Environments Section

Potsdam

Germany

Seventh Author

Sophie Opfergelt

Email: sophie.opfergelt@uclouvain.be

UCLouvain, Earth and Life Institute

Earth and Life Institute

Croix du Sud, 1

Louvain-la-Neuve 1348

Belgium