

PROBABILITY EQUIVALENT LEVEL FOR COVAR AND VAR

Patricia Ortega-Jiménez, Franco Pellerrey,
Miguel Sordo, Alfonso Suárez-Llorens

REPRINT | 2024 / 17


ISBA

Voie du Roman Pays 20 - L1.04.01

B-1348 Louvain-la-Neuve

Email : lidam-library@uclouvain.be

<https://uclouvain.be/en/research-institutes/lidam/isba/publication.html>



Probability equivalent level for CoVaR and VaR

Patricia Ortega-Jiménez^{a,b,*}, Franco Pellerey^c, Miguel A. Sordo^b, Alfonso Suárez-Llorens^b

^a Institute of Statistics, Biostatistics and Actuarial Science-ISBA, Louvain Institute of Data Analysis and Modeling-LIDAM, UC Louvain, Louvain-la-Neuve, Belgium

^b Departamento de Estadística e Investigación Operativa, Facultad de Ciencias, Universidad de Cádiz, Spain

^c Dipartimento di Scienze Matematiche, Politecnico di Torino, Italy

ARTICLE INFO

JEL classification:
G220

Keywords:
Systematic risk
Contagion risk measure
Value at risk
Conditional value at risk
Stochastically increasing

ABSTRACT

For a given risk, the well-known classical definition of Value-at-Risk (VaR) does not take into account possible interactions with other observable risks. For this reason, conditional VaRs that capture contagion effects and tail dependence among risks, such as the Co-Value-at-Risk (CoVaR), have been defined and studied in recent literature. In this paper we study conditions that guarantee, in the bivariate setting, the ordering between VaR and CoVaR, allowing to understand which, among the two measures, is more or less conservative than the other. By doing this, we introduce the notion of Probability Equivalent Level of CoVaR-VaR (PELCoV), which is the VaR value of the observable variable for which VaR and CoVaR coincide, and we study some of its properties such as uniqueness and boundedness. In particular, we show that its properties are entirely explained by the copula that describes the dependence between risks, and we provide a list of copulas for which PELCoV is explicitly available, and for which it is or not bounded. A practical applicative example is also presented.

1. Introduction

The study of risk measures to determine capital requirements, the amount of cash that a company should hold in reserve to cover unforeseen losses, has generated considerable attention in the economic and actuarial literature. Capital requirements can be calculated using a value-at-risk approach. Since the Basel Committee on Banking Supervision (BCBS) adopted it in the nineties of the previous century, the value-at-risk (VaR) has become the predominant measure of risk and has been employed in regulatory frameworks of banking (Basel III/IV) and insurance (Solvency II). Representing “the worst expected loss over a given horizon under normal market conditions at a given level of confidence”, the popularity of VaR is mainly due to its simplicity and ease of computation. The interested reader may consult Artzner et al. (1999), Jorion (2000), Kupiec (2002) and Denuit et al. (2005) for overviews on the topic.

Formally, given a random risk Y with distribution function F_Y the value-at-risk (VaR) of Y at level $v \in (0, 1)$, denoted by $\text{VaR}_v[Y]$, is just the inverse (or quantile function) of F_Y at v , given by

$$\text{VaR}_v[Y] = F_Y^{-1}(v) = \inf \{x : F_Y(x) \geq v\}.$$

However, the value-at-risk approach is focused on the individual risk of an institution and does not capture the systemic risk, which is the risk of failures caused by interactions with other financial institutions (see Bisias et al. (2012) and Benoit et al. (2017) for general surveys on systemic risk). As reported, among others, by Drehmann and Tarsheev (2011), Acharya et al. (2017) and Brownlees and Engle (2017), the strong interconnection within the financial system has contributed, especially since the 2007–2009 financial crisis, to the spreading of systemic risk, with adverse consequences in terms of credit supply to the real economy.

One important source of systemic risk is the risk of financial contagion, which occurs when losses in one financial institution spillover to another institution that is linked with the first one (see Glasserman and Young (2016)). As a result, since the first version of the paper by Adrian and Brunnermeier (2016), an active area of research has been devoted to the quantification, management and comparison of contagion risk measures. See, for example, Chen et al. (2013), Girardi and Ergün (2013), Mainik and Schaanning (2014), Sordo et al. (2015), Acharya et al. (2017), Sordo et al. (2018), Das and Fasen-Hartmann (2018a), Ortega-Jiménez et al. (2021) and Dhaene et al. (2022). In particular, Adrian and Brunnermeier (2016) considered a dependence-adjusted version of the VaR called CoVaR that measures the institution

* Corresponding author.

E-mail addresses: patricia.ortega@uclouvain.be (P. Ortega-Jiménez), franco.pellerey@polito.it (F. Pellerey), mangel.sordo@uca.es (M.A. Sordo), alfonso.suarez@uca.es (A. Suárez-Llorens).

<https://doi.org/10.1016/j.insmatheco.2023.12.004>

Received November 2022; Received in revised form December 2023; Accepted 21 December 2023

Available online 10 January 2024

0167-6687/© 2024 Elsevier B.V. All rights reserved.

risk, conditional on other institutions being in distress. Formally, the co-value-at-risk (CoVaR) of Y at level $v \in (0, 1)$, given that X is at level $u \in (0, 1)$, denoted by $\text{CoVaR}_{v,u}[Y|X]$, is defined by the VaR of the conditional variable $[Y|X = \text{VaR}_u[X]]$ at level risk v as follows¹

$$\text{CoVaR}_{v,u}[Y|X] = \text{VaR}_v[Y|X = \text{VaR}_u[X]] = F_{Y|X=\text{VaR}_u[X]}^{-1}(v).$$

Standard financial regulations determine capital requirements based on isolated institutions without considering the risk of systemic institutional failure. However, as pointed out by Acharya (2009), “the goal of prudential regulation should be to ensure the financial stability of the system as a whole, i.e., of an institution not only individually but also as a part of the overall financial system”. To address this shortcoming, some authors (including Lehar (2005), Acharya (2009), Huang et al. (2009) and Acharya et al. (2017)) have proposed different prudential regulation models that consider both individual and systemic failure risk. Successful implementation of these models relies on effective methods for monitoring systemic risk. This paper is a step towards achieving this goal by examining a strategy based on the combination of conditional and unconditional VaR to capture the dimension of systemic risk. In particular, we study conditions that guarantee the ordering between VaR and CoVaR, allowing us to understand which, among the two measures, is more or less conservative than the other.

For a random variable Y representing a financial risk, this strategy requires the presence of an adequate covariate X (we discuss in Section 3 some guidelines on choosing it). Suppose that, initially, the risk-capital calculation is based on the VaR at level v of Y , for some $v \in (0, 1)$. If, given a risk X and $u \in [0, 1]$, $\text{CoVaR}_{v,u}[Y|X] < \text{VaR}_v[Y]$ holds, replacing VaR by CoVaR does not make sense as CoVaR contemplates a more optimistic scenario. However, if $\text{CoVaR}_{v,u}[Y|X] > \text{VaR}_v[Y]$ holds, then $\text{VaR}_v[Y]$ underestimates the spillover effect and it is prudent to replace it with $\text{CoVaR}_{v,u}[Y|X]$ as a more conservative provision.

Given $v \in (0, 1)$, the point in question here is the existence of a risk-level threshold $u_v \in (0, 1)$, as a function of v , such that $\text{CoVaR}_{v,u_v}[Y|X] = \text{VaR}_v[Y]$. Although this u_v might not be unique or even exist, its study is especially interesting in situations where it is unique. As we show in Section 2, we can guarantee its existence and uniqueness under reasonable assumptions on the dependence structure of the vector (X, Y) . The primary purpose of this paper is to provide a plausible interpretation of the level u_v and discuss its implications under such assumptions. The findings of this study may enable decision-makers to understand better when “to sound the alarm” and implement adequate response actions.

Another motivation for the study of u_v is the recent work by Li and Wang (2022), who studied a similar balancing point for the equivalence between the value-at-risk and the expected shortfall, called PELVE (Probability Equivalent Level of value-at-risk and expected shortfall). Specifically, for a chosen level ϵ close to 0 and a loss random variable Y , the PELVE is a constant multiplier c such that $\text{ES}_{1-c\epsilon}(Y) = \text{VaR}_{1-\epsilon}(Y)$, where

$$\text{ES}_v[Y] = \frac{1}{1-v} \int_v^1 \text{VaR}_t[Y] dt,$$

is the expected shortfall at level $v \in (0, 1)$. The value of PELVE addresses the issue of regulatory capital change if one replaces VaR with ES (see Li and Wang (2022) for details). Generalizations of PELVE have been proposed in Fiori and Rosazza Gianin (2022) and Barczy et al. (2023).

Throughout the paper, we assume the following regularity conditions hold: the random vector (X, Y) has absolutely continuous strictly increasing marginal distribution functions $F_X(\cdot)$ and $F_Y(\cdot)$ and strictly increasing conditional distribution $F_{Y|X=x}(\cdot)$, for all x , on their respective supports. For a random vector satisfying these conditions, we define

the probability equivalent level of CoVaR-VaR at risk level v (in short, PELCoV_v) as follows.

Definition 1. Let (X, Y) be a random vector satisfying regularity conditions. Given $v \in (0, 1)$, we say that $u_v \in (0, 1)$ is a probability equivalent level of CoVaR-VaR at risk level v (PELCoV_v) for X if $\text{CoVaR}_{v,u_v}[Y|X] = \text{VaR}_v[Y]$.

We model the dependence structure between X and Y using copulas. If K is the joint distribution of the random vector (X, Y) , by the Sklar's theorem, we can write

$$K(x, y) = C(F_X(x), F_Y(y)),$$

where C is the joint distribution function of the vector-copula (U, V) given by $U = F_X(X)$ and $V = F_Y(Y)$. The function C is the copula and collects the information on the inner dependence of the vector, independently from the marginal behaviour of the components. Under the regularity conditions, C is unique and differentiable (see Nelsen, 2007). For a random vector (X, Y) satisfying regularity conditions and $v \in (0, 1)$, we show below that a PELCoV_v only depends on the copula, a result that places our research in the context of other recent studies that model systemic risk using a copula-CoVaR approach, including Reboredo and Ugolini (2015), Bernardi et al. (2017), Jaworski (2017), Karimalis and Nomikos (2018), Jin (2018), Di Clemente (2018) and Zhao (2022), among others.

Decision-makers in finance and insurance are often interested in positive dependence structures among risks. We mean by positive dependence that X and Y are likely to be large or small together. The interested reader can find an introduction to dependence notions in the monograph by Joe (1997) and a recent review of concepts in Navarro et al. (2020). In particular, we focus on the positive regression dependence (PRD) concept introduced by Lehmann (1966), also called stochastically increasing (SI). We first recall the definition of the usual stochastic order. Given two random variables X_1 and X_2 , X_1 is said to be smaller than X_2 in the usual stochastic order, denoted by $X_1 \leq_{st} X_2$, if the survival probabilities satisfy

$$\Pr\{X_1 > x\} \leq \Pr\{X_2 > x\}, \forall x \in \mathbb{R}. \quad (1)$$

Roughly speaking, X_1 is less likely than X_2 to take on large values. It is straightforward to check the following equivalence,

$$X_1 \leq_{st} X_2 \iff \text{VaR}_u[X_1] \leq \text{VaR}_u[X_2], \forall u \in [0, 1]. \quad (2)$$

Further details about the usual stochastic order can be found in Shaked and Shanthikumar (2007). Now we define the SI concept in terms of the $\leq_{st}$ order between conditional distributions. Let (X, Y) be a random vector with copula C . We will say that Y is SI in X , in short $Y \uparrow_{SI} X$, if

$$[Y|X = x_1] \leq_{st} [Y|X = x_2], \forall x_1 \leq x_2. \quad (3)$$

In some sense, if $Y \uparrow_{SI} X$ we expect that Y is likely to take on large values when the conditional random variable X increases. It is clear from (1) that (3) holds if and only if the survival probability $\Pr\{Y > y | X = x\}$ is an increasing function of x , for all y . The SI concept is characterized by the copula: $Y \uparrow_{SI} X$ is equivalent to checking that the partial derivative

$$\partial_1 C(u, v) = \Pr\{V \leq v | U = u\} \quad (4)$$

is a decreasing function of u , for all v , where (U, V) is defined as above, (see page 41 in Nelsen, 2007). Consequently, the vector (X, Y) is SI if and only if its copula is SI. If $Y \uparrow_{SI} X$ and $X \uparrow_{SI} Y$, we say that the vector (X, Y) is positively dependent through stochastic ordering (PDS). Note that (X, Y) is PDS if and only if its copula C is PDS (see Cai and Wei, 2012). Throughout the paper, we will only consider the case where Y is SI in X in a strict sense, i.e., where the conditional distribution

¹ A modified definition of CoVaR was given in Girardi and Ergün (2013) and Mainik and Schaanning (2014).

$\Pr\{Y > y \mid X = x\}$ is a strict increasing function of x , for all y . We will denote this property by $Y \uparrow_{SSI} X$ (strict SI property). Note that the SSI property implies that the conditional distribution $[Y \mid X = x]$ varies when $X = x$ does.

The rest of the paper is organized as follows. In Section 2, we show that under some regularity conditions on the copula and a particular dependence structure of the vector, the PELCoV_v exists and is unique. In Section 3, we interpret this notion as a signal that the VaR of Y at level v is “starting to underestimate” the spillover risk effect.

The issue is appealing when we have a unique signal for all $v \in (0, 1)$, which takes us to study the existence of an upper bound for the set of PELCoV_v s, with $v \in (0, 1)$. This is why, in Section 4, we focus on the monotonicity and boundedness properties of the PELCoV_v . In Section 5, we determine the PELCoV_v and study its boundedness for several parametric families of copulas. In Section 6 we provide an example with real data to illustrate the methodology application.

2. Existence and uniqueness of PELCoV_v

In this section we establish conditions to guarantee the existence and uniqueness of the PELCoV_v . To begin with, we observe that the relative position of VaR and CoVaR only depends on the copula.

Lemma 2. *Let (X, Y) be a random vector satisfying regularity conditions with copula C . Given $u, v \in (0, 1)$, $\text{CoVaR}_{v,u}[Y \mid X] \geq \text{VaR}_v[Y]$ (respectively $\leq, =$) if, and only if $\partial_1 C(u, v) \leq v$ (respectively $\geq, =$).*

Proof. Rewriting (4) as $\partial_1 C(u, v) = F_{Y \mid X=\text{VaR}_u[X]}(\text{VaR}_v[Y])$ the result follows easily. $\square$

Given $v \in (0, 1)$, Lemma 2 says that a PELCoV_v only depends on the copula. The next result shows that the continuity of $\partial_1 C(u, v)$ determines the continuity of $\text{CoVaR}_{v,u}[Y \mid X]$.

Lemma 3. *Let (X, Y) be a random vector satisfying regularity conditions with copula C . Then, $\text{CoVaR}_{v,u}[Y \mid X]$ is continuous in $u \in (0, 1)$, for all $v \in (0, 1)$ if, and only if, $\partial_1 C(u, v)$ is continuous in $u \in (0, 1)$, for all $v \in (0, 1)$.*

Proof. We only prove the “if” part (the “only if” part is proved in the same way). Given an $\epsilon > 0$ and $u_0 \in (0, 1)$, the inequality

$$|\text{CoVaR}_{v,u}[Y \mid X] - \text{CoVaR}_{v,u_0}[Y \mid X]| < \epsilon,$$

is equivalent to

$$\partial_1 C(u, F_Y(y_1)) < v < \partial_1 C(u, F_Y(y_2)), \quad (5)$$

where $y_1 = \text{CoVaR}_{v,u_0}[Y \mid X] - \epsilon$ and $y_2 = \text{CoVaR}_{v,u_0}[Y \mid X] + \epsilon$. By the assumptions, using that $\partial_1 C(u, v)$ is strict increasing in v ,

$$\lim_{u \rightarrow u_0} \partial_1 C(u, F_Y(y_1)) = \partial_1 C(u_0, F_Y(y_1)) < v \quad (6)$$

and

$$\lim_{u \rightarrow u_0} \partial_1 C(u, F_Y(y_2)) = \partial_1 C(u_0, F_Y(y_2)) > v. \quad (7)$$

From (6) and (7), we see that (5) holds in an interval $|u - u_0| < \delta$, with δ small enough. $\square$

Proposition 4. *Let (X, Y) be a random vector satisfying regularity conditions with copula C . Given a risk-level $v \in (0, 1)$, there exist $u_v^+, u_v^- \in (0, 1)$ such that $\text{CoVaR}_{v,u_v^-}[Y \mid X] \leq \text{VaR}_v[Y] \leq \text{CoVaR}_{v,u_v^+}[Y \mid X]$.*

Proof. Let $v \in (0, 1)$. Reasoning by contradiction, assume that u_v^+ or u_v^- do not exist. Then, the conditional quantile curve $u \mapsto \text{CoVaR}_{v,u}[Y \mid X]$ lies strictly either above or below the $\text{VaR}_v[Y]$. Suppose that the curve

lies strictly above (the other case is analogous). From Lemma 2, the inequality $\text{CoVaR}_{v,u}[Y \mid X] > \text{VaR}_v[Y]$, for all $u \in (0, 1)$, is equivalent to checking that $v > \partial_1 C(u, v)$, for all $u \in (0, 1)$. The latter implies that

$$v = \int_0^1 v du > \int_0^1 \partial_1 C(u, v) du = C(1, v) = v, \quad (8)$$

which is a contradiction and concludes the proof. $\square$

We remark that Proposition 4 holds for $0 < v < 1$. This result says that, under regularity conditions, the curves $u \mapsto \text{VaR}_v[Y]$ and $u \mapsto \text{CoVaR}_{v,u}[Y \mid X]$ necessarily cut each other. Using (2), this is equivalent to saying that there always exists a range of values of u such that the random variables Y and $[Y \mid X = \text{VaR}_u[X]]$ fail to be stochastically ordered. Consider, for example, $v_1 < v_2$ such that the probability equivalent levels u_1, u_2 verify $u_{v_1} < u_{v_2}$. For $u \in (u_{v_1}, u_{v_2})$ one has $\text{CoVaR}_{v_1,u}[Y \mid X] > \text{VaR}_{v_1}[Y]$ and $\text{CoVaR}_{v_2,u}[Y \mid X] < \text{VaR}_{v_2}[Y]$. Thus, by (2), neither $Y \leq_{st} [Y \mid X = \text{VaR}_u[X]]$ nor $Y \geq_{st} [Y \mid X = \text{VaR}_u[X]]$ can hold true. Regarding this issue, we will provide in Section 4 sufficient conditions for the existence of a $u^* \in (0, 1)$ such that Y and $[Y \mid X = \text{VaR}_u[X]]$ are stochastically ordered for all $u \in [u^*, 1]$.

The next theorem gives sufficient conditions for existence and uniqueness of $u_v = \text{PELCoV}_v$.

Corollary 5. *Let (X, Y) be a random vector satisfying regularity conditions with a copula C and let $v \in (0, 1)$. Then:*

- (a) *If $\partial_1 C(u, v)$ is continuous in $u \in (0, 1)$, there exists at least a $u_v \in (0, 1)$ such that $\text{CoVaR}_{v,u_v}[Y \mid X] = \text{VaR}_v[Y]$.*
- ii) *If $Y \uparrow_{SSI} X$, then $u_v = \text{PELCoV}_v$ is unique.*

Proof. The proof of (a) follows from Lemma 3 and Proposition 4 by application of Darboux property to the continuous function $u \mapsto \text{CoVaR}_{v,u}[Y \mid X]$ in the closed interval given by the values u_v^- and u_v^+ . To prove (b), note that $Y \uparrow_{SSI} X$ if, and only if, $\text{CoVaR}_{v,u}[Y \mid X]$ is strictly increasing in u , for all v (see Fernández-Ponce et al., 2011). $\square$

The following example shows that the continuity of $\partial_1 C(u, v)$ in $u \in (0, 1)$ is necessary for the existence of u_v .

Example 6. Let C be a copula having density $c(u, v)$ as follows

$$c(u, v) = \begin{cases} 2 & \text{if } (u - 1/2)(v - 1/2) > 0, \\ 0 & \text{if } (u - 1/2)(v - 1/2) \leq 0. \end{cases}$$

A straightforward computation shows that

$$\partial_1 C(u, v) = \begin{cases} 2v & \text{if } 0 \leq u, v \leq 1/2, \\ 1 & \text{if } 0 \leq u \leq 1/2, 1/2 < v \leq 1, \\ 0 & \text{if } 1/2 < u \leq 1, 0 \leq v \leq 1/2, \\ 2v - 1 & \text{if } 1/2 < u, v \leq 1. \end{cases}$$

By construction, C is absolutely continuous and satisfies the regularity conditions. However, $\partial_1 C(u, v)$ is not continuous in $u \in (0, 1)$, for all $v \in (0, 1)$. In particular, it is always discontinuous at $u = 0.5$. It is clear that $\partial_1 C(u, v) \neq v$ for all $u, v \in (0, 1)$. Hence the PELCoV_v does not exist, for all $v \in (0, 1)$.

3. Further discussion on the PELCoV_v

In what follows, in order to guarantee existence and uniqueness, we consider a random vector (X, Y) that satisfies the regularity conditions with copula C , such that $Y \uparrow_{SSI} X$ and $\partial_1 C(u, v)$ is continuous in u .




Fig. 1. PELCoV_v values for $v_1 = 0.4, v_2 = 0.7$ and $v_3 = 0.95$ for a normal vector (X, Y) with $\mu_X = 10000, \sigma_X = 1000, \mu_Y = 40000, \sigma_Y = 2500$ and $\rho = 0.9$. (For interpretation of the colours in the figure(s), the reader is referred to the web version of this article.)

To clarify the exposition, we present in Fig. 1 some computations² of $u_v = \text{PELCoV}_v$ for a bivariate normal vector (X, Y) with $\mu_X = 10000, \sigma_X = 1000, \mu_Y = 40000, \sigma_Y = 2500$ and $\rho = 0.9$ (the positive correlation coefficient guarantees that $Y \uparrow_{S,SI} X$). Note that u_v only depends on the copula (that is, on ρ). The graphic plots $u_{v_i}, i = 1, 2, 3$, for $v_1 = 0.4, v_2 = 0.7$ and $v_3 = 0.95$, where u_{v_i} is the abscissa of the intersection point between the conditional quantile curve $u \mapsto \text{CoVaR}_{v_i,u}[Y|X]$ and the line $y = \text{VaR}_{v_i}[Y]$. Given $v \in [0, 1]$, we see from the graphic that

$$u_v = \min_{u \in [0,1]} \{ \text{CoVaR}_{v,u'}[Y|X] \geq \text{VaR}_v[Y], \text{ for all } u' > u \}, \quad (9)$$

representing the lowest risk level u of X from which $\text{CoVaR}_{v,u}[Y|X]$ will be a more conservative reserve position than $\text{VaR}_v[Y]$. For the bivariate normal distribution shown in Fig. 1, the PELCoV_v is an increasing function of v and tends to 1 as v goes to 1. However, this is not always the case: the PELCoV_v is not necessarily increasing in v , as we show in Section 5. The existence of an upper bound for the PELCoV_v deserves further discussion. If there exists a $u^* < 1$ such $u_v \leq u^*$ for all v , then

$$\begin{aligned} \text{CoVaR}_{v,u}[Y|X] &\geq \text{CoVaR}_{v,u^*}[Y|X] \\ &\geq \text{CoVaR}_{v,u_v}[Y|X] \\ &= \text{VaR}_v[Y], \end{aligned}$$

for all $v \in (0, 1)$ and for all $u \in [u^*, 1]$, where the inequalities follow from the fact that $\text{CoVaR}_{v,u}[Y|X]$ is strictly increasing in u (due to $Y \uparrow_{S,SI} X$) and the equality is by definition of u_v . Using (2), this can be rewritten as³

$$Y \leq_{SI} [Y|X = \text{VaR}_u[X]], \quad \forall u \in [u^*, 1]. \quad (10)$$

We interpret (10) as follows. Given $v \in (0, 1)$, if X exceeds its quantile u^* , $\text{VaR}_v[Y]$ underestimates the risk spillover effect, and it is more prudent to determine reserves in terms of $\text{CoVaR}_{v,u}[Y|X]$, for all $u > u^*$. Returning to the interpretation of (9), the condition $Y \uparrow_{S,SI} X$ implies that the vector (X, Y) is PQD (positive quadrant dependent, see Nelsen, 2007), that is,

$$\Pr\{Y > \text{VaR}_v[Y] | X > \text{VaR}_u[X]\} \geq 1 - v, \quad \text{for all } u, v \in (0, 1), \quad (11)$$

where $1 - v = \Pr\{Y > \text{VaR}_v[Y]\}$. This means that the assumption $Y \uparrow_{S,SI} X$, by itself, ensures that the conditional information that X exceeds

its quantile u (for any u) increases the probability of Y exceeding its v quantile. The additional information is that, given $v \in (0, 1)$, u_v is the smallest probability-level such that

$$\text{CoVaR}_{v,u}[Y|X] \geq \text{VaR}_v[Y], \quad \forall u \geq u_v. \quad (12)$$

In our approach, this means that if the institution X is becoming riskier and exceeds its u_v quantile, $\text{CoVaR}_{v,u}[Y|X]$ is more conservative than $\text{VaR}_v[Y]$. Thus, u_v acts as a signal or “alert” that indicates that the unconditional VaR is “starting to underestimate” the risk. If we define the regression expected shortfall (RES) at level v as

$$\text{RES}_v[Y|X] = \frac{1}{1 - u_v} \int_{u_v}^1 \text{CoVaR}_{v,u}[Y|X] du, \quad (13)$$

it follows from (12) that $\text{RES}_v[Y|X] \geq \text{VaR}_v[Y]$. We can interpret (13) as a modified expected shortfall, where the conditional event $\{Y > \text{VaR}_v[Y]\}$ is replaced by $\{X > \text{VaR}_{u_v}[X]\}$, a stress scenario intended to capture the contagion effect.⁴

We now return to the initial problem of determining provisions for Y using a VaR-CoVaR-based approach. To choose a good covariate X for Y , we suggest taking into consideration the following criteria.

- The dependence structure between X and Y should be easily observable (it determines both u_v and the monotonicity of $\text{CoVaR}_{v,u}[Y|X]$ when u exceeds u_v) but not necessarily too strong. Under comonotonicity, for example, $u_v = v$ and the PELCoV is not helpful. We should also mention that a greater positive dependence between X and Y does not imply a higher PELCoV_v . In general, the PELCoV_v is not monotonic with respect to the concordance ordering of copulas.
- Given $v \in (0, 1)$, a good “alert” level u_v should be reasonably smaller than v . The higher the PELCoV_v , the less likely the event

⁴ The risk measure (13) should not be confused with the conditional expected shortfall (or CoES) introduced by Mainik and Schaanning (2014) as

$$\text{CoES}_{v,u}[Y|X] = \frac{1}{1 - v} \int_v^1 \text{CoVaR}_{t,u}[Y|X] dt.$$

While the latter averages the quantiles $F_{Y|X=\text{VaR}_u[X]}^{-1}(t)$ with u fixed and t varying from v on, the regression expected shortfall given by (13) averages the quantiles $F_{Y|X=\text{VaR}_u[X]}^{-1}(v)$, with v fixed and u varying from u_v on.

² Computations are based on formulas (14) to (16).

³ Comparisons of the form (10) are discussed in Navarro and Sordo (2018).

$[X > \text{VaR}_{u_v}[X]]$ to occur. If the strategy consists of replacing $\text{VaR}_{0.99}[Y]$ with $\text{CoVaR}_{0.99,0.99}[Y|X]$, the alert level $u_v = 0.99$ may come too late to be effective.

- (c) Given $v \in (0, 1)$, the regression expected shortfall (13) quantifies the impact of X exceeding its u_v -quantile on Y . Therefore, the higher $\text{RES}_v[Y|X]$, the more effective the replacement of $\text{VaR}_v[Y]$ with $\text{CoVaR}_{v,u_v}[Y|X]$ in the sense that it protects the company from a worse scenario.

For example, suppose that Y is normally distributed with parameters μ_Y and σ_Y and we base risk calculations on $\text{VaR}_v[Y]$, for some $0 < v < 1$. It is well-known that

$$\text{VaR}_v[Y] = \mu_Y + \text{VaR}_v[Z]\sigma_Y, \quad (14)$$

where $Z \sim N(0, 1)$. Let X be a random variable such that (X, Y) is a bivariate normal vector with correlation coefficient ρ and let μ_X and σ_X be the parameters of X . It is well-known that the conditional distribution $[Y|X = x]$ is a normal distribution with parameters $\mu_Y + \rho \frac{\sigma_Y}{\sigma_X}(x - \mu_X)$ and $\sigma_Y \sqrt{1 - \rho^2}$. According to (14) we have

$$\begin{aligned} \text{CoVaR}_{v,u}[Y|X] = & \mu_Y + \rho \frac{\sigma_Y}{\sigma_X}(\text{VaR}_u[X] - \mu_X) \\ & + \text{VaR}_v[Z]\sigma_Y \sqrt{1 - \rho^2}, \end{aligned} \quad (15)$$

and, by solving the equation $\text{CoVaR}_{v,u}[Y|X] = \text{VaR}_v[Y]$, we obtain

$$u_v = F_Z\left(\frac{1 - \sqrt{1 - \rho^2}}{\rho} \text{VaR}_v[Z]\right). \quad (16)$$

Now suppose that $v = 0.95$ and $\rho = 0.4$. We can check that $u_{0.95} = 0.6344$. Since the event $[X > \text{VaR}_{0.6344}[X]]$ is likely to happen, we expect the “alarm” to sound often enough to be effective. Combining (13) and (15), we evaluate the impact on Y caused by the contagion effect by

$$\text{RES}_v[Y|X] = \mu_Y + k_{\rho,v}\sigma_Y, \quad (17)$$

where $k_{\rho,v} = \rho \text{ES}_{u_v}[Z] + \sqrt{1 - \rho^2} \text{VaR}_v[Z]$ (here we have used that $\text{ES}_u[X] = \mu_X + \text{ES}_u[Z]\sigma_X$). It follows from comparing (14) and (17) that the difference $k_{0.4,0.95} - \text{VaR}_{0.95}[Z]$ quantifies the severity of the spillover effect. In this example, $k_{0.4,0.95} = 1.9189 > 1.6448 = \text{VaR}_{0.95}[Z]$, which reasonably justifies the replacement of $\text{VaR}_{0.95}[Y]$ with $\text{CoVaR}_{0.95,0.6344}[Y|X]$.

4. Monotonicity and boundedness

As in Section 3, we consider a random vector (X, Y) with copula C that satisfies the regularity conditions, such that $Y \uparrow_{SSI} X$ and $\partial_1 C(u, v)$ is continuous in u for all $v \in (0, 1)$. In this section, we discuss monotonicity and boundedness properties of the PELCoV_v .

4.1. Monotonicity

The growth of the PELCoV_v as a function of v is meaningful and easy to interpret: as the capital requirements increase (moving, for example, from $\text{VaR}_{0.95}[Y]$ to $\text{VaR}_{0.99}[Y]$), the risk-level u for X at which CoVaR becomes more conservative than VaR increases accordingly (that is, $u_{0.95} \leq u_{0.99}$). From Fig. 1 and the discussion in Section 3 (where we said that the PELCoV_v of a bivariate normal distribution with $\rho > 0$ is increasing in v), we can think that the map $v \mapsto u_v$ is increasing whenever $Y \uparrow_{SSI} X$. This is a false intuition, as the following example shows.

Example 7. Let X and W be two independent random variables, $X \sim \text{Exp}(1)$ and $W \sim \text{Weibull}(0.5, 1)$. The vector (X, Y) , where $Y = X + W$, satisfies the regularity conditions (in particular, $Y \uparrow_{SSI} X$). Given $v \in (0, 1)$, the PELCoV_v for this vector is the unique u such that




Fig. 2. Function $v \mapsto u_v$ for the vector (X, Y) , where $Y = X + W$, where $X \sim \text{Exp}(1)$, $W \sim \text{Weibull}(0.5, 1)$ and X and W are independent.

$\text{CoVaR}_{v,u}[Y|X] = \text{VaR}_v[Y]$. Using that $\text{VaR}_v[W + c] = \text{VaR}_v[W] + c$ for any constant c , we get $u = F_X((\text{VaR}_v[Y] - \text{VaR}_v[W]))$. We can compute numerically these values taking into account that the marginal distribution function of Y is given by

$$\begin{aligned} F_Y(x) &= \int_0^x F_W(x - y)f_X(y)dy = \int_0^x (1 - e^{-\sqrt{x-y}}) \cdot e^y dy = \\ &= 1 - e^{-\sqrt{x}} - e^{-x} \cdot DF\left(\frac{1}{2}\right) + e^{-\sqrt{x}} DF\left(\frac{1}{2} - x\right), \end{aligned}$$

where $DF(x)$ is the Dawson's integral $DF(x) = e^{-x^2} \int_0^x e^{-y^2} dy$. Fig. 2 plots the PELCoV_v as a function of v . The maximum is attained at $v = u_0 = 0.74816$ and $u^* = u_{u_0} = 0.764213$.

Example 7 shows that the function $v \mapsto u_v$ is not necessarily increasing under the assumption $Y \uparrow_{SSI} X$. It is natural, therefore, to ask for additional conditions under which u_v increases as a function of v . Observe that, by Lemma 2, under the regularity conditions, the PELCoV_v is the unique value $u_v \in (0, 1)$ such that $\partial_1 C(u_v, v) = v$. Assuming differentiability of $\partial_1 C(u, v)$ and using the implicit function theorem we see that

$$\frac{du_v}{dv} = \frac{1 - c(u_v, v)}{\partial_{11} C(u_v, v)}, \quad (18)$$

where $c(u, v) = \frac{\partial^2}{\partial u \partial v} C(u, v)$ is the copula density. In particular, if $\partial_{11} C(u, v) < 0$ for all $u, v \in (0, 1)^2$ (which is equivalent to $Y \uparrow_{SSI} X$), we see that

$$\frac{du_v}{dv} > 0 \text{ if and only if } c(u_v, v) > 1. \quad (19)$$

Direct verification of the growth of u_v as a function of v using (19) requires the analytical expression of u_v . Next we present a sufficient (but not necessary) condition, that does not require it, based on the Bickel and Lehmann (1979) dispersive order. Recall that X_1 is said to be smaller than X_2 in the dispersive order, denoted by $X_1 \leq_{disp} X_2$, if the distances between their VaRs satisfy

$$\text{VaR}_{u_1}[X_1] - \text{VaR}_{u_2}[X_1] \leq \text{VaR}_{u_1}[X_2] - \text{VaR}_{u_2}[X_2] \text{ for all } 0 \leq u_1 < u_2 \leq 1.$$

Although the PELCoV is based only on the copula, we can give sufficient conditions for its monotonicity in terms of both the marginals and copula behaviour.

Proposition 8. Let (X, Y) be a random vector satisfying regularity conditions with a copula C , such that $Y \uparrow_{SSI} X$ and $\partial_1 C(u, v)$ is continuous in both arguments. If $[Y|X = \text{VaR}_u[X]] \leq_{disp} Y$ for all $u \in (0, 1)$, then the PELCoV_v increases with respect to v .

Proof. Given $v_1 < v_2$, for all $u \in (0, 1)$ we have

$$\begin{aligned} \text{VaR}_{v_2}[Y|X = \text{VaR}_u[X]] - \text{VaR}_{v_1}[Y|X = \text{VaR}_u[X]] \\ \leq \text{VaR}_{v_2}[Y] - \text{VaR}_{v_1}[Y] \end{aligned}$$

By Corollary 5, there exists a u_{v_2} such that $\text{CoVaR}_{v_2, u_{v_2}}[Y|X] = \text{VaR}_{v_2}[Y]$. Therefore, taking $u = u_{v_2}$ in the above inequality it follows that

$$\text{CoVaR}_{v_1, u_{v_2}}[Y|X] - \text{VaR}_{v_1}[Y] \geq 0, \text{ for all } v_1 < v_2. \quad (20)$$

Since $Y \uparrow_{SSI} X$, the function $u \mapsto \text{CoVaR}_{v, u}[Y|X]$ is strictly increasing. This fact, together with (20), implies $u_{v_1} \leq u_{v_2}$. $\square$

In Section 3, we said without proof that the PELCoV_v of a bivariate normal distribution with $\rho > 0$ is increasing in v (see Fig. 1). Now we can prove it as follows. If (X, Y) is a bivariate normal vector with correlation coefficient ρ , the univariate random variables $[Y|X = F^{-1}(u)]$ and Y are normally distributed with variances $\sigma_Y^2(1 - \rho^2)$ and σ_Y^2 , respectively. Using that a normal distribution is smaller than another one in the dispersive order if its variance is smaller (see, Table 1.1 in Müller and Stoyan, 2002), the result follows as a consequence of Proposition 8.

Let (X^C, Y^C) denote a random vector with marginals F and G and copula C . Belzunce et al. (2008) say that (X^C, Y^C) is less in the conditional dispersive order⁵ than $(X^{C'}, Y^{C'})$ if $[Y^C|X^C = F^{-1}(u)] \leq_{disp} [Y^{C'}|X^{C'} = F^{-1}(u)]$, for all $u \in (0, 1)$. Observe that $[Y|X = F^{-1}(u)] \leq_{disp} Y$, for all $u \in (0, 1)$, is a particular case for C' , the independence copula. This order is also a necessary condition for the growth of u_v as a function of v in some particular models, as we now show.

Proposition 9. Let $\mathbf{X} = (X, Y)$ be a random vector such that $Y = \psi(X) + W$, where X and W are independent random variables and ψ is a strictly increasing function. The PELCoV_v , given by

$$u_v = F_X(\psi^{-1}(\text{VaR}_v[Y] - \text{VaR}_v[W])),$$

increases with respect to $v \in (0, 1)$ if and only if $W \leq_{disp} \psi(X) + W$. Moreover, this last inequality is always true when W has a logconcave density function.

Proof. First, $Y \uparrow_{SSI} X$ trivially holds. Given $v \in (0, 1)$, u_v is determined using the same argument as in Example 7 and the fact that $\psi(\text{VaR}_u[X]) = \text{VaR}_u[\psi(X)]$ for a strictly increasing ψ . Clearly, the growth of u_v as a function of v is equivalent to the growth of $\text{VaR}_v[\psi(X) + W] - \text{VaR}_v[W]$ in v , which is the same as $\psi(X) + W \geq_{disp} W$. By Theorem 3.B.7 in Shaked and Shanthikumar (2007), this inequality is always true when W has a logconcave density function. $\square$

Note that Proposition 9 does not apply to the vector (X, Y) in Example 7 because the density function of the Weibull distribution is not log-concave.

4.2. Boundedness

As we explained in Section 3, the existence of a $u^* < 1$ such $u_v \leq u^*$ for all v , ensures that $\text{VaR}_v[Y]$ will underestimate, for all v , the risk spillover effect as soon as X exceeds its quantile u^* . The aim of this section is to provide conditions to ensure the existence of an upper bound $u^* < 1$ for the set $\{u_v : v \in (0, 1)\}$.

If the function that maps v to u_v is increasing, then the supremum of the set $\{u_v : v \in (0, 1)\}$ is $\lim_{v \rightarrow 1} u_v$. In this case, the existence of an

upper bound $u^* < 1$ reduces to the study of this limit. In general, when the function $v \mapsto u_v$ is not monotonic, the supremum may be attained in some $v_0 \in (0, 1)$, as in Example 7 (where $v_0 = 0.74816$). The following result establishes conditions under which the existence of an upper bound $u^* < 1$ for the set $\{u_v : v \in (0, 1)\}$ can be proved by computing the limits of the PELCoV_v when v tends to 0 and to 1.

Corollary 10. Let (X, Y) be a random vector satisfying regularity conditions with a copula C , such that $Y \uparrow_{SSI} X$ and $\partial_1 C(u, v)$ is continuous in u . The following conditions are equivalent:

- (1) There exist $u^* < 1$ such that $u_v \leq u^*$ for all $v \in (0, 1)$.
- (2) $\lim_{v \rightarrow 0} u_v < 1$ and $\lim_{v \rightarrow 1} u_v < 1$.

Proof. First we show, by contradiction, that there does not exist a $v \in (0, 1)$ such that $v = \partial_1 C(1, v)$. Assume that there exists $v^* \in (0, 1)$ such that $v^* = \partial_1 C(1, v^*)$. It follows from $Y \uparrow_{SSI} X$ that $\partial_1 C(1, v)$ decreases in u . Then, $v^* < \partial_1 C(u, v^*)$, for all $u \in (0, 1)$, which is a contradiction as in (8). Consequently, if the supremum of the set $\{u_v : v \in (0, 1)\}$ is attained at some $v_0 \in (0, 1)$, then it is necessarily less than one. This implies that a necessary and sufficient condition for the supremum of $\{u_v : v \in (0, 1)\}$ to be 1 is $\lim_{v \rightarrow 0} u_v = 1$ or $\lim_{v \rightarrow 1} u_v = 1$. $\square$

The following theorem gives sufficient conditions for the existence of an upper bound $u^* < 1$ for the set $\{u_v : v \in (0, 1)\}$ in terms of the copula density $c(u, v)$.

Theorem 11. Let (X, Y) be a random vector satisfying regularity conditions with copula C , such that $Y \uparrow_{SSI} X$ and $\partial_1 C(u, v)$ is continuous in u . Let $c(u, v)$ be the copula density. If

- (i) there exists $m < 1$ such that $c(u, v)$ increases in v for all $u \geq m$ and $v \leq u$,
 - (ii) the limit $\lim_{(u, v) \rightarrow (1, 1)} c(u, v)$ exists and it is strictly greater than one,
- then there exist $u^* < 1$ such that $u_v \leq u^*$ for all $v \in (0, 1)$.

Proof. The idea of the proof is to show that the assumptions imply $\lim_{v \rightarrow 0} u_v < 1$ and $\lim_{v \rightarrow 1} u_v < 1$ and then to apply Corollary 10.

We first show that (ii) implies $\lim_{v \rightarrow 1} u_v < 1$. From (ii), there exists $\delta > 0$ such that $c(u, v) > 1$ for all $(u, v) \in (1 - \delta, 1)^2$. This implies that $\frac{\int_v^1 c(u, z) dz}{1 - v} > 1$ for all $(u, v) \in (1 - \delta, 1)^2$ or, equivalently, that

$$\frac{\int_0^v c(u, z) dz}{v} = \frac{\partial_1 C(u, v)}{v} < 1, \text{ for all } (u, v) \text{ in } (1 - \delta, 1)^2. \quad (21)$$

Using Lemma 2, it follows that $u_v < 1 - \delta$ for $v > 1 - \delta$ and, consequently, $\lim_{v \rightarrow 1} u_v < 1$. Now observe, taking into account the equality in (21), that

(i) implies that

$$\frac{\partial_1 C(u, v)}{v} \text{ increases in } v \text{ for all } u \geq m \text{ and } v \leq u. \quad (22)$$

To prove that $\lim_{v \rightarrow 0} u_v < 1$, we assume, by contradiction, that $\lim_{v \rightarrow 0} u_v = 1$. Then, given $\epsilon > \min(1 - m, \delta)$, there exists $\alpha > 0$ such that for $v_1 < \alpha$, $u_{v_1} > 1 - \epsilon$. For such v_1 there exists $v_2 > v_1$ such that $1 - \delta < v_2 < u_{v_1}$. For these v_1 and v_2 we get

$$1 = \frac{\partial_1 C(u_{v_1}, v_1)}{v_1} \leq \frac{\partial_1 C(u_{v_1}, v_2)}{v_2} < 1,$$

a contradiction, where the equality follows from the definition of PELCoV_v , the first inequality follows from (22) and the second (strict) inequality follows from (21). $\square$

5. Parametric examples of PELCoV_v s

In this section, we determine the PELCoV_v and study the existence of an upper bound $u^* < 1$ for several parametric families of copulas. Recall

⁵ For further details on multivariate dispersive orderings, see Fernandez-Ponce and Suárez-Llorens (2003) and Arias-Nicolás et al. (2005).




Fig. 3. Function $v \mapsto u_v$ for the Frank copula with different values of the dependence parameter.

that given a copula $C(u, v)$ satisfying the assumptions of Corollary 5, the PELCoV_v is the unique value $u_v \in (0, 1)$ such that $\partial_1 C(u_v, v) = v$. For the existence of an upper bound, it is sufficient to verify the conditions of Theorem 11.

5.1. Families of copulas with bounded PELCoV_v s

The result of this section are summarized in Table 1.

5.1.1. Farlie–Gumbel–Morgenstern family of copulas

Let us consider (X, Y) with the Farlie–Gumbel–Morgenstern copula, given by $C(u, v) = uv(1 + \theta(1 - u)(1 - v))$, for $\theta \in [-1, 1]$, $u, v \in [0, 1]$. Since $\partial_1 C(u, v) = v + \theta v(2u - 1)(1 - v)$ if $\theta \in (0, 1]$, $Y \uparrow_{SSI} X$. For $\theta > 0$ the copula density $c(u, v) = 1 + \theta(1 - 2u)(1 - 2v)$ increases in v for all $u \geq 1/2$. Moreover, $\lim_{(u,v) \rightarrow (1,1)} c(u, v) = 1 + \theta > 0$ and it follows from Theorem 11 that there exists an upper bound $u^* < 1$ for the PELCoV_v s.

Observe that $\partial_1 C(u, v) = v$ if and only if $u = 1/2$. Hence, the PELCoV_v is $u_v = 1/2$ for all $v \in (0, 1)$. Moreover, under the SSI condition ($\theta > 0$), $u^* = 1/2$ and $Y \leq_{st} [Y | X = x]$ for all $x \geq F^{-1}(1/2)$. Consequently, for all $v \in (0, 1)$, $\text{VaR}_v[Y]$ underestimates the risk spillover effect whenever X exceeds its median.

5.1.2. Frank family of copulas

The Frank copula $C(u, v) = -\frac{1}{\theta} \ln \left(1 + \frac{(e^{-u\theta} - 1)(e^{-v\theta} - 1)}{e^{-\theta} - 1} \right)$ (see (4.1.5) in Nelsen, 2007) is SSI for $\theta > 0$. The PELCoV_v , given by

$$u_v = \frac{1}{\theta} \ln \left(\frac{e^\theta (1 - e^{\theta v}) (1 - v)}{v (e^{\theta v} - e^\theta)} \right), \quad (23)$$

increases in v . Observe that $v \geq u_v$ if and only if $v \geq \frac{1}{2}$ (see Fig. 3). The copula density, given by

$$c(u, v) = \frac{e^{(1+u+v)\theta} (e^\theta - 1) \theta}{(e^{(u+v)\theta} - e^\theta (e^{u\theta} + e^{v\theta} - 1))^2},$$

satisfies the conditions of Theorem 11 for $\theta > 0$ and $m = 1/2$. The upper bound⁶ is given by $u^* = \lim_{v \rightarrow 1} u_v = \frac{1}{\theta} \ln \left(\frac{e^\theta - 1}{\theta} \right)$.

5.1.3. Clayton family of copulas

Let consider the Clayton copula (see (4.1.1) in Nelsen, 2007), given by $C(u, v) = [\max(u^{-\theta} + v^{-\theta} - 1, 0)]^{-1/\theta}$. This copula is SSI for $\theta > 0$. The copula density is given by

$$c(u, v) = u^{-1-\theta} v^{-1-\theta} (u^{-\theta} + v^{-\theta} - 1)^{-2-\frac{1}{\theta}} (1 + \theta).$$

⁶ See Appendix A for details on the computation.




Fig. 4. Function $v \mapsto u_v$ for the Clayton copula with different values of the dependence parameter.

It can be shown that $c(u, v)$ increases in v if $u > \phi_\theta(v)$, where

$$\phi_\theta(v) = \left(1 - \frac{\theta}{\theta + v^\theta(1 + \theta)} \right)^{1/\theta}$$

is a function that increases in v such that

$$\lim_{v \rightarrow 1} \phi_\theta(v) = \left(\frac{1 + \theta}{1 + 2\theta} \right)^{1/\theta} = m_\theta.$$

For this m_θ we see that $c(u, v)$ satisfies (i) in Theorem 11. Since

$$\lim_{(u,v) \rightarrow (1,1)} c(u, v) = 1 + \theta,$$

condition (ii) also holds for $\theta > 0$ and it follows that the PELCoV_v s are bounded. The PELCoV_v and the upper bound u^* are given by

$$u_v = \left(\frac{v^{-\theta/(\theta+1)} - 1}{v^{-\theta} - 1} \right)^{1/\theta}; \quad u^* = \lim_{v \rightarrow 1} u_v = (1 + \theta)^{-\frac{1}{\theta}}.$$

Fig. 4 plots the function $v \mapsto u_v$ for different values of θ .

5.1.4. Ali–Mikhail–Haq family of copulas

Let us consider the Ali–Mikhail–Haq copula (see (4.3.1) in Nelsen, 2007), given by $C(u, v) = \frac{uv}{(1 - \theta(1 - u)(1 - v))}$ for $\theta \in [-1, 1]$. The copula is SSI for $\theta \in (0, 1]$. Given $\theta \in (0, 1)$, it can be shown that the copula density, given by

$$c(u, v) = \frac{1 + \theta(u + v + uv - 2 + (1 - u)(1 - v)\theta)}{(1 - (1 - u)(1 - v)\theta)^3},$$

increases in v for $u \geq \frac{2}{3}$. Moreover, for $\theta > 0$

$$\lim_{(u,v) \rightarrow (1,1)} c(u, v) = 1 + \theta > 1,$$

and it follows from Theorem 11 that there exists an upper bound u^* for the PELCoV_v s. It can be checked that

$$u_v = \frac{\theta(1 - v) - 1 + \sqrt{1 - \theta(1 - v)}}{\theta(1 - v)}; \quad u^* = \lim_{v \rightarrow 1} u_v = \frac{1}{2}.$$

Fig. 5 plots the function $v \mapsto u_v$ for different values of θ .

5.1.5. Other families of copulas

For the following families of copulas, the conditions (i) and (ii) in Theorem 11 and the bound u^* have been numerically investigated.

(a) Family of copulas (4.1.13) in Nelsen (2007). The copula

$$C(u, v) = \exp \left(1 - [(1 - \ln u)^\theta + (1 - \ln v)^\theta - 1]^{1/\theta} \right)$$

is SSI for $\theta > 1$. The function $v \mapsto u_v$ increases in v and the limit $u^* = \lim_{v \rightarrow 1} u_v$ exists for all $\theta \in (1, 1000)$ (see Fig. 6). (b) Family of copulas (4.1.17) in Nelsen (2007). The copula




Fig. 5. Function $v \mapsto u_v$ for the Ali-Mikhail-Haq copula with different values of the dependence parameter.




Fig. 6. Function $\theta \mapsto u^* = \lim_{v \rightarrow 1} u_v$ for the copula (4.1.13) in Nelsen (2007).




Fig. 7. Function $\theta \mapsto u^* = \lim_{v \rightarrow 1} u_v$ for the copula (4.1.17) in Nelsen (2007).

$$C(u, v) = \left(1 + \frac{[(1+u)^{-\theta} - 1][(1+v)^{-\theta} - 1]}{2^{-\theta} - 1} \right)^{-1/\theta} - 1, \quad \theta \neq 0,$$

is SSI for $\theta > -1$ ($\theta \neq 0$). The PELCoV_v increases in v and the limit $u^* = \lim_{v \rightarrow 1} u_v$ exists for all $\theta \in (-1, 1000) \setminus \{0\}$ (see Fig. 7). (c) Family of copulas 19 in Table 4.1 in Nelsen (2007). The copula

$$C(u, v) = \frac{\theta}{\ln(e^{\theta/u} + e^{\theta/v} - e^{\theta})}, \quad \theta > 0,$$

is SSI. The PELCoV_v increases in v and the limit $u^* = \lim_{v \rightarrow 1} u_v$ exists for all $\theta \in (0, 1000)$ (see Fig. 8).




Fig. 8. Function $\theta \mapsto u^* = \lim_{v \rightarrow 1} u_v$ for the copula 19 in Table 4.1 in Nelsen (2007).




Fig. 9. Function $v \mapsto u_v$ for the Gaussian copula with different values of the dependence parameter.

5.2. Families of copulas with unbounded PELCoV_v s

This section shows several families of copulas such that the PELCoV_v s do not have an upper bound. We start with the gaussian family, which has appeared before in several examples joining a bivariate random vector. To be consistent with such examples, for the gaussian family we also consider the marginals.

5.2.1. Gaussian family of copulas

Let $\mathbf{X} = (X, Y) \sim \mathbf{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ be a bivariate normal random vector with $\boldsymbol{\mu} = (\mu_1, \mu_2)$ and $\boldsymbol{\Sigma} = \begin{pmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{pmatrix}$. For $\rho > 0$ the vector is SSI. We saw in Section 4 that the PELCoV_v increases in terms of v , with u_v given by (16). Observe that $u_v < v$ for all $v \in (\frac{1}{2}, 1)$ (see Fig. 9). Moreover, since $\lim_{v \rightarrow 0} u_v = 0$ and $\lim_{v \rightarrow 1} u_v = 1$, it follows from Corollary 10, that there does not exist a $u^* < 1$ such that $u_v \leq u^*$ for all $v \in (0, 1)$.

5.2.2. Gumbel Hougaard family of copulas

The Gumbel Hougaard family of copulas (see 4.1.4 in Nelsen, 2007) has the form $C(u, v) = \exp\left(-\left[(-\ln u)^\theta + (-\ln v)^\theta\right]^{1/\theta}\right)$. It is SSI for $\theta \geq 1$. The density is given by

$$c(u, v) = \frac{e^{-((-\ln u)^\theta + (-\ln v)^\theta)^{1/\theta}} (\ln[v] \ln[u])^{\theta-1} h(u, v)}{uv}$$

where

$$h(u, v) = \left(\theta - 1 + ((-\ln u)^\theta + (-\ln v)^\theta)^{1/\theta} \right) \left((-\ln u)^\theta + (-\ln v)^\theta \right)^{-2 + 1/\theta}.$$

Since $\lim_{v \rightarrow 1} c(u, v) = 0$, for all $u \in (0, 1)$, $\lim_{u \rightarrow 1} c(u, v) = 0$, for all $v \in (0, 1)$ and $\lim_{u \rightarrow 1} c(u, u) = \infty$, we conclude that $\lim_{(u,v) \rightarrow (1,1)} c(u, v)$ does not exist and Theorem 11 does not apply. The plot of u_v as a function of u is obtained numerically for different values of $\theta \geq 1$. Fig. 10 shows

Table 1

PELCoVs and their bounds for vectors (X, Y) such that $Y \uparrow_{SSI} X$.

Copulas with bounded PELCoV	PELCoV	Bound
$(X, \psi(X) + W)$ with ψ strictly increasing and X, W independent	$u_v = F_X \left(\psi^{-1} \left(F_{\psi(X)+W}^{-1}(v) - F_W^{-1}(v) \right) \right)$	If W has a logconcave density function, $u^* = \lim_{v \rightarrow 1} u_v$
FGM copula: for $\theta \in (0, 1]$ $C(u, v) = uv(1 + \theta(1-u)(1-v))$	$u_v = 1/2$	$u^* = 1/2$
Frank copula: for $\theta > 0$, $C(u, v) = -\frac{1}{\theta} \ln \left(1 + \frac{(e^{-\theta u} - 1)(e^{-\theta v} - 1)}{e^{-\theta} - 1} \right)$	$u_v = \frac{1}{\theta} \ln \left(\frac{e^{\theta(1-u)(1-v)} - 1}{v(e^{\theta u} - 1)} \right)$	$u^* = \frac{1}{\theta} \ln \left(\frac{e^{\theta} - 1}{\theta} \right)$
Clayton copula: for $\theta > 0$, $C(u, v) = [\max(u^{-\theta} + v^{-\theta} - 1, 0)]^{-1/\theta}$	$u_v = \left(\frac{v^{-\theta/(1-v)} - 1}{v^{-\theta} - 1} \right)^{1/\theta}$	$u^* = (1 + \theta)^{-\frac{1}{\theta}}$
AMH copula: for $\theta \in (0, 1]$, $C(u, v) = \frac{uv}{(1-\theta(1-u)(1-v))}$	$u_v = \frac{\theta(1-v)-1+\sqrt{1-\theta(1-v)}}{\theta(1-v)}$	$u^* = 1/2$
Copula 13 in Table 4.1 in Nelsen (2007): for $\theta > 0$, $C(u, v) = \exp \left(1 - [(1 - \ln u)^\theta + (1 - \ln v)^\theta - 1]^{1/\theta} \right)$	Numerically computed	Corollary 10 applies. Numerically computed
Copula 17 in Table 4.1 in Nelsen (2007): for $\theta > 0$, $C(u, v) = \left(1 + \frac{[(1+u)^{-\theta} - 1][(1+v)^{-\theta} - 1]}{2^{-\theta} - 1} \right)^{-1/\theta} - 1$	Numerically computed	Corollary 10 applies. Numerically computed
Copula 19 in Table 4.1 in Nelsen (2007): for $\theta > 0$, $C(u, v) = \frac{\theta}{\ln(e^{\theta/2} + e^{\theta/2} - e^{\theta})}$	Numerically computed	Corollary 10 applies. Numerically computed




Fig. 10. Function $v \mapsto u_v$ for the Gumbel Hougard copula with different values of the dependence parameter.




Fig. 11. Function $v \mapsto u_v$ for the copula 4.1.6 in Nelsen (2007) with different values of the dependence parameter.

that there does not exist an upper bound $u^* < 1$ for the sequence of u_v , $v \in (0, 1)$.

5.3. Other families of copulas

For the following families of copulas, the property SSI and the PELCoVs have been numerically investigated for all $\theta \in (1, 1000)$. In general, for these copulas, there does not exist a $u^* < 1$ such that $u_v \leq u^*$ for all $v \in (0, 1)$.

(a) The family of copulas 4.1.6 in Nelsen (2007), given by $C(u, v) = 1 - [(1-u)^\theta + (1-v)^\theta - (1-u)^\theta(1-v)^\theta]^{1/\theta}$, for $\theta > 1$, see Fig. 11.




Fig. 12. Function $v \mapsto u_v$ for the copula 4.1.12 in Nelsen (2007) with different values in the dependence parameter.




Fig. 13. Function $v \mapsto u_v$ for the copula 4.1.14 in Nelsen (2007) with different values of the dependence parameter.

(b) The family of copulas 4.1.12 in Nelsen (2007) given, for $\theta \geq 1$, by $C(u, v) = \left(1 + [(u^{-1} - 1)^\theta + (v^{-1} - 1)^\theta]^{1/\theta} \right)^{-1}$, see Fig. 12.

(c) The family of copulas 4.1.14 in Nelsen (2007), given, for $\theta \geq 1$, by $C(u, v) = \left(1 + [(u^{-1/\theta} - 1)^\theta + (v^{-1/\theta} - 1)^\theta]^{1/\theta} \right)^{-\theta}$, see Fig. 13.

6. Example

This section provides a real-data-based example to illustrate the methodology. We consider the negative log-returns (or losses) of stock

prices of two companies that operate in highly dependent sectors: International Consolidated Airlines Group S.A. (or simply, IAG), a holding company for several airlines including British Airways, Iberia and Air Europa, and Melia Hotels International S.A. (or MHI) a Spanish chain that operates more than 400 hotels throughout more than 40 countries. The example shows that the VaR of MHI asset log returns underestimated the risk compared to CoVaR (conditional on VaR for IAG asset log returns) during periods of the COVID-19 financial crisis.

6.1. The model

The copula models discussed so far assume a dependence structure constant over time. However, empirical studies of multivariate time series have revealed the existence of time-varying dependence among asset returns. We use the copula time series model introduced by Patton (2006b) to capture a possible dynamic dependence structure. This model assumes that the functional form of the copula remains fixed over the sample while the parameters vary according to some evolution equation. In order to estimate the parameters in this equation, we first need to specify the marginal distributions for the asset log returns. This model has been widely used to study the co-movements of asset returns in the framework of time-varying dependence structures (see, for example, Patton (2006a) Reboredo (2011, 2013) and Ji et al. (2019)).

We denote the negative log returns of IAG as the variable X_t and the negative log returns of MHI as the variable Y_t . Since log-returns may present the characteristics of autocorrelation and conditional heteroscedasticity, two ARMA-GARCH models are employed to describe their marginal distributions. The analyzes described in Section 6.3 support the following models for the two marginal series. The marginal distribution of IAG is specified as an MA(1)-GARCH(1, 1) model with skew t-innovations:

$$\begin{aligned} X_t &= \eta_t - \theta_1 \eta_{t-1} \\ \eta_t &= \sigma_{x,t} a_t, \quad \sigma_{x,t}^2 = \omega_x + \alpha_x \eta_{t-1}^2 + \beta_x \sigma_{x,t-1}^2, \\ \sqrt{\frac{v_x}{\sigma_{x,t}^2(v_x - 2)}} \eta_t &\sim \text{iid } t_{v_x} \end{aligned}$$

where a_t is a sequence of iid random variables with mean 0 and variance 1. Similarly, the marginal distribution of MHI is specified as a AR(5)-GARCH (1, 1) model with skew t-innovations as follows:

$$\begin{aligned} Y_t &= \phi_4 Y_{t-4} + \phi_5 Y_{t-5} + \varepsilon_t \\ \varepsilon_t &= \sigma_{y,t} b_t, \quad \sigma_{y,t}^2 = \omega_y + \alpha_y \varepsilon_{t-1}^2 + \beta_y \sigma_{y,t-1}^2, \\ \sqrt{\frac{v_y}{\sigma_{y,t}^2(v_y - 2)}} \varepsilon_t &\sim \text{iid } t_{v_y} \end{aligned}$$

where b_t is a sequence of iid random variables with mean 0 and variance 1.

Denote by $F_t(\cdot; \lambda_c)$ the conditional distribution function of the bivariate time series $\mathbf{X}_t = (X_t, Y_t)$, given the information set at time $t - 1$. By applying Sklar's theorem to the joint conditional distribution function, we have

$$F_t(x, y; \lambda) = C_t(F_t(x; \lambda_1), G_t(y; \lambda_2); \lambda_c), \quad (24)$$

where λ_1 and λ_2 are the parameters for the marginal conditional distributions, λ_c are the parameters for the conditional copula and $\lambda = (\lambda_1, \lambda_2, \lambda_c)$ are the parameters for the joint conditional distribution. For the applicability of the copula models studied in this paper, we considered a model with bounded PELCoV (the Frank model) and another with unbounded PELCoV (the Gaussian model) to perform the fits. The goodness-of-fit test for bivariate copulas based on White's information matrix equality (White (1982)) gave a p -value equal to 0.12 for the Frank copula and that based on Kendall process, as proposed by Wang and Wells (2000), gave a p -value equal to 0.14 for the Gaussian

copula. Although AIC showed that the Gaussian performed better than the Frank copula, we do not have enough evidence to reject these models at usual significance levels under the assumption of time-invariant dependence.

Following Patton (2006b), for each model, we specify the time-varying copula as follows. For the Gaussian copula, the linear correlation coefficient ρ_t evolves according to an ARMA(1,10)-type process:

$$\rho_t = \Lambda_1 \left(v_0 + v_1 \rho_{t-1} + v_2 \frac{1}{10} \sum_{j=1}^{10} \Phi^{-1}(u_{t-j}) \cdot \Phi^{-1}(v_{t-j}) \right)$$

where Φ^{-1} is the inverse of the standard normal distribution function, $\Lambda_1(x) = (1 - e^{-x}) / (1 + e^{-x})$ is the modified logistic transformation, designed to keep ρ_t in $(-1, 1)$ and $\lambda_c = (v_0, v_1, v_2)$. The model includes ρ_{t-1} as a regressor to capture any persistency in the dependence parameter and incorporates the intuition that the correlation should increase (decrease) if the transformed marginals have the same (opposite) sign. For the Frank copula, the driving mechanism for the dependence parameter is:

$$\theta_t = \Lambda_2 \left(\omega_0 + \omega_1 \theta_{t-1} + \omega_2 \frac{1}{10} \sum_{j=1}^{10} |u_{t-j} - v_{t-j}| \right)$$

where $\Lambda_2(x) = (1 + e^{-x})$ and $\lambda_c = (\omega_0, \omega_1, \omega_2)$. The choice of Λ_2 ensures that C_t is SSI. The dependence parameter equation includes θ_{t-1} as a regressor and a forcing variable based on the closeness of the most recent u_t and v_t .

6.2. Estimation and testing

The set of parameters λ_c of the copula is estimated by maximum likelihood, where the log-likelihood function of a random sample $(x_t, y_t)_{t=1}^n$ is, according to (24), given by

$$l(\lambda) = \sum_{t=1}^n \left\{ \log f_t(x_t; \lambda_1) + \log g_t(y_t; \lambda_2) + \log c_t(F_t(x_t; \lambda_1), G_t(y_t; \lambda_2); \lambda_c) \right\},$$

where f_t, g_t and c_t are the marginal conditional density functions and the conditional copula density function, respectively. We use a two-stage maximum likelihood estimation procedure based on maximizing the parameters separately for the marginals and the copula, see Joe and Xu (1996). At the first stage only the parameters from the marginal distributions are estimated. At the second stage, the dependence parameter is estimated from the copula likelihood by solving the following:

$$\hat{\lambda}_c = \arg \max_{\lambda_c} \sum_{t=1}^n \log c_t(\hat{u}_t, \hat{v}_t; \lambda_c),$$

where $\hat{u}_t = F_t(x_t; \hat{\lambda}_1)$ and $\hat{v}_t = G_t(y_t; \hat{\lambda}_2)$ are pseudo-sample observations from the copula. Under standard conditions, this procedure gives consistent and asymptotically normal estimates (see Joe (1997)).

6.3. Data analysis

We used adjusted weekly closing prices for IAG and MHI obtained from the public website <http://es.finance.yahoo.com> covering seven years, from January 8, 2016, to December 30, 2022, yielding $n = 363$ observations for each group. We considered negative log returns computed as $r_t = \log(p_{t-1}/p_t)$, where p_t and p_{t-1} are stock prices at weeks t and $t - 1$, respectively. Table 2 provides descriptive statistics for log returns series. The Shapiro-Wilk test strongly rejected the normality, and the assets log returns exhibit a positive excess kurtosis. The empirical correlation coefficient is 0.7034.

The parameters of the ARMA(p,q)-tGARCH(r,s) models were empirically determined by identifying the optimal models among alternative values using the Akaike Information Criterion (AIC). The Jarque-Bera test strongly rejected the normality of residuals. We used the Ljung-Box test of the squared standardized residuals to test the validity of

Table 2

Descriptive statistics for log returns. Ljung–Box test for the squared returns is computed with 20 lags.

	MHI	IAG
Mean	0.002081	0.003845
Std dev	0.057062	0.076719
Max	0.327792	0.513617
Min	-0.345877	-0.369389
Skewness	-0.020238	1.017383
Kurtosis	8.145324	10.414437
Shapiro-Wilk p -val	0.00000	0.000000
Pearson's r	0.7034985	
number observed	363	




Fig. 14. Scatter-plot of the empirical copula of the bivariate time series (X_t, Y_t) , given by $(F_t(x_t; \hat{\lambda}_1), G_t(y_t; \hat{\lambda}_2))$.

Table 3

Maximum likelihood estimates with asymptotic standard errors in parentheses of the parameters of the marginal distribution models for MHI and IAG log returns. Ljung–Box test for the squared residuals (R^2) and for standardized residuals (RS) are computed with 20 lags. Jarque–Bera tests the normality of residuals.

Mean equation			
MHI		IAG	
$\hat{\phi}_4$	-0.1604 (0.0520)	$\hat{\theta}_1$	0.1668(0.0481)
$\hat{\phi}_5$	0.0893 (0.0521)		
Variance equation			
$\hat{\alpha}_x$	0.1129(0.03869)	$\hat{\alpha}_y$	0.1129 (0.03869)
$\hat{\beta}_x$	0.8649 (0.03999)	$\hat{\beta}_y$	0.8649 (0.03999)
Ljung-Box R^2 p -val	0.88279		0.8827
Jarque-Bera p -val	0		0
Marginal models			
d.f. $\hat{u}_x$	2.96	d.f. $\hat{u}_y$	3.01
KS test p -val	0.5598		0.4357
Ljung-Box (RS) p -val	0.39		0.53

the volatility equation. The sample autocorrelation function (ACF) and p -values of Kolmogorov–Smirnov and Ljung–Box tests suggest that the models are adequate. Table 3 provides the parameter estimates for the marginal models. Fig. 14 shows the scatterplot of the empirical copula $(F_t(x_t; \hat{\lambda}_1), G_t(y_t; \hat{\lambda}_2))$.

6.4. Results

Given the bivariate time series (X_t, Y_t) , where X_t and Y_t describe the losses (negative log returns) of IAG and MHI, respectively, we have obtained the time-varying PELCoV $_v$ for the series X_t , $v \in (0, 1)$, under different dynamic copula models. Fig. 15 plots the time series X_t to-

gether with the $u_v(t)$ -quantiles for different values of v ($v_1 = 0.99, v_2 = 0.95$ and $v_3 = 0.8$). Given t , the $u_v(t)$ -quantile is $F_{X_t}^{-1}(u_v(t))$ (called PELCoV $_v$ -IAG-quantile in the plots). The left-hand plots show the quantiles associated with the PELCoV $_v$ s under the fitted Gaussian copula and the right-hand plots show the quantiles associated to the PELCoV $_v$ s and to their upper bounds u_t^* under the fitted Frank copula. The upper bounds depicted in the plots are given by $F_{X_t}^{-1}(u_t^*)$, where u_t^* is obtained from equation (23), where θ is replaced with θ_t and v takes the probability levels 0.99, 0.95 and 0.80, respectively. If X_t exceeds $F_{X_t}^{-1}(u_t^*)$, it is more prudent to determine reserves in terms of CoVaR for all $u_t > u_t^*$. The plots allow us to compare, for each t , the actual IAG loss (in terms of negative log returns) and the minimum loss above which CoVaR $_{v_i, u}$ [MHI|IAG] had been a more prudential reserve provision than VaR $_v$ [MHI].

X_t was always closer to its u_v -quantile (whatever the probability level u) under the Frank copula than the Gaussian copula. All the plots, except one (the plot in the top left corner), show that VaR $_v$ [MHI] underestimated the risk compared to CoVaR $_{v_i, u}$ [MHI|IAG] during the pandemic period.⁷ The COVID-19 pandemic significantly impacted many sectors of the economy, especially in the airlines and hotel industry, and materialized as a significant drop in the demand for passenger flights and accommodations. In these circumstances, it is not surprising that the value at risk underestimated the spillover effect.

7. Discussion

We have explored a method for monitoring systemic risk that can be used to determine provisions in the framework of a prudential regulation model. In particular, we have provided conditions that guarantee the ordering between VaR and CoVaR, allowing us to understand which, among the two measures, is more or less conservative than the other.

The prudential regulation strategy described is based on determining the PELCoV to decide whether to replace the VaR by the CoVaR. It is important to remark that, even if the PELCoV acts as a signal warning that the unconditional VaR underestimates the spillover effect, it does not capture itself such effect. In case of extreme positive dependence (comonotonicity), since $Y = F_Y^{-1}(F_X(X))$, it is clear that the PELCoV at level v (understood as in (9)) is equal to v . Therefore, when considering families of copulas where the convergence to comonotonicity is determined by the convergence of a parameter, we would expect that the PELCoV at level v converges to v in terms of the dependence parameter. This is in concordance with what we can observe in the families of copulas considered in this paper. For instance, considering the Gaussian, Gumbel Hoogard or Frank families of copulas, the PELCoV at level v tends to v when the underlying parameter tends to the comonotonicity case. However, in such a case, the PELCoV at level v is smaller (larger) than v and tends increasingly (decreasingly) to v when $v > \frac{1}{2}$ ($v < \frac{1}{2}$). On the other hand, the PELCoV is constant for the Farlie–Gumbel–Morgenstern and decreases in the dependence parameter considering the Ali–Mikhail–Haq family of copulas. The different behaviours of the PELCoV considering these families of copulas serve to illustrate that the monotonicity of the PELCoV is influenced by the dependence structure itself (not only by the intensity of the dependence) and the considered level of risk v .

An important problem that remains to be addressed concerns the non-parametric inference on the PELCoV. Reasoning as in Li and Wang (2022), a natural candidate to estimate the PELCoV, given $v \in (0, 1)$, is the u that solves the equation

$$\widehat{\text{CoVaR}}_{v, u}[Y|X] = \widehat{\text{VaR}}_v[Y], \quad (25)$$

⁷ Reversing the roles between MHI and IAG leads to a similar result in this period.




Fig. 15. Given the bivariate time series (X_t, Y_t) , the left-hand plots show the time series X_t (log-returns of IAG) together with their u_v -quantiles for $v = 0.99, 0.95$ and 0.80 , respectively, under the assumption of a time-varying Gaussian copula. The right-hand plots show the time series X_t , together with their u_v -quantiles for $v = 0.99, 0.95$ and 0.80 , respectively, and with their upper bounds (given by the u^* -quantiles) under the assumption of a time-varying Frank copula.

where $\widehat{\text{VaR}}_v[Y]$ and $\widehat{\text{CoVaR}}_{v,u}$ are the empirical VaR and CoVaR, respectively. The empirical VaR is the sample quantile, given by $\widehat{\text{VaR}}_v[Y] = X_{[i]}$, for $v \in \left(\frac{i-1}{n}, \frac{i}{n}\right]$, $i = 1, \dots, n$, where $X_{[1]} \leq X_{[2]} \leq \dots \leq X_{[n]}$, are the order statistics of the data $X_1, \dots, X_n$. Regarding CoVaR, there are in the literature different ways to estimate it. It can be obtained, for example, from models with time-varying second moments, by using maximum likelihood estimation or, as in Adrian and Brunnermeier (2016), using quantile regression. Unfortunately, the curve $u \mapsto \text{CoVaR}_{v,u}(Y|X)$ is not, in general, strictly increasing. Fixed v , the equation $\text{CoVaR}_{v,u}(Y|X) = \text{VaR}_v[Y]$ may not have a solution or have more than one and we cannot guarantee the existence or uniqueness of the solution of equation (25).

An alternative way to estimate the PELCoV is by solving the equation $\partial_1 \widehat{C}(u, v) = u$. The most straightforward way to estimate $\partial_1 C(u, v)$ would be by means of the Markov kernel, as in Fuchs and Trutschnig (2020). However, as before, we need additional assumptions to ensure that the curve $u \mapsto \partial_1 C(u, v)$ is strictly increasing. This is our theoretical SSI property, an assumption whose empirical verification is open to further study.

As suggested by one of the referees, in a similar way to the Probability Equivalent Level between VaR and CoVaR (PELCoV) defined and studied here, a probability equivalent level could be also considered between the risk measures ES (Expected Shortfall) and CoES (Conditional

Expected Shortfall); see, e.g., Sordo et al. (2018), for their definition and applications. Despite its analogous interpretation as an alert level, there is a significant difference between the new concept and the PELCoV: a probability equivalent level between ES and CoES does not only depend on the copula, but also on the marginals. Another difference is that, under the SSI assumption, we interpret the existence of an upper bound for this probability equivalent level in terms of the increasing convex order (instead of the stochastic order, as in the case of the PELCoV) between the random variables $[Y|X = \text{VaR}_u[X]]$ and Y . This will be a topic for future research.

CRedit authorship contribution statement

Patricia Ortega-Jiménez: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Supervision, Visualization, Writing – original draft, Writing – review & editing. **Franco Pellerey:** Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology, Supervision, Validation, Writing – original draft, Writing – review & editing, Project administration. **Miguel A. Sordo:** Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Resources, Supervision, Validation, Writing – original draft, Writing – review & editing. **Alfonso Suárez-Llorens:** Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology, Supervision, Visualization, Writing – original draft, Writing – review & editing.

sition, Investigation, Methodology, Project administration, Resources, Supervision, Validation, Writing – original draft, Writing – review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

We thank three anonymous reviewers for constructive and very helpful comments. POJ, MAS and AA acknowledge support received by Ministerio de Economía y Competitividad of Spain under grant PID2020-116216GB-I00 and from the 2014-2020 ERDF Operational Programme and the Department of Economy, Knowledge, Business and University of the Regional Government of Andalusia under grant FEDER-UCA18-107519. POJ acknowledges University of Cádiz for the PhD grant (call 2018) linked to the project MTM2017-89577-P and funding from the FWO and F.R.S.-FNRS and under the Excellence of Science (EOS) programme, project ASTERISK (40007517). FP is partially supported by the GNAMPA research group of INDAM (Istituto Nazionale Di Alta Matematica), Italy, and by the grant Progetto di Eccellenza, CUP: E11G18000350001, Italy.

Appendix A. Additional computations

This appendix gives the details in the computation of $u^* = \lim_{v \rightarrow 1} u_v$ for the Frank family of copulas, whose density function is given by

$$c(u, v) = \frac{e^{(1+u+v)\theta} (e^\theta - 1) \theta}{(e^{(u+v)\theta} - e^\theta (e^{u\theta} + e^{v\theta} - 1))^2}.$$

To find the analytical form of the PELCoV_v and the corresponding upper bound, let us first note that

$$\partial_1 C(u, v) = \frac{e^\theta (1 - e^{\theta v})}{e^{\theta(u+v)} + e^\theta (1 - e^{\theta u} - e^{\theta v})},$$

and that $\partial_1 C(u, v) > 0$ for $\theta > 0$, so that the copula is SSI for $\theta > 0$. Moreover, we can see that $\partial_1 C(u, v) \leq v$ if and only if

$$\frac{e^\theta (1 - e^{\theta v})}{v} \leq e^{\theta u} e^{\theta v} + e^\theta - e^{\theta u} e^\theta - e^{\theta(1+v)},$$

that is, if $\frac{e^\theta (1 - e^{\theta v})}{v} - e^\theta + e^{\theta(1+v)} \leq e^{\theta u} (e^{\theta v} - e^\theta)$, which holds when

$$\frac{1}{\theta} \ln \left(\frac{e^\theta (1 - e^{\theta v}) - v e^\theta + v e^{\theta(1+v)}}{v (e^{\theta v} - e^\theta)} \right) \geq u.$$

Therefore, the PELCoV_v, u_v , is given by:

$$u_v = \frac{1}{\theta} \ln \left(\frac{e^\theta (1 - e^{\theta v}) (1 - v)}{v (e^{\theta v} - e^\theta)} \right).$$

Then, for $\theta > 0$, u_v increases in v if $\frac{(1 - e^{\theta v})(1 - v)}{v(e^{\theta v} - e^\theta)}$ increases in v . Considering $f(v) = (1 - e^{\theta v})(1 - v)$ and $g(v) = v(e^{\theta v} - e^\theta)$, by L'Hôpital's Rule of Monotonicity (Anderson et al., 1997), it would suffice to observe that $\frac{f'(v)}{g'(v)}$ increases in $v \in (0, 1)$. To this aim, note that, given $\theta > 0$,

$$\frac{d}{dv} \left(\frac{f'(v)}{g'(v)} \right) = \frac{e^{\theta v} \theta (2 + e^{\theta v} \theta + v \theta + e^\theta (-2 + \theta - v \theta))}{(e^\theta - e^{\theta v} (1 + v \theta))^2}$$

is positive for all $v \in (0, 1)$ if

$$h(v) = 2 + e^{\theta v} \theta + v \theta + e^\theta (-2 + \theta - v \theta)$$

is positive for all $v \in (0, 1)$. One can easily check that $h(v)$ is continuous and derivable and that $h'(v) = \theta - e^\theta \theta + e^{\theta v} \theta^2$, which is zero for $v_0 = \frac{1}{\theta} \ln \left(\frac{e^\theta - 1}{\theta} \right)$. Since $h''(v_0) = (e^\theta - 1) \theta^2 > 0$, v_0 is a minimum. Therefore, it suffices to verify that $h(0)$, $h(1)$ and $h(v_0)$ are positive when $\theta > 0$. It is easy to see that $h(0) = h(1) = 2 + e^\theta (\theta - 2) + \theta > 0$ and that $h(v_0) = 1 + e^\theta (\theta - 1) + (e^\theta - 1) \ln \left(\frac{\theta}{e^\theta - 1} \right)$. As $\ln(x) > 1 - x$ for all $x \in (0, 1)$, letting $x = \frac{\theta}{e^\theta - 1}$, one gets that

$$h(v_0) > 1 + e^\theta (\theta - 1) + (e^\theta - 1) \left(1 - \frac{\theta}{e^\theta - 1} \right) = (e^\theta - 1) \theta.$$

Thus, for the Frank copula, u_v is increasing with respect to v . In order to find u^* , one should only consider the limit when v tends to 1.

Using continuity and L'Hôpital's result, as $u_v = \frac{1}{\theta} \ln \left(\frac{e^\theta (1 - e^{\theta v})(1 - v)}{v(e^{\theta v} - e^\theta)} \right)$, it follows

$$\begin{aligned} \lim_{v \rightarrow 1} u_v &= \frac{1}{\theta} \cdot \ln \left(\lim_{v \rightarrow 1} \frac{e^\theta (1 + e^{\theta v} (-1 + \theta - v \theta))}{e^\theta - e^{\theta v} (1 + v \theta)} \right) \\ &= \frac{1}{\theta} \ln \left(\frac{e^\theta (1 - e^\theta)}{e^\theta (1 - 1 - \theta)} \right) \\ &= \frac{1}{\theta} \ln \left(\frac{e^\theta - 1}{\theta} \right). \end{aligned}$$

Therefore, $u^* = \frac{1}{\theta} \ln \left(\frac{e^\theta - 1}{\theta} \right)$.

References

- Acharya, V., Pedersen, L., Philippon, T., Richardson, M., 2017. Measuring systemic risk. *The Review of Financial Studies* 30 (1), 2–47.
- Acharya, V.V., 2009. A theory of systemic risk and design of prudential bank regulation. *Journal of Financial Stability* 5 (3), 224–255.
- Adrian, T., Brunnermeier, M.K., 2016. CoVaR. *American Economic Review* 106 (7), 1705.
- Anderson, G.D., Vamanamurthy, M.K., Vuorinen, M., 1997. Conformal Invariants, Inequalities and Quasiconformal maps. John Wiley and Sons, New York.
- Arias-Nicolás, J.P., Fernández-Ponce, J.M., Luque-Calvo, P., Suárez-Llorens, A., 2005. Multivariate dispersion order and the notion of copula applied to the multivariate t-distribution. *Probability in the Engineering and Informational Sciences* 19 (3), 363–375.
- Artzner, P., Delbaen, F., Eber, J.M., Heath, D., 1999. Coherent measures of risk. *Mathematical Finance* 9 (3), 203–228.
- Barczy, M., Nedényi, F.K., Sütő, L., 2023. Probability equivalent level of value at risk and higher-order expected shortfalls. *Insurance: Mathematics and Economics* 108, 107–128.
- Belzunce, F., Ruiz, J.M., Suárez-Llorens, A., 2008. On multivariate dispersion orderings based on the standard construction. *Statistics & Probability Letters*, 271–281.
- Benoit, S., Colliard, J.E., Hurlin, C., Pérignon, C., 2017. Where the risks lie: a survey on systemic risk. *Review of Finance* 21 (1), 109–152.
- Bernardi, M., Durante, M., Jaworski, P., 2017. CoVaR of families of copulas. *Statistics & Probability Letters* 120, 8–17.
- Bickel, P.J., Lehmann, E.L., 1979. Descriptive Statistics for Nonparametric Models (IV. Spread). Jureckova, Prague, Czech Republic.
- Bisias, D., Flood, M., Lo, A.W., Valavanis, S., 2012. A survey of systemic risk analytics. *Annual Review of Financial Economics* 4 (1), 255–296.
- Brownlees, C., Engle, R.F., 2017. SRISK: a conditional capital shortfall measure of systemic risk. *The Review of Financial Studies* 30 (1), 48–79.
- Cai, J., Wei, W., 2012. On the invariant properties of notions of positive dependence and copulas under increasing transformations. *Insurance: Mathematics and Economics* 50 (1), 43–49.
- Chen, C., Iyengar, G., Moallemi, C.C., 2013. An axiomatic approach to systemic risk. *Management Science* 59 (6), 1373–1388.
- Das, B., Fassen-Hartmann, V., 2018a. Risk contagion under regular variation and asymptotic tail independence. *Journal of Multivariate Analysis* 165, 194–215.
- Denuit, M., Dhaene, J., Goovaerts, M., Kaas, R., 2005. Actuarial Theory for Dependent Risks: Measures, Orders and Models. John Wiley & Sons.
- Dhaene, J., Laeven, R.J.A., Zhang, Y., 2022. Systemic risk: conditional distortion risk measures. *Insurance: Mathematics and Economics* (ISSN 0167-6687) 102, 126–145.
- Di Clemente, A., 2018. Estimating the marginal contribution to systemic risk by a CoVaR-model based on copula functions and extreme value theory. *Economic Notes: Review of Banking, Finance and Monetary Economics* 47 (1), 69–112.

- Drehmann, M., Tarashev, N.A., 2011. Systemic importance: some simple indicators. *BIS Quarterly Review*, March. Available at SSRN: <https://ssrn.com/abstract=1785264>.
- Fernandez-Ponce, J.M., Suárez-Llorens, A., 2003. A multivariate dispersion ordering based on quantiles more widely separated. *Journal of Multivariate Analysis* 85 (1), 40–53.
- Fernández-Ponce, J.M., Pellerey, F., Rodríguez-Griñolo, M.R., 2011. A characterization of the multivariate excess wealth ordering. *Insurance: Mathematics and Economics* (ISSN 0167-6687) 49 (3), 410–417.
- Fiori, A.M., Rosazza Gianin, E., 2022. Generalized PELVE and applications to risk measures. *European Actuarial Journal*.
- Fuchs, S., Trutschnig, W., 2020. On quantile based co-risk measures and their estimation. *Dependence Modeling* 8 (1), 396–416.
- Girardi, G., Ergün, A.T., 2013. Systemic risk measurement: multivariate GARCH estimation of CoVaR. *Journal of Banking & Finance* 37, 3169–3180.
- Glasserman, P., Young, H.P., 2016. Contagion in financial networks. *Journal of Economic Literature* 54 (3), 779–831.
- Huang, X., Zhou, H., Zhu, H., 2009. A framework for assessing the systemic risk of major financial institutions. *Journal of Banking & Finance* 33 (11), 2036–2049.
- Jaworski, P., 2017. On Conditional Value at Risk (CoVaR) for tail-dependent copulas. *Dependence Modeling* 5, 1–17.
- Ji, Q., Liu, B.Y., Fan, Y., 2019. Risk dependence of covar and structural change between oil prices and exchange rates: a time-varying copula model. *Energy Economics* 77, 80–92.
- Jin, X., 2018. Downside and upside risk spillovers from China to Asian stock markets: a CoVaR-copula approach. *Finance Research Letters* 25, 202–212.
- Joe, H., 1997. *Multivariate Models and Multivariate Dependence Concepts*. CRC Press.
- Joe, H., Xu, J.J., 1996. The Estimation Method of Inference Functions for Margins for Multivariate Models.
- Jorion, P., 2000. *Value at Risk*. McGraw-Hill Professional Publishing.
- Karimalis, E.N., Nomikos, N.K., 2018. Measuring systemic risk in the European banking sector: a copula CoVaR approach. *European Journal of Finance* 24 (11), 944–975.
- Kupiec, P., 2002. Stress testing in a value at risk framework. In: Dempster, M. (Ed.), *Risk Management: Value at Risk and Beyond*, pp. 76–99.
- Lehar, A., 2005. Measuring systemic risk: a risk management approach. *Journal of Banking & Finance* 29 (10), 2577–2603.
- Lehmann, E.L., 1966. Some concepts of dependence. *The Annals of Mathematical Statistics* 37 (5), 1137–1153.
- Li, H., Wang, R., 2022. PELVE: probability equivalent level of VaR and ES. *Journal of Econometrics* (ISSN 0304-4076).
- Mainik, G., Schaanning, E., 2014. On dependence consistency of CoVaR and some other systemic risk measures. *Statistics & Risk Modeling* 31 (1), 49–77.
- Müller, A., Stoyan, D., 2002. *Comparison Methods for Stochastic Models and Risks*. Wiley Series in Probability and Statistics. New York.
- Navarro, J., Sordo, M.A., 2018. Stochastic comparisons and bounds for conditional distributions by using copula properties. *Dependence Modeling* 6 (1), 156–177.
- Navarro, J., Pellerey, F., Sordo, M.A., 2020. Weak dependence notions and their mutual relationships. *Mathematics* 9 (1), 81.
- Nelsen, R.B., 2007. *An Introduction to Copulas*. Springer Science & Business Media.
- Ortega-Jiménez, P., Sordo, M.A., Suárez-Llorens, A., 2021. Stochastic orders and multivariate measures of risk contagion. *Insurance: Mathematics and Economics* 96, 199–207.
- Patton, A.J., 2006a. Estimation of multivariate models for time series of possibly different lengths. *Journal of Applied Econometrics* 21 (2), 147–173.
- Patton, A.J., 2006b. Modelling asymmetric exchange rate dependence. *International Economic Review* 47 (2), 527–556.
- Reboredo, J.C., 2011. How do crude oil prices co-move?: a copula approach. *Energy Economics* 33 (5), 948–955.
- Reboredo, J.C., 2013. Is gold a safe haven or a hedge for the us dollar? Implications for risk management. *Journal of Banking & Finance* 37 (8), 2665–2676.
- Reboredo, J.C., Ugolini, A., 2015. Systemic risk in European sovereign debt markets: a CoVaR-copula approach. *Journal of International Money and Finance* 51, 214–244.
- Shaked, M., Shanthikumar, J.G., 2007. *Stochastic Orders*. Springer Science & Business Media.
- Sordo, M.A., Suárez-Llorens, A., Bello, A., 2015. Comparison of conditional distributions in portfolios of dependent risks. *Insurance: Mathematics and Economics* 61, 62–69.
- Sordo, M.A., Bello, A., Suárez-Llorens, A., 2018. Stochastic orders and co-risk measures under positive dependence. *Insurance: Mathematics and Economics* 78, 105–113.
- Wang, W., Wells, M.T., 2000. Model selection and semiparametric inference for bivariate failure-time data. *Journal of the American Statistical Association* 95 (449), 62–72.
- White, H., 1982. Maximum likelihood estimation of misspecified models. *Econometrica*, 1–25.
- Zhao, S., 2022. Systemic risk measurement: a limiting threshold copula approach to CoVaR. *Computers & Industrial Engineering* 171, 108464.