

HISTOLOGY AND HISTOPATHOLOGY

ISSN: 0213-3911
e-ISSN: 1699-5848

Submit your article to this Journal (<http://www.hh.um.es/Instructions.htm>)

Optimal human ovarian follicle isolation: A review focused on enzymatic digestion

Authors: Farnaz Tajbakhsh, Somayeh Tavana, Mohammad Kazemi Ashtiani, Christiani Andrade Amorim and Rouhollah Fathi

DOI: 10.14670/HH-18-747

Article type: REVIEW

Accepted: 2024-04-12

Epub ahead of print: 2024-04-12

This article has been peer reviewed and published immediately upon acceptance.
Articles in "Histology and Histopathology" are listed in Pubmed.
Pre-print author's version

Optimal human ovarian follicle isolation: A review focused on enzymatic digestion

Farnaz Tajbakhsh^{1,2}, Somayeh Tavana¹, Mohammad Kazemi Ashtiani³, Christiani Andrade Amorim⁴, Rouhollah Fathi¹

1. Department of Embryology, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran

2. Department of Developmental Biology, University of Science and Culture, Tehran, Iran

3. Department of Cell Engineering, Cell Science Research Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, Iran

4. Pôle de Recherche en Gynécologie, Institut de Recherche Expérimentale et Clinique, Université Catholique de Louvain, Brussels, Belgium

* **The Corresponding Email:** rfathi79@royaninstitute.org

Keywords: human ovarian tissue, follicle isolation, enzymatic isolation, artificial ovary, live/dead.

Short title: Enzymatic isolation of the human ovarian follicle

Abstract

The damage or depletion of ovarian reserves due to aging or cancer treatment can increase the need for fertility preservation techniques. One of the most common ways of supporting fertility in prepubertal girls and women who require immediate cancer treatment is through ovarian tissue cryopreservation and re-transplantation following cancer treatment. However, a more appropriate method should be employed in diseases such as leukemia, where malignant cells may be present in cryopreserved tissue, instead of ovarian tissue transplantation. Human ovarian follicle isolation for *in vitro* culture or the use of artificial ovaries for their growth can decrease the risk of reintroducing cancer cells into these individuals. Here we review the methods for the isolation of human ovarian follicles.

List of abbreviations

POI: Premature Ovarian Insufficiency

BM: Basement Membrane

GMP: Good Manufacturing Practices

DH: Dispace High

TM: Thermolysin Medium

TDE: Tumor Dissociation Enzyme

Introduction

Over the last few decades, there has been an increase in cancer incidence among women, with around 20 percent of women of childbearing age being affected (Maltaris et al., 2006; Akahori et al., 2019). Although anticancer treatments, including chemotherapy, radiation therapy, and other methods, have increased survival rates, the potential complications of these treatments were not fully understood until around 20 years ago (Meirow and endocrinology, 2000; Donnez et al., 2006; Resetkova et al., 2013; Smith et al., 2014). The ovaries are sensitive organs, and cytotoxic drugs such as alkylating agents and radiotherapy can damage a patient's reproductive ability and increase the risk of premature ovarian insufficiency (POI) (Delbaere, and Englert, 2009; Peddie et al., 2012; Demeestere et al., 2017; Weners et al., 2017; Buckenmeyer et al., 2020).

Fertility preservation is recommended when there is a high risk of POI in cancer patients. Among the different strategies available, ovarian tissue cryopreservation is the only one indicated for prepubertal girls or women who need to start cancer treatment as soon as possible and may not have sufficient time to undergo embryo or oocyte cryopreservation protocols (Wang et al., 2016; Donnez and Dolmans, 2017; Andersen et al., 2019; Silvestris et al., 2020).

Primordial and primary follicles found in the cortex area are preserved during ovarian tissue cryopreservation. Primordial follicles make up the majority of follicles in the ovarian cortex and are referred to as the ovarian reserve (Wallace and Kelsey, 2010). Furthermore, the small size, low metabolic rate, and absence of zona pellucida and cortical granules make primordial follicles more resistant to cryopreservation injuries (Gerritse et al., 2008). These small follicles are surrounded by collagen IV, laminin, and fibronectin (Heeren et al., 2015; Mouloungui et al., 2018).

In recent years, more than 200 live births have been achieved after the transplantation of cryopreserved ovarian tissue (Dolmans et al., 2020). However, the most critical concern about ovarian tissue transplantation is the possibility of cancer cells in the tissue and the recurrence of cancer. This issue is particularly relevant for cancers such as leukemia, Burkitt lymphoma, and neuroblastoma, which pose a higher risk (Dolmans, 2021). Therefore, researchers have been developing effective methods to isolate

viable follicles from cancer cells, either to grow them *in vitro* or to transplant them into an artificial ovary, to minimize the risk of cancer recurrence.

The purpose of this study was to review and compare the human ovarian follicle isolation methods.

Isolation of human preantral follicles

Why should we isolate ovarian follicles?

Acute lymphoblastic and myeloid leukemias are the most commonly occurring cancers in prepubertal girls (Ward et al., 2014). For these patients, ovarian tissue transplantation can increase the risk of cancer recurrence (Rosendahl et al., 2010). However, there are techniques available to identify cancer cells in the ovarian tissue, including real-time quantitative polymerase chain reaction, multicolor flow cytometry, and ovarian tissue xenografts (Mouloungui et al., 2018). Moreover, the basement membrane around the follicles serves as a physical barrier that protects them against malignant cells (Irving-Rodgers and Rodgers, 2006). Therefore, follicle isolation using approaches such as artificial ovaries (Soares et al., 2017; Chiti et al., 2018; Chen et al., 2021) or *in vitro* culture (Dolmans et al., 2006; McLaughlin et al., 2018; Wiweko et al., 2019) can be helpful for this group of patients.

How should we isolate ovarian follicles?

Numerous studies have investigated different techniques for isolating preantral follicles in humans, which are generally classified as either mechanical or enzymatic methods. In mechanical processes, ovarian fragments are cut using a tissue chopper or a sharp tool (e.g., scissors or scalpel) followed by dissection with needles. Although mechanical isolation is a safe method, human ovaries have a dense and fibrous structure, and using mechanical methods alone can be a hard and time-consuming approach. In one study, Langbeen et al. applied an Ultra Turrax T18 basic homogenizer instead of a tissue chopper in three ruminant species (bovine, caprine, and ovine) to decrease the length of the isolation procedure (Langbeen et al., 2015). In another study, some mechanical isolation methods including micro-dissection, tissue chopping followed by filtration, mincing, trypsinization and sonication were combined (Sharma et al., 2009).

However, this technique was not used for human follicle isolation. Therefore, it seems that a combination of mechanical and enzymatic methods is typically employed to successfully isolate a substantial number of follicles. To the best of our knowledge, the only study comparing mechanical isolation followed by enzymatic digestion with that of mechanical isolation alone was performed by Dong et al. (Dong et al., 2014). The authors reported that the mechanical/enzymatic isolation method yielded a higher survival rate for primordial follicles after ten days of *in vitro* culture, while the survival rate of secondary follicles was higher with mechanical isolation alone. Interestingly, they found that culturing ovarian tissue for six days before follicle isolation led to larger follicles and higher levels of estradiol than with mechanical isolation alone. They explained that this was due to the ovarian tissue becoming looser after the six-day culture, allowing for easier follicle isolation with the mechanical method (Dong et al., 2014).

Considering the composition of preantral follicles, the basement membrane (BM) can be important in choosing the best enzyme. The follicular BM is made of laminin, collagenase IV, and fibronectin, and it maintains the structure of the developing follicle, protects the cellular structure, and regulates follicle growth and development by also regulating hormones and growth factors. These BM can be remodeled during follicle development (Heeren et al., 2015). On the other hand, BM can support capillaries, white blood cells, and nerve processes from the follicle compartment (Donnez et al., 2006). Therefore, it is essential to use an appropriate enzyme to isolate more viable follicles with a healthy BM. Enzymes, including various types of collagenases and liberases, and more recently, tumor dissociation enzyme (TDE), have been utilized over the years in numerous studies aimed at isolating human preantral follicles. During enzymatic digestion, sticky DNA ends released from damaged cells act as glue, causing isolated follicles to attach to other cells or remaining fragments of ovarian tissue, which can prevent the harvesting of fully isolated follicles (Wang et al., 2007). To circumvent this problem, DNase I is added to the enzymatic solution. All enzymes are produced under good manufacturing practices (GMP) for clinical practice.

Which materials could be useful to isolate human ovarian follicles?

Collagenase

Various types of collagenase (Ia, II, IX, XI) derived from *Clostridium histolyticum* can be used for isolating follicles, with or without DNase I (Lierman et al., 2015).

In 1993, Roy and Treacy (Roy and Treacy, 1993) reported the first successful isolation of primordial follicles from human ovaries. They dissected the ovarian tissue with a razor blade and then digested the fragments using a combination of collagenase type I and DNase. The authors reported that this protocol yielded a large number of isolated follicles, which were able to grow *in vitro*, as evidenced by their increased size and synthesis of estradiol, progesterone, and androstenedione after five days of culture (Roy and Treacy, 1993). It should be mentioned that shorter exposure to the enzyme, with lower or higher amounts of enzyme was not effective. Oktay et al. (Oktay et al., 1997) achieved high efficiency and viability rates by using collagenase type IA in combination with the mechanical method for partial disaggregation of ovarian tissue before or after cryopreservation. However, electron microscopy revealed some damage following isolation and cryopreservation (Oktay et al., 1997). Several studies have utilized various types of collagenases for partial or complete isolation of human preantral follicles (Figueiredo et al., 1993; Hovatta et al., 1999; Martinez-Madrid et al., 2004; Xu et al., 2009; Chen et al., 2021). For example, collagenase NB6, which is a combination of collagenase I and II frequently employed for isolating human umbilical cord cells, was used by Mouloungui et al. to isolate human ovarian follicles (Mouloungui et al., 2018). They isolated a high number of viable primordial follicles with an intact BM. Belen Martinez-Madrid et al. also applied collagenase IA for primordial follicle isolation (Martinez-Madrid et al., 2004). They used a Ficoll density gradient to recover isolated follicles and discovered that this gradient can minimize the manipulation time and maintain high follicular viability.

While collagenase is a commonly used enzyme for ovarian follicle isolation, it contains a large amount of endotoxin that can negatively impact follicle quality, in addition to the batch-to-batch variation that can affect reproducibility (Lierman et al., 2015). Furthermore, it has been reported that collagenase can cause damage to the follicle's basal lamina (Hovatta et al., 1999; Dong et al., 2014). To address these concerns, a

purified collagenase blend or Liberase were used as alternatives for follicle isolation. Dolmans et al. compared Liberase Blendzyme 3 and collagenase type IA and found that the use of the former could be a suitable alternative to the latter (Dolmans et al., 2006, 2007, 2008). Due to using Liberase Blendzyme 3, not only was the follicular ultrastructure preserved but more than 86% of isolated follicles were viable, with sufficient capacity to grow *in vitro* or *in vivo* (Dolmans et al., 2006, 2008).

Liberase DH

Liberase DH (Dispase High) is a type of enzyme blend that contains collagenase I, collagenase II, a high concentration of Dispase, and a non-clostridial neutral protease. Studies comparing Liberase and collagenase for isolating preantral follicles have shown that Liberase DH produces better results in terms of the number and quality of preantral follicles, survival rates, *in vitro* culture outcomes, and structure of follicles (Vanacker et al., 2011). Mouloungui et al. (Mouloungui et al., 2018) conducted a comparison of collagenases NB6 and IA with Liberase DH for follicle isolation. However, their results did not show any significant difference in the number and survival rate of the isolated follicles using the three enzymes (Mouloungui et al., 2018).

Wang et al. (Wang et al., 2016) used a combination of Liberase DH and DNase I to achieve the highest quality isolation and *in vitro* culture of follicles. Chiti et al. (Chiti et al., 2017) also used Liberase DH and DNase I in their study to enhance follicle isolation. Interestingly, they adapted the duration of the incubation in the enzymatic solution and reported that interrupting the enzymatic digestion every 30 min to retrieve the isolated follicles resulted in a greater number and better quality of isolated follicles compared with incubating the tissue in the enzymatic solution for 75 min. Moreover, this modification allows for the length of the isolation procedure to be adjusted based on the fibrous nature of the ovarian tissue (Chiti et al., 2017).

Liberase TM

Liberase TM (Thermolysin Medium) is a type of Liberase that contains collagenase I, collagenase II, and a medium concentration of Thermolysin (Schmidt et al., 2018). Two studies by Kristensen et al. (Kristensen

et al., 2011) and Yin et al. (Yin et al., 2016) used Liberase TM to isolate human preantral follicles from the ovarian medulla. While both groups used a mixture of 0.04 mg/ml Liberase TM and 0.2 mg/ml collagenase IV in warm McCoy's medium to isolate the follicles, Yin et al. added a 0.2 mg/ml DNase I to the solution. A comparison study by Lierman et al. shows that there was no significant difference between the number and type of follicles isolated with the low concentration of Liberase DH and Liberase TM combined with collagenase IV if they ignored the maturation status of the isolated follicles. However, Liberase DH and Liberase TM with collagenase IV work better than collagenase IV alone, while using collagenase IV is sufficient to isolate high-quality secondary follicles (Lierman et al., 2015).

Tumor Dissociation Enzyme (TDE)

Studies aimed at improving the isolation of human preantral follicle isolation have continued, with researchers such as Schmidt et al. (Schmidt et al., 2018) and Chen et al. (Chen et al., 2021) using a commercial enzyme cocktail called TDE to digest ovarian tissue and comparing it to Liberase TM and Liberase DH combined with DNase I, respectively. TDE is a highly purified enzyme that contains collagenase, protease, Dispase, and DNase, which aid the isolation of many follicles and prevent the digested tissue suspensions from becoming sticky (Chen et al., 2021; Schmidt et al., 2018). According to these studies, TDE is an excellent alternative to different Liberases and is effective in reducing oxidative stress while maintaining follicle morphology. Additionally, the survival rate of follicles isolated with TDE can be higher than with Liberase and it decreases the risk of cancer cell reseeding (Chen et al., 2021; Schmidt et al., 2018). Furthermore, the apoptosis rate does not increase with TDE (Schmidt et al., 2018).

Discussion

Based on various studies, using a mechanical method to isolate secondary follicles in a two-step culture is sufficient, however, in the case of isolated follicle culture or transplantation, it is better to obtain both enzymatic and mechanical procedures (McLaughlin et al., 2018; Schmidt et al., 2018).

Promising results have been achieved in the continued study of various enzymes for the isolation of human ovarian follicles. Among the first enzymes used for this purpose were those of the collagenase family (Figueiredo et al., 1993; Oktay et al., 1997; Martinez-Madrid et al., 2004). In addition, the Liberase family and TDE (Schmidt et al., 2018) were employed for follicle isolation. Although, maintaining follicular structure is a crucial subject, obtaining healthy follicles with minimum damage to granulosa cells (GCs) and oocytes is very important (Dolmans et al., 2006). Therefore, choosing the best enzyme is essential. For example, the BM contains collagen IV (Heeren et al., 2015), and the enzyme collagenase IV can harm this barrier and reduce follicle quality (Kristensen et al., 2011). On the other hand, primordial follicles have a tighter contact with surrounding tissues but this connection will become looser over follicle development (Hovatta et al., 1999). It seems that complete separation of primordial follicles during an isolation procedure may need a stronger enzyme, like collagenase IA; however, to recover the isolated follicles and select the viable ones, a Ficoll density gradient can be helpful (Kristensen et al., 2011). Most human studies used the Liberase family for preantral follicle isolation (Dolmans et al., 2008; Paulini et al., 2016).

In all studies, whether follicle isolation was for *in vitro* culture or for transplantation, assessing the viability of isolated follicles is crucial. To check this issue, Neutral Red is the best option to help isolate healthy viable follicles faster (Nemes et al., 1979). Nevertheless, using calcein AM and ethidium homodimer I can help classify isolated follicles based on their cell and oocyte damage by exhibiting a green and red fluorescent color (Paulini et al., 2016). Viability assessment of human follicles isolated by the Liberase enzyme blend (Dolmans et al., 2006) or Liberase DH shows these enzymes did not impair the isolated follicles (Paulini et al., 2016). Our study (unpublished data) also confirmed the viability of isolating human follicles by collagenase IA. Although, collagenase I is a common enzyme for follicle isolation, its effect has not been thoroughly investigated (Gosden, 1990; Telfer and Gosden, 1990). It seems that partial follicle isolation by collagenase can lead to premature oocyte extrusion (Hovatta et al., 1999). Comparison between Liberases and collagenase confirms that although the number of fully isolated follicles is lower than with collagenase, it can protect the ultrastructure and morphology of the follicles. Because collagenase is a stronger enzyme and TEM analysis shows that it can harm follicle structures such as the follicular wall

(Dolmans et al., 2006). Even so, there are different comparisons between various Liberase enzymes and it seems that Liberase TM is a better option for follicle isolation (Buckenmeyer et al., 2020). Also, TDE can be used instead of Liberase TM and Liberase DH; apoptosis and necrosis were not observed after follicle isolation (Schmidt et al., 2018; Chen et al., 2021). However, there is speculation that the isolation process, regardless of the enzyme type, can impact cell-cell interactions within the follicle (Smith et al., 2014). Furthermore, comparison of some enzymatic protocols for follicular isolation shows that the maturation status of follicles plays a key role in response to enzymatic isolation (Lierman et al., 2015). As a result, there is a lack of comprehensive studies regarding the effects of common enzymes on isolated follicle structure and their consequences on further follicular *in vitro* culture or transplantation.

Conclusion

To conclude, in dense ovaries, such as human, Liberase TM is the best option for human ovarian follicular isolation. The Liberase family can maintain the follicular ultrastructure and reduce damage to GCs. On the other hand, TDE can be an alternative enzyme to decrease the apoptosis and necrosis of human ovarian follicles in enzymatic digestion.

Acknowledgment

All authors in the Cryobioengineering team of the Royan Institute acknowledge financial support from the Council for Development of Regenerative Medicine and Stem Cells (Grant. 11/54216).

Conflict of interest statement

The authors of this article have no conflict of interest.

References:

- Akahori T., Woods D.C. and Tilly J.L. (2019). Female fertility preservation through stem cell-based ovarian tissue reconstitution *in vitro* and ovarian regeneration *in vivo*. Clin. Med. Insights Reprod. Health 13, 1179558119848007.
- Andersen C.Y., Mamsen L.S. and Kristensen S.G. (2019). FERTILITY PRESERVATION: Freezing of ovarian tissue and clinical opportunities. Reproduction 158, F27-F34.
- Buckenmeyer M.J., Sukhwani M., Iftikhar A., Nolfi A.L., Xian Z., Dadi S., Case Z.W., Steimer S.R., D'Amore A., Orwig K.E. and Brown B.N. (2023). A bioengineered in situ ovary (ISO) supports follicle engraftment and live-births post-chemotherapy. J. Tissue Eng. 14, 20417314231197282.
- Chen J., Isachenko E., Wang W., Du X., Wang M., Rahimi G., Mallmann P. and Isachenko V. (2022). Optimization of follicle isolation for bioengineering of human artificial ovary. Biopreserv. Biobank. 20, 529-539.
- Chiti M.C., Dolmans M.-M., Hobeika M., Cernogoraz A., Donnez J. and Amorim C.A.J. (2017). A modified and tailored human follicle isolation procedure improves follicle recovery and survival. J. Ovarian Res. 10, 1-9.
- Chiti M.C., Dolmans M.-M., Mortiaux L., Zhuge F., Ouni E., Shahri P.A.K., Ruymbeke E.V., Champagne S.-D., Donnez J. and Amorim C.A. (2018). A novel fibrin-based artificial ovary prototype resembling human ovarian tissue in terms of architecture and rigidity. J. Assist. Reprod. Genet. 35, 41-48.
- Demeestere I., Simon P., Emiliani S., Delbaere A. and Englert Y. (2009). Orthotopic and heterotopic ovarian tissue transplantation. Hum. Reprod. Update 15, 649-665.
- Dolmans M.-M., Michaux N., Camboni A., Martinez-Madrid B., Van Langendonck A., Annarita Nottola S. and Donnez J. (2006). Evaluation of Liberase, a purified enzyme blend, for the isolation of human primordial and primary ovarian follicles. Hum. Reprod. 21, 413-420.
- Dolmans M.M., Martinez-Madrid B., Gadisseux E., Guiot Y., Yuan W.Y., Torre A., Camboni A., Van Langendonck A. and Donnez J. (2007). Short-term transplantation of isolated human ovarian follicles and cortical tissue into nude mice. Reproduction 134, 253-262.
- Dolmans M.-M., Yuan W.Y., Camboni A., Torre A., Van Langendonck A., Martinez-Madrid B. and Donnez J. (2008). Development of antral follicles after xenografting of isolated small human preantral follicles. Reprod. Biomed. Online 16, 705-711.
- Dolmans M.-M., Falcone T. and Patrizio P. (2020). Importance of patient selection to analyze in vitro fertilization outcome with transplanted cryopreserved ovarian tissue. Fertil. Steril. 114, 279-280.
- Dolmans M.-M. (2021). From isolated follicles to the artificial ovary: why and how?. Curr. Opin. Endocr. Metab. Res. 18, 62-68.
- Dong F.-L., Ma L., Shi S.-L., Dai S.-J., Liu X.-G., Su Y.P., Guo Y.-H., Wang F. and Sun Y.-P. (2014). An research on the isolation methods of frozen-thawed human ovarian preantral follicles. Int. J. Clin. Exp. Med. 7, 2298-2303.
- Donnez J. and Dolmans, M.-M. (2017). Fertility preservation in women. N. Engl. J. Med. 377, 1657-1665.
- Donnez J., Martinez-Madrid B., Jadoul P., Van Langendonck A., Demylle D. and Dolmans M.-M. (2006). Ovarian tissue cryopreservation and transplantation: a review. Hum. Reprod. Update 12, 519-535.
- Figueiredo J.R., Hulshof S.C., Van den Hurk R., Ectors F.J., Fontes R.S., Nusgens B., Bevers M.M. and Beckers J.F. (1993). Development of a combined new mechanical and enzymatic method for the isolation of intact preantral follicles from fetal, calf and adult bovine ovaries. Theriogenology 40, 789-799.
- Gerritse R., Beerendonk C., Tijink M., Heetkamp A., Kremer J., Braat D. and Westphal J. (2008). Optimal perfusion of an intact ovary as a prerequisite for successful ovarian cryopreservation. Hum. Reprod. 23, 329-335.
- Gosden R. (1990). Restitution of fertility in sterilized mice by transferring primordial ovarian follicles. Hum. Reprod. 5, 117-122.

- Heeren A.M., Van Iperen L., Klootwijk D.B., de Melo Bernardo A., Roost M.S., Gomes Fernandes M.M., Louwe L.A., Hilders C.G., Helmerhorst F.M., Van Der Westerlaken L.A. and Chuva de Sousa Lopes S.M. (2015). Development of the follicular basement membrane during human gametogenesis and early folliculogenesis. *BMC Dev. Biol.* 15, 1-13.
- Hovatta O., Wright C., Krausz T., Hardy K. and Winston R.M. (1999). Human primordial, primary and secondary ovarian follicles in long-term culture: effect of partial isolation. *Hum. Reprod.* 14, 2519-2524.
- Irving-Rodgers H.F. and Rodgers R. (2006). Extracellular matrix of the developing ovarian follicle. *Semin. Reprod. Med.* 24, 195-203.
- Kristensen S.G., Rasmussen A., Byskov A.G. and Andersen C.Y. (2011). Isolation of pre-antral follicles from human ovarian medulla tissue. *Hum. Reprod.* 26, 157-166.
- Langbein A., Jorssen E., Fransen E., Rodriguez A., García M.C., Leroy J. and Bols P. (2015). Characterization of freshly retrieved preantral follicles using a low-invasive, mechanical isolation method extended to different ruminant species. *Zygote* 23, 683-694.
- Lierman S., Tilleman K., Cornelissen M., De Vos W.H., Weyers S., T'Sjoen G., Cuvelier C.A. and De Sutter P. (2015). Follicles of various maturation stages react differently to enzymatic isolation: a comparison of different isolation protocols. *Reprod. Biomed. Online* 30, 181-190.
- Maltaris T., Boehm D., Dittrich R., Seufert R. and Koelbl H. (2006). Reproduction beyond cancer: a message of hope for young women. *Gynecol. Oncol.* 103, 1109-1121.
- Martinez-Madrid B., Dolmans M.-M., Van Langendonck A., Defrère S., Van Eyck A.-S. and Donnez J. (2004). Ficoll density gradient method for recovery of isolated human ovarian primordial follicles. *Fertil. Steril.* 82, 1648-1653.
- McLaughlin M., Albertini D., Wallace W., Anderson R. and Telfer E. (2018). Metaphase II oocytes from human unilaminar follicles grown in a multi-step culture system. *MHR: Mol. Hum. Reprod.* 24, 135-142.
- Meirow D. (2000). Reproduction post-chemotherapy in young cancer patients. *Mol. Cell. Endocrinol.* 169, 123-131.
- Mouloungui E., Zver T., Roux C. and Amiot C.J. (2018). A protocol to isolate and qualify purified human preantral follicles in cases of acute leukemia, for future clinical applications. *J. Ovarian Res.* 11, 1-15.
- Nemes Z., Dietz R., Lüth J., Gomba S., Hackenthal E. and Gross F. (1979). The pharmacological relevance of vital staining with neutral red. *Experientia* 35, 1475-1476.
- Oktay K., Nugent D., Newton H., Salha O., Chatterjee P. and Gosden R.G. (1997). Isolation and characterization of primordial follicles from fresh and cryopreserved human ovarian tissue. *Fertil. Steril.* 67, 481-486.
- Paulini F., Vilela J.M., Chiti M.C., Donnez J., Jadoul P., Dolmans M.-M. and Amorim C.A. (2016). Survival and growth of human preantral follicles after cryopreservation of ovarian tissue, follicle isolation and short-term xenografting. *Reprod. BioMed. Online* 33, 425-432.
- Peddie V., Porter M.A., Barbour R., Culligan D., MacDonald G., King D., Horn J. and Bhattacharya S. (2012). Factors affecting decision making about fertility preservation after cancer diagnosis: a qualitative study. *BJOG.* 119, 1049-1057.
- Resetkova N., Hayashi M., Kolp L.A. and Christianson M.S. (2013). Fertility preservation for prepubertal girls: update and current challenges. *Curr. Obstet. Gynecol. Rep.* 2, 218-225.
- Rosendahl M., Andersen M.T., Ralfkiær E., Kjeldsen L., Andersen M.K. and Andersen C.Y. (2010). Evidence of residual disease in cryopreserved ovarian cortex from female patients with leukemia. *Fertil. Steril.* 94, 2186-2190.
- Roy S.K. and Treacy B.J. (1993). Isolation and long-term culture of human preantral follicles. *Fertil. Steril.* 59, 783-790.
- Schmidt V.M., Isachenko E., Rappl G., Rahimi G., Hanstein B., Morgenstern B., Mallmann P. and Isachenko V. (2018). Construction of human artificial ovary from cryopreserved ovarian tissue:

- Appearance of apoptosis and necrosis after enzymatic isolation of follicles. *Cryobiology*. 84, 10-14.
- Schmidt V.M., Isachenko V., Rappl G., Rahimi G., Hanstein B., Morgenstern B., Mallmann P. and Isachenko E. (2018). Comparison of the enzymatic efficiency of Liberase TM and tumor dissociation enzyme: effect on the viability of cells digested from fresh and cryopreserved human ovarian cortex. *Reprod. Biol. Endocrinol.* 16, 1-14.
- Sharma G.T., Dubey P.K. and Meur S. (2009). Effect of different mechanical isolation techniques on developmental competence and survival of buffalo ovarian preantral follicles. *Livestock Sci.* 123, 300-305.
- Silvestris E., De Palma G., Canosa S., Palini S., Dellino M., Revelli A. and Paradiso A.V. (2020). Human ovarian cortex biobanking: a fascinating resource for fertility preservation in cancer. *Int. J. Mol. Sci.* 21, 3245.
- Smith M.A., Altekruze S.F., Adamson P.C., Reaman G.H. and Seibel N.L. (2014). Declining childhood and adolescent cancer mortality. *Cancer* 120, 2497-2506.
- Soares M., Saussoy P., Maskens M., Reul H., Amorim C.A., Donnez J. and Dolmans M.M. (2017). Eliminating malignant cells from cryopreserved ovarian tissue is possible in leukaemia patients. *Br. J. Haematol.* 178, 231-239.
- Telfer E., Torrance C. and Gosden R. (1990). Morphological study of cultured preantral ovarian follicles of mice after transplantation under the kidney capsule. *J. Reprod. Fertil.* 89, 565-571.
- Vanacker J., Camboni A., Dath C., Van Langendonck A., Dolmans M.-M., Donnez J. and Amorim C.A. (2011). Enzymatic isolation of human primordial and primary ovarian follicles with Liberase DH: protocol for application in a clinical setting. *Fertil. Steril.* 96, 379-383.e3.
- Wallace W.H.B. and Kelsey T.W. (2010). Human ovarian reserve from conception to the menopause. *PLoS One*. 5, e8772.
- Wang W., Upshaw L., Zhang G., Strong D.M. and Reems J.-A. (2007). Adjustment of digestion enzyme composition improves islet isolation outcome from marginal grade human donor pancreata. *Cell Tissue Bank* 8, 187-194.
- Wang T.R., Yan J., Lu C.L., Xia X., Yin T.L., Zhi X., Zhu X.H., Ding T., Hu W.H., Guo H.Y. and Li R. (2016). Human single follicle growth in vitro from cryopreserved ovarian tissue after slow freezing or vitrification. *Hum. Reprod.* 31, 763-773.
- Ward E., DeSantis C., Robbins A., Kohler B. and Jemal A. (2014). Childhood and adolescent cancer statistics, 2014. *CA Cancer J. Clin.* 64, 83-103.
- Wenners A., Grambach J., Koss J., Maass N., Jonat W., Schmutzler A. and Mundhenke C. (2017). Reduced ovarian reserve in young early breast cancer patients: preliminary data from a prospective cohort trial. *BMC Cancer*. 17, 1-9.
- Wiweko B., Soebijanto S., Boediono A., Mansyur M., Siregar N.C., Suryandari D.A., Aulia A., Djuwantono T. and Affandi B. (2019). Survival of isolated human preantral follicles after vitrification: Analyses of morphology and Fas ligand and caspase-3 mRNA expression. *Clin. Exp. Reprod. Med.* 46, 152-165.
- Xu M., Barrett S.L., West-Farrell E., Kondapalli L.A., Kiesewetter S.E., Shea L.D. and Woodruff T.K. (2009). *In vitro* grown human ovarian follicles from cancer patients support oocyte growth. *Hum. Reprod.* 24, 2531-2540.
- Yin H., Kristensen S., Jiang H., Rasmussen A. and Andersen C.Y. (2016). Survival and growth of isolated pre-antral follicles from human ovarian medulla tissue during long-term 3D culture. *Hum. Reprod.* 31, 1531-1539.

Table 1. Summary of studies on the isolation of human preantral follicles using enzymatic digestion

Sample (age)	Mechanical method	Enzymatic procedure	Diameter of the isolated follicles (µm)	Aim	Main outcomes	Reference
Premenopausal women with regular menstrual cycles who were undergoing hysterectomy-oophorectomy (range 16 to 32 years of age)	Razor blade before and agitation during and at 4°C in a refrigerator for 36 hours after the enzymatic process for slow digestion	The mixture of collagenase and deoxyribonuclease (DNase) for 1 hour at 37°C and for 36 hours at 4°C	<90 µm	Long-term culture of isolated follicles	Successful follicle isolation and long culture	Roy and Treacy, 1993
Biopsies of women undergoing laparoscopic surgery (range 25 to 32 years of age)	27-gauge insulin needles after enzymatic digestion	DNase and Collagenase IA	30 - 50 µm	Follicle isolation before and after cryopreservation	Successful preantral isolation from fresh and frozen ovarian tissue	Oktay et al., 1997
Women undergoing elective	Tissue was chopped	Collagenase type IA and DNase I	-	Ontogeny FSHr expression	FSHr expression in the early stages of	Oktay et al., 1997

cesarean section (range 5 to 33 years of age)				on in isolated follicles	follicle development	
Biopsies of women undergoing laparoscopic surgery (range 25–43 years of age)	Sliced into 0.3–1 mm before and using a 27- gauge needle after enzymatic digestion or just partial isolation for the mechanical isolation group	Collagenase II at 1 or 0.5 mg/ml for the enzymatic group	20 - 80 μ m	Culture of the ovarian follicle by two enzymati c and mechani cal methods	Culturing the small tissue slices is better than partial isolation	Hovatta et al., 1999
Biopsies of women undergoing laparoscopic surgery (range 26 and 31 years of age)	Using the tissue sectioner before, gentle agitation to mechanically disrupt digested tissue	Collagenase type IA	30 -120 μ m	Isolation of preantral follicles by a ficoll density gradient method	A Ficoll density gradient can reduce the manipulation time and increase the recovery of isolated follicles	Martinez- Madrid et al. 2004

Biopsies of women who underwent laparoscopic surgery for gynecological disease (range 20 to 32 years of age)	Using the tissue sectioner, and using a 130- μ m micropipette to transfer isolated follicles	Collagenase type IA versus Liberase blendzyme 3	40 - 80 μ m	Comparison between Liberase blend enzyme and collagenase IA	Liberase shows high viability in isolated follicles and is a good alternative to collagenase IA	Dolmans et al., 2006
Biopsies of women undergoing laparoscopic surgery (range 23 to 32 years of age)	Using the tissue sectioner before, aspirated up and down during enzymatic digestion and aspirated up and down with a 10 ml pipette to pick up follicles	Liberase blendzyme 3	-	Short-term xenografting human isolated follicles	Isolated follicles can survive after 7 days of xenotransplantation by plasma clot	Dolmans et al., 2007

Biopsies of women undergoing laparoscopic surgery (range 26 to 29 years of age)	Using the tissue sectioner before and using a 130 μ m micropipette after enzymatic digestion	Collagenase IA versus Liberase blendzyme 3 (PCB)	30 -100 μ m	Long-term xenograf ting human isolated follicles	Isolated follicles can survive after long-term xenotransplantation by plasma clot	Dolmans et al., 2008
Patients were seeking fertility-sparing options in the oncofertility program (range 16 to 39 years of age)	Ovarian cortical strips were cut into 1 mm ³ pieces before and 25-gauge needles in dissection media after enzymatic digestion	Collagenase and DNase	175 $\pm$ 10 μ m	<i>In vitro</i> culture of secondary follicles	Successful culture of the secondary follicle in a bio-engineered system for 30 days	Xu et al., 2009
Biopsies of women undergoing laparoscopic	Using the tissue sectioner, and using a 130-	Collagenase IA versus Liberase DH	30 - 80 μ m	Comparison between collagenase IA and	The high number of follicles were isolated by Liberase DH. Also, they can	Vanacker et al., 2011

surgery (range 21 to 35 years of age)	mm micropipette for the transfer of isolated follicles			Liberase DH	maintain their viability structure after <i>in vitro</i> culture	
Biopsies of girls/women undergoing cryopreservation of the ovarian cortex (range 3 to 35 years of age)	Using the tissue sectioner before, and using a thin Pasteur pipette after enzymatic digestion	The mixture of Collagenase IV, Liberase TM, and DNase I	40 - 160 μm	Follicle isolation from the ovarian medulla	The medulla is full of preantral follicles, especially in young girls that can used for investigation and clinical approaches	Kristensen et al., 2011
Fresh and frozen-thawed ovarian tissue (range not mentioned)	1 ml syringe under an inverted microscope	Collagenase type I	The mean of preantral follicles isolated by collagenase IA combined with mechanical digestion after 6 days was $84.9 \pm 4.3 \mu\text{m}$, mechanical alone was $84.9 \pm 4.3 \mu\text{m}$	Comparison of isolation methods on frozen-thawed ovarian follicles	There is no difference in the stage of isolated follicles before and after vitrification and the survival rate of secondary follicles is higher with mechanical isolation after 6 days of culture	Dong et al., 2014


Frozen ovarian tissue of the transsexual person	Not mentioned	Liberase DH and Liberase TM alone or combined with collagenase IV versus collagenase IV alone	>60 μm (primordial follicles) to >200 μm (secondary follicles)	Comparison of five group enzymes for follicle isolation	The maturation status of follicles has a key role in response to enzymatic isolation	Lierman et al., 2015
Biopsies of women undergoing laparoscopic surgery (range 22 and 36 years of age)	Using the tissue chopper, manually pipetting and transferring the follicles to fresh droplets using a 130- μm micropipette after enzymatic digestion	Liberase DH	-	Safe follicle isolation	Washing can eliminate malignant cells	Soares et al., 2015

Biopsies of women who underwent laparoscopic surgery for gynecological disease (mean 31.8 ± 8.36 years of age)	Using the tissue sectioner before and 29-gauge after enzymatic digestion	Liberase DH and DNase I	Mean $77.7 \pm 2.7 \mu\text{m}$ for fresh tissue, $77.2 \pm 3.2 \mu\text{m}$, and $78.3 \pm 2.4 \mu\text{m}$, from slow-frozen and vitrified tissue, respectively	Comparison of slow freezing, vitrification, and fresh tissue after single follicle <i>in vitro</i> culture	There is no difference between slow freezing and vitrification	Wang et al., 2016
Tiny amounts of cortical tissues of cancer patients for cryopreservation (range 9 to 37 years of age)	Using the tissue chopper before and aspirating the suspension up and down to release the follicles into the media. Isolated follicles with red coloration	The mixture of Liberase TM, collagenase IV, and DNase I	$\leq 60 \mu\text{m}$ to $120\text{--}250 \mu\text{m}$	Culture of medulla-isolated follicles in a 3-D system	Secondary isolated follicles had better growth in comparison with smaller follicles after 7 days	Yin et al., 2016

	were removed and washed twice in 37°C DPBS supplemented with 10% FBS after enzymatic digestion					
Freeze-thawed biopsies of women undergoing laparoscopic surgery (range 21 to 37 years of age)	Using the tissue sectioner before, and using a 130- µm micropipette after enzymatic digestion	Liberase DH	Between 38.8±6.3 µm to 48.8±2.5 µm	Survival and growth of xenotran splanted follicles by fibrin	It seems that fibrin can be used as a human artificial ovary	Paulini et al., 2016
Using vitrified ovarian strips (range not mentioned)	Using the tissue chopper before, filtration every	Liberase DH and DNase I	30 to 129 µm	Design a modified protocol for	Increase the number and quality of isolated follicles	Chiti et al., 2017

	30 min for complete follicle isolation during, and 1 hour using a 130-µm micropipette after enzymatic digestion			follicle isolation		
Biopsies of women undergoing laparoscopic surgery (range 25 to 37 years of age)	Manually pipetting and transferring the follicles using a 130-µm micropipette after enzymatic digestion	Collagenase IA, a GMP-grade purified collagenase NB6 or Liberase DH	Mean for collagenase NB6 was 31.66 ± 6.79 µm, for collagenase IA 36.77 ± 7.69 µm and for liberase DH 36.77 ± 7.69 µm	Follicle isolation without leukemic cells presentin g	Follicle isolation was made by GMP for cell therapy	Mouloungui et al., 2018
Fresh and frozen ovarian cortex fragments	Mechanically cut	TDE versus Liberase TM	-	Compari son of TDE with	Isolated follicles with TDE had no necrosis and apoptosis	Schmidt et al., 2018


(range 29 to 39 years of age)				Liberase TM		
Donated ovarian cortex from patients with various malignant diseases (range 15 to 37 years of age)	Using the tissue chopper	A mixture of Liberase TM and Collagenase IV	54.5–162.2 μm	Human ovary reconstruction	Xenotransplantation of decellularized ovarian cortical tissue can support the growth of human follicles	Pors et al., 2019
Biopsies of women undergoing laparoscopic undergoing laparoscopic surgery (range 18 to 35 years of age)	Mechanically cut into $\sim 0.5 \times 1$ to 1×1 mm^3 pieces by quick simultaneous movements of two scalpels no. 22, and using 125 μm V-denuded capillaries after enzymatic digestion	TDE-cocktail versus Liberase DH/ /DNase	The mean TDE was 40.5 ± 4.1 μm , and the mean Liberase DH was 42 ± 3.5 μm	Optimizing the follicle isolation protocol	TDE can be used instead of Liberase DH	Chen et al., 2021



Ovarian strips of:

- Transgender patients;
- Biopsies of women undergoing laparoscopic surgery for gynecological disease;
- Women undergoing elective cesarean section;
- Patients were seeking fertility-sparing options in the oncofertility program.

Fresh or frozen-warmed/thawed ovarian strips





or



Mechanical process for chopped ovarian strips before follicle isolation

Enzymatic digestion such as:

- Collagenase (I, II, IV or IX);
- collagenase NB6
- Liberase blendzyme 3;
- Liberase DH;
- Liberase TM;
- Tumor Dissociation Enzyme (TDE).



Mechanically follicle isolation by using an insulin syringe under a stereomicroscope