

# Biophysical mechanisms governing large-scale brain network dynamics underlying individual-specific variability of perception

Vinodh G. Kumar  | Shrey Dutta  | Siddharth Talwar | Dipanjan Roy  | Arpan Banerjee 

Cognitive Brain Dynamics Lab, National Brain Research Centre, Gurgaon, India

## Correspondence

Dipanjan Roy and Arpan Banerjee, Cognitive Brain Dynamics Lab, National Brain Research Centre, NH-8, Manesar, Gurgaon 122052, Haryana, India. Emails: dipanjan.nbrc@gov.in (DR); arpan@nbrc.ac.in (AB)

## Funding information

Department of Biotechnology, Government of India, Grant/Award Number: BT/RLF/Re-entry/07/2014, BT/RLF/Re-entry/31/2011 and BT/07/IYBA/2013; Department of Science and Technology (DST) Ministry of Science and Technology, Government of India, Grant/Award Number: SR/CSRI/21/2016

## Abstract.

Perception necessitates interaction among neuronal ensembles, the dynamics of which can be conceptualized as the emergent behavior of coupled dynamical systems. Here, we propose a detailed neurobiologically realistic model that captures the neural mechanisms of inter-individual variability observed in cross-modal speech perception. From raw EEG signals recorded from human participants when they were presented with speech vocalizations of McGurk-incongruent and congruent audio–visual (AV) stimuli, we computed the global coherence metric to capture the neural variability of large-scale networks. We identified that participants' McGurk susceptibility was negatively correlated to their alpha band global coherence. The proposed biophysical model conceptualized the global coherence dynamics emerge from coupling between the interacting neural masses—representing the sensory-specific auditory/visual areas and modality nonspecific associative/integrative regions. Subsequently, we could predict that an extremely weak direct AV coupling results in a decrease in alpha band global coherence—mimicking the cortical dynamics of participants with higher McGurk susceptibility. Source connectivity analysis also showed decreased connectivity between sensory-specific regions in participants more susceptible to McGurk effect, thus establishing an empirical validation to the prediction. Overall, our study provides an outline to link variability in structural and functional connectivity metrics to variability of performance that can be useful for several perception and action task paradigms.

**Abbreviations:** AAL, automated anatomical labeling; AV, audio–visual; BEM, boundary element method; DICS, dynamic imaging of coherent sources; EEG, electroencephalogram; fMRI, functional magnetic resonance imaging; HR, Hindmarsh–Rose; IOC, inferior occipital cortex; LGN, lateral geniculate nucleus; MEG, magnetoencephalogram; MGN, medial geniculate nucleus; MOC, medial occipital cortex; SOC, superior occipital cortex; STG, superior temporal gyrus; STS, superior temporal sulcus.

Vinodh G. Kumar and Shrey Dutta contributed equally to this work.

Edited by Christoph M. Michel

The peer review history for this article is available at <https://publons.com/publon/10.1111/ejn.14747>

© 2020 Federation of European Neuroscience Societies and John Wiley & Sons Ltd



## KEYWORDS

EEG, global coherence, large-scale networks, McGurk effect, multisensory, perception, source connectivity, variability

## 1 | INTRODUCTION

Neurobiologically realistic dynamical models of perception have provided attractive explanation of behavior (Cuppini, Shams, Magosso, & Ursino, 2017; Moreno-Bote, Rinzel, & Rubin, 2007; Thakur, Mukherjee, Sen, & Banerjee, 2016). Concurrently, they also provide insights into brain dynamics observed at different scales, for example, microscopic level of single neurons (Deco, Pérez-Sanagustín, de Lafuente, & Romo, 2007) and macroscopic level of field recordings (David & Friston, 2003; Onojima, Goto, Mizuhara, & Aoyagi, 2018; Spiegler, Knösche, Schwab, Haueisen, & Atay, 2011). How does this framework adapt to perceptual variability across individuals during the perception of complex signals? One intuitive approach to explain perceptual variability involves characterization of stochastic fluctuations in component neural systems. Thereafter, probability distributions set by internal processes that generate prediction can follow Bayes' optimization principles to set up perceptual object identification (Körding et al., 2007; Ma, Beck, & Pouget, 2008). One drawback of these descriptive Bayesian brain hypotheses is that, unlike predictive coding formulations, they are agnostic to the neurobiological mechanisms and systems-level interactions that govern perception (Ursino, Crisafulli, di Pellegrino, Magosso, & Cuppini, 2017). In this article, we employ McGurk effect (McGurk & Macdonald, 1976) to demonstrate how the systems-level interactions in multisensory and unisensory subsystems can explain differences in McGurk susceptibility across individuals.

The fact that speech perception is fundamentally driven by visual cues is demonstrated in this paradigm—an auditory */bal* dubbed onto visual lip movements of */gal* is perceived by the listener as a completely different syllable */dal* (McGurk & Macdonald, 1976). Variability in the perception of the illusion has enabled researchers to unravel the neural correlates of illusory/cross-modal percept in terms of cortical activations and neural oscillations (Keil, Muller, Ihssen, & Weisz, 2012; Nath & Beauchamp, 2011). However, these studies have overlooked the substantial variability of McGurk susceptibility across individuals that could influence the objective behavioral and neural estimates.

From a neurophysiological standpoint, how do we reconcile the neural correlates of McGurk percept under the purview of inter-individual variability or vice versa? Moreover, it raises a fundamental question on what constitutes an appropriate neural correlate. For instance, activity in superior temporal sulcus (STS) and its functional connectedness to sensory-specific and fronto-parietal regions has been shown to subserve multisensory speech perception and McGurk susceptibility (Keil et al., 2012;

Nath & Beauchamp, 2012) as well. Moreover, other research areas like face processing (Haxby, Hoffman, & Gobbini, 2000), motion perception (Puce & Perrett, 2003), and social perception (Saxe, 2006) attribute activations in the STS to the particular behavior being studied. This suggests that the STS is a prime nexus for information processing; however, it may not be a neural correlate of any specific processing component.

Emerging studies indicate complex cognitive capabilities to emerge as a result of interactions among large-scale networks of cortical areas (Bressler & Kelso 2001; Bressler & Menon, 2010; Varela, Lachaux, Rodriguez, & Martinerie, 2001). Therefore, a measure that could capture the “dynamic” and “preferential” interactions among distributed neuronal assemblies/brain regions will offer the most pertinent representation of the distinct neural machinery supporting McGurk susceptibility.


Recent studies have indicated that beyond a specific region of interest, a large-scale network of oscillatory brain networks underwrite the McGurk effect (Kumar et al., 2016; Kumar, Kumar, Roy, & Banerjee, 2017). A key question emerges, how robust is this network across individuals and whether the organization of such networks is contingent on the stimulus configurations. Secondly, what are the neural mechanisms that give rise to such network level correlates? While the first question needs to be answered empirically using a detailed neurophysiological study of underlying brain networks, the more broader question of systems-level insight requires a neurobiologically inspired large-scale computational model.

We hypothesized that variability in McGurk susceptibility across individuals stems from implicit differences in their large-scale functional connectivity. In order to test that, we conducted psychophysical experiments in tandem with EEG recordings to characterize the large-scale oscillatory modes in whole-brain dynamics. Subsequently, we constructed a neural mass model for multisensory perception—an abstraction of summed synapto-dendritic activity of thousands of neurons, comprising three nodes representing sensory and multisensory areas. Finally, we validated the plausible neural mechanisms made by the model using source-space analysis (Figure 1), thus providing a link between functional metrics and underlying neurophysiological complexity.

## 2 | MATERIALS AND METHODS

### 2.1 | Participants

Eighteen healthy right-handed volunteers (8 females, mean age 24.9,  $SD = 2.8$ ) gave written informed consent under an



**FIGURE 1** Pipeline of the study: We address an ongoing challenge of characterizing the neural mechanisms that govern the variability during multisensory speech perception. Employing the McGurk paradigm, we identify the neural signatures of the variability in the global coherence metric. We then turn to test our hypothesis—“dynamic and preferential coupling between sensory-specific and multisensory regions underlies the observed variability,” using a neural mass model. Finally, we validate the inferences drawn from the model by analyzing pairwise coherence in the source space between concordant cortical generators of neural activity [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]


experimental protocol approved by the Institutional Human Ethics Committee of the National Brain Research Centre, Gurgaon, which is in agreement with the 2008 Declaration of Helsinki.

## 2.2 | Stimuli and behavioral task


Digital videos of a native Hindi-speaking male volunteer articulating the syllables */pal/*, */kal/*, and */tal/* were recorded and edited using the audio editing software Audacity ([www.audacityteam.org](http://www.audacityteam.org)) and the video editing software VideoPad Editor ([www.nchsoftware.com](http://www.nchsoftware.com)). The duration of each video clip ranged from 1.5 to 1.7 s to include the neutral, mouth-closed position, and all the mouth movements of articulation to closing. The duration of auditory syllables in the videos ranged from 0.4 to 0.5 s. The stimuli (Figure 2) consisted of four kinds of videos: three congruent (auditory and visual matching) syllables (*/pal/*, */kal/*, and */tal/*) and one McGurk/incongruent (auditory and visual mismatch) syllable (auditory */pal/* + visual */kal/*) producing the McGurk percept of */tal/*.

The experiment contained five blocks—each block consisting of 120 trials (30 trials of each video presented in random). Inter-stimulus intervals were pseudo-randomly varied between 1.2 and 2.8 s to minimize expectancy effects. Using a forced-choice task, on every trial, participant reported their

### (a) McGurk stimulus



### (b) Congruent stimuli



**FIGURE 2** Stimuli: (a) Example trial with 3 video frames from McGurk stimulus (audio */pal/* + video */kal/*) used in this experiment (top row), the audio trace of the syllable */pal/* presented simultaneously to the video (middle row), and the onset and offset time of the audio. (b) The congruent AV stimuli: Each block represents a video with audio */pal/*, */tal/*, and */kal/* dubbed onto a video of a person articulating */pal/*, */tal/*, and */kal/*, respectively [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

percept by pressing a specified key on the keyboard corresponding to */pal/*, */kal/*, */tal/* or something else (others) while watching the videos.



## 2.3 | EEG recording and preprocessing

Continuous EEG scans were acquired using a Neuroscan system (Synamps 2, Compumedics, Inc.) with 64 Ag/AgCl scalp electrodes placed on an elastic cap in a 10–20 montage. Individual electrode locations were registered using the FASTRAK 3D digitizing system (Polhemus Inc.). Recordings were made against the center (near Cz) reference electrode on the Neuroscan cap and digitized at a sampling rate of 1,000 Hz. Channel impedances were monitored to be at values <10 kΩ.

Preprocessing and offline data analysis were performed using EEGLAB (Delorme & Makeig, 2004), FieldTrip (Oostenveld, Fries, Maris, & Schoffelen, 2011), and custom-made MATLAB scripts. Continuous data were high-passed at 0.1 Hz with finite impulse response (FIR) filters, low-passed at 80 Hz FIR, and notch-filtered between 46 and 54 Hz to remove line noise (9th-order 2-pass Butterworth filter). The noisy channels were removed and interpolated, following which the data were average re-referenced. For the data analysis, epochs of 0.8 s after the onset of the sound were extracted and categorized according to the type of AV stimuli—congruent AV stimuli, *lta*, *lpa*, *lka*, and incongruent McGurk stimulus. The extracted epochs were then baseline-corrected by removing the temporal mean of the EEG signal 0.2 s before the onset of the sound, on an epoch-by-epoch basis. Finally, epochs with amplitudes above or below  $\pm 75 \mu\text{V}$  were removed to minimize contamination from ocular (eyeblinks and movements) and muscle-related artifacts. To ensure comparable signal-to-noise ratio, an equal number of trials were picked randomly from each participant for every kind of AV stimulus presented.

## 2.4 | Large-scale functional connectivity analysis

Neuronal synchronization was quantified using global coherence (Cimenser et al. 2011). The metric quantifies the strength of covariation of neural oscillations at the global scale in specific frequency bands (Railo, Koivisto, & Revonsuo, 2011) that corresponds to specific functional state of the brain. We employed the Chronux (Mitra & Bokil, 2008) function CrossSpecMatc.m to obtain trialwise global coherence on the 0.8-s epochs sorted based on stimuli and responses. We considered 5 orthogonal discrete prolate spheroidal sequences (dpss), also known as Slepian tapers to restrict the leakage of spectral estimates into nearby frequency bins. The output variable “Ctot” of the function yielded global coherence at frequency  $f$ .

The function initially multiplies the epochs with the specified number of Slepian tapers before performing the fast Fourier transform (FFT). The resulting FFT values are

averaged, and the cross-spectrum for all sensor combinations at frequency  $f$  is computed. The cross-spectral density between two sensors was computed from the tapered Fourier transforms using the following equation

$$C_{ij} = \text{conj}(X_i(f) \cdot X_j(f)) \quad (1)$$

where  $C_{ij}$  represents the cross-spectrum;  $X_i$  and  $X_j$  represent the tapered Fourier transforms from the sensors  $i$  and  $j$ . Subsequently, singular value decomposition (SVD) is applied on the cross-spectral density matrix for every frequency  $f$  which yields the following

$$D(f) = VCV^T \quad (2)$$

The diagonal matrix  $D$  comprises the values proportional to the variance explained by the orthogonal set of eigenvectors ( $V$ ,  $V^T$ ). The global coherence  $C_{\text{Global}}(f)$  at frequency bin  $f$  is subsequently computed by normalizing the largest eigenvalue (first entry of the  $D(f)$  at frequency  $f$  on a trial-by-trial basis for each participant employing the following equation

$$C_{\text{Global}}(f) = \frac{D_1(f)}{\sum_{i=1}^n D_i(f)} \quad (3)$$

We initially analyzed if changes in global coherence values at specific frequency bands ( $\alpha$ : 8–13 Hz,  $\beta$ : 14–30 Hz,  $\gamma$ : 31–45 Hz) correlated with the participants' susceptibility of McGurk perception. Participant-wise mean of the global coherence in specific frequency bands was computed and was statistically analyzed using Spearman rank correlation and  $t$  tests.


## 2.5 | Architecture of large-scale dynamical model of perception

To account the neuronal interactions that could facilitate changes in global coherence, we constructed a large-scale dynamical model. Striking a balance between biological realism and model parsimony, we proposed a neural mass model comprising three nodes representing the auditory, visual, and the multisensory areas (Figure 3a). We followed closely a previously established practice and convention in computational modeling by treating each cortical region as an individual node comprising excitatory and inhibitory population of Hindmarsh–Rose (HR) neurons (Stefanescu & Jirsa, 2008). Rationally, we incorporate the following factors to reflect the biological realism in the proposed model architecture -

1. Concurring with previous evidences, auditory node was assumed to be most sensitive to ambient temporal fluctuations operating with a fast time scale (Rosen &



## Model architecture



**FIGURE 3** Model architecture: (a) Representation of the model comprising three nodes signifying auditory (fast time constant), visual (slow time constant), and higher order multisensory regions (intermediate time constant). Excitatory influences are balanced by a reciprocal inhibitory connection. (b) Representation of each node comprising of network of 150 Hindmarsh–Rose excitatory and 50 inhibitory neurons. Each neuron can exhibit isolated spiking, periodic spiking, and bursting behavior. The detailed model parameters are presented in Table 1 [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

Howell, 2011), visual node was assumed to be slowest in terms of sensitivity (Williams, Mechler, Gordon, Shapley, & Hawken, 2004), and the multisensory node was assumed to operate at an intermediate time scale.

- Inspired by the anatomical evidences, in our model, inputs from visual node were directed to the auditory cortex in two ways: (a) direct influence in a feedforward manner (Falchier, Clavagnier, Barone, & Kennedy, 2002; Rockland & Ojima, 2003; Wallace, Ramachandran, & Stein, 2004) and (b) in feedback manner via the multisensory association node (Bizley & King, 2011).
- Excitatory and inhibitory afferents on the spatially extended and geometrically aligned pyramidal neurons result in post-synaptic potentials (Kirschstein & Köhling, 2009). These communications emerge as EEG patterns. Hence, we used a population of excitatory and inhibitory neurons in each node (Figure 3b). Following the approach used by Stefanescu and Jirsa (2008), we employed a total of 150 excitatory ( $N_E = 150$ ) and 50 inhibitory ( $N_I = 50$ ) HR neurons, maintaining a ratio of 3:1. Importantly, inhibitory neurons within an individual node were not coupled directly with other inhibitory neurons based on the logic that these connections in microcircuits are typically sparse in nature (Stefanescu & Jirsa, 2008; Wilson & Cowan, 1972).

Incorporating the aforementioned factors, we defined a neural mass model comprising three sets of ordinary differential equations for each excitatory and inhibitory HR neuron. The three variables  $x_n^L$ ,  $y_n^L$ , and  $z_n^L$  account for the membrane potential dynamics of a neuron  $n$ , fast gating current, and slow gating current, respectively. Thus, the entire network can be represented as a network of coupled nonlinear differential equations containing –

### 2.5.1 | Excitatory subpopulations

$$\begin{aligned} \tau_L \frac{dx_{n_E}^L}{dt} = & y_{n_E}^L - ax_{n_E}^{L3} - z_{n_E}^L + K_{EE} \left( E \left( x_{n_E}^L \right) - x_{n_E}^L \right) \\ & - K_{IE} \left( E \left( x_{n_I}^L \right) - x_{n_I}^L \right) + I_{n_E}^L \\ & + \sum_{u \neq L} W_{Lu} E \left( x_{n_E}^M \right) + \epsilon(0, \epsilon dt) \end{aligned}$$

$\forall u = V, M, A;$   
 $u \neq L$

$$\tau_L \frac{dy_{n_E}^L}{dt} = c - dx_{n_E}^{L2} - y_{n_E}^L$$

$$\tau_L \frac{dz_{n_E}^L}{dt} = r \left( s \left( x_{n_E}^L - x_0 \right) - z_{n_E}^L \right); \quad n_E = 1, \dots, N_E; L = A, V, M$$

### 2.5.2 | Inhibitory subpopulations

$$\tau_L \frac{dx_{n_I}^L}{dt} = y_{n_I}^L - ax_{n_I}^{L3} + bx_{n_I}^{L2} - z_{n_I}^L + K_{EI} \left( E \left( x_{n_E}^L \right) - x_{n_E}^L \right) + I_{n_I}^L$$

$$\tau_L \frac{dy_{n_I}^L}{dt} = c - dx_{n_I}^{L2} - y_{n_I}^L$$

$$\tau_L \frac{dz_{n_I}^L}{dt} = r \left( s \left( x_{n_I}^L - x_0 \right) - z_{n_I}^L \right); \quad n_I = 1, \dots, N_I \quad (4)$$

where  $L$ :  $A$ ,  $V$ , and  $M$  represent auditory, visual, and multisensory nodes, respectively. The nodes were driven by a noise of normal distribution  $\epsilon(0, \epsilon dt)$ , where  $\epsilon = 0.1$ .

**TABLE 1** Description of the model parameters and their corresponding values used in the model

| Parameter                 | Value                                                           | Description of parameters                                             |
|---------------------------|-----------------------------------------------------------------|-----------------------------------------------------------------------|
| $a, b, c, d$              | 1, 3, 1, 5                                                      | Constants for faster variables $x$ and $y$                            |
| $r$                       | 0.006                                                           | Lower value of $r$ is responsible for slower time scale of $z$        |
| $S$                       | 4                                                               | Constant affecting slower variable $z$                                |
| $x_0$                     | −1.6                                                            | $(x_0, y_0, 0)$ is a stable equilibrium point of Hindmarsh–Rose model |
| $N_E, N_I$                | 150, 50                                                         | Number of excitatory and inhibitory neurons in each node              |
| $\mu, \sigma$             | 2.8, 0.4                                                        | Mean and dispersion of input current in each node                     |
| $I_{n_E}, I_{n_I}$        | Computed from $\mu, \sigma$                                     | Input current for excitatory and inhibitory neurons in each node      |
| $\tau_A, \tau_V, \tau_M$  | 0.05, 2.5, 1 in milliseconds                                    | Time-scales for auditory, visual and multisensory nodes               |
| $\varepsilon$             | 0.1                                                             | Constant affecting the dispersion of noise                            |
| Within-node connectivity  |                                                                 |                                                                       |
| $K_{EE}$                  | 0.5                                                             | Coupling between excitatory neurons                                   |
| $K_{EI}$                  | 0.5                                                             | Excitatory to inhibitory coupling                                     |
| $n$                       | 3.39                                                            | $K_{IE}$ to $K_{EE}$ ratio                                            |
| $K_{IE}$                  | Computed from $n$ and $K_{EE}$                                  | Inhibitory to excitatory coupling                                     |
| Between-node connectivity |                                                                 |                                                                       |
| $W_{AV}, W_{AM}, W_{VM}$  | Explored in the range 0–1 (or −1 to 0)                          |                                                                       |
| State variables           |                                                                 |                                                                       |
| $x_{n_E}, x_{n_I}$        | Membrane potentials for excitatory and inhibitory subpopulation |                                                                       |
| $y_{n_E}, y_{n_I}$        | Spiking variables for excitatory and inhibitory subpopulation   |                                                                       |
| $z_{n_E}, z_{n_I}$        | Bursting variables for excitatory and inhibitory subpopulation  |                                                                       |

Auditory node had the fastest ( $\tau_A \sim 0.05$  ms), and visual node had the slowest time constant ( $\tau_V \sim 2.5$  ms). Time constant of the multisensory node was chosen to be in between the two ( $\tau_M \sim 1$  ms) as it integrates information from both the modalities. The mean activity of excitatory neurons in a given node ( $E(x_{n_E}) = \frac{1}{N_E} \sum_{n_E=1}^{N_E} x_{n_E}$ ) influencing the neuronal activity of other nodes was governed by coupling parameters:  $W_{AV}$  (auditory–visual coupling),  $W_{AM}$

(auditory–multisensory coupling), and  $W_{VM}$  (visual–multisensory coupling). Positive and negative values of the coupling parameters reflect excitatory and inhibitory coupling, respectively. Inhibitory couplings were chosen to maintain a balance with excitation. For instance, visual node's excitatory coupling of  $+W_{AV}$  on auditory node was balanced with inhibitory coupling of the same strength ( $-W_{AV}$ ) from the auditory node. Also, in such a configuration, visual node is being the source of excitation and is referred as the source, whereas auditory node is referred as the sink as all excitatory couplings are directed toward it. The multisensory node in such configuration behaves as both source and sink.

We place each individual neuron in a dynamical regime where both spiking and bursting behaviors are possible depending on the external input current ( $I$ ) that enters the neuron when other parameters are held constant at the following values (Stefanescu & Jirsa, 2008):

$$a = 1; b = 3; c = 1; d = 5; s = 4; r = 0.006; x_0 = -1.6.$$

The coupling between neurons within a node was linear, and its strength was governed by the following parameters:  $K_{EE}$  for excitatory–excitatory coupling,  $K_{EI}$  for excitatory–inhibitory coupling, and  $K_{IE}$  for inhibitory–excitatory coupling. Within a node, the relation between  $K_{EE}$  and  $K_{IE}$  was captured by the ratio  $n = \frac{K_{IE}}{K_{EE}}$ . Electrophysiological evidences indicate

enhanced oscillatory power in delta (1–4 Hz) and alpha (8–12 Hz) bands during rest (Gold, Henze, Koch, & Buzsáki, 2006). To emulate the characteristics of the large-scale network during rest, we chose the value of the ratio,  $n = 3.39$  where the average oscillatory activity of the nodes in a disconnected network, that is, in the absence of stimulus ( $\mu(I_{A,V,M}) = 0.1$ ; baseline) was higher in delta and alpha bands.

The external currents to both the excitatory and inhibitory subpopulations were drawn from a Gaussian distribution where  $\mu$  and  $\sigma$  represent the mean and standard deviation. As the input stimulus relays to auditory, visual, and multisensory regions via thalamus, we interpret lateral geniculate nucleus (LGN) and medial geniculate nucleus (MGN) to be the source of external current ( $I_A$ ,  $I_V$ , and  $I_M$ )—a pulse of 450 ms in the nodes when the model was simulated for 1 s. In rhesus monkey, the projections of MGN to multisensory areas like STS were found to be sparse (Yeterian & Pandya, 1989). Therefore, we choose lower mean value of external current to the multisensory node ( $\mu(I_M) = 0.85$ ) in comparison with visual ( $\mu(I_V) = 2.8$ ) and auditory ( $\mu(I_V) = 2.8$ ) nodes. All parameters are presented together in Table 1.

We hypothesized variability in global coherence patterns to emerge from differences in the coupling among the auditory, visual and multisensory nodes operating at fast, slow,



and intermediate time scales, respectively. Conceptually, the auditory and visual nodes are sensory-specific brain regions while multisensory regions include areas of associative cortex that are non-specific to a particular sensory stream. Thereafter, we sought to explore the global coherence while varying the coupling strength among the nodes in our model.

In summary, we considered different architectures or connectivity motifs, under the constraint that any reciprocal coupling comprised a forward excitatory set of connections and a reciprocal (and balanced) inhibitory set of connections. In what follows, we will designate the source of the excitatory connections as a source and the source of recurrent inhibitory connections as a target.

## 2.6 | Source-space connectivity

To validate the inferences gathered from our model, we turned to the empirical data again and undertook source-space connectivity analysis (see Figure 1). We employed dynamic imaging of coherent sources (DICS; Gross et al., 2001)—a frequency domain adaptive spatial filtering algorithm—to identify the sources of global coherence in specific frequency bands. This algorithm has proven to be particularly powerful in localizing oscillatory sources of cortical activity with well-established hypothesis (Halder, Talwar, Jaiswal, & Banerjee, 2019). Customized MATLAB codes employing FieldTrip toolbox were used. Using `ft_prepare_leadfield.m` and employing the boundary element method (BEM), we generated the leadfield matrix for the Colin 27 brain template. The leadfield matrix corresponds to the tissue and geometrical properties of the brain represented as discrete grids or voxels. Then, we employed `ft_freqanalysis.m` to evaluate the cross-spectral matrices of the epochs pooled over all the participants. Leadfield matrix and the cross-spectral matrices were used to compute the common spatial filters that regulate the amplitude of brain electrical activity passing from a specific location while attenuating activity originating at other locations. The distribution of the output amplitude of the spatial filters provides the metric for source localization.

To reduce the dimensionality in our source-space connectivity analysis, we adopted a parcellation scheme before generating source time series. We combined the source estimates of the voxels using the labeling according to automated anatomical labeling (AAL) atlas (Tzourio-Mazoyer et al., 2002). Then, for reconstructing the parcel-wise time series, we projected single-trial epoched time series to the source-space by multiplying them with spatial filter of the respective parcel obtained employing DICS. Subsequently, we computed the coherence between relevant parcels for rare and frequent perceivers explicitly.

As the creation of common spatial filter required multiple trials and participants to reconstruct one reliable estimate of source time series, we resort to bootstrapping to compare the

differences in pairwise source-level coherence between rare and frequent perceivers. Initially for each relevant parcel, from the trials pooled over participants in each group (rare and frequent), 500 trials were randomly picked to form two sets of random partition. Subsequently, pairwise coherence between the parcels was computed on a trial-by-trial basis and averaged for rare and frequent perceivers separately.

The aforementioned steps were repeated 500 times to generate a distribution of pairwise coherence values for frequent and rare perceivers. Significance of pairwise coherence difference values between the rare and frequent perceivers was evaluated from bootstrapped estimates.

## 3 | RESULTS


### 3.1 | Variability in multisensory perception

Incongruent McGurk stimulus, visual */kal* paired with audio */pal* also referred as */pal-/kal*, and three kinds of congruent AV stimuli—*/tal-/tal*, */pal-/pal*, and */kal-/kal*—were presented to all participants. The sequence of presentation was randomized. As the participants observed each of the stimuli, they were instructed to report with a set of four keys if they had heard */tal*, */pal*, */kal*, or “something else.”

Response tendency (*inter-trial variability*), computed as the relative proportion of illusory (*/tal*) responses in all McGurk trials across all participants, showed that */tal* percept was reported in 58.19% ( $SD = 32.33$ ) of trials, whereas unisensory */pal* percept was reported in 37.56% ( $SD = 31.79$ ) trials (Figure 4a). However, paired two-sample *t* tests comparing the percentage of illusory */tal* and unisensory */pal* percept over all the participants revealed no significant difference ( $t_{17} = 1.36$ ,  $p = .19$ ). Congruent AV stimuli (*/tal*, */pal*, and */kal*) were correctly identified in 96.56% ( $SD = 2.88$ ) trials by all the participants.

A high degree of *inter-individual* variability in McGurk susceptibility was observed (Figure 4b) while analyzing participant-wise */tal* percept during McGurk stimulus. The participants were classified based on their McGurk susceptibility in the incongruent trials into two groups—rare perceivers ( $<50\%$  */tal* percept) and frequent perceivers ( $>50\%$  */tal* percept). Eight of the eighteen participants elicited  $<50\%$  propensity of perceiving the McGurk effect. A two-way analysis of variance with congruent AV stimuli (*/pal*, */tal*, and */kal*) and participant groups (frequent and rare) as factors revealed that only the type of congruent stimulus affects the hit rate,  $F(2, 53) = 4.44$ ,  $p = .02$ , and the effect of participant group (frequent or rare) on the hit rate was not significant,  $F(1, 53) = 4.18$ ,  $p = .05$ . Also, there was no evidence of interaction between the congruent AV stimulus type and the participant group  $F(2, 53) = 0.07$ ,  $p = .937$ . We further performed paired two-sample *t* test to compare





**FIGURE 4** Behavior: Behavior: (a) *Inter-trial variability*—Percentage of */ta/* (illusory) and */pa/* (unisensory) percept during the presentation of McGurk stimulus and the hit rate during the presentation of congruent AV stimuli (*/pa/*, */ta/*, and */ka/*) averaged over all the participants. The red mark in each box plot indicates the median. (b) *Inter-individual variability*—Propensity of McGurk effect for each of the 18 participants expressed as the percentage of */ta/* percept during the presentation of McGurk stimulus. The participants were categorized as frequent perceivers (blue diamonds, >50%) and rare perceivers (red diamonds, ≤50%) [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

the differences in the hit rate during the congruent stimuli. We observed that hit rate during congruent AV stimulus */pa/* was significantly lower than hit rate during congruent AV stimuli: */ta/* ( $t_{17} = -2.06$ ,  $p = .047$ ) and */ka/* ( $t_{17} = -2.97$ ,  $p = .005$ ).

### 3.2 | Large-scale functional connectivity in EEG sensor space

The emergence of specific cognitive trait is subserved by the coordination of distributed functionally specialized neuronal assemblies or brain regions (Bressler, 1995; Bressler & Menon, 2010; Varela et al., 2001). Based on our previous study (Kumar et al., 2016), we hypothesized that neural signatures of inter-individual can be quantified using the global coherence metric that captures the degree of neuronal synchronization over the entire brain from EEG scalp-level recordings.


We observed a significant negative correlation between each participant's alpha band global coherence and their McGurk susceptibility during incongruent McGurk condition (*/pa/-/ka/*),  $r_s(18) = -0.49$ ,  $p = .04$  (Figure 5a), congruent */ta/-/ta/*,  $r_s(18) = -0.51$ ,  $p = .03$  (Figure 5b),

and */ka/-/ka/*,  $r_s(18) = -0.49$ ,  $p = .04$  (Figure 5c) stimuli. However, alpha modulation among participants during congruent */pa/-/pa/* condition was not correlated with their McGurk susceptibility,  $r_s(18) = -0.18$ ,  $p = .47$  (Figure 5d). Evidently, there was also no correlation between global coherence in any other frequency bands and McGurk susceptibility during incongruent McGurk (*/pa/-/ka/*),  $\theta$ :  $r_s(18) = -0.29$ ,  $p = .22$ ,  $\beta$ :  $r_s(18) = -0.21$ ,  $p = .39$ ,  $\gamma$ :  $r_s(18) = -0.10$ ,  $p = .45$ ; congruent */pa/*,  $\theta$ :  $r_s(18) = -0.38$ ,  $p = .12$ ,  $\beta$ :  $r_s(18) = -0.14$ ,  $p = .57$ ,  $\gamma$ :  $r_s(18) = -0.15$ ,  $p = .53$ ; congruent */ta/*,  $\theta$ :  $r_s(18) = -0.36$ ,  $p = .14$ ,  $\beta$ :  $r_s(18) = -0.17$ ,  $p = .50$ ,  $\gamma$ :  $r_s(18) = -0.19$ ,  $p = .42$ , and congruent */ka/* ( $\theta$ :  $r_s(18) = -0.31$ ,  $p = .20$ ,  $\beta$ :  $r_s(18) = -0.22$ ,  $p = .37$ ,  $\gamma$ :  $r_s(18) = -0.16$ ,  $p = .50$  stimuli).

### 3.3 | A simple interacting neural mass model with fast, intermediate, and slow time constants explains de-synchronization of alpha band global coherence

Global coherence dynamics may reflect the potential operation of a large-scale network that supports multisensory speech

**FIGURE 5** Large-scale functional connectivity: Mean alpha band coherence in each participant plotted against that participant's susceptibility of McGurk effect during stimulus: (a) McGurk, (b) congruent */ta/*, (c) congruent */ka/*, and (d) congruent */pa/* [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



perception. We hypothesized the variability in McGurk percept across participants to be grounded in specific interactions between neural assemblies (Figure 6). In order to account for the observed global coherence patterns, we employed a neural mass model comprising three nodes broadly representing the auditory and visual areas in the sensory cortices, and the multisensory node representing the collection of multisensory (or modality nonspecific) areas. Each node comprised a population of excitatory and inhibitory Hindmarsh–Rose (HR) neurons (Hindmarsh & Rose, 1984) operating at different physiologically motivated time scales (see Section 2 for more details). Model parameter details are presented in Table 1.

We sought to explore the global coherence while varying the configuration and strength of coupling among the nodes in our model. For instance, in Figure 6a auditory node's excitatory coupling of  $+W_{AV}$  on multisensory node was balanced with inhibitory coupling of the same strength ( $-W_{AV}$ ) from the multisensory node. In such a configuration, auditory node is being the source of excitation and is referred as the source, whereas multisensory node is referred as the sink as all excitatory couplings are directed toward it. The visual node in such configuration behaves as both source and sink.

The coexistence of coherence in alpha and gamma band facilitated by the two time scales of visual (slow) and auditory (fast) nodes was observed in network configuration wherein visual node operates as the source and auditory node acts as a sink (Figure 6g). Moreover, the concurrence of the coherence in alpha and gamma bands emerged in the coupling between auditory and visual regions (coupling parameter ranging between 0.35 and 0.7) via two routes—(a) direct coupling ( $W_{AV}$ ) and (b) indirect coupling ( $W_{AM}$  &  $W_{VM}$ ) involving multisensory node (M). We assumed coupling strength  $<0.35$  to be weak coupling (WC), coupling strength between 0.35 and 0.7 to be moderate coupling (MC), and coupling greater than 0.7 to be strong coupling (SC). In SC range, we observed high coherence in alpha band and gamma bands; however, no peak was observed in alpha band. In the MC range, none of the model configurations were able to generate concurrent coherent states in alpha and gamma bands (Figure 6a–h), except Figure 6g, where visual node acts as the source of excitation. Furthermore, direct coupling between auditory and visual nodes alone ( $W_{AV}$  ranges from 0 to 1,  $W_{AM}$  &  $W_{VM} = 0$ ) results only in alpha band coherence (Figure 6j). Interestingly, emergence of only gamma band coherence was observed when the indirect coupling in the model was varied between A–V nodes via multisensory node ( $W_{AM}$  &  $W_{VM}$  range from 0 to 1,  $W_{AV} = 0$ ) in MC range (Figure 6k).

### 3.3.1 | Direct audio–visual interaction underpins inter-individual variability

Our empirical findings indicate a significant negative correlation between participant-wise alpha band coherence and


McGurk susceptibility. We hypothesized mediation of interaction between auditory (A) and visual (V) nodes via the multisensory node (M; i.e., stronger A–M and V–M coupling than A–V coupling) to be characteristic of participants with higher susceptibility to the McGurk effect. To test this hypothesis, a balanced network coupling state,  $W_{AM} = W_{VM} = W_{AV} = 0.35$  where alpha and gamma band coherences coexist was considered, following which the change in the coherence peaks with decreasing direct A–V coupling ( $W_{AV}$ ) was studied. A suppression of alpha coherence peak with decrease in  $W_{AV}$  was observed (Figure 7); however, gamma coherence peak remained more or less unchanged. Only when A–V coupling in the computational model became negligible ( $W_{AV} < 0.05$ ), disappearance of alpha coherence peak was observed. We use this basic observation to revisit the empirical data and ask whether there was evidence for a disconnection or uncoupling between auditory and visual sources in subjects who evinced differential alpha coherence.

### 3.4 | Validation of model predictions using source connectivity analysis


Finally, we obtained source-level time series for each brain parcel by projecting single-trial time series to the source-space by multiplying them with spatial filter of the respective parcel using voxel to parcel mapping provided by AAL atlas (see Section 2 for details). Subsequently, the pairwise coherence—a measure of functional connectedness—between the parcels associated with auditory and visual areas was computed. Auditory-specific parcel, superior temporal gyrus (STG), and the visual parcels, superior occipital cortex (SOC), medial occipital cortex (MOC), and inferior occipital cortex (IOC), were identified. A significant increase of 0.8% and 0.5% in the functional connectivity in rare perceivers than frequent perceivers was observed between left STG and left SOC (mean coherence increase = 0.0071,  $SD = 0.0015$ ), and between left STG and left MOC (mean coherence increase = 0.0053,  $SD = 0.0020$ ), respectively (Figure 8 and Table 2). Percentage increase in the aforementioned coherence differences in rare perceivers compared to frequent perceivers between the auditory and visual parcels in both hemispheres is summarized in Table 2. Achieved significance levels were estimated using normal intervals as the bootstrapped estimates of coherence differences followed normal distribution (Wasserman, 2005). Normality of each coherence distribution was tested using Lilliefors test where the null hypothesis that bootstrapped coherence values come from a normal distribution could not be rejected at 95% significance. To cross-check the results, we ran several permutations of the comparison, and the main pattern of left STG–SOC and left STG–MOC connectivity being significantly higher in rare perceivers did not change.



## Plausible coupling scenarios



## Prediction of alpha and gamma band coherence



→ Excitatory coupling (EC)

**WC** Weak coupling range

**MC** Moderate coupling range

**SC** Strong coupling range

○ Source of EC

● Sink of EC

◐ Both source and sink of EC

**V<sub>S</sub>** Visual node with slow time constant

**A<sub>F</sub>** Auditory node with fast time constant

**M<sub>I</sub>** Multisensory node with intermediate time constant



## 4 | DISCUSSION

An ongoing challenge in multisensory speech perception community is to accurately identify and characterize the neural mechanisms that govern the perceptual variability across individuals. Traditionally, studies have focused on identifying the neural correlates in terms of candidate brain regions or interactions among specific regions of interest that are responsible for the observed heterogeneity (Nath & Beauchamp, 2012). A more emerging understanding, however, suggests the existence of large-scale networks of brain regions facilitating perceptual processing (Beauchamp, 2015; Keil et al., 2012; Kumar et al., 2017). Neuronal oscillations have been identified as a key marker of neuronal information processing in the relevant brain regions that needs to be fully explored to shape an individual's perceptual experience (Keil et al., 2012; Keil & Senkowski, 2018; Kumar et al., 2016, 2017). Robust oscillations observed from macroscopic recordings such as EEG/MEG are an outcome of network interactions among local subpopulations of excitatory and inhibitory neurons (Becker, Knock, Ritter, & Jirsa, 2015; Deco, Rolls, & Romo, 2010; Wilson & Cowan, 1972). Empirically, such interactions result in global coherence dynamics (Kumar et al., 2017). Our study demonstrates how distinct global coherence patterns correlate with inter-individual heterogeneity. Subsequently, using a computational model of interacting large-scale brain networks, we explain the plausible neural mechanisms that generate the coherence dynamics associated with the observed heterogeneity. Finally, we validate the plausible mechanism that emerged from the model about the connection parameters among interacting neural subsystems using source-space functional connectivity analysis. Put together, we present a biologically plausible mechanistic proposal that underlies the observed inter-individual variability in multisensory speech perception.

The key observations in our study are as follows: (a) Susceptibility of individuals' McGurk effect is negatively correlated to their alpha band global coherence—indicating that the degree of synchronization of large-scale neural assemblies in the alpha band underlies inter-individual variability; (b) weakened direct A–V coupling in the neural mass model leads to the decrease in alpha band coherence; and (c) individuals with higher McGurk susceptibility elicited decreased pairwise connectivity between the sensory-specific (audio and visual) areas than individuals with lower McGurk susceptibility.

### 4.1 | Global coherence as the index of perceptual variability


Inter-individual differences observed in our study replicate (including reports from our group) and extend previous studies. 44% of our participants had more than 80% McGurk percept which lies within the range of 26%–100% reported in previous studies (Benoit, Raij, Lin, Jääskeläinen, & Stufflebeam, 2009; MacDonald, Andersen, & Bachmann, 2000; McGurk & Macdonald, 1976; Nath & Beauchamp, 2012; Figure 4a). Also, our mean McGurk percept (collapsed across participants) of 58% is within the literature range of 24%–100% (Benoit et al., 2009; MacDonald et al., 2000; McGurk & Macdonald, 1976; Nath & Beauchamp, 2012; Strand, Cooperman, Rowe, & Simenstad, 2014; Figure 4a,b). Hence, the behavioral results give confidence that the corresponding brain patterns will be of robust significance and general applicability. Our results show that the extent of de-synchronization of the cortical network in the alpha band indexes an individual's McGurk susceptibility (Figure 5a–c). This finding has two key elements. First, the inverse relationship between McGurk susceptibility and alpha band global coherence was consistent for McGurk stimulus and the congruent AV stimuli (*lta* and *kla*) as well. This highlights that our observations indeed reflect inter-individual differences/perceptual categorization and not necessarily stimulus configurations. Second, it pinpoints that inter-individual variability is contingent on the functional state of the brain which can be captured by exploring the dynamics of the large-scale cortical network. The syllables *lta* and *kla* can be difficult to distinguish with visual cues alone, but acoustically are different. Therefore, coincidence of AV cues allows both the group of perceivers to reap the benefits of AV integration during congruent AV stimuli *lta* and *kla*. However, the bilabial syllable *pta* is very different visually from *lta* and *kla*. Hence, identification of the congruent AV syllable *pta* does not necessarily require AV integration. Therefore, we do not see any significant correlation during congruent *pta* stimulus (Figure 5d).

Previous evidences accentuate the modulations in alpha band coherence to central executive processes (Klimesch, 1999; Sauseng, Klimesch, Schabus, & Doppelmayr, 2005) that are postulated to be involved in allocating working memory storage to phonological loop


**FIGURE 6** Prediction of alpha and gamma coherences from neural mass model: (a–h) Exploration of various coupling scenarios to identify whether it is possible to generate alpha and gamma coherence in moderate coupling range. Only the model configuration (g) exhibits concurrent coherence in the alpha and gamma bands in the moderate coupling range. (i) Alpha and gamma band coherences coexist in moderate coupling range when visual node is source and auditory node is the sink (j) Only direct A–V coupling generates alpha coherence independently. (k) Indirect A–V coupling via multisensory node generates gamma coherence at the limit case scenario of weak direct coupling. Note: (reciprocal) inhibitory connections have been omitted for clarity [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

**FIGURE 7** Mechanistic understanding of inter-individual variability: Alpha desynchronization characteristic of frequent perceivers resulted due to negligible A–V coupling [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

### Inter-individual variability



**FIGURE 8** Pairwise connectivity between the parcels: Pairwise coherence between: (a) left superior temporal gyrus (STG) and left superior occipital cortex (SOC), (b) right STG and right SOC, (c) left STG and left medial occipital cortex (MOC), (d) right STG and right MOC, (e) left STG and left inferior occipital cortex (IOC), and (f) right STG and right IOC in rare and frequent perceivers. The brain images represent parcels between which the pairwise coherence is computed [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]





**TABLE 2** Summary of the source-level coherence difference between the parcels from rare and frequent perceivers (bootstrapped estimates, see Section 2)

| Connectivity | Left hemisphere |                 |        |                                            | Right hemisphere |                 |        |                                            |
|--------------|-----------------|-----------------|--------|--------------------------------------------|------------------|-----------------|--------|--------------------------------------------|
|              | % Increase      | Mean difference | SD     | Achieved significance level (ASL) <i>p</i> | % Increase       | Mean difference | SD     | Achieved significance level (ASL) <i>p</i> |
| STG-SOC      | 0.8             | 0.0071          | 0.0015 | <.0001*                                    | 0.2              | 0.0016          | 0.0017 | .53                                        |
| STG-MOC      | 0.5             | 0.0053          | 0.0020 | <.0001*                                    | 0.4              | 0.0038          | 0.0015 | .12                                        |
| STG-IOC      | 0.3             | 0.0035          | 0.0010 | .16                                        | 0.3              | 0.0028          | 0.0016 | .09                                        |

\*Significance.

that maintains verbal information, and the visuo-spatial sketchpad that maintains transient visuo-spatial information (Baddeley, 1992). Such processes are involved in cross-modal perception that requires the integration of information from distributed neuronal assemblies. Taken together, communication via alpha coherence may be the most plausible neuromarker of inter-group variability at the network level bearing the signature of integration.

#### 4.2 | Multisensory node operational at intermediate time scale renders communication effective between sensory-specific nodes operating at slow and fast time scales

Neural mass models that operate to explain mesoscopic and macroscopic features of dynamics in EEG/MEG data offer an attractive tool to understand the underlying neural mechanisms (David & Friston, 2003; Jansen & Rit, 1995; Lopes da Silva, Hoeks, Smits, & Zetterberg, 1974). The time scale of processing in our computational model is most disparate for the auditory and visual nodes, with auditory the fastest and visual the slowest (Rosen & Howell, 2011; Williams et al., 2004). Without the presence of an intermediate time scale, only alpha coherence is sustained by the neural mass model within biologically relevant parameter regimes (Figure 6). On introduction of another neural mass operating at intermediate time scale, synchronization in the gamma band emerges because higher dimensionality of the resultant dynamical system allows creation of another mode of communication exchange. Hence, our model suggests that emergence of gamma band coherence could be attributed to the communication between areas of auditory and visual cortices gated through multisensory areas such as STS (Figure 7). Our suggestion is in line with earlier observations of visual stimuli modulating auditory perception either directly resulting in alpha band coherence (Kayser, Petkov, & Logothetis, 2008) or indirectly via higher order regions resulting primarily in gamma band coherence (Kayser & Logothetis, 2009; Maier, Chandrasekaran, & Ghazanfar, 2008).

Drawing from the model, inter-individual differences primarily emerge from the differences in the A–V coupling. A stronger A–V coupling implies a lesser mediation via the multisensory node (stronger A–M and V–M) leading to weaker integration which is akin to rare perceivers. Conversely, lesser A–V coupling implies a larger involvement of the multisensory node (stronger A–M and V–M) in mediating the integration of the sensory inputs that is characteristic of frequent perceivers. Findings from our model corroborate with the earlier findings that pinpoint the significance of functional connectedness of the multisensory regions like STS to sensory-specific, frontal, and parietal regions in facilitating cross-modal perception (Keil et al., 2012).

There are a number of qualifications that attend this work and its interpretation. In brief, our exploration of different connectivity architectures was limited to extrinsic (i.e., between nodes) coupling differences. These differences were both in terms of the arrangement of the connections and connection strengths. However, we did not consider intrinsic connectivity, namely within-node connections. The role of intrinsic connectivity in accounting for inter-individual differences is something that we hope to explore in the future. This is a potentially important avenue of enquiry, given the sensitivity of global dynamics to subtle changes in intrinsic connectivity and excitability (Freyer, Roberts, Ritter, & Breakspear, 2012; Gilson, Moreno-Bote, Ponce-Alvarez, Ritter, & Deco, 2016; Roy et al., 2014). This is particularly prescient from the point of view of predictive coding models of multisensory integration. In this setting, the precision assigned to various sources of information (i.e., prediction errors) is thought to be mediated by intrinsic excitability (or self-inhibition; Bhatt et al., 2016; Brown & Friston, 2012; Pinotsis et al., 2014). We also note, in contrast to the neural mass models used in dynamic causal modeling (Bastos et al., 2012, 2015; Litvak, Garrido, Zeidman, & Friston, 2015), that the designation of forward and backward is based purely upon the distinction between excitatory and inhibitory connectivity. In biologically plausible models, this designation usually rests upon laminar specificity (Markov et al., 2013; Shaw et al., 2019). However,



our neural mass model did not distinguish between different excitatory and inhibitory populations in different cortical layers. It would be interesting to reproduce the analysis reported above using nuanced models of the sort used in dynamic causal models based on the canonical microcircuit. Provisional work in this area has already considered A/V sensory integration (Li, Xi, Zhang, Liu, & Tang, 2019); however, it has not yet addressed the cardinal issue of inter-individual variability.

### 4.3 | Dynamic coupling between audio and visual cortices underlies perceptual variability

Most importantly, we validated our predictions from the dynamical model to the empirical estimates of source-level functional connectivity. Our pairwise connectivity between the parcels (Figure 8) shows increased connectivity between the auditory and visual parcels in rare than frequent perceivers. Interestingly, we observed this increase only in the left hemisphere which corroborates the earlier findings accentuating left laterality in multisensory speech perception (Beauchamp, 2010; Keil et al., 2012; Nath & Beauchamp, 2012). Notably, for our source reconstruction, we used a form of beamforming (DICS) to identify sources that were likely to contribute the most power in sensor space. While this was sufficient for our purposes, this kind of source reconstruction has to be used carefully when subsequently measuring coherence among sources. This is because the scheme used to identify or estimates sources suppresses coherent activity between sources. It will be interesting to perform a more detailed source reconstruction that embodies our prior knowledge about the key nodes or sources involved in the McGurk effect, for example, using equivalent current dipole reconstructions that are guided by the functional magnetic resonance imaging (fMRI) literature in this area.

The analysis pipeline (Figure 1) can be generalized easily to other studies of complex perceptual phenomena. For example, alpha and/or gamma coherences have been observed in AV perception studies involving speech phrases (Doesburg, Emberson, Rahi, Cameron, & Ward, 2008), natural scenes (Kayser et al., 2008), and also artificially generated A–V looming signals (Maier et al., 2008). Modulations in A–V interactions that explain differences in McGurk susceptibility in our study could also explain the increase in alpha phase consistency observed during natural AV scenes in rhesus monkeys (Kayser et al., 2008). Our model demonstrates that increase in the interaction between neural nodes operating at fast and slow time scales via a node operating at an intermediate time scale results in increased gamma and decreased alpha and beta band coherence. However, a slight tweak in the model topology elicits coherence modulations in varied frequency bands. A basic understanding that multifrequency

coherent states of the network emerge from their difference in time scales can be used to explain a large repertoire of empirical observations.

Although our model of three interacting neural masses with different time constants that generate band-specific coherences can come across as a canonical model of neuronal network dynamics, certain limitations exist. Multi-parametric and unbounded nature of the parameter space results in myriads of dynamics including chaos (Stefanescu & Jirsa, 2008). Therefore, such models cannot be used to directly fit the data using optimization techniques and should be limited to use as a phenomenological or minimalistic model in providing mechanistic insights into many findings (Engel, Senkowski, & Schneider, 2011; Fries, 2015)—both in normal and in clinical population. Also, this kind of neural mass model can be contrasted with the neural mass models used in dynamic causal modeling, which have the fixed point attractors and can be used to fit observed EEG time series or spectra.

### ACKNOWLEDGEMENTS

This study was supported by NBRC Core funds, Ramalingaswami Fellowships (Department of Biotechnology, Government of India) to DR (BT/RLF/Re-entry/07/2014) and AB (BT/RLF/Re-entry/31/2011), and Innovative Young Biotechnologist Award (IYBA) to AB (BT/07/IYBA/2013). DR was supported by SR/CSRI/21/2016 extramural grant from the Department of Science and Technology (DST) Ministry of Science and Technology, Government of India.

### CONFLICT OF INTEREST

Authors declare no conflict of interest.

### DATA AVAILABILITY STATEMENT

Raw and processed data used for generation of figures from empirical data as well as large-scale dynamical model will be shared on request to corresponding author.

### ORCID

Vinodh G. Kumar  <https://orcid.org/0000-0002-4604-7379>  
 Shrey Dutta  <https://orcid.org/0000-0001-9692-1645>  
 Dipanjan Roy  <https://orcid.org/0000-0002-1669-1083>  
 Arpan Banerjee  <https://orcid.org/0000-0002-3725-0463>

### REFERENCES

- Baddeley, A. (1992). Working memory. *Science*, 255(5044), 556–559. <https://doi.org/10.1126/science.1736359>
- Bastos, A. M., Litvak, V., Moran, R., Bosman, C. A., Fries, P., & Friston, K. J. (2015). A DCM study of spectral asymmetries in feedforward and feedback connections between visual areas V1 and V4 in the monkey. *NeuroImage*, 108, 460–475. <https://doi.org/10.1016/j.neuroimage.2014.12.081>
- Bastos, A. M., Usrey, W. M., Adams, R. A., Mangun, G. R., Fries, P., & Friston, K. J. (2012). Canonical microcircuits for

- predictive coding. *Neuron*, 76(4), 695–711. <https://doi.org/10.1016/j.neuron.2012.10.038>
- Beauchamp, M. S. (2010). fMRI-guided TMS reveals that the STS is a cortical locus of the McGurk effect. *Journal of Neuroscience*, 30(7), 2414–2417. <https://doi.org/10.1523/JNEUROSCI.4865-09.2010>. fMRI-guided
- Beauchamp, M. S. (2015). The social mysteries of the superior temporal sulcus. *Trends in Cognitive Sciences*, 19(9), 489–490. <https://doi.org/10.1016/j.tics.2015.07.002>
- Becker, R., Knock, S., Ritter, P., & Jirsa, V. (2015). Relating alpha power and phase to population firing and hemodynamic activity using a Thalamo-cortical neural mass model. *PLoS Computational Biology*, 11(9), e1004352. <https://doi.org/10.1371/journal.pcbi.1004352>
- Benoit, M. M., Raji, T., Lin, F.-H., Jääskeläinen, I. P., & Stufflebeam, S. (2009). Primary and multisensory cortical activity is correlated with audiovisual percepts. *Human Brain Mapping*, 31(4). <https://doi.org/10.1002/hbm.20884>
- Bhatt, M. B., Bowen, S., Rossiter, H. E., Dupont-Hadwen, J., Moran, R. J., Friston, K. J., & Ward, N. S. (2016). Computational modelling of movement-related beta-oscillatory dynamics in human motor cortex. *NeuroImage*, 133, 224–232. <https://doi.org/10.1016/j.neuroimage.2016.02.078>
- Bizley, J. K., & King, A. J. (2011). What can multisensory processing tell us about the functional organization of auditory cortex? In M. M. Murray & M. T. Wallace (Eds.), *The neural bases of multisensory processes*, (pp. 31–48). Boca Raton, FL: CRC Press/Taylor & Francis. Retrieved from <https://doi.org/10.1201/b11092-5>
- Bressler, S. L. (1995). Large-scale cortical networks and cognition. *Brain Research Reviews*, 20(3), 288–304. [https://doi.org/10.1016/0165-0173\(94\)00016-1](https://doi.org/10.1016/0165-0173(94)00016-1)
- Bressler, S. L., & Kelso, J. A. S. (2001). Cortical coordination dynamics and cognition. *Trends in Cognitive Sciences*, 5(1), 26–36. [https://doi.org/10.1016/S1364-6613\(00\)01564-3](https://doi.org/10.1016/S1364-6613(00)01564-3)
- Bressler, S. L., & Menon, V. (2010). Large-scale brain networks in cognition: Emerging methods and principles. *Trends in Cognitive Sciences*, 14(6), 277–290. <https://doi.org/10.1016/j.tics.2010.04.004>
- Brown, H. R., & Friston, K. J. (2012). Dynamic causal modelling of precision and synaptic gain in visual perception – An EEG study. *NeuroImage*, 63(1), 223–231. <https://doi.org/10.1016/j.neuroimage.2012.06.044>
- Cimenser, A., Purdon, P. L., Pierce, E. T., Walsh, J. L., Salazar-Gomez, A. F., Harrell, P. G., ... Brown, E. N. (2011). Tracking brain states under general anesthesia by using global coherence analysis. *Proceedings of the National Academy of Sciences of the United States of America*, 108(21), 8832–8837. [https://doi.org/10.1073/pnas.108\(21\).8832-8837](https://doi.org/10.1073/pnas.108(21).8832-8837)
- Cuppini, C., Shams, L., Magosso, E., & Ursino, M. (2017). A biologically inspired neurocomputational model for audiovisual integration and causal inference. *European Journal of Neuroscience*, 46(9), 2481–2498. <https://doi.org/10.1111/ejn.13725>
- David, O., & Friston, K. J. (2003). A neural mass model for MEG/EEG: Coupling and neuronal dynamics. *NeuroImage*, 20(3), 1743–1755. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/14642484>
- Deco, G., Pérez-Sanagustín, M., de Lafuente, V., & Romo, R. (2007). Perceptual detection as a dynamical bistability phenomenon: A neurocomputational correlate of sensation. *Proceedings of the National Academy of Sciences of the United States of America*, 104(50), 20073–20077. <https://doi.org/10.1073/pnas.0709794104>
- Deco, G., Rolls, E. T., & Romo, R. (2010). Synaptic dynamics and decision making. *Proceedings of the National Academy of Sciences of the United States of America*, 107(16), 7545–7549. <https://doi.org/10.1073/pnas.1002333107>
- Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, 134(1), 9–21. <https://doi.org/10.1016/j.jneumeth.2003.10.009>
- Doesburg, S. M., Emberson, L. L., Rahi, A., Cameron, D., & Ward, L. M. (2008). Asynchrony from synchrony: Long-range gamma-band neural synchrony accompanies perception of audiovisual speech asynchrony. *Experimental Brain Research*, 185(1), 11–20. <https://doi.org/10.1007/s00221-007-1127-5>
- Engel, A. K., Senkowski, D., & Schneider, T. R. (2011). Multisensory integration through neural coherence. In M. M. Murray & M. T. Wallace (Eds.), *The neural bases of multisensory processes*, (pp. 115–130). Boca Raton, FL: CRC Press/Taylor & Francis. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/22593880>
- Falchier, A., Clavagnier, S., Barone, P., & Kennedy, H. (2002). Anatomical evidence of multimodal integration in primate striate cortex. *The Journal of Neuroscience*, 22(13), 5749–5759. <https://doi.org/10.1523/JNEUROSCI.22-13-05749.2002>
- Freyer, F., Roberts, J. A., Ritter, P., & Breakspear, M. (2012). A canonical model of multistability and scale-invariance in biological systems. *PLoS Computational Biology*, 8(8), e1002634. <https://doi.org/10.1371/journal.pcbi.1002634>
- Fries, P. (2015). Rhythms for cognition: Communication through coherence. *Neuron*, 88(1), 220–235. <https://doi.org/10.1016/j.neuron.2015.09.034>
- Gilson, M., Moreno-Bote, R., Ponce-Alvarez, A., Ritter, P., & Deco, G. (2016). Estimation of directed effective connectivity from fMRI functional connectivity hints at asymmetries of cortical connectome. *PLoS Computational Biology*, 12(3), e1004762. <https://doi.org/10.1371/journal.pcbi.1004762>
- Gold, C., Henze, D. A., Koch, C., & Buzsáki, G. (2006). On the origin of the extracellular action potential waveform: A modeling study. *Journal of Neurophysiology*, 95(5), 3113–3128. <https://doi.org/10.1152/jn.00979.2005>
- Gross, J., Kujala, J., Hamalainen, M., Timmermann, L., Schnitzler, A., & Salmelin, R. (2001). Dynamic imaging of coherent sources: Studying neural interactions in the human brain. *Proceedings of the National Academy of Sciences of the United States of America*, 98(2), 694–699. <https://doi.org/10.1073/pnas.98.2.694>
- Halder, T., Talwar, S., Jaiswal, A. K., & Banerjee, A. (2019). Quantitative evaluation in estimating sources underlying brain oscillations using current source density methods and beamformer approaches. *Eneuro*, 6(4), ENEURO.0170-19.2019. <https://doi.org/10.1523/ENEURO.0170-19.2019>
- Haxby, J. V., Hoffman, E. A., & Gobbini, M. I. (2000). The distributed human neural system for face perception. *Trends in Cognitive Sciences*, 4(6), 223–233. [https://doi.org/10.1016/S1364-6613\(00\)01482-0](https://doi.org/10.1016/S1364-6613(00)01482-0)
- Hindmarsh, J. L., & Rose, R. M. (1984). A model of neuronal bursting using three coupled first order differential equations. *Proceedings of the Royal Society of London. Series B, Biological Sciences*, 221(1222), 87–102. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/6144106>
- Jansen, B. H., & Rit, V. G. (1995). Electroencephalogram and visual evoked potential generation in a mathematical model of coupled cortical columns. *Biological Cybernetics*, 73(4), 357–366. <https://doi.org/10.1007/BF00199471>



- Kayser, C., & Logothetis, N. K. (2009). Directed interactions between auditory and superior temporal cortices and their role in sensory integration. *Frontiers in Integrative Neuroscience*, 3, 7. <https://doi.org/10.3389/neuro.07.007.2009>
- Kayser, C., Petkov, C. I., & Logothetis, N. K. (2008). Visual modulation of neurons in auditory cortex. *Cerebral Cortex*, 18(7), 1560–1574. <https://doi.org/10.1093/cercor/bhm187>
- Keil, J., Muller, N., Ihssen, N., & Weisz, N. (2012). On the variability of the McGurk effect: Audiovisual integration depends on pre-stimulus brain states. *Cerebral Cortex*, 22(1), 221–231. <https://doi.org/10.1093/cercor/bhr125>
- Keil, J., & Senkowski, D. (2018). Neural oscillations orchestrate multi-sensory processing. *The Neuroscientist*, 24(6), 609–626. <https://doi.org/10.1177/1073858418755352>
- Kirschstein, T., & Köhling, R. (2009). What is the source of the EEG? *Clinical EEG and Neuroscience*, 40(3), 146–149. <https://doi.org/10.1177/155005940904000305>
- Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: A review and analysis. *Brain Research Reviews*, 29, 169–195. [https://doi.org/10.1016/S0165-0173\(98\)00056-3](https://doi.org/10.1016/S0165-0173(98)00056-3)
- Körding, K. P., Beierholm, U., Ma, W. J., Quartz, S., Tenenbaum, J. B., & Shams, L. (2007). Causal inference in multisensory perception. *PLoS ONE*, 2(9), e943. <https://doi.org/10.1371/journal.pone.0000943>
- Kumar, G. V., Halder, T., Jaiswal, A. K., Mukherjee, A., Roy, D., & Banerjee, A. (2016). Large scale functional brain networks underlying temporal integration of audio-visual speech perception: An EEG study. *Frontiers in Psychology*, 7, 1558. <https://doi.org/10.3389/fpsyg.2016.01558>
- Kumar, G. V., Kumar, N., Roy, D., & Banerjee, A. (2017). Segregation and integration of cortical information processing underlying cross-modal perception. *Multisensory Research*, 31(5), 481–500. <https://doi.org/10.1163/22134808-00002574>
- Li, Q., Xi, Y., Zhang, M., Liu, L., & Tang, X. (2019). Distinct mechanism of audiovisual integration with informative and uninformative sound in a visual detection task: A DCM study. *Frontiers in Computational Neuroscience*, 13, 59. <https://doi.org/10.3389/fncom.2019.00059>
- Litvak, V., Garrido, M., Zeidman, P., & Friston, K. (2015). Empirical Bayes for group (DCM) studies: A reproducibility study. *Frontiers in Human Neuroscience*, 9(DEC), 670. <https://doi.org/10.3389/fnhum.2015.00670>
- Lopes da Silva, F. H., Hoeks, A., Smits, H., & Zetterberg, L. H. (1974). Model of brain rhythmic activity. The alpha-rhythm of the thalamus. *Kybernetik*, 15(1), 27–37. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/4853232>
- Ma, W. J., Beck, J. M., & Pouget, A. (2008). Spiking networks for Bayesian inference and choice. *Current Opinion in Neurobiology*, 18(2), 217–222. <https://doi.org/10.1016/j.conb.2008.07.004>
- MacDonald, J., Andersen, S., & Bachmann, T. (2000). Hearing by eye: How much spatial degradation can be tolerated? *Perception*, 29(10), 1155–1168. <https://doi.org/10.1068/p3020>
- Maier, J. X., Chandrasekaran, C., & Ghazanfar, A. A. (2008). Integration of bimodal looming signals through neuronal coherence in the temporal lobe. *Current Biology*, 18(13), 963–968. <https://doi.org/10.1016/j.cub.2008.05.043>
- Markov, N. T., Ercsey-Ravasz, M., Van Essen, D. C., Knoblauch, K., Toroczkai, Z., & Kennedy, H. (2013). Cortical high-density counterstream architectures. *Science*, 342(6158), 1238406–1238406. <https://doi.org/10.1126/science.1238406>
- McGurk, H., & Macdonald, J. (1976). Hearing lips and seeing voices. *Nature*, 264, 691–811. <https://doi.org/10.1038/264746a0>
- Mitra, P., & Bokil, H. (2008). *Observed brain dynamics*. New York, NY: Oxford Univ Press.
- Moreno-Bote, R., Rinzel, J., & Rubin, N. (2007). Noise-induced alternations in an attractor network model of perceptual bistability. *Journal of Neurophysiology*, 98(3), 1125–1139. <https://doi.org/10.1152/jn.00116.2007>
- Nath, A. R., & Beauchamp, M. S. (2011). Dynamic changes in superior temporal sulcus connectivity during perception of noisy audiovisual speech. *The Journal of Neuroscience*, 31(5), 1704–1714. <https://doi.org/10.1523/JNEUROSCI.4853-10.2011>
- Nath, A. R., & Beauchamp, M. S. (2012). A neural basis for interindividual differences in the McGurk effect, a multisensory speech illusion. *NeuroImage*, 59(1), 781–787. <https://doi.org/10.1016/j.neuroimage.2011.07.024>
- Onojima, T., Goto, T., Mizuhara, H., & Aoyagi, T. (2018). A dynamical systems approach for estimating phase interactions between rhythms of different frequencies from experimental data. *PLOS Computational Biology*, 14(1), e1005928. <https://doi.org/10.1371/journal.pcbi.1005928>
- Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, 2011, 156869. <https://doi.org/10.1155/2011/156869>
- Pinotsis, D. A., Brunet, N., Bastos, A., Bosman, C. A., Litvak, V., Fries, P., & Friston, K. J. (2014). Contrast gain control and horizontal interactions in V1: A DCM study. *NeuroImage*, 92, 143–155. <https://doi.org/10.1016/j.neuroimage.2014.01.047>
- Puce, A., & Perrett, D. (2003). Electrophysiology and brain imaging of biological motion. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 358(1431), 435–445. <https://doi.org/10.1098/rstb.2002.1221>
- Railo, H., Koivisto, M., & Revonsuo, A. (2011). Tracking the processes behind conscious perception: A review of event-related potential correlates of visual consciousness. *Consciousness and Cognition*, 20(3), 972–983. <https://doi.org/10.1016/j.concog.2011.03.019>
- Rockland, K. S., & Ojima, H. (2003). Multisensory convergence in calcarine visual areas in macaque monkey. *International Journal of Psychophysiology*, 50(1–2), 19–26. [https://doi.org/10.1016/S0167-8760\(03\)00121-1](https://doi.org/10.1016/S0167-8760(03)00121-1)
- Rosen, S., & Howell, P. (2011). *Signals and systems for speech and hearing/S. Rosen and P. Howell – Details – Trove*. Retrieved from <https://trove.nla.gov.au/work/6339833>
- Roy, D., Sigala, R., Breakspear, M., McIntosh, A. R., Jirsa, V. K., Deco, G., & Ritter, P. (2014). Using the virtual brain to reveal the role of oscillations and plasticity in shaping brain's dynamical landscape. *Brain Connectivity*, 4(10), 791–811. <https://doi.org/10.1089/brain.2014.0252>
- Sauseng, P., Klimesch, W., Schabus, M., & Doppelmayr, M. (2005). Fronto-parietal EEG coherence in theta and upper alpha reflect central executive functions of working memory. *International Journal of Psychophysiology*, 57(2), 97–103. <https://doi.org/10.1016/j.ijpsycho.2005.03.018>
- Saxe, R. (2006). Uniquely human social cognition. *Current Opinion in Neurobiology*, 16(2), 235–239. <https://doi.org/10.1016/J.CONB.2006.03.001>

- Shaw, A. D., Hughes, L. E., Moran, R., Coyle-Gilchrist, I., Rittman, T., & Rowe, J. B. (2019). In vivo assay of cortical microcircuitry in frontotemporal dementia: A platform for experimental medicine studies. *Cerebral Cortex*, 1–11. <https://doi.org/10.1093/cercor/bhz024>
- Spiegler, A., Knösche, T. R., Schwab, K., Haueisen, J., & Atay, F. M. (2011). Modeling brain resonance phenomena using a neural mass model. *PLoS Computational Biology*, 7(12), e1002298. <https://doi.org/10.1371/journal.pcbi.1002298>
- Stefanescu, R. A., & Jirsa, V. K. (2008). A low dimensional description of globally coupled heterogeneous neural networks of excitatory and inhibitory neurons. *PLoS Computational Biology*, 4(11), e1000219. <https://doi.org/10.1371/journal.pcbi.1000219>
- Strand, J., Cooperman, A., Rowe, J., & Simenstad, A. (2014). Individual differences in susceptibility to the McGurk effect: Links with lip-reading and detecting audiovisual incongruity. *Journal of Speech, Language, and Hearing Research*, 57(6), 2322–2331. [https://doi.org/10.1044/2014\\_JSLHR-H-14-0059](https://doi.org/10.1044/2014_JSLHR-H-14-0059)
- Thakur, B., Mukherjee, A., Sen, A., & Banerjee, A. (2016). A dynamical framework to relate perceptual variability with multisensory information processing. *Scientific Reports*, 6(1), 31280. <https://doi.org/10.1038/srep31280>
- Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., ... Joliot, M. (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage*, 15(1), 273–289. <https://doi.org/10.1006/NIMG.2001.0978>
- Ursino, M., Crisafulli, A., di Pellegrino, G., Magosso, E., & Cuppini, C. (2017). Development of a Bayesian estimator for audio-visual integration: A neurocomputational study. *Frontiers in Computational Neuroscience*, 11, 89. <https://doi.org/10.3389/fncom.2017.00089>
- Varela, F., Lachaux, J.-P., Rodriguez, E., & Martinerie, J. (2001). The brainweb: Phase synchronization and large-scale integration. *Nature Reviews Neuroscience*, 2(4), 229–239. <https://doi.org/10.1038/35067550>
- Wallace, M. T., Ramachandran, R., & Stein, B. E. (2004). A revised view of sensory cortical parcellation. *Proceedings of the National Academy of Sciences of the United States of America*, 101(7), 2167–2172. <https://doi.org/10.1073/pnas.0305697101>
- Wasserman, L. (2005). *All of non-parametric statistics*. New York, NY: Springer.
- Williams, P. E., Mechler, F., Gordon, J., Shapley, R., & Hawken, M. J. (2004). Entrainment to video displays in primary visual cortex of macaque and humans. *Journal of Neuroscience*, 24(38), 8278–8288. <https://doi.org/10.1523/JNEUROSCI.2716-04.2004>
- Wilson, H. R., & Cowan, J. D. (1972). Excitatory and inhibitory interactions in localized populations of model neurons. *Biophysical Journal*, 12(1), 1–24. [https://doi.org/10.1016/S0006-3495\(72\)86068-5](https://doi.org/10.1016/S0006-3495(72)86068-5)
- Yeterian, E. H., & Pandya, D. N. (1989). Thalamic connections of the cortex of the superior temporal sulcus in the rhesus monkey. *The Journal of Comparative Neurology*, 282(1), 80–97. <https://doi.org/10.1002/cne.902820107>

**How to cite this article:** Kumar VG, Dutta S, Talwar S, Roy D, Banerjee A. Biophysical mechanisms governing large-scale brain network dynamics underlying individual-specific variability of perception. *Eur J Neurosci*. 2020;52:3746–3762. <https://doi.org/10.1111/ejn.14747>