

Structure and pH Dependence of Membranolytic Mechanisms by Truncated Oxidized Phospholipids

Min Xie, Maik G. N. Derks, Eveline H. W. Koch, C. Bjorn van Boven, Minchakarn Janlad, Behnaz Bagheri, Zexi Xu, Daria Kovryzhenko, Cornelis A. van Walree, Ana Sobota, Markus Weingarth, Jirasak Wong-ekabut, Mikko Karttunen, Eefjan Breukink, J. Antoinette Killian, Andreas F. P. Sonnen, and Joseph H. Lorent*



Cite This: <https://doi.org/10.1021/jacs.4c12543>



Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: Membrane lipid oxidation is a universal process that occurs in situations of oxidative stress and is encountered in numerous physiological and pathological situations. Oxidized truncated phospholipids make up a large part of the oxidation products and alter the membrane properties in a way that can lead to cell death. However, the underlying mechanisms are not well understood nor is it clear whether environmental factors, such as pH, can modulate these effects. Using model membranes, we investigate how individual lipid aldehydes and carboxylic acids with truncated acyl chains alter the membrane structure. Our data shows that lipid aldehydes and carboxylic acids have different permeabilization efficiencies toward molecules of varying charge and size and that $\Delta C9$ truncated lipids are usually more efficient in permeabilizing membranes than $\Delta C5$. In terms of physical mechanisms, the $\Delta C9$ truncated lipid carboxylic acid induces permeabilization and membrane curvature in a pH-dependent fashion. This is explained by ionization-dependent exposure of the carboxyl group to the water–bilayer interface, which increases the intrinsic molecular curvature of the oxidized lipid. Conversely, $\Delta C9$ truncated lipid aldehydes and nonionized carboxyls do not induce curved structures but are more efficient in increasing permeability toward larger molecules. We further show that truncated lipids can escape the bilayer and accumulate at interfaces, implying that they might act on neighboring cells. This study indicates that oxidized phospholipids with truncated acyl chains disrupt membrane structure, depending on their specific molecular structure and the pH of the environment, opening a possible route for the design of lipid nanoparticles with pH-dependent drug release.



INTRODUCTION

Lipid oxidation not only is involved in various basic cellular reaction programs such as cellular senescence¹ but also contributes to pathological dysfunctions, including Parkinson's disease,^{2–4} cardiovascular diseases,^{5,6} diabetes,⁷ and Alzheimer's disease.⁸ Phospholipid bilayers are fundamental structural units of cellular membranes which are susceptible to short and long-lived reactive oxygen species (ROS).⁹ The sources of ROS derive from either endogenous production, such as the mitochondrial respiratory chain in mammalian cells,¹⁰ or exogenous stimuli, such as UV light and heat.^{11,12} Generally, the majority of formed oxidation products are lipid hydroperoxides and various truncated lipid oxidation products such as aldehydes and carboxylic acids.^{9,13–15} Oxidative truncation produces further smaller volatile products, such as malondialdehyde (MDA) and 4-hydroxynonenal (4-HNE), that derive from their parental lipids. Their electrophilic properties render these molecules very reactive toward biomolecules which supposedly leads to pathological as well as physiological processes.¹⁶ The diversity and the physiological

roles of these secondary products has been reviewed extensively.^{11,17,18} While these small volatile products can diffuse passively through membrane bilayers, lipid aldehydes and lipid carboxylic acids with truncated acyl chains can alter structure and function of membrane bilayers.¹⁹ Bilayers or lamellar phases are formed by amphipathic phospholipids that bury their hydrophobic tails in the membrane interior and expose their hydrophilic head groups to the aqueous phase.^{20,21} Lipid oxidation can disrupt the bilayer structure due to increased hydrophilicity or a radical change in lipid shape. Molecular dynamics (MD) simulations show that in some cases lipid aldehydes and carboxylic acids might expose their chains

Received: September 10, 2024

Revised: February 18, 2025

Accepted: February 19, 2025

59 with the oxidized hydrophilic functions to the membrane
60 surface.^{22–25} This change of conformation results supposedly in
61 conically shaped lipids which increase the membrane surface
62 area, reduce bilayer thickness,^{26–28} and induce spontaneous
63 positive curvature, which results in the formation of micelles or
64 pores.^{28,29}

65 The resulting defects can significantly influence the
66 biophysical and physiological functions of the membrane
67 bilayers. Studies show that the presence of oxidized lipids alter
68 lipid packing differentially, depending on their structure.^{9,23,28}
69 Experimental and MD simulations have demonstrated that
70 especially lipid aldehydes increase the membrane permeability.
71 Free energy profile analyses indicate that the activation free
72 energy requirement for diffusion of small hydrophilic molecules,
73 such as H₂O, H₂O₂, and O₂, across the bilayer is much lower
74 when lipid aldehydes are present.^{28,30} Studies have further
75 shown that lipid oxidation leads to the formation of pores in
76 model membranes either by artificially oxidizing membranes of
77 unsaturated lipids^{9,27,31} or by incorporating certain oxidized
78 lipids.^{32,33}

79 Although impairment of the membrane bilayer structure
80 induced by oxidized lipids has been reported in many studies, a
81 systematic characterization of the permeabilization mechanisms
82 upon structural differences of truncated oxidized lipids is still
83 lacking and there is a need for experimental data that could
84 confirm the conclusions that have been obtained from MD
85 simulations.^{22–25} In addition, there are almost no data on
86 possible additive effects of different OxPCs, nor the influence of
87 pH. A pH dependent effect could arise from ionization of the
88 COOH group of carboxylated lipids and has been proposed in
89 MD simulations.³⁴ We therefore characterized the membranolytic
90 mechanisms by complementary techniques using model
91 membranes of different truncated carboxylic acids and
92 aldehydes (structures in Figure 1) upon different pHs and we

investigated the additivity of these effects. We studied 93
permeabilization toward different types of molecules in giant 94
unilamellar vesicles (GUVs) and large unilamellar vesicles 95
(LUVs). We further investigated lipid packing by polarity 96
sensitive probes, induction of curvature and order parameter by 97
³¹P NMR, and the formation of micelles and the release of 98
oxidized lipids from the membrane by dynamic light scattering 99
and surface tension measurements. We measured the zeta (ζ)- 100
potential to determine exposure of the ionized carboxyl group to 101
the membrane interface and used MD simulations to explain 102
some of the observed results and to help attribute permeabiliza- 103
tion mechanisms to the different OxPCs. 104

RESULTS

Truncated Aldehydes Produce Stronger Influx of 4
kDa FITC-Dextran and Sulforhodamine B than Carboxy-
yls. We first used GUVs to evaluate permeabilization induced by 108
lipid aldehydes and carboxylic acids with truncated acyl chains. 109
We observed that 20 mol % of PC(16:0/9:0(CHO@C9)) or 110
PC(16:0/9:0(COOH@C9)) induced permeabilization toward 111
FITC-Dextran (4 kDa) in GUVs made of PC(16:0/18:1), with a 112
larger effect induced by PC(16:0/9:0(CHO@C9)) (Figure 113
2A,B). This result is further supported by a sulforhodamine B 114
(SRB) leakage assay in GUVs containing 10 mol % of PC(16:0/ 115
9:0(CHO@C9)) or PC(16:0/9:0(COOH@C9)) (Figure S2). 116
We observed a larger release of SRB in GUVs containing the 117
aldehyde truncated form. 118

Dithionite Quenching of Inner Leaflet NBD-PE Occurs Only above a Threshold of Truncated OxPCs but the Effect Is Additive.

To quantify the permeabilization or 121
possibly lipid flip-flop induced by oxidized lipids in membrane 122
bilayers, we used a dithionite quenching assay in LUVs (Figure 123
2C). OxPCs were integrated into PC(16:0/18:1) LUV 124
containing headgroup labeled NBD-PE(18:1/18:1) at 0.5% 125
(mol/mol) and dithionite was further used to chemically quench 126
the fluorescence of NBD irreversibly. This results in 50% 127
quenching if dithionite does not diffuse to the luminal 128
monolayer of the LUV, and if there is no flopping of NBD-PE 129
to the outside monolayer. Our results show that only around 130
40–45% of quenching was attained in the controls (Figure S1). 131
This is because of some remaining MLVs in the sample (~4%; 132
see Figure S3). Since the same ratio of MLVs was present 133
throughout all conditions, we concluded that the observed 134
effects did not result from a change of lamellarity. Since the same 135
ratio of MLVs was present throughout all conditions, we 136
concluded that the observed effects did not result from a change 137
of lamellarity. Triton X-100 was used to solubilize the 138
membranes, leading to complete fluorescence quenching. To 139
quantify the quenching by dithionite ions, we fitted the 140
quenching curves to an exponential plateau model. The 141
equation of this model is % quenching = $Y_M - (Y_M - Y_0)^{-kt}$, 142
with k reflecting the rate constant (min⁻¹). Y_M and Y_0 indicate 143
the maximum and minimum percentages of quenching, 144
respectively. As expected, the quenching rate increased upon 145
higher incorporated fractions of ΔC9 and ΔC5 OxPCs in LUVs, 146
with carboxylated acyl chains being slightly more efficient in 147
promoting quenching of NBD-PE than lipid aldehydes (Figure 148
2D,E, and Figure S1A,B,C,D). Interestingly, a significant 149
increase of quenching was only observed when truncated 150
OxPCs were present above a threshold of ~12–15 mol % 151
(Figure 2D,E, and Figure S1A,B,C,D). However, when both 152
PC(16:0/9:0(CHO@C9)) and PC(16:0/9:0(COOH@C9)) 153
were mixed at different ratios but at a constant total molar 154




Figure 1. Chemical structures of lipids and fluorescent probes used in this study. (A) Host lipid for LUV or GUV. (B) Oxidized lipids. (C) Fluorescent lipids. (D) Fluorescent probes.




Figure 2. Incorporation of OxPCs disrupts PC(16:0/18:1) membranes. (A) Confocal microscopy images of GUVs containing no oxidized lipids (control) or 20 mol % PC(16:0/9:0<CHO@C9>) or PC(16:0/9:0<COOH@C9>) incubated for 10 min in the presence of 40 μM of FITC-Dextran (4 kDa). Red arrows indicate GUVs that are permeabilized. (B) Quantification of the FITC-Dextran intensity inside versus outside of GUVs. (C) Schematic diagram of the dithionite quenching assay. Quenching speed (k values, min^{-1}) induced by (D) ΔC9 and (E) ΔC5 truncated OxPCs and (F) varied ratios of ΔC9 truncated OxPCs at a total fraction of 20 mol % based on curve fitting with an exponential plateau model as described in Materials and Methods in the SI. One-way ANOVA analysis was used to compare fluorescence intensity in GUVs toward controls. Two-way ANOVA analysis was used to compare the k values between different incorporation fractions and the negative controls (colored stars) and the k values between PC(16:0/9:0<CHO@C9>) and PC(16:0/9:0<COOH@C9>) at the same incorporation fractions (black stars). At least two independent experiments were performed for the NBD-PE fluorescence quenching assay in triplicates ($n \geq 2$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$).

fraction of 20 mol % of total oxidized lipids, quenching slightly increased with higher PC(16:0/9:0<COOH@C9>) ratios (Figure 2F and Figure S1E). Curiously, even when one of both lipids dropped below the 12–15 mol % hurdle, quenching was still observed, meaning that the observed threshold for one

OxPCs is not valid if two different OxPCs are present and that somehow the effect is additive (Figure 2F and Figure S1E).

Truncated Aldehydes Induce Quicker Permeation toward Sulforhodamine B (SRB) than Carboxyls while a Low pH Enhances the Effect of Carboxyls. In addition to the dithionite assay, we examined permeabilization by adding




Figure 3. Permeabilization of PC(16:0/18:1) LUV induced by the addition of PC(16:0/9:0<CHO@C9>) and PC(16:0/9:0<COOH@C9>) to the suspension. (A) Schematic diagram of permeabilization assay via sulforhodamine B leakage. Release constants (k value, min^{-1}) of sulforhodamine B after addition of (B) $\Delta C9$ and (C) $\Delta C5$ truncated OxPCs, and (E) varied ratios of $\Delta C9$ truncated OxPCs at 20 mol % based on curve fitting as described in materials and methods. Percentage (%) of escaped sulforhodamine B at 20 min after addition of (D) $\Delta C9$ and (E) $\Delta C5$ truncated OxPCs and (G) variable ratios of $\Delta C9$ aldehydes and carboxyls at a total concentration of 20 mol %. (H) Comparison of release constants (k value, min^{-1}) of sulforhodamine B between buffers at pH = 7.4 and pH = 4.5. One-way ANOVA analysis was used to compare the k values and the leakage at 20 min between different fractions of each OxPCs and the corresponding controls (colored stars). Unpaired t -test analysis (two-tailed) was used to compare the k values between PC(16:0/9:0<CHO@C9>) and PC(16:0/9:0<COOH@C9>) at the same concentrations or different pHs (black stars). At least three independent experiments were performed ($n \geq 3$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$).




Figure 4. Membrane packing characterized by polarity sensitive probes. (A) Di-4-ANEPPDHQ GP and (B) Laurdan GP upon increasing fractions of PC(16:0/9:0<CHO@C9>) and PC(16:0/9:0<COOH@C9>) integrated into PC(16:0/18:1) LUV. Two-way ANOVA analysis was used to compare different incorporated fractions to the corresponding controls (colored stars) and between PC(16:0/9:0<CHO@C9>) and PC(16:0/9:0<COOH@C9>) (black stars) at the same incorporated fractions ($n \geq 6$, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$).




Figure 5. DLS analysis of PC(16:0/18:1) LUV containing PC(16:0/9:0<CHO@C9>) (A, C) or PC(16:0/9:0<COOH@C9>) (B, C). Size distribution of LUVs containing different amounts of (A) PC(16:0/9:0<CHO@C9>) or (B) PC(16:0/9:0<COOH@C9>). The arrow indicates the average hydrodynamic diameters in nanometers based on the second peak (smaller peak). (C) Average hydrodynamic diameters in nanometers based on the main peak (larger peak). The values are derived from 6 measurements with error bars indicating the standard deviation (SD). One-way ANOVA was used to compare the size difference between LUVs with different incorporated fractions of OxPCs toward the corresponding controls ($n \geq 6$, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$).

different OxPCs to the suspension containing PC(16:0/18:1) LUVs which enclosed sulforhodamine B at a self-quenching concentration of 20 mM (Figure 3A). Triton X-100 was used as a positive control to induce complete permeabilization. However, it induced a constant increase of SRB fluorescence that was accounted for in the calculation of release (Figure S4E). The addition of OxPCs to the suspension induced concentration-dependent permeabilization (Figure S4A,B,C,D, arrows indicate the addition of OxPCs). As controls, equal amounts of ethanol and PC(16:0/18:1) lipid were added, and neither of them induced permeabilization (Figure S4A,B,C,D). As for the quenching assay, we fitted the release curves with the exponential plateau model. The k value showed that $\Delta C9$ and $\Delta C5$ lipid aldehydes induced faster permeabilization than $\Delta C9$ and $\Delta C5$ carboxylic OxPCs, respectively (Figure 3B,C and Figure S4A,B,C,D). Curiously, the opposite was observed for dithionite quenching (Figure 2D,E). However, after 20 min, PC(16:0/9:0<CHO@C9>) and PC(16:0/9:0<COOH@C9>) induced a similar level of release at 91.5% and 87.0%, respectively (Figure 3D,E). Further, $\Delta C9$ truncated lipids were slightly more effective in releasing SRB than $\Delta C5$ truncated lipids (Figure 3B,C and Figure S4A,B,C,D). SRB release induced by varying ratios of PC(16:0/9:0<COOH@

C9>) upon PC(16:0/9:0<CHO@C9>) at a total of 20 mol % was not significantly altered (Figure 3F,G), consolidating the hypothesis that the effect on permeability of truncated lipids with different chemical functions is additive.

To investigate the probable pH dependence of SRB release because of the protonatable carboxyl function, we also assessed SRB leakage under acidic conditions (pH = 4.5). Importantly, PC(16:0/9:0<COOH@C9>) induced a quicker permeabilization in acidic (pH = 4.5) medium, whereas no significant changes were observed with PC(16:0/9:0<CHO@C9>) upon pHs (Figure 3H and Figure S4G). This suggests that nonionized carboxyl and aldehyde functions are somehow more efficient in releasing SRB.

Truncated Lipid Aldehydes Alter Membrane Polarity and Membrane Fluidity Differently from Carboxyls. Polarity sensitive probes, such as Laurdan and di-4-ANEPPDHQ, have recently been shown to report different trends upon increasing membrane oxidation.⁹ In theory, these probes should be able to sense increased water insertion or a change of polarity in their microenvironment that is quantified by the generalized polarization (GP).³⁵ The fluorophore of di-4-ANEPPDHQ is assumed to localize at the membrane interface and determine properties of only the outer leaflet of the




Figure 6. Surface tension determination with pendant drop tensiometry. Interfacial tension of PBS suspensions containing PC(16:0/18:1) LUVs that have been incorporated with different amounts of (A) PC(16:0/9:0<CHO@C9>) and (B) PC(16:0/9:0<COOH@C9>) upon inspection time. The log₂ T_{50%} values of surface tension reduction in the suspension containing (C) LUVs with different amounts of OxPCs (LUVs suspension) or (D) different concentrations of free OxPCs that have been added to the buffer and that do not contain LUVs (free lipids, see also Figure S6). The log₂ T_{50%} values were obtained by fitting the curves with a sigmoidal model using GraphPad Prism as described in the main text. (E) Minimal surface tension (γ_{min}, mN/m) at critical micellar concentration (CMC) and beyond reached by the different oxidized lipids. At least two independent experiments were performed for surface tension measurements (*n* ≥ 3). The values are the mean ± SD (*n* ≥ 3). One-way ANOVA was used to compare the γ_{min} (mN/m) values between different OxPCs (**p* < 0.05, ***p* < 0.01, ****p* < 0.001, *****p* < 0.0001).

membrane bilayer due to its positive charge and the absence of flipping,^{36,37} while Laurdan is supposedly localized in between headgroups and acyl chains and probes both leaflets of the membrane bilayer.³⁸ We investigated here how defined molar ratios of truncated carboxylic and aldehyde OxPCs alter the GP of these probes with the aim of investigating changes in packing or membrane water incursion (Figure 4). In addition, we measured the effect on DPH fluorescence anisotropy to determine the environmental fluidity of the lipid acyl chains.³⁹ We first observed that the incorporation of both OxPCs decreased di-4-ANEPPDHQ GP upon their present concentrations, with PC(16:0/9:0<CHO@C9>) causing larger effects than PC(16:0/9:0<COOH@C9>) (Figure 4A). Less membrane packing alterations were observed by Laurdan GP, although PC(16:0/9:0<COOH@C9>) decreased Laurdan GP significantly at concentrations higher than 10 mol % (Figure 4B). The fluidity of the hydrophobic core, probed by DPH anisotropy, seemed to be only slightly affected by both OxPCs (Figure S5). Taken together, we observed that carboxylated and aldehyde OxPCs affect membrane packing differently.

OxPCs Reduce the Vesicle Size but Only Carboxylated Lipids Induce the Formation of a Population of Small Aggregates. To investigate if OxPCs alter the vesicle structure in LUVs by inducing the formation of micelles or other potential

amphiphilic aggregates, we measured LUV size by dynamic light scattering (DLS) upon increasing fraction of truncated OxPCs. We observed that both PC(16:0/9:0<CHO@C9>) and PC(16:0/9:0<COOH@C9>) reduced the size of PC(16:0/18:1) LUVs in a concentration dependent manner (Figure SA,B,C). In addition, at 30 mol % of PC(16:0/9:0<COOH@C9>), we observed the formation of small aggregates with an average hydrodynamic diameter of 19 nm (Figure SB, arrow indicates the second peak formation). To confirm a size reduction of vesicles, we did some cryo-EM measurements at pH = 7.4 and 4.5 and obtained a reduction of vesicles size specifically with PC(16:0/9:0<COOH@C9>) at pH = 7.4 (Figure S3G).

Truncated OxPCs Are Released from the Lipid Bilayer with Different Kinetics and Effects on Surface Tension while pH Influences Strongly PC(16:0/9:0<COOH@C9>) Behavior. We recently showed that complex oxidation of lipids leads to a decrease of surface tension at the air–water interface because of the release of certain oxidized species from the membrane.⁹ Since truncated lipids should have a higher water solubility and possibly form micelles because of their high intrinsic molecular curvature, we wanted to measure the capacity of these lipids to leave the lipid bilayer, which also might explain the reduction of vesicle size (Figure 5). Pure PC(16:0/18:1) LUVs did not significantly affect the surface tension of the




Figure 7. ^{31}P NMR analysis and Zeta (ζ) potential determination. Static ^{31}P NMR spectra of PC(16:0/18:1) MLVs incorporated with different fractions of (A) PC(16:0/9:0<CHO@C9>) and (B) PC(16:0/9:0<COOH@C9>) hydrated with Milli-Q- H_2O (top panel) or NH_4HCO_3 (bottom panel) or (C) PC(16:0/9:0<COOH@C9>) neutralized with NaOH at an equivalent amount. (D) Zeta (ζ) potential determination in LUVs composed of PC(16:0/18:1) with or without 15 mol % of PC(16:0/9:0<COOH@C9>) upon pH changes.

suspension within the inspected period while the incorporation of both OxPCs into LUVs reduced the surface tension of the buffer at both neutral (pH 7.4) and acidic (pH 4.5) pHs (Figure 6A,B and Figure S6D). This indicates that a certain amount of OxPCs left the bilayer phase of LUVs and adsorbed at the air–water interface. The rate of surface tension reduction was strongly influenced by the nature of the truncated species and the respective molar ratio inside the vesicles (Figure 6A,B). To compare these effects to a system where no lipid bilayer was present, we measured surface tension reduction of free lipids

added to the buffer (Figure S6A,B,C,E). To quantify the rate of surface tension reduction of the different concentrations of OxPCs in both data sets, we applied a four-parameter sigmoidal regression model:

$$\gamma = \gamma_{\min} + \frac{(\gamma_{\max} - \gamma_{\min})}{1 + 10^{(\log T_{50\%} - t) \times a}}$$

Here, γ is the surface tension upon time, $\gamma_{\min}$ is the minimal surface tension obtained (Figure 6E), $\gamma_{\max}$ is the initial surface




Figure 8. MD simulations of PC(16:0/18:1) membrane bilayer in the presence of OxPCs. Snapshots at the end of the 1 μ s simulation of (A) 100 mol % concentration of PC(16:0/18:1), at the end of the 2 μ s simulations of (B) 50 mol % PC(16:0/9:0(COOH@C9)), ionized and deprotonated, and (C) 50 mol % PC(16:0/9:0(CHO@C9)). The light blue region is water. The cyan spheres are water molecules within 3.5 Å of the phosphate atom of PC(16:0/18:1) and oxidized lipids. The green spheres are phosphate atoms of the lipid headgroups, and the green, yellow, and white lines represent PC(16:0/18:1), PC(16:0/9:0(COOH@C9)), and PC(16:0/9:0(CHO@C9)) molecules, respectively. Calculations of (D) area of per lipid, (E) thickness, and (F) permeability to water molecules in the presence of 50 mol % PC(16:0/9:0(CHO@C9)) and PC(16:0/9:0(COOH@C9)), respectively. The table shows the area compressibility modulus (K_A) and bending modulus (K_C) in the presence of OxPCs.

tension, $T_{50\%}$ is the time in seconds necessary to reach the inflection point of the sigmoid, t is the time in seconds, and a is the slope.

We observed that free lipids added to the buffer reduced surface tension quicker than when they were incorporated into LUVs (Figure 6 and Figure S6A,B) at the same molar fractions. In addition, PC(16:0/9:0(COOH@C9)) reduced surface tension quicker than PC(16:0/9:0(CHO@C9)) ($\sim 3\text{--}5 \log_2$), especially when OxPCs were incorporated into LUV at 20–30 mol % at pH = 7.4 (Figure 6C). This implies that PC(16:0/9:0(COOH@C9)) is easily released from the membrane bilayer either because its ionized carboxyl residue renders it more soluble in the aqueous phase and therefore can quicker migrate to the interface or because it forms more efficiently micelles or other lipid aggregates that can quickly interchange lipids with the interface. Interestingly, the time to reach the inflection point ($T_{50\%}$) drastically decreased from 403.4 to 10.15 s for the carboxylated lipid when the pH increased from 4.5 to 7.4 (Figure 6C and Figure S6D), while the kinetics of the CHO lipid were not strongly altered (Figure 6C and Figure S6D). This indicates that the protonated carboxylated lipid remained longer in the LUVs than in the ionized version.

Based on the curve regressions made from concentrations that reached the minimum surface tension, $\gamma_{\min}$ (mN/m) at CMC (Figure 6E), we observed that PC(16:0/9:0(CHO@C9)) reduced surface tension to a lower value than PC(16:0/9:0(COOH@C9)) at pH 7.4. Importantly, at pH = 4.5, the surface tension reduction of PC(16:0/9:0(COOH@C9))

became similar to that of PC(16:0/9:0(CHO@C9)) (Figure 6E). In addition, we observed that OxPCs with the same functional groups, the ones with shorter acyl chains (truncation at $\Delta C5$), had a slightly weaker capacity to reduce surface tension (Figure 6E). In lipid monolayers, the maximum reduction of surface pressure obtained with a lipid at complete coverage of the water/air interface⁴⁰ or to the critical micellar concentration (CMC),²⁰ although the CMC might not always correspond to complete coverage of the monolayer at collapse.⁴¹ The different capacities of reducing surface tension of both OxPCs and the pH dependence that is observed for PC(16:0/9:0(COOH@C9)) suggests different behaviors which strongly depend on the molecular structure and the pH.⁴² Since the only difference between these two OxPCs is the functional group at the truncated tail, it is reasonable to assume that PC(16:0/9:0(CHO@C9)) can form more compressed monolayers than PC(16:0/9:0(COOH@C9)) at pH = 7.4, which might depend on the fact that the tail of lipid aldehydes might not protrude the same way to the interface as lipid carboxyls.^{24,25} A collapse at a lower coverage or surface pressure, corresponding to a higher surface tension, might also be induced by higher solubility of the carboxyl truncated lipid or the shorter acyl chain ($\Delta C5$) in the aqueous phase.⁴³ Since a low pH increases the capacity of PC(16:0/9:0(COOH@C9)) to reduce surface tension to a similar amount as PC(16:0/9:0(CHO@C9)), we can assume that both molecules adopt a similar conformation at low pH.

³¹P NMR Analysis Reveals a Different Effect of Truncated Carboxyls and Aldehydes on the Bilayer Structure. Membrane bilayer structure in the presence of OxPCs with truncated acyl chains was further analyzed by phosphorus NMR (³¹P NMR), which provides information on dynamics and structure at the phosphate group of phospholipids.⁴⁴ The static ³¹P NMR spectra for PC(16:0/18:1) in the presence of different fractions of PC(16:0/9:0<CHO@C9>) and PC(16:0/9:0<COOH@C9>) are displayed in Figure 7. In the presence of both OxPCs, the membranes remained mostly in a liquid crystalline lamellar phase upon increasing fractions of OxPCs (Figure 7A,B), but a small isotropic peak at lower fractions of PC(16:0/9:0<COOH@C9>) emerged (Figure 7B, top panel). Since concentrations of lipids were quite high in this MLV assay and since the protonated PC(16:0/9:0<COOH@C9>) might hence exceed the buffer capacity, we used an ammonium bicarbonate solution to increase ionization of the carboxyl function beforehand (Figure 7B, bottom panel). Under these conditions, we observed a concentration-dependent shift of the right shoulder but still not a concentration-dependent increase of the isotropic peak at ~−2 ppm, even though it was strongest at 30 mol % (Figure 7B, bottom panel).

The chemical shift anisotropy gives an indication of rotational diffusion of phospholipids and the size of the vesicles.⁴⁵ A much larger effect on chemical shift anisotropy was reached with PC(16:0/9:0<COOH@C9>) than PC(16:0/9:0<CHO@C9>) which might indicate the formation of smaller structures (increased averaging by rotational diffusion, due to a slightly different orientation) and/or a change in headgroup dynamics (Figure S7A).

In addition to the effects from individual OxPCs, we also measured the effects from the mix of PC(16:0/9:0<CHO@C9>) and PC(16:0/9:0<COOH@C9>) at a ratio of 1:1 and a total fraction of 20 mol % (Figure S7C). The increased fraction of PC(16:0/9:0<COOH@C9>) led to a larger chemical shift anisotropy (Figure S7D), as expected.

Ionization of the Carboxyl Group in PC(16:0/9:0<COOH@C9>) Induces Highly Curved Structures in a pH Dependent Way. The ³¹P NMR results point out that the pH plays a major role in the effects induced by PC(16:0/9:0<COOH@C9>) and that the carboxyl functional group is able to significantly alter the structure of the membrane if it is ionized, possibly by producing micelles or other curved lipid aggregates. To further investigate the effect of pH on ionization of the carboxyl group and membrane structure, we used NaOH at an equivalent amount to PC(16:0/9:0<COOH@C9>) to completely ionize the carboxyl group. Under these conditions, we observed a large disruption of the lamellar structure and formation of an isotropic phase, pointing to the possibility that the intrinsic curvature of the ionized PC(16:0/9:0<COOH@C9>) increases largely with pH (Figure 7C). This effect was temperature dependent (Figure S7B), with high temperatures favoring highly curved structures. To monitor the importance of pH on the ionization of PC(16:0/9:0<COOH@C9>), we measured the zeta (ζ) potential of LUVs containing PC(16:0/9:0<COOH@C9>) upon pH changes. We observed that the ζ-potential decreased strongly upon increasing pH, indicating that the ionization of the carboxyl function in the membrane depended on pH (Figure 7D). This result further suggests that the ionized carboxyl function contributes to the ζ-potential and is probably localized at the membrane–water interface.

MD Simulations Reveal Differences in Positioning and Effects on Bilayer Structure of Truncated Carboxyls and

Aldehydes. To further investigate the positions of the carboxyl and aldehyde groups in the bilayer plane, we performed MD simulations (Figure 8). We observed that the ionized carboxyl group pointed to the bilayer interface (Figure 8B), while the aldehyde function remained in the hydrophobic core (Figure 8C). This resulted in a slightly larger area/molecule for the truncated carboxylic acid, while the aldehyde function effectively reduced this parameter (Figure 8D). The reduction of the area per lipid by the lipid aldehyde could be in line with the reduction of surface tension that was more efficient with the aldehyde lipid. A smaller molecular volume of the CHO molecule would mean a higher density of molecules at the water–air interface and hence a lower surface tension.

Thickness was more reduced with the ionized COOH lipid (Figure 8E), meaning it can probably more efficiently abolish the bilayer structure, as observed in NMR studies (Figure 7). In terms of water permeability, we observed a slightly higher value for the CHO truncated lipid (Figure 8F). This could be in line with the permeability observed toward SRB (Figure 3 and Figure S2).

The mechanical properties of the bilayer (i.e., the area compressibility modulus (K_A) and the bending modulus (K_C)), as from our MD trajectories, were strongly reduced by PC(16:0/9:0<COOH@C9>), while PC(16:0/9:0<CHO@C9>) had only little effects (table in Figure 8). These values reinforce the current picture that PC(16:0/9:0<COOH@C9>) has a stronger tendency to impair the bilayer structure of the membrane.

DISCUSSION

Our results show that truncated oxidized lipids have different mechanisms to destabilize lipid membranes. Permeabilization depends on the position of the acyl chain break, with ΔC9 lipids being more effective in permeabilizing membranes than ΔC5, and on the functional group at the truncation site (carboxyl or aldehyde). For truncated lipids with carboxyl groups, the pH largely influences ionization of the carboxyl group, and ionization at high pH promotes the disruption of the bilayer structure, release of carboxylated lipids from the bilayer, and formation of highly curved isotropic lipid aggregates while sulforhodamine B and dextran-FITC leakage is higher for nonionized residues.

Permeabilization driven by oxidized lipid aldehydes and carboxylic acids has been investigated by many groups but no clear permeabilization mechanisms have been established.^{28,32,47,48} Our data suggest different permeabilization mechanisms for truncated oxidized lipids containing aldehyde and carboxyl termini and pH dependence of the mechanism for carboxyls. Lipid aldehydes were more efficient in increasing permeability toward larger molecules, including sulforhodamine B (Figure 3, Figure S2) and FITC-Dextran (Figure 2A,B), which has a hydrodynamic radius above 1 nm.⁴⁹ Nonionized carboxyl at ΔC9 was also more efficient in releasing SRB than its ionized version (Figure 3H).

Ionized lipid carboxylic acids seemed to be more efficient in inducing permeabilization toward dithionite ions or increasing lipid-flip-flop of NBD-PE than aldehydes (Figure 2D,E,F). Similarly, in another study, an increase in flip-flop of NBD-PS was strongly induced by PC(16:0/9:0<COOH@C9>) but less with PC(16:0/9:0<CHO@C9>) at around 10% mol/mol in a PC(16:0/18:1) bilayer.⁵⁰ At these concentrations and because of the setup of their assay,⁵⁰ they could exclude permeabilization toward dithionite. Time scales in these experiments were far




Figure 9. Schematic diagram of effects in membrane bilayers induced by OxPCs with truncated acyl chains. (A) Representations of PC(16:0/18:1), PC(16:0/9:0<CHO@C9>), and PC(16:0/9:0<COOH@C9>) (Top panel). In PC(16:0/9:0<CHO@C9>), the aldehyde function points to the hydrophobic core. At physiological pH (I), the carboxylic acid of PC(16:0/9:0<COOH@C9>) is to a large part ionized, and the truncated acyl chain points to the bilayer interface, while at low pH (II), the carboxyl function is protonated and points probably to the hydrophobic core in an angle that does not induce high intrinsic molecular curvature.⁴⁶ (B) In the presence of PC(16:0/9:0<CHO@C9>), the membrane bilayer becomes more permeable probably because the hydrophilicity of the membrane core is increased by the polar aldehyde groups. At higher concentrations, oxidized lipids might also aggregate and induce diffusion spots. Further, PC(16:0/9:0<CHO@C9>) is probably released from the membrane bilayer and can adsorb at the air–water interface, reducing interfacial tension. (C) In the presence of PC(16:0/9:0<COOH@C9>), (I) at high (or neutral) pH, the presence of charged interfacial COO[−] functions reduces the ζ-potential and induces high curvature that could induce toroidal pores or transient instabilities due to the formation of isotropic aggregates (probably micelles). The formation of these aggregates could enhance the lipid exchange between bilayer phases and the air–water interface, thereby reducing surface tension more rapidly. (II) At low pH, the truncated sn-2 acyl chains point probably to the membrane core which inhibits curvature induction and the formation of isotropic structures, while permeabilization toward SRB is increased, probably by a similar effect observed for PC(16:0/9:0<CHO@C9>).

greater (hours-weeks) than in our assay, but a strong increase of NBD-PS flip-flopping occurred at 14% mol of PC(16:0/9:0<COOH@C9>) and 16% mol of PC(16:0/9:0<CHO@C9>), reducing half-life to 1–2 h. At concentrations higher than 20% mol, they observed permeabilization of the membrane toward dithionite. In our slightly different quenching assay, we observed a threshold of around 12–15% mol for all truncated lipids tested (Figure 2D,E and Figure S1A,B,C,D). Since our time scales are in the minutes range, a half-life of 1–2 h for flip-flopping would not be detectable, but since we used NBD-PE instead of NBD-PS, flip-flopping might be faster because PE is zwitterionic and has an uncharged small headgroup. We therefore cannot exclude that flip-flopping and/or permeabilization is occurring in the dithionite quenching assay.

In terms of permeabilization thresholds, other studies have seen similar effects for truncated lipids. Runas et al. showed three permeability regimes for PEG12-NBD, a small uncharged hydrophilic molecule of 821 Da, in GUV containing PC(16:0/9:0<CHO@C9>). From 2.5% to 10%, permeability increased clearly toward the control, while at concentrations >12.5%, the bilayer seemed to be disrupted or pores were formed.⁵¹ Thus, the mechanism of permeabilization, and possibly flip-flopping, is dependent on the concentration and the type of oxidized lipids. When oxidized lipids were added from the outside to the LUVs, there seemed to be no clear threshold at 12–15% mol since even smaller concentrations induced sulforhodamine B

release (Figure 3B,C and Figure S4A,B,C,D). It is possible that the truncated oxidized lipids added from the outside of the LUVs are more concentrated at certain spots of the suspension just after addition and thereby induce permeabilization of a subpopulation of LUVs, although the experiment on GUVs (Figure S2) confirmed that SRB leakage also occurs when the oxidized lipid is present in the membrane at a concentration of 10% which is below the threshold observed for the NBD-PE quenching assay.

At a pH of 7.4, the reduction of surface tension is faster with PC(16:0/9:0<COOH@C9>) than PC(16:0/9:0<CHO@C9>) when the lipid is added to the suspension (Figure 6C,D) and also when the oxidized lipid is released from LUVs (Figure 6C). This means that ionized PC(16:0/9:0<COOH@C9>) is quicker at entering and leaving the monolayer/bilayer than PC(16:0/9:0<CHO@C9>), probably because a higher water solubility of the ionized PC(16:0/9:0<COOH@C9>) goes ahead with a shorter residence time in the bilayer/monolayer.⁹ This seems to be confirmed by the fact that a low pH drastically reduces the speed of surface tension reduction by PC(16:0/9:0<COOH@C9>) (Figure S6). Similarly, it has been shown that, at a pH of 3.0, a monolayer composed of PC(16:0/9:0<COOH@C9>) collapses at a much higher surface pressure than at a pH of 10, meaning that the ionized form of PC(16:0/9:0<COOH@C9>) solubilizes easier in the aqueous phase.⁵²

Grauby-Heywang et al. observed the formation of aggregates in the suspension when the monolayer collapsed.⁵² We obtained a population of small aggregates, similar in size to micellar structures with PC(16:0/9:0(COOH@C9)) (Figure 5B), which might correspond to the isotropic peak observed in ³¹P NMR (Figure 7B). Lipid exchange of bilayers or monolayers with micellar aggregates containing PC(16:0/9:0(COOH@C9)) might also be more rapid. However, permeabilization was faster with PC(16:0/9:0(CHO@C9)) and nonionized PC(16:0/9:0(COOH@C9)), meaning that the permeabilization rate toward sulforhodamine B relied probably not essentially upon insertion of the lipid into the membrane but on a process that occurs in the membrane.

The fact that oxidized lipids can leave the membrane bilayer and insert at the air–water interface has been observed earlier by our group.⁹ This implies that oxidized lipids can not only exert their effects on the cell in which they are produced but also have long-range effects on neighboring cells. Since a reduction of surface tension is observed quite quickly already at small percentages of oxidized lipids (5 mol %), it is possible that the same phenomenon is observed in cells with low amounts of oxidized lipids. The reduction of surface tension is more efficient with PC(16:0/9:0(CHO@C9)) than PC(16:0/9:0(COOH@C9)) at pH = 7.4, probably because the ionized form of PC(16:0/9:0(COOH@C9)) is not likely to accumulate at a high coverage density (surface pressure) and solubilizes more easily in the aqueous phase.⁵³ Further, PC(16:0/9:0(CHO@C9)) should have a smaller hydrophilic head because its two tails point into the membrane core or air phase (Figure 8). This might result in a larger coverage area at the air–water phase compared to ionized PC(16:0/9:0(COOH@C9)) that points its ionized carboxyl tail to the water phase, thereby increasing the headgroup area (Figure 8 and Figure 9A).⁵⁴ This is further supported by the increased surface tension reduction capacity of PC(16:0/9:0(COOH@C9)) at pH 4.5 where both tails should point to the membrane core or air phase, reducing the headgroup size and increasing coverage (Figure 6E). At acidic pH (4.5), the carboxylic acid functional group is hence less ionized and tends to adopt a molecular conformation similar to PC(16:0/9:0(CHO@C9)) (Figure 9A).

In terms of membranolytic mechanisms, we observed a different picture for aldehyde and carboxylic acids truncated at ΔC5 or ΔC9 (Figure 9). PC(16:0/9:0(COOH@C9)) was able to induce highly curved structures at a pH of 7.4 and beyond (Figure 5B and Figure 7C). In the absence of a high buffer capacity in MLVs and at low pH, these structures were not observed (Figure 7B). Further, the ζ-potential of LUVs containing PC(16:0/9:0(COOH@C9)) decreased with pH (Figure 7D), meaning that PC(16:0/9:0(COOH@C9)) becomes ionized and that its carboxyl function points to the membrane–water interface as was also indicated by our MD simulations (Figure 8B). This behavior has been proposed earlier, but pH dependence has not been shown in liposomes but suggested by MD simulations.⁵⁵ The fact that the ionized carboxyl group points to the membrane interface should increase intrinsic molecular curvature of PC(16:0/9:0(COOH@C9)), favoring nonbilayer isotropic aggregates. Permeabilization could hence be induced by the formation of toroidal pores through positive intrinsic molecular curvature, although formation of these pores might require quite high concentrations of PC(16:0/9:0(COOH@C9)).^{47,56} The membrane might also simply be destabilized by the formation of micelles and the continuous release of PC(16:0/9:0(COOH@

C9)) from the lipid bilayer which might lead to transient instabilities allowing molecules such as dithionite to pass through or increase flip-flopping of NBD-PE. MD simulations show further that the thickness of the membrane is reduced (Figure 8E) with the ionized form of PC(16:0/9:0(COOH@C9)) at high concentrations. This might favor the formation of transient pores and reduce the thickness of the hydrophobic barrier.

When PC(16:0/9:0(COOH@C9)) was less ionized, we observed less curved structures (Figures 7B and S3G), suggesting that the intrinsic molecular curvature of the molecule is reduced.⁵⁶ It has been shown in earlier studies that the protonated form of PC(16:0/9:0(COOH@C9)) does not have its truncated acyl chain being pointed to the membrane interface, but rather being in the membrane bilayer, although it is not sure at what membrane depth the protonated COOH group would accumulate.⁵⁵ Since there is less formation of isotropic structures, we assume that the intrinsic molecular curvature of the lipid is reduced (Figure 9), suggesting either that the loss of a negative charge at the interface leads to less repulsion of headgroups and carboxyl groups at the membrane interface or the truncated sn-2 acyl chain adopts a conformation that produces less membrane curvature, or both.²⁰ The permeabilization mechanism might hence be altered and become similar to PC(16:0/9:0(CHO@C9)) whose aldehyde function also remains in the hydrophobic membrane core (Figure 8C) and where no curved structures were observed (Figure 5A and Figure 7A). Ouchi et al. argued that the hydrophobic barrier of the membrane becomes more polar and, hence, more permeable for small hydrophilic molecules, only if the hydrophilic group at the truncation site is pointing to the membrane interior.⁴⁸ This might explain why permeabilization toward SRB at low pH was quicker with PC(16:0/9:0(COOH@C9)) than PC(16:0/9:0(CHO@C9)), maybe because of the slightly higher polarity of the COOH function. For the passage of larger and/or charged molecules such as FITC-Dextran, the formation of larger pores would be necessary. It is possible that aggregation of several OxPCs with their aldehyde/nonionized carboxyl function pointing to the membrane interior leads to the formation of larger pores.⁴⁸ We did not observe them in our MD simulations, although full pore formation occurs in time scales that are generally not accessible by MD simulations.

Interestingly, high temperatures favored the formation of curved aggregates over bilayers (Figure S7B). It has been proposed for truncated ΔC6 and ΔC12 phospholipids bearing a charged nitrobenzoxadiazole (NBD) group at the truncation site that the conformation of the charged truncated chain depends on the packing of the surrounding bilayer. In a densely packed bilayer environment at low temperature, the NBD group is rather aligned with the acyl chains of other phospholipids, while lower packing at higher temperature promotes the outward conformation.^{57,58} A similar behavior of PC(16:0/9:0(COOH@C9)) in membranes would explain the temperature dependent curvature induction. Since headgroup packing is reduced in the presence of truncated OxPCs (Figure 4 and Figure S7A), an increase in their concentrations might thus facilitate the interfacial position of the COOH group.

It seems that some of the effects of PC(16:0/9:0(CHO@C9)) and PC(16:0/9:0(COOH@C9)) are additive (Figure 2F, Figure 3F, Figure S1E, Figure S4F, and Figure S7C,D). This was especially interesting regarding NBD-PE quenching by dithionite since the threshold for individual OxPCs to induce permeabilization or flip-flopping was abolished when a

combination of both OxPCs at different molar ratios was used (Figure 2F and Figure S1E). As discussed above, packing defects induced by either of the OxPCs might facilitate the effect of the other OxPCs.^{55,59} It is also not clear if both molecules could aggregate to form pores or if mixed lipid aggregates, such as mixed micelles, are formed.

We conclude that the different mechanisms of permeabilization induced by oxidized lipid aldehydes and carboxylic acids depend on a series of parameters (Figure 9), including the pH, ionization state, and the length of the truncated acyl chains.

Physiologically speaking, these results show that permeabilization by truncated carboxylated OxPCs is pH dependent. Upon immune response, macrophages produce ROS at pH 4–5 to kill phagocytosed microorganisms. Since truncated carboxylated lipids are produced in this acid environment, the membrane permeability of these microorganisms might be enhanced and increase the killing efficiency. We recently observed in bacterial membranes that permeabilization at pH = 4.5 upon oxidative stress was stronger.⁶⁰ This could indicate that different oxidation products were formed, that acidity itself influences permeabilization, or that permeabilization by OxPCs is enhanced, as we observed here. The release of OxPCs from the membrane is physiologically relevant since certain OxPCs, such as lipid hydroperoxides, have also signaling functions in cells.⁶¹ Truncated OxPCs would thus not need specific transport mechanisms and could simply diffuse to neighboring cells and exert paracrine or membrane destabilizing effects.

Finally, truncated carboxylic OxPCs might be interesting in lipid-based pharmaceutical delivery systems. The ζ -potential of lipid vesicles can be a strong indicator of the stability of the system.⁶² In pH-dependent liposomal delivery systems that are used to increase content release in the stomach (pH = 1–4), the intestine (pH = 7–8), or lysosomes (pH 4.5), truncated carboxyls could diminish the negative ζ -potential of vesicles and increase release of pharmacologically active substances at low pH.

CONCLUSION

We report a systematic investigation on how oxidized lipids with one truncated acyl chain alter properties of membrane bilayers depending on their present concentrations, functional groups, lengths of truncations, and pH. Oxidized lipids with longer truncated tails ($\Delta C9$) are more effective in permeabilizing membranes than those with shorter ones ($\Delta C5$). Lipid aldehydes are likely to permeabilize membrane toward molecules with larger size (e.g., FITC-Dextran and sulforhodamine B) while lipid carboxylic acids tend to permeabilize membrane toward molecules of smaller size or increase lipid flip-flop at neutral pH. At neutral pH, carboxylic acids also induce the formation of highly curved isotropic lipid aggregates and possibly toroidal pores because the ionized carboxyl group protrudes to the membrane interface and induces a high intrinsic molecular curvature. Ionization also seems to favor the release of the lipids from the lipid bilayer and results in a quicker accumulation at the air–water interface. At low pH, the truncated carboxylic acid is nonionized and remains probably inside the hydrophobic core at an angle that prevents the formation of highly curved isotropic structures. However, a low pH of 4.5 increases the capacity of the lipid carboxyl to permeabilize membranes toward larger molecules, similarly to the truncated aldehyde. The mechanism of permeabilization probably depends here on the reduction of the effectiveness of the hydrophobic barrier because of the presence of hydrophilic

functions inside the membrane or the formation of nontoroidal pores.

Our work provides a whole picture of membranolytic effects of truncated oxidized lipids, which could help to understand the effects of these lipids on biological cell membranes. We further describe the possibility of a paracrine effect of these lipids on neighboring cells and the usefulness of truncated carboxylated lipids in the design of pH-dependent drug delivery systems.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.4c12543>.

Computational and experimental materials and methods; additional data for membrane permeabilization, Cryo-EM imaging, membrane packing, surface tension reduction, and ³¹P NMR analysis (PDF)

AUTHOR INFORMATION

Corresponding Author

Joseph H. Lorent – Membrane Biochemistry & Biophysics, Bijvoet Center for Biomolecular Research, Department of Chemistry, Utrecht University, 3584 CH Utrecht, The Netherlands; Cellular and Molecular Pharmacology, Translational Research from Experimental and Clinical Pharmacology to Treatment Optimization, Louvain Drug Research Institute, UCLouvain, B-1200 Brussels, Belgium; orcid.org/0000-0002-7537-8521; Email: joseph.lorent@uclouvain.be

Authors

Min Xie – Membrane Biochemistry & Biophysics, Bijvoet Center for Biomolecular Research, Department of Chemistry, Utrecht University, 3584 CH Utrecht, The Netherlands; orcid.org/0009-0002-2828-8096

Maik G. N. Derks – Membrane Biochemistry & Biophysics, Bijvoet Center for Biomolecular Research, Department of Chemistry, Utrecht University, 3584 CH Utrecht, The Netherlands; Nuclear Magnetic Resonance Spectroscopy, Bijvoet Center for Biomolecular Research, Department of Chemistry, Utrecht University, 3584 CH Utrecht, The Netherlands; orcid.org/0000-0001-8962-5362

Eveline H. W. Koch – Membrane Biochemistry & Biophysics, Bijvoet Center for Biomolecular Research, Department of Chemistry, Utrecht University, 3584 CH Utrecht, The Netherlands; orcid.org/0009-0004-3432-2910

C. Bjorn van Boven – Membrane Biochemistry & Biophysics, Bijvoet Center for Biomolecular Research, Department of Chemistry, Utrecht University, 3584 CH Utrecht, The Netherlands

Minchakarn Janlad – Department of Physics, Faculty of Science, Kasetsart University, Bangkok 10900, Thailand; Computational Biomodelling Laboratory for Agricultural Science and Technology (CBLAST), Faculty of Science, Kasetsart University, Bangkok 10900, Thailand

Behnaz Bagheri – Department of Applied Physics and Science Education, Technical University of Eindhoven, Eindhoven 5600 MB, The Netherlands; Institute for Complex Molecular Systems, Eindhoven 5600 MB, The Netherlands

Zexi Xu – Membrane Biochemistry & Biophysics, Bijvoet Center for Biomolecular Research, Department of Chemistry, Utrecht University, 3584 CH Utrecht, The Netherlands

754 **Daria Kovryzhenko** – Cellular and Molecular Pharmacology,
755 Translational Research from Experimental and Clinical
756 Pharmacology to Treatment Optimization, Louvain Drug
757 Research Institute, UCLouvain, B-1200 Brussels, Belgium
758 **Cornelis A. van Walree** – Membrane Biochemistry &
759 Biophysics, Bijvoet Center for Biomolecular Research,
760 Department of Chemistry, Utrecht University, 3584 CH
761 Utrecht, The Netherlands; University College Utrecht, 3584
762 ED Utrecht, The Netherlands; orcid.org/0000-0002-6178-1424
763
764 **Ana Sobota** – Atmospheric Pressure Non-Thermal Plasmas and
765 Their Interaction with Targets, Applied Physics Department,
766 Eindhoven University of Technology, 5600 MB Eindhoven,
767 The Netherlands
768 **Markus Weingarth** – Nuclear Magnetic Resonance
769 Spectroscopy, Bijvoet Center for Biomolecular Research,
770 Department of Chemistry, Utrecht University, 3584 CH
771 Utrecht, The Netherlands; orcid.org/0000-0003-0831-8673
772
773 **Jirasak Wong-ekkabut** – Department of Physics, Faculty of
774 Science, Kasetsart University, Bangkok 10900, Thailand;
775 Computational Biomodelling Laboratory for Agricultural
776 Science and Technology (CBLAST), Faculty of Science,
777 Kasetsart University, Bangkok 10900, Thailand
778 **Mikko Karttunen** – Department of Chemistry, Western
779 University, London, Ontario N6A 5B7, Canada; Department
780 of Physics and Astronomy, Western University, London,
781 Ontario N6A 3K7, Canada; orcid.org/0000-0002-8626-3033
782
783 **Eefjan Breukink** – Membrane Biochemistry & Biophysics,
784 Bijvoet Center for Biomolecular Research, Department of
785 Chemistry, Utrecht University, 3584 CH Utrecht, The
786 Netherlands; orcid.org/0000-0002-7311-0660
787
788 **J. Antoinette Killian** – Membrane Biochemistry & Biophysics,
789 Bijvoet Center for Biomolecular Research, Department of
790 Chemistry, Utrecht University, 3584 CH Utrecht, The
791 Netherlands
792 **Andreas F. P. Sonnen** – Membrane Biochemistry & Biophysics,
793 Bijvoet Center for Biomolecular Research, Department of
794 Chemistry, Utrecht University, 3584 CH Utrecht, The
795 Netherlands; Pathology Department, University Medical
Center Utrecht, 3508 GA Utrecht, The Netherlands

796 Complete contact information is available at:

797 <https://pubs.acs.org/10.1021/jacs.4c12543>

798 Notes

799 The authors declare no competing financial interest.

800 ■ ACKNOWLEDGMENTS

801 We are grateful to Bonny W. M. Kuipers for support in DLS
802 analysis and Martin F. Haase for providing pendant drop
803 tensiometer equipment. We further thank the Alliance of
804 Utrecht University and Eindhoven University of Technology
805 (grant number BG.180191.4.09) for financing this project. M.X.
806 was supported by a scholarship from the China Scholarship
807 Council (CSC), <https://www.csc.edu.cn/>, file no.
808 201809370077. M.K. thanks the Canada Research Chairs
809 Program and the Natural Sciences and Engineering Research
810 Council of Canada (NSERC) for financial support. J.W. is
811 supported by the National Science Research and Innovation
812 fund (NSRF) via the Program Management Unit for Human
813 Resources and Institutional Development, Research and

Innovation [grant numberB42G670041]. J.H.L. thanks the
Fédération Wallonie-Bruxelles (grant number 24/29-143) for
financial support. The authors also thank the Digital Research
Alliance of Canada for computational resources.

■ REFERENCES

- (1) Byrns, C. N.; Perlegos, A. E.; Miller, K. N.; Jin, Z.; Carranza, F. R.; Manchandra, P.; Beveridge, C. H.; Randolph, C. E.; Chaluvadi, V. S.; Zhang, S. L.; Srinivasan, A. R.; Bennett, F. C.; Sehgal, A.; Adams, P. D.; Chopra, G.; Bonini, N. M. Senescent Glia Link Mitochondrial Dysfunction and Lipid Accumulation. *Nature* **2024** *630* (8016), 475–483.
- (2) Yoritaka, A.; Hattori, N.; Uchida, K.; Tanaka, M.; Stadtman, E. R.; Mizuno, Y. Immunohistochemical Detection of 4-Hydroxynonenal Protein Adducts in Parkinson Disease. *Proc. Natl. Acad. Sci. U. S. A.* **1996**, *93* (7), 2696–2701.
- (3) Sanyal, J.; Bandyopadhyay, S. K.; Banerjee, T. K.; Mukherjee, S. C.; Chakraborty, D. P.; Ray, B. C.; Rao, V. R. Plasma Levels of Lipid Peroxides in Patients with Parkinson's Disease. *Eur. Rev. Med. Pharmacol. Sci.* **2009**, *13* (2), 129–132.
- (4) Navarro, A.; Boveris, A.; Bández, M. J.; Sánchez-Pino, M. J.; Gómez, C.; Muntané, G.; Ferrer, I. Human Brain Cortex: Mitochondrial Oxidative Damage and Adaptive Response in Parkinson Disease and in Dementia with Lewy Bodies. *Free Radic Biol. Med.* **2009**, *46* (12), 1574–1580.
- (5) Valko, M.; Leibfritz, D.; Moncol, J.; Cronin, M. T. D.; Mazur, M.; Telser, J. Free Radicals and Antioxidants in Normal Physiological Functions and Human Disease. *Int. J. Biochem. Cell Biol.* **2007**, *39* (1), 44–84.
- (6) Negre-Salvayre, A.; Auge, N.; Ayala, V.; Basaga, H.; Boada, J.; Brenke, R.; Chapple, S.; Cohen, G.; Feher, J.; Grune, T.; Lengyel, G.; Mann, G. E.; Pamplona, R.; Poli, G.; Portero-Otin, M.; Riahi, Y.; Salvayre, R.; Sasson, S.; Serrano, J.; Shamni, O.; Siems, W.; Siow, R. C. M.; Wiswedel, I.; Zarkovic, K.; Zarkovic, N. Pathological Aspects of Lipid Peroxidation. *Free Radic Res.* **2010**, *44* (10), 1125–1171.
- (7) Wang, C.-H.; Chang, R.-W.; Ko, Y.-H.; Tsai, P.-R.; Wang, S.-S.; Chen, Y.-S.; Ko, W.-J.; Chang, C.-Y.; Young, T.-H.; Chang, K.-C. Prevention of Arterial Stiffening by Using Low-Dose Atorvastatin in Diabetes Is Associated with Decreased Malondialdehyde. *PLoS One* **2014**, *9* (3), No. e90471.
- (8) Cristalli, D. O.; Arnal, N.; Marra, F. A.; de Alaniz, M. J. T.; Marra, C. A. Peripheral Markers in Neurodegenerative Patients and Their First-Degree Relatives. *J. Neurol Sci.* **2012**, *314* (1–2), 48–56.
- (9) Xie, M.; Koch, E. H. W.; van Walree, C. A.; Sobota, A.; Sonnen, A. F. P.; Breukink, E.; Killian, J. A.; Lorent, J. H. Two Separate Mechanisms Are Involved in Membrane Permeabilization during Lipid Oxidation. *Biophys. J.* **2023**, *122* (23), 4503.
- (10) Murphy, M. P. How Mitochondria Produce Reactive Oxygen Species. *Biochem. J.* **2009**, *417* (1), 1–13.
- (11) Ayala, A.; Muñoz, M. F.; Argüelles, S. Lipid Peroxidation: Production, Metabolism, and Signaling Mechanisms of Malondialdehyde and 4-Hydroxy-2-Nonenal. *Oxid Med. Cell Longev* **2014**, *2014*, 360438.
- (12) Altan, O.; Pabuçcuoğlu, A.; Altan, A.; Konyalıoğlu, S.; Bayraktar, H. Effect of Heat Stress on Oxidative Stress, Lipid Peroxidation and Some Stress Parameters in Broilers. *Br Poult Sci.* **2003**, *44* (4), 545–550.
- (13) Spickett, C. M. Formation of Oxidatively Modified Lipids as the Basis for a Cellular Epilipidome. *Front Endocrinol (Lausanne)* **2020**, *11*, 1–8.
- (14) Hutchins, P. M.; Moore, E. E.; Murphy, R. C. Electrospray MS/MS Reveals Extensive and Nonspecific Oxidation of Cholesterol Esters in Human Peripheral Vascular Lesions[S]. *J. Lipid Res.* **2011**, *52* (11), 2070–2083.
- (15) Guo, S.; Lu, J.; Zhuo, Y.; Xiao, M.; Xue, X.; Zhong, S.; Shen, X.; Yin, C.; Li, L.; Chen, Q.; Zhu, M.; Chen, B.; Zhao, M.; Zheng, L.; Tao, Y.; Yin, H. Endogenous Cholesterol Ester Hydroperoxides Modulate

- 880 Cholesterol Levels and Inhibit Cholesterol Uptake in Hepatocytes and
881 Macrophages. *Redox Biol.* **2019**, *21*, 101069.
- 882 (16) Guéraud, F.; Atalay, M.; Bresgen, N.; Cipak, A.; Eckl, P. M.; Huc,
883 L.; Jouanin, L.; Siems, W.; Uchida, K. Chemistry and Biochemistry of
884 Lipid Peroxidation Products. *Free Radic Res.* **2010**, *44* (10), 1098–
885 1124.
- 886 (17) Sousa, B. C.; Pitt, A. R.; Spickett, C. M. Chemistry and Analysis of
887 HNE and Other Prominent Carbonyl-Containing Lipid Oxidation
888 Compounds. *Free Radic Biol. Med.* **2017**, *111*, 294–308.
- 889 (18) Reis, A.; Spickett, C. M. Chemistry of Phospholipid Oxidation.
890 *Biochim Biophys Acta Biomembr* **2012**, *1818* (10), 2374–2387.
- 891 (19) Jurkiewicz, P.; Cwiklik, L.; Jungwirth, P.; Hof, M. Lipid
892 Hydration and Mobility: An Interplay between Fluorescence Solvent
893 Relaxation Experiments and Molecular Dynamics Simulations.
894 *Biochimie*. **2012**, *94* (1), 26–32.
- 895 (20) Israelachvili, J. *Intermolecular and Surface Forces*, Third ed.;
896 Academic Press, 2011; DOI: 10.1016/C2009-0-21560-1.
- 897 (21) Singer, S. J.; Nicolson, G. L. The Structure and Chemistry of
898 Mammalian Cell Membranes. *Am. J. Pathol.* **1971**, *65* (2), 427–437.
- 899 (22) Rosa, R. De; Spinozzi, F.; Itri, R. Hydroperoxide and Carboxyl
900 Groups Preferential Location in Oxidized Biomembranes Experimen-
901 tally Determined by Small Angle X-Ray Scattering: Implications in
902 Membrane Structure. *Biochim Biophys Acta Biomembr* **2018**, *1860* (11),
903 2299–2307.
- 904 (23) Wong-Ekkabut, J.; Xu, Z.; Triampo, W.; Tang, I. M.; Tieleman,
905 D. P.; Monticelli, L. Effect of Lipid Peroxidation on the Properties of
906 Lipid Bilayers: A Molecular Dynamics Study. *Biophys. J.* **2007**, *93* (12),
907 4225–4236.
- 908 (24) Khandelia, H.; Mouritsen, O. G. Lipid Gymnastics: Evidence of
909 Complete Acyl Chain Reversal in Oxidized Phospholipids from
910 Molecular Simulations. *Biophys. J.* **2009**, *96* (7), 2734–2743.
- 911 (25) Bagheri, B.; Boonnoy, P.; Wong-Ekkabut, J.; Karttunen, M. Effect
912 of Oxidation on POPC Lipid Bilayers: Anionic Carboxyl Group Plays a
913 Major Role. *Phys. Chem. Chem. Phys.* **2023**, *25* (27), 18310–18321.
- 914 (26) Bour, A.; Kruglik, S. G.; Chabanon, M.; Rangamani, P.; Puff, N.;
915 Bonneau, S. Lipid Unsaturation Properties Govern the Sensitivity of
916 Membranes to Photoinduced Oxidative Stress. *Biophys. J.* **2019**, *116*
917 (5), 910–920.
- 918 (27) Parra-Ortiz, E.; Browning, K. L.; Damgaard, L. S. E.; Nordström,
919 R.; Micciulla, S.; Bucciarelli, S.; Malmsten, M. Effects of Oxidation on
920 the Physicochemical Properties of Polyunsaturated Lipid Membranes.
921 *J. Colloid Interface Sci.* **2019**, *538*, 404–419.
- 922 (28) Boonnoy, P.; Jarerattanachai, V.; Karttunen, M.; Wong-Ekkabut,
923 J. Bilayer Deformation, Pores, and Micellation Induced by Oxidized
924 Lipids. *J. Phys. Chem. Lett.* **2015**, *6* (24), 4884–4888.
- 925 (29) Heuvingh, J.; Bonneau, S. Asymmetric Oxidation of Giant
926 Vesicles Triggers Curvature-Associated Shape Transition and Per-
927 meabilization. *Biophys. J.* **2009**, *97* (11), 2904–2912.
- 928 (30) Yusupov, M.; Van der Paal, J.; Neyts, E. C.; Bogaerts, A.
929 Synergistic Effect of Electric Field and Lipid Oxidation on the
930 Permeability of Cell Membranes. *Biochim Biophys Acta Gen Subj* **2017**,
931 *1861* (4), 839–847.
- 932 (31) Mertins, O.; Bacellar, I. O. L.; Thalmann, F.; Marques, C. M.;
933 Baptista, M. S.; Itri, R. Physical Damage on Giant Vesicles Membrane as
934 a Result of Methylene Blue Photoirradiation. *Biophys. J.* **2014**, *106* (1),
935 162–171.
- 936 (32) Tsubone, T. M.; Junqueira, H. C.; Baptista, M. S.; Itri, R.
937 Contrasting Roles of Oxidized Lipids in Modulating Membrane
938 Microdomains. *Biochim Biophys Acta Biomembr* **2019**, *1861* (3), 660–
939 669.
- 940 (33) Runas, K. A.; Acharya, S. J.; Schmidt, J. J.; Malmstadt, N.
941 Addition of Cleaved Tail Fragments during Lipid Oxidation Stabilizes
942 Membrane Permeability Behavior. *Langmuir* **2016**, *32* (3), 779–786.
- 943 (34) Bagheri, B.; Boonnoy, P.; Wong-Ekkabut, J.; Karttunen, M. Effect
944 of Oxidation on POPC Lipid Bilayers: Anionic Carboxyl Group Plays a
945 Major Role. *Phys. Chem. Chem. Phys.* **2023**, *25* (27), 18310–18321.
- 946 (35) Parasassi, T.; De Stasio, G.; d'Ubaldo, A.; Gratton, E. Phase
947 Fluctuation in Phospholipid Membranes Revealed by Laurdan
948 Fluorescence. *Biophys. J.* **1990**, *57* (6), 1179–1186.
- (36) Lorent, J. H.; Levental, K. R.; Ganesan, L.; Rivera-Longworth, 949
G.; Sezgin, E.; Doktorova, M.; Lyman, E.; Levental, I. Plasma 950
Membranes Are Asymmetric in Lipid Unsaturation, Packing and 951
Protein Shape. *Nat. Chem. Biol.* **2020**, *16* (6), 644–652. 952
- (37) Suhaj, A.; Gowland, D.; Bonini, N.; Owen, D. M.; Lorenz, C. D. 953
Laurdan and Di-4-ANEPPDHQ Influence the Properties of Lipid 954
Membranes: A Classical Molecular Dynamics and Fluorescence Study. 955
J. Phys. Chem. B **2020**, *124* (50), 11419–11430. 956
- (38) Bacalum, M.; Zorila, B.; Radu, M. Fluorescence Spectra 957
Decomposition by Asymmetric Functions: Laurdan Spectrum 958
Revisited. *Anal. Biochem.* **2013**, *440* (2), 123–129. 959
- (39) Sousa, T.; Castro, R. E.; Coutinho, A.; Rodrigues, C. M. P.; 960
Prieto, M.; Fernandes, F. *Measuring the Impact of Bile Acids on the* 961
Membrane Order of Primary Hepatocytes and Isolated Mitochondria by 962
Fluorescence Imaging and Spectroscopy BT - Experimental Cholestasis 963
Research; Vinken, M., Ed.; Springer New York: New York, NY, 2019; pp 964
99–115; DOI: 10.1007/978-1-4939-9420-5_7. 965
- (40) Brown, R. E.; Brockman, H. L. Using Monomolecular Films to 966
Characterize Lipid Lateral Interactions. *Lipid Rafts* **2007**, *398*, 41–58. 967
- (41) Mukherjee, I.; Moulik, S. P.; Rakshit, A. K. Tensiometric 968
Determination of Gibbs Surface Excess and Micelle Point: A Critical 969
Revisit. *J. Colloid Interface Sci.* **2013**, *394* (1), 329–336. 970
- (42) Zhao, J.; Dai, C.; Ding, Q.; Du, M.; Feng, H.; Wei, Z.; Chen, A.; 971
Zhao, M. The Structure Effect on the Surface and Interfacial Properties 972
of Zwitterionic Sulfobetaine Surfactants for Enhanced Oil Recovery. 973
RSC Adv. **2015**, *5* (18), 13993–14001. 974
- (43) Qiao, L.; Ge, A.; Liang, Y.; Ye, S. Oxidative Degradation of the 975
Monolayer of 1-Palmitoyl-2-Oleoyl-Sn-Glycero-3-Phosphocholine 976
(POPC) in Low-Level Ozone. *J. Phys. Chem. B* **2015**, *119* (44), 977
14188–14199. 978
- (44) Gröbner, G.; Williamson, P. Solid-State NMR Methods for 979
Studying Membrane Systems. In *Solid-State NMR*; IOP Publishing, 980
2020; pp 1-1–1-30; DOI: 10.1088/978-0-7503-2532-Sch1. 981
- (45) McLaughlin, A. C.; Cullis, P. R.; Berden, J. A.; Richards, R. E. 31P 982
NMR of Phospholipid Membranes: Effects of Chemical and Anisotropy 983
at High Magnetic Field Strengths. *Journal of Magnetic Resonance* (1969) 984
1975, *20* (1), 146–165. 985
- (46) Mendes Ferreira, T.; Bärenwald, R.; Sood, R.; Volinsky, R.; 986
Drescher, S.; Karlström, G.; Topgaard, D.; Kinnunen, P.; Saalwächter, 987
K.; Ollila, S. Oxidized Lipids in Model Membranes: Atomistic Details 988
from Solid-State NMR Experiments and MD Simulations. *Biophys. J.* 989
2016, *110* (3), 584a. 990
- (47) Jurkiewicz, P.; Olżyńska, A.; Cwiklik, L.; Conte, E.; Jungwirth, P.; 991
Megli, F. M.; Hof, M. Biophysics of Lipid Bilayers Containing 992
Oxidatively Modified Phospholipids: Insights from Fluorescence and 993
EPR Experiments and from MD Simulations. *Biochimica et Biophysica* 994
Acta - Biomembranes **2012**, *1818*, 2388–2402. 995
- (48) Ouchi, Y.; Unoura, K.; Nabika, H. Role of Oxidized Lipids in 996
Permeation of H₂O₂ Through a Lipid Membrane: Molecular 997
Mechanism of an Inhibitor to Promoter Switch. *Sci. Rep* **2019**, *9* (1), 998
1–4. 999
- (49) Sandrin, D.; Wagner, D.; Sitta, C. E.; Thoma, R.; Felekyan, S.; 1000
Hermes, H. E.; Janiak, C.; De Sousa Amadeu, N.; Kühnemuth, R.; 1001
Löwen, H.; Egelhaaf, S. U.; Seidel, C. A. M. Diffusion of Macro- 1002
molecules in a Polymer Hydrogel: From Microscopic to Macroscopic 1003
Scales. *Phys. Chem. Chem. Phys.* **2016**, *18* (18), 12860–12876. 1004
- (50) Volinsky, R.; Cwiklik, L.; Jurkiewicz, P.; Hof, M.; Jungwirth, P.; 1005
Kinnunen, P. K. J. Oxidized Phosphatidylcholines Facilitate Phospho- 1006
lipid Flip-Flop in Liposomes. *Biophys. J.* **2011**, *101* (6), 1376–1384. 1007
- (51) Runas, K. A.; Malmstadt, N. Low Levels of Lipid Oxidation 1008
Radically Increase the Passive Permeability of Lipid Bilayers. *Soft Matter* 1009
2015, *11* (3), 499–505. 1010
- (52) Grauby-Heywang, C.; Moroté, F.; Mathelié-Guinlet, M.; 1011
Gammoudi, I.; Faye, N. R.; Cohen-Bouhacina, T. Influence of Oxidized 1012
Lipids on Palmitoyl-Oleoyl-Phosphatidylcholine Organization, Con- 1013
tribution of Langmuir Monolayers and Langmuir-Blodgett Films. 1014
Chem. Phys. Lipids **2016**, *200*, 74–82. 1015
- (53) Grauby-Heywang, C.; Moroté, F.; Mathelié-Guinlet, M.; 1016
Gammoudi, I.; Faye, N. R.; Cohen-Bouhacina, T. Influence of Oxidized 1017

- 1018 Lipids on Palmitoyl-Oleoyl-Phosphatidylcholine Organization, Con-
1019 tribution of Langmuir Monolayers and Langmuir-Blodgett Films.
1020 *Chem. Phys. Lipids* **2016**, *200*, 74–82.
- 1021 (54) Czajka, A.; Hazell, G.; Eastoe, J. Surfactants at the Design Limit.
1022 *Langmuir* **2015**, *31* (30), 8205–8217.
- 1023 (55) Mendes Ferreira, T.; Sood, R.; Bärenwald, R.; Carlström, G.;
1024 Topgaard, D.; Saalwächter, K.; Kinnunen, P. K. J.; Ollila, O. H. S. Acyl
1025 Chain Disorder and Azelaoyl Orientation in Lipid Membranes
1026 Containing Oxidized Lipids. *Langmuir* **2016**, *32* (25), 6524–6533.
- 1027 (56) Lorent, J.; Le Duff, C. S.; Quetin-Leclercq, J.; Mingeot-Leclercq,
1028 M.-P. Induction of Highly Curved Structures in Relation to Membrane
1029 Permeabilization and Budding by the Triterpenoid Saponins, Alpha-
1030 and Delta-Hederin. *J. Biol. Chem.* **2013**, *288* (20), 14000–14017.
- 1031 (57) Stockl, M.; Plazzo, A. P.; Korte, T.; Herrmann, A. Detection of
1032 Lipid Domains in Model and Cell Membranes by Fluorescence
1033 Lifetime Imaging Microscopy of Fluorescent Lipid Analogues. *J. Biol.*
1034 *Chem.* **2008**, *283* (45), 30828–30837.
- 1035 (58) Filipe, H. A. L.; Pokorná, Š.; Hof, M.; Amaro, M.; Loura, L. M. S.
1036 Orientation of Nitro-Group Governs the Fluorescence Lifetime of
1037 Nitrobenzoxadiazole (NBD)-Labeled Lipids in Lipid Bilayers. *Phys.*
1038 *Chem. Chem. Phys.* **2019**, *21* (4), 1682–1688.
- 1039 (59) Megli, F. M.; Conte, E.; Russo, L. Comparative 5-
1040 Doxylstearoyllecithin and 3-Doxylcholestane EPR Spin Labeling
1041 Study of Phospholipid Bilayer Perturbation by Different Oxidized
1042 Lecithin Species. *Biochim Biophys Acta Biomembr* **2010**, *1798* (1),
1043 1886–1898.
- 1044 (60) Xie, M.; Koch, E. H. W.; van Walree, C. A.; Sobota, A.; Sonnen,
1045 A. F. P.; Killian, J. A.; Breukink, E.; Lorent, J. H. Synergistic Effects of
1046 Oxidative and Acid Stress on Bacterial Membranes of Escherichia Coli
1047 and Staphylococcus Simulans. *Commun. Biol.* **2024**, *7* (1), 1161.
- 1048 (61) Korytowski, W.; Basova, L. V.; Pilat, A.; Kernstock, R. M.; Girotti,
1049 A. W. Permeabilization of the Mitochondrial Outer Membrane by Bax/
1050 Truncated Bid (TBid) Proteins as Sensitized by Cardiolipin Hydro-
1051 peroxide Translocation: Mechanistic Implications for the Intrinsic
1052 Pathway of Oxidative Apoptosis. *J. Biol. Chem.* **2011**, *286* (30), 26334–
1053 26343.
- 1054 (62) Smith, M. C.; Crist, R. M.; Clogston, J. D.; McNeil, S. E. Zeta
1055 Potential: A Case Study of Cationic, Anionic, and Neutral Liposomes.
1056 *Anal Bioanal Chem.* **2017**, *409* (24), 5779–5787.