



# Invasive fountain grass (*Pennisetum setaceum* (Forssk.) Chiov.) increases its potential area of distribution in Tenerife island under future climatic scenarios

Daniele Da Re · Enrico Tordoni · Federico De Pascalis · Zaira Negrín-Pérez · José María Fernández-Palacios · José Ramón Arévalo · Duccio Rocchini · Félix Manuel Medina · Rüdiger Otto · Eduardo Arlé · Giovanni Bacaro

Received: 26 October 2019 / Accepted: 4 June 2020  
© Springer Nature B.V. 2020

**Abstract** Mapping the distribution of invasive species under current and future climate conditions is crucial to implement sustainable and effective conservation strategies. Several studies showed how invasive species may benefit from climate change fostering their invasion rate and, consequently,

affecting the native species community. In the Canary Islands and on Tenerife in particular, previous research mostly focused on climate change impacts on the native communities, whereas less attention has been paid on alien species distribution under climate change scenarios. In this study, we modelled the habitat distribution of *Pennisetum setaceum*, one of the most invasive alien species on Tenerife. In addition, we described the species' potential distribution shift in the light of two climate change scenarios (RCP2.6, RCP8.5), highlighting the areas that should be prioritized during management and eradication

Communicated by Paul M. Ramsay.

**Electronic supplementary material** The online version of this article (doi:<https://doi.org/10.1007/s11258-020-01046-9>) contains supplementary material, which is available to authorized users.

D. Da Re · E. Tordoni · F. De Pascalis · G. Bacaro  
Department of Life Sciences, University of Trieste, Via L. Giorgieri 10, 34127 Trieste, Italy

*Present Address:*

D. Da Re (✉)  
George Lemaitre Center for Earth and Climate Research,  
Earth and Life Institute, UCLouvain, Place Louis Pasteur  
3, 1348 Louvain-la-Neuve, Belgium  
e-mail: daniele.dare@uclouvain.be

*Present Address:*

F. De Pascalis  
Department of Environmental Sciences and Policy,  
University of Milan, via Celoria 26, 20133 Milan, Italy

Z. Negrín-Pérez · J. M. Fernández-Palacios ·  
J. R. Arévalo · R. Otto  
Departamento de Botánica, Ecología y Fisiología Vegetal,  
Universidad de La Laguna, Avenida Astrofísico Francisco  
Sánchez s/n, 38206 La Laguna, Tenerife, Spain

D. Rocchini  
Alma Mater Studiorum University of Bologna,  
Department of Biological, Geological and Environmental  
Sciences, via Irnerio 42, 40126 Bologna, Italy

D. Rocchini  
Department of Applied Geoinformatics and Spatial  
Planning, Faculty of Environmental Sciences, Czech  
University of Life Sciences Prague, Kamýcka 129,  
Suchbát, 16500 Praha, Czech Republic

F. M. Medina  
Servicio de Medio Ambiente, Cabildo Insular de La  
Palma, Avenida Los Indianos, 20,  
38700 Santa Cruz de La Palma, Canary Islands, Spain

E. Arlé  
German Centre for Integrative Biodiversity Research  
(iDiv), Leipzig Halle-Jena-Leipzig, Deutscher Platz 5e,  
04103 Leipzig, Germany

programs. *P. setaceum*'s suitable areas are located in the coastal area, with higher habitat suitability near cities and below 800 m asl. In both future climate change scenarios, the geographic distribution of *P. setaceum* suitable areas is characterized by an elevational shift, which is more pronounced in the RCP8.5 scenario. Despite being drought resistant, water supply is crucial for the species' seed germination, thus supporting future species' shift to higher elevation and in the north–north–west part of the island, where it could benefit from the combined effect of orographic precipitations and humidity carried by trade winds.

**Keywords** Habitat suitability models · Invasive alien species · Canary Archipelago · Global warming

## Introduction

Climate change is having strong ecological impacts on biodiversity from the polar regions to the tropics, and predicting the response of biodiversity to future climate change scenarios has become a primary field of research (Pereira et al. 2010; Bellard et al. 2012; Matías and Jump 2013; Dyderski et al. 2018). Global warming, along with the reduction in precipitation during the growing season, may strongly alter species distribution (Kleinbauer et al. 2010; Stocker 2014). Furthermore, the increase in CO<sub>2</sub> atmospheric concentration and Nitrogen deposition promotes the presence of invasive species and enhances the risk of biological invasions (Dukes and Mooney 1999). In this context, special attention should be paid to island systems: islands are, in fact, at the forefront against global changes such as sea-level rise and biological invasions (Bellard et al. 2014; Pyšek et al. 2017). Islands are well known to host habitats rich in rare and endemic species; hence, more than one third of biodiversity hotspots in the world are entirely, or largely, within islands (Bellard et al. 2014).

Biological invasions are dramatically threatening island biodiversity and ecosystem integrity (Hulme 2009; Scalera et al. 2012) since invasive alien species (hereafter IAS, see Pyšek et al. 2004 for comprehensive definitions) may directly reduce local plant species diversity (Tordoni et al. 2019). However, the deleterious effect of plant invasion is not limited to competition with native species, but it could affect the

whole ecosystem, especially enclosed and fragile ones such as islands. Firstly, changes in carbon and nitrogen soil dynamics, resulting from alien plants invasions (Vilà and López-Darias 2006; Vilà et al. 2011; Qian and Ricklefs 2006), impact soil biotic community and potentially alter important processes, such as mutualism. In nutrient-enriched soils, for example, mycorrhizal populations can become antagonistic to hosts (Toby Kiers et al. 2010). The balance in macroinvertebrate communities can be disrupted (Gremmen et al. 1998), with potential effects on the whole ecosystems. Secondly, invasive alien plant species could promote alien insect presence (Morales and Aizen 2002), which has the potential to affect entire habitats and ultimately alter ecosystem functioning and services (Kenis et al. 2009). Finally, recent studies suggest that IAS may benefit from climate change (e.g. Kleinbauer et al. 2010; Brundu and Richardson 2016; Dyderski et al. 2018), which foster their invasion rate and, consequently, invasion-related risks.

Due to the high rate of endemism (Whittaker and Fernández-Palacios 2007; Fernández-Palacios et al. 2016), the Canary archipelago represents an extremely vulnerable area for alien species invasions (Courchamp et al. 2003; Millennium Ecosystem Assessment 2005; Kueffer et al. 2010; Bacaro et al. 2015). Particularly, the island of Tenerife hosts a total of 9325 species, 1468 of them being vascular plants with 279 local endemic plant species (Arechavaleta et al. 2010). This island has been affected by an intense human activity that altered the natural ecosystems, causing fragmentation and the introduction of invasive species in protected areas (Delgado et al. 2004), such as fountain grass (*Pennisetum setaceum* (Forssk.) Chiov.).

*Pennisetum setaceum* is a perennial, wind-dispersed, apomictic, C4 bunch grass native to North Africa and the Middle East (Williams et al. 1995; Poulin et al. 2007). It was introduced in the Canary Islands as an ornamental plant and its presence has been reported since 1940s (Hansen et al. 1970; de Paz et al. 1999). Currently, it is considered one of the most invasive species on Tenerife (Arechavaleta et al. 2010; Francisco-Ortega et al. 2009) as well as in other areas (e.g. California, Hawaii, South Africa; Williams et al. 1995; Poulin et al. 2007; Rahlao et al. 2010), constantly expanding its range along roads, from urbanized areas toward natural ones (Martín Esquivel et al. 1995; González-Rodríguez et al. 2010). Elevated phenotypic plasticity, characterised by variation in functional trait value according to the environment, and

its resilience, makes *P. setaceum* well adapted to different ecological conditions (Poulin et al. 2007; González-Rodríguez et al. 2010). Moreover, it can establish symbiosis with different local mycorrhizal fungi (AMF) communities, further promoting a successful establishment and spread (Rodríguez-Caballero et al. 2018). It is an aggressive invader of arid and semiarid coastal habitats such as thermo-xerophilous grasslands and shrublands. Here, it establishes almost monospecific stands and causes long-lasting ecological consequences (Cordell and Sandquist 2008; González-Rodríguez et al. 2010). The species is known to have negative impacts on resources acquisitions in the dry forest plant communities and it can interact with the soil bacterial community, shifting its structure and composition, which can result in severe alterations of the N cycle (Rodríguez-Caballero et al. 2017). Moreover, it is a fire promoting species that can cause dramatic environmental changes, since fire is one of the most important drivers of land use and atmospheric changes globally (D'Antonio and Vitousek 1992). Finally, it can promote the presence of the alien invasive leafhopper *Balclutha brevis* (Bella et al. 2012). It has been estimated that approximately 30% of all protected areas of the Canary Islands have been invaded by fountain grass (Martín Esquivel et al. 1995; González-Rodríguez et al. 2010). Considering the dramatic effects that an uncontrolled increase in its distribution could create on Tenerife, understanding *P. setaceum* future habitat distribution is particularly important both from a conservational and ecological perspective, and it is a time-sensitive task.

Correlative models such as habitat suitability models (HSMs) have extensively been used so far to estimate the geographic distribution of a species based on an index of environmental similarity (Kearney 2006; Peterson et al. 2011; Guisan et al. 2017) which, in turn, is estimated starting from its occurrences in relation to its environmental determinants (Thuiller 2007; Steiner et al. 2008; Jiménez-Valverde et al. 2011). HSMs became a central tool in invasion biology, providing both interesting insight on species' ecology and practical suggestions for eradication management. For invasive species, the correct projection in space and time of HSMs estimates depends on different assumptions, which are defined by the research questions. It is fundamental to consider niche conservatism among native and invasive range when the research objective is to project the suitability index estimated in the native range into the invasive one. When using the species


invaded range to train a HSM, the invasive species must be at quasi-equilibrium with the environment in which it occurs (Guisan and Thuiller 2005; Gallien et al. 2012). However, these assumptions are not always met, since the naturalized climatic niche of invasive species may differ from the natives climatic niches (e.g. Medley 2010; Early and Sax 2014) and since an invasive species is not at equilibrium with its environment until the latest stage of invasion (Barbet-Massin et al. 2018). Both cases result in a likely underestimate on the species predicted habitat suitability distribution. However, in situations where assumptions cannot be met for practical reasons (e.g. impossibility to compare native and invasive niche), HSMs may still provide predictions which are useful for invasive species management, such as highlighting the areas of likely expansion in the near future, which should be targeted and prioritized by conservation efforts (West et al. 2016).

Considering the importance of predictive tools to halt the loss of biodiversity, especially in relation to the spread of IAS fostered by climate change, here we aimed at (i) estimating *P. setaceum*'s current habitat distribution through HSM, across its' invaded range in Tenerife, and (ii) assessing the effect of climate change in shaping *P. setaceum*'s habitat distribution under two climate change scenarios (RCP2.6, RCP8.5). We hypothesize that *P. setaceum* may benefit from climate warming, increasing its current habitat distribution by spreading toward more humid zone of the island (e.g. at higher elevation), hence potentially increasing its impact on native habitats. Final goal of this study is to disclose a clearer image of those portions of Tenerife which will be more prone to the invasion process and therefore in urgent need of eradication and control activities.

## Materials and methods

### Study area

The study was carried out in Tenerife (27°–29°N, 13°–18°W; Fig. 1), the largest island of the Canary archipelago and the one with the highest elevation within Macaronesia (Mount Teide, 3718 m asl). Strong variations in elevation and aspect define local mesoclimatic zones and land uses that are primary factors in structuring both native and alien plant communities on the Canary Islands (Whittaker and



**Fig. 1** *Pennisetum setaceum* thinned occurrences distribution on Tenerife

Fernández-Palacios 2007). Mesoclimate is affected by the trade winds that create a contrast between the northern or windward aspect (more humid and cloudier) and the southern or leeward aspect (more arid and cloudless).

#### Response variable

Data of *P. setaceum* occurrences were obtained from ATLANTIS (Gobierno de Canarias 2015), a regional database containing IAS occurrences within a grid of  $500 \times 500$  m square cells covering the entire archipelago. Species records spanned in time from

1970 to 2017 but only those from 2005 to 2014 were considered for the analysis in order to maintain data consistency with the climatic data available. Only the grid cells covering Tenerife landmass were selected from the grid covering the whole Canary archipelago (5515 selected out of 8519 total cells). This dataset was complemented with records obtained by the Teno Rural park internal database (Suppl. Mat.).

## Predictors

Three abiotic variables were selected to predict species' habitat distribution based on previous knowledge on the ecology and biology of *P. setaceum*.

Mean winter precipitation and mean spring precipitation were chosen since the period October–March concentrates the 87.3 % of Canary Islands' annual precipitation (Sánchez-Benítez et al. 2017). Moreover, it has been reported that in Mediterranean-type climate areas, *P. setaceum* germination occurs primarily in the Winter–Spring period (especially in December–January), followed by a constant decline towards the Summer due to moisture reduction (Hernandez and Sandquist 2011). Mean spring temperature was also selected, since it is known to be a limiting factor for alien species colonization along an elevational gradient, being usually correlated to elevation (Barni et al. 2012; Bacaro et al. 2015; Steinbauer et al. 2017) and because the species has a maximum growth in warmer conditions (Sweet and Holt 2015).

Finally, road kernel density was selected as a proxy of propagule pressure and species dispersion, especially for *P. setaceum* (Foxcroft et al. 2019). Indeed anthropogenic activities enhance alien species' spread (Pyšek et al. 2010; Tordoni et al. 2017), and roads are known to be primary introduction pathways. Particularly, roadside may serve as invasion epicentres for annual species with high reproductive rates (Pauchard and Alaback 2004; Bacaro et al. 2015; Da Re et al. 2019).

Monthly precipitation of October–March periods and monthly temperature of March–May periods, spanning from 2005 to 2014, were obtained from Agencia Estatal de Meteorología (AEMET, accessed on April 2016). The methodology presented in Da Re et al. (2019) was used for filtering and the processing of the meteorological data in order to obtain continuous climatic interpolated surfaces through a co-Kriging procedure, using elevation as covariate (Myers 1984; Garzón-Machado et al. 2014; Wilson and Silander 2014) and the R-package “geoR” (Ribeiro et al. 2001). Road kernel density, calculated on the road network using 10 km regularly distributed sample points, was used here as a proxy of human disturbance and as a source of propagule pressure (Bacaro et al. 2015; Da Re et al. 2019). The digital elevation model (DEM, 10 m of spatial resolution) and road network of

Tenerife Island were downloaded from Cartográfica de Canarias S.A. (GRAFCAN, <https://www.grafcan.es/>, accessed on March 2016), all the derived variables were calculated using the grid spatial resolution as for the IAS occurrences (500 m) (Table 1).

## Climate change predictions

Temperature time series analysis of the Canary Islands reported a temperature increase tendency, showing the greatest increases on island summit (above 2000 m asl) and above the cloud layer of windward mid-altitude areas (Martín et al. 2012; Luque et al. 2014). With regard to precipitation, there is no clear and significant change (García-Herrera et al. 2003; Cropper and Hanna 2014).

Cropper (2013) estimates an increase in surface air temperature of 1 °C and a decrease in water availability of 15–20% due to more than 30% reduction in precipitation (Giorgi 2006; Somot et al. 2008). In order to get a reliable estimate of climate change effects on the climatic variables considered (though deemed approximate), we apply a constant correction on interpolated climatic surfaces. Specifically, we considered the two Representative Concentration Pathways (RCPs), namely RCP2.6 and RCP8.5, as described by the 5th Intergovernmental Panel on Climate Change (IPCC) report representing possible ranges of radiative forcing values in the year 2100 relative to pre-industrial values (+ 2.6 and + 8.5 W/m<sup>2</sup>, respectively). Winter and Spring precipitations were corrected by – 10.6% and – 36.7% (RCP2.6 and RCP8.5) according to Winter (December–January–February, DJF) precipitation reduction prevision made by Cropper (2013), while for Spring, temperature values 0.96 °C (RCP2.6) and 2.68 °C (RCP8.5) were added.

## Habitat suitability modelling

HSMs were performed using Maxent v3.4.1 (Phillips et al. 2017) through the R-package “dismo” (Hijmans et al. 2017). Maxent estimates the relationship between species occurrences and various environmental predictors, identifying areas with suitable environmental conditions for the species, given a background uniform probability of 0.5 (Elith et al. 2011). To reduce the effects of sampling bias and thus avoiding a possible source of model inaccuracy (Phillips et al.

**Table 1** Predictors summary statistics at the occurrences locations

|                                                 | Mean   | SD    | Median | Min    | Max    |
|-------------------------------------------------|--------|-------|--------|--------|--------|
| October–March mean monthly precipitation        | 266.48 | 87.42 | 263.22 | 140.47 | 460.07 |
| Mean spring temperature                         | 18.00  | 1.00  | 18.24  | 15.54  | 19.93  |
| Road kernel density                             | 0.06   | 0.07  | 0.04   | 0.00   | 0.40   |
| RCP2.6 October–March mean monthly precipitation | 223.85 | 73.43 | 221.10 | 118.00 | 386.46 |
| RCP2.6 mean spring temperature                  | 18.96  | 1.00  | 19.20  | 16.50  | 20.89  |
| RCP8.5 October–March mean monthly precipitation | 168.68 | 55.33 | 166.62 | 88.92  | 291.22 |
| RCP8.5 mean spring temperature                  | 20.68  | 1.00  | 20.92  | 18.22  | 22.61  |


2006, 2009; Syfert et al. 2013), spatial filtering with a thinning distance of 2 km was applied to the *P. setaceum* occurrence dataset using the R-package “spThin” (Aiello-Lammens et al. 2015). The thinned occurrence data were then randomly divided into a training set and testing dataset applying a spatial block approach (Guevara et al. 2018; Roberts et al. 2017). Models with different combinations of feature classes (FC: Linear; Quadratic; Linear and Quadratic; Hinge; Linear, Quadratic, and Hinge) and beta regularization multipliers (RM: 0.5–5.0 with 0.5 intervals) were computed in order to avoid overfitting and to approximate optimal levels of complexity. The best model was then chosen based on Akaike Information Criterion corrected for small sample sizes (AICc, Warren and Seifert 2011; Muscarella et al. 2014). Following Guevara et al. (2018), we used all 8134-island pixel as background point to ensure a full representation of environments available for the species. The optimal settings obtained from above were used to train the final models using maxent.jar software. Model outputs were set as logistic response of the predicted distribution. The logistic output was then interpreted conservatively as a suitability index rather than as a probability (Merow et al. 2013) and it was evaluated using both the area under the curve (AUC) provided for the test data (Phillips et al. 2006), and the Boyce’s Index (Hirzel et al. 2006). The Boyce’s Index, computed through the “ecospat” R-package (Di Cola et al. 2017), ranges between  $-1$  (the model predict areas where presences are more frequent as being highly suitable for the species) and  $+1$  (the model predictions are consistent with the distribution of presences in the evaluation dataset), and values close to zero mean that the model is not different from random expectations (Hirzel et al. 2006). Finally, the

model was projected onto current, RCP2.6, and RCP8.5 climatic scenarios using unconstrained extrapolation (Guevara et al. 2018). The R codes used are available in the gitLab repository presented in the Supplementary Materials.

## Results

*Pennisetum setaceum* is mainly present in coastal areas, especially on the North-Western side of the island and near the largest cities (Santa Cruz de Tenerife, San Cristobal de La Laguna and Güímar on the N–E coast; Bajamar, Punta del Hidalgo, and Puerto de La Cruz on the N–W coast, Los Gigantes and Las Americas on the south; Fig. 2 and 2a).

48 occurrences (out of the original 227) were produced using the spatial thinning approach and then these were used to train the model. Model selection based on the AICc criteria pointed out that the most appropriate model was the one having linear, quadratic and product features and a beta regularization multiplier equal to 0.5 (Table 2). For this model, both Boyce’s Index and AUC suggested good performance in predicting species’ environmental suitability (AUC = 0.757; Boyce’s Index = 0.855, respectively). Temperature was the most important variable (88.6% of variable importance) followed by precipitation and roads density (8.7 and 2.7% of variable importance, respectively). Current suitable areas for *P. setaceum* mirrored occurrences’ distribution appearing prevalently below 800 m (Fig. 2). Model prediction based on the RCP2.6 scenario substantially confirmed as highly suitable areas, the current suitable ones, though a shift to higher values of habitat suitability can also be observed at higher elevation (Fig. 2b). An elevational



**Fig. 2** *Pennisetum setaceum* HSM predictions: **a** current climatic conditions, **b** RCP2.6 climatic conditions, **c** RCP8.5 climatic conditions

shift is emphasized by the model prediction based on the RCP8.5 scenario (Fig. 2c), where *P. setaceum* seems to climb up along elevation (especially in the northern part of the island on the Anaga mountains).

In warmer and drier climatic conditions, *P. setaceum* is likely to shift from lower to higher elevations, increasing the total area occupied by the species (Fig. 3). Indeed, estimates of habitat suitability indicate the preference of the species for climatic conditions with a mean spring temperature higher than 10 °C (Fig. 4).

## Discussion

We investigated here the present and future spread of the invasive *P. setaceum* under different climatic scenarios, highlighting the current suitability of areas below 800 m asl (mirroring its current distribution; Supplementary Material Fig. 5), and predicting an increase in habitat suitability at higher elevations (~ 1000 m asl) for both future scenarios considered. Specifically, the predicted increase resulted particularly pronounced for the RCP8.5 scenario. Our predictions are quite alarming, considering that 800 m asl represents the current upper distribution limit of *P. setaceum* in Tenerife (Hobi 2008). *P. setaceum* is one of the most invasive species in Tenerife and it is currently spreading around the Mediterranean basin (Devesa Alcaraz et al. 2006; Pasta et al. 2010; Saavedra and Alcántara 2017). Our work is therefore particularly valuable to halt its spread via an effective monitoring of these areas that are under high invasion risk in the future.

### Invaded habitats

*Pennisetum setaceum* spreads below 800 m asl is likely to be enhanced by anthropic disturbance (e.g. land-cover change and consequent habitat fragmentation), rather than global warming. However, due to the year-by-year increase in anthropic disturbance and temperature at higher elevations, its present distribution is still expanding above 800 m asl (see for instance Kalwij et al. 2015). It has also been observed that *P. setaceum* can reach up to 2000 m asl in other islands contexts such as Hawaii (Williams et al. 1995). Moreover, *P. setaceum* has been recently observed at altitudes of 1500 m in the Barranco Risco Liso, located

**Table 2** Results of the Maxent model selection via AICc

|    | Settings | Features | rm   | train.AUC | avg.test.<br>AUC | var.test.<br>AUC | avg.diff.<br>AUC | var.diff.<br>AUC | avg.test.<br>orMTP | var.test.<br>orMTP | avg.test.<br>or10pct | var.test.<br>or10pct | AICc   | delta.AICc | w.AIC | Parameters |
|----|----------|----------|------|-----------|------------------|------------------|------------------|------------------|--------------------|--------------------|----------------------|----------------------|--------|------------|-------|------------|
| 1  | L_0.5    | L        | 0.50 | 0.76      | 0.77             | 0.02             | 0.04             | 0.01             | 0.05               | 0.01               | 0.13                 | 0.05                 | 787.33 | 3.22       | 0.05  | 3.00       |
| 2  | LQ_0.5   | LQ       | 0.50 | 0.78      | 0.76             | 0.02             | 0.05             | 0.02             | 0.09               | 0.03               | 0.16                 | 0.07                 | 785.73 | 1.62       | 0.10  | 5.00       |
| 3  | LQP_0.5  | LQP      | 0.50 | 0.78      | 0.77             | 0.02             | 0.05             | 0.02             | 0.02               | 0.00               | 0.16                 | 0.04                 | 784.11 | 0.00       | 0.23  | 4.00       |
| 4  | LQPH_0.5 | LQPH     | 0.50 | 0.81      | 0.75             | 0.02             | 0.08             | 0.02             | 0.02               | 0.00               | 0.28                 | 0.02                 | 946.93 | 162.82     | 0.00  | 30.00      |
| 5  | L_1      | L        | 1.00 | 0.76      | 0.77             | 0.01             | 0.04             | 0.01             | 0.02               | 0.00               | 0.16                 | 0.05                 | 787.47 | 3.36       | 0.04  | 3.00       |
| 6  | LQ_1     | LQ       | 1.00 | 0.77      | 0.76             | 0.02             | 0.04             | 0.02             | 0.07               | 0.02               | 0.16                 | 0.07                 | 786.63 | 2.52       | 0.07  | 4.00       |
| 7  | LQP_1    | LQP      | 1.00 | 0.76      | 0.77             | 0.02             | 0.04             | 0.01             | 0.02               | 0.00               | 0.16                 | 0.05                 | 789.02 | 4.91       | 0.02  | 4.00       |
| 8  | LQPH_1   | LQPH     | 1.00 | 0.80      | 0.76             | 0.02             | 0.07             | 0.02             | 0.02               | 0.00               | 0.16                 | 0.02                 | 865.59 | 81.48      | 0.00  | 23.00      |
| 9  | L_1.5    | L        | 1.50 | 0.76      | 0.76             | 0.01             | 0.03             | 0.01             | 0.02               | 0.00               | 0.16                 | 0.05                 | 787.71 | 3.60       | 0.04  | 3.00       |
| 10 | LQ_1.5   | LQ       | 1.50 | 0.76      | 0.76             | 0.01             | 0.04             | 0.01             | 0.02               | 0.00               | 0.16                 | 0.05                 | 790.11 | 6.00       | 0.01  | 4.00       |
| 11 | LQP_1.5  | LQP      | 1.50 | 0.76      | 0.76             | 0.01             | 0.04             | 0.01             | 0.02               | 0.00               | 0.18                 | 0.04                 | 788.25 | 4.14       | 0.03  | 3.00       |
| 12 | LQPH_1.5 | LQPH     | 1.50 | 0.79      | 0.75             | 0.02             | 0.06             | 0.02             | 0.02               | 0.00               | 0.16                 | 0.02                 | 803.79 | 19.68      | 0.00  | 11.00      |
| 13 | L_2      | L        | 2.00 | 0.76      | 0.76             | 0.01             | 0.03             | 0.01             | 0.02               | 0.00               | 0.16                 | 0.05                 | 788.03 | 3.92       | 0.03  | 3.00       |
| 14 | LQ_2     | LQ       | 2.00 | 0.76      | 0.76             | 0.01             | 0.03             | 0.01             | 0.02               | 0.00               | 0.18                 | 0.04                 | 788.26 | 4.15       | 0.03  | 3.00       |
| 15 | LQP_2    | LQP      | 2.00 | 0.76      | 0.75             | 0.01             | 0.03             | 0.01             | 0.02               | 0.00               | 0.18                 | 0.04                 | 789.79 | 5.68       | 0.01  | 3.00       |
| 16 | LQPH_2   | LQPH     | 2.00 | 0.78      | 0.74             | 0.01             | 0.05             | 0.02             | 0.02               | 0.00               | 0.16                 | 0.02                 | 810.26 | 26.15      | 0.00  | 12.00      |
| 17 | L_2.5    | L        | 2.50 | 0.76      | 0.76             | 0.01             | 0.03             | 0.01             | 0.02               | 0.00               | 0.18                 | 0.04                 | 788.45 | 4.34       | 0.03  | 3.00       |
| 18 | LQ_2.5   | LQ       | 2.50 | 0.76      | 0.75             | 0.01             | 0.03             | 0.01             | 0.02               | 0.00               | 0.18                 | 0.04                 | 788.81 | 4.70       | 0.02  | 3.00       |
| 19 | LQP_2.5  | LQP      | 2.50 | 0.75      | 0.75             | 0.01             | 0.03             | 0.01             | 0.02               | 0.00               | 0.16                 | 0.02                 | 788.86 | 4.75       | 0.02  | 2.00       |
| 20 | LQPH_2.5 | LQPH     | 2.50 | 0.77      | 0.74             | 0.01             | 0.05             | 0.02             | 0.05               | 0.00               | 0.18                 | 0.02                 | 803.32 | 19.21      | 0.00  | 9.00       |
| 21 | L_3      | L        | 3.00 | 0.76      | 0.76             | 0.01             | 0.03             | 0.01             | 0.02               | 0.00               | 0.18                 | 0.04                 | 788.96 | 4.85       | 0.02  | 3.00       |
| 22 | LQ_3     | LQ       | 3.00 | 0.76      | 0.75             | 0.01             | 0.03             | 0.01             | 0.02               | 0.00               | 0.18                 | 0.04                 | 789.47 | 5.36       | 0.02  | 3.00       |
| 23 | LQP_3    | LQP      | 3.00 | 0.75      | 0.74             | 0.00             | 0.03             | 0.00             | 0.02               | 0.00               | 0.13                 | 0.01                 | 789.99 | 5.88       | 0.01  | 2.00       |
| 24 | LQPH_3   | LQPH     | 3.00 | 0.75      | 0.73             | 0.01             | 0.05             | 0.01             | 0.05               | 0.00               | 0.13                 | 0.01                 | 797.87 | 13.76      | 0.00  | 6.00       |
| 25 | L_3.5    | L        | 3.50 | 0.76      | 0.75             | 0.01             | 0.03             | 0.01             | 0.02               | 0.00               | 0.18                 | 0.04                 | 789.55 | 5.45       | 0.02  | 3.00       |
| 26 | LQ_3.5   | LQ       | 3.50 | 0.75      | 0.75             | 0.01             | 0.03             | 0.01             | 0.05               | 0.00               | 0.18                 | 0.04                 | 790.26 | 6.15       | 0.01  | 3.00       |
| 27 | LQP_3.5  | LQP      | 3.50 | 0.75      | 0.74             | 0.00             | 0.03             | 0.00             | 0.02               | 0.00               | 0.11                 | 0.01                 | 788.79 | 4.68       | 0.02  | 1.00       |
| 28 | LQPH_3.5 | LQPH     | 3.50 | 0.75      | 0.73             | 0.01             | 0.04             | 0.01             | 0.02               | 0.00               | 0.11                 | 0.01                 | 788.79 | 4.68       | 0.02  | 1.00       |
| 29 | L_4      | L        | 4.00 | 0.75      | 0.75             | 0.01             | 0.03             | 0.01             | 0.02               | 0.00               | 0.18                 | 0.04                 | 790.24 | 6.13       | 0.01  | 3.00       |
| 30 | LQ_4     | LQ       | 4.00 | 0.75      | 0.74             | 0.00             | 0.03             | 0.01             | 0.02               | 0.00               | 0.16                 | 0.02                 | 791.17 | 7.06       | 0.01  | 3.00       |
| 31 | LQP_4    | LQP      | 4.00 | 0.75      | 0.75             | 0.00             | 0.02             | 0.00             | 0.02               | 0.00               | 0.09                 | 0.01                 | 789.09 | 4.98       | 0.02  | 1.00       |
| 32 | LQPH_4   | LQPH     | 4.00 | 0.75      | 0.74             | 0.00             | 0.03             | 0.00             | 0.02               | 0.00               | 0.11                 | 0.01                 | 789.05 | 4.94       | 0.02  | 1.00       |
| 33 | L_4.5    | L        | 4.50 | 0.75      | 0.75             | 0.01             | 0.03             | 0.01             | 0.05               | 0.00               | 0.18                 | 0.04                 | 791.02 | 6.91       | 0.01  | 3.00       |
| 34 | LQ_4.5   | LQ       | 4.50 | 0.75      | 0.74             | 0.00             | 0.03             | 0.00             | 0.02               | 0.00               | 0.13                 | 0.01                 | 789.75 | 5.64       | 0.01  | 2.00       |
| 35 | LQP_4.5  | LQP      | 4.50 | 0.75      | 0.75             | 0.00             | 0.02             | 0.00             | 0.02               | 0.00               | 0.09                 | 0.01                 | 789.33 | 5.22       | 0.02  | 1.00       |
| 36 | LQPH_4.5 | LQPH     | 4.50 | 0.75      | 0.75             | 0.00             | 0.02             | 0.00             | 0.02               | 0.00               | 0.09                 | 0.01                 | 789.39 | 5.29       | 0.02  | 1.00       |


Table 2 continued

| Settings | Features | rm   | train.AUC | avg.test.<br>AUC | var.test.<br>AUC | avg.diff.<br>AUC | var.diff.<br>AUC | avg.test.<br>orMTP | var.test.<br>orMTP | avg.test.<br>or10pct | var.test.<br>or10pct | AICc   | delta.AICc | w.AIC | Parameters |
|----------|----------|------|-----------|------------------|------------------|------------------|------------------|--------------------|--------------------|----------------------|----------------------|--------|------------|-------|------------|
| 37       | L_5      | L    | 5.00      | 0.75             | 0.00             | 0.03             | 0.01             | 0.02               | 0.00               | 0.16                 | 0.02                 | 791.88 | 7.78       | 0.00  | 3.00       |
| 38       | LQ_5     | LQ   | 5.00      | 0.75             | 0.00             | 0.03             | 0.00             | 0.02               | 0.00               | 0.13                 | 0.01                 | 790.37 | 6.26       | 0.01  | 2.00       |
| 39       | LQP_5    | LQP  | 5.00      | 0.75             | 0.00             | 0.02             | 0.00             | 0.02               | 0.00               | 0.09                 | 0.01                 | 789.63 | 5.52       | 0.01  | 1.00       |
| 40       | LQPH_5   | LQPH | 5.00      | 0.75             | 0.00             | 0.02             | 0.00             | 0.02               | 0.00               | 0.09                 | 0.01                 | 789.63 | 5.52       | 0.01  | 1.00       |

RM beta regularization multipliers; AUC Area Under the Curve; MTP the minimum training presence

within the Caldera de Taburiente in La Palma island (Walentowitz et al. 2019). Around Tenerife island, *P. setaceum* has rapidly colonized mainly areas that were already experiencing anthropic disturbance, such as agricultural and rural areas, spreading at higher altitudinal belts using the road network as the main propagation pathway like other IAS (Arteaga et al. 2009; Arévalo et al. 2010; Benedetti and Morelli 2017; Follak et al. 2018). In fact, in the area near Arafo, which has a long land-use history for agriculture, the species has climbed up approximately 100 m in elevation in three years (from 655 to 740 m asl, personal observation of DDR and ZNP in April 2019). *P. setaceum* established firstly along the roadside, and then it spread deeper into the surrounding landscapes, confirming that roads remain a crucial dispersion pathway for this species and could promote its spread into protected areas (Foxcroft et al. 2019; Walentowitz et al. 2019). However, its ability to compete and outstand native plant communities could be questioned in areas characterized by a high degree of biotic resistance and where the native community has sufficient propagule pressure for new colonization (Schuster et al. 2018).

*Pennisetum setaceum* also spread into many ravines within the lowlands in the South of Tenerife as well as into the lower parts of the National Park Caldera de Taburiente (Barranco de las Angustias), settling in semi natural communities (personal observations of RO). In ravines, called *barrancos* in Tenerife, temporary water flow during the rainy season facilitates the spread of seeds downriver, improving also the water supply in semiarid areas. However, in the southernmost part of Tenerife, *P. setaceum* probably is still limited by soil water conditions and at the highest sites of its distribution (currently at about 1000 m asl) low temperature might limit its growth. A still unanswered question is if *P. setaceum* will be able to grow and spread in open Canary pine forests growing above 800 to 1000 m asl, which might have serious consequences for natural regeneration and fire dynamics. However, according to Walentowitz et al. (2019), the dense canopy of forest ecosystems should prevent *P. setaceum* spread due to its heliophilous nature.




**Fig. 3** Relationships between *P. setaceum* HSM predictions and elevation: blue dots and line, current climatic conditions; orange dots and line, RCP2.6 climatic conditions; red dots and line, RCP8.5 climatic conditions

### Limiting factors, functional traits, and competitive advantages

Generally, rainfall and temperature patterns are among the most important limiting factors determining plant distribution and abundance (Stephenson 1990), affecting reproduction and other life-history traits. Temperature is a strong limiting factor for the species, which usually prefers areas with temperature higher than 10 °C (Fig. 4). Under the current conditions, these areas are located under 1000 m asl, but they are expected to rise under both the climate change scenarios analysed (RCP2.6 and RCP8.5). The geographic distribution of the species in both scenarios showed an elevation shift, particularly pronounced in the RCP8.5 scenario. The species could move to higher elevation probably benefitting from the effect of orographic precipitations, and it could migrate in the north-western part of the island, where the trade winds enhance water availability (Figs. 2, 3). The ecophysiological traits of *P. setaceum* support its large

size, extensive canopy, shorter leaf senescence period, and the capacity to buffer seasonal and local water shortages (González-Rodríguez et al. 2010). Indeed, they confer considerable competitive advantages to the invader, and this can partially explain its invasion success in the Canary archipelago, considering local environmental and climatic conditions. On the other hand, seedlings' growth could be a limiting factor of *P. setaceum* invasion success: despite their relatively broad germination requirements, seedling cannot cope with low soil moisture or drought conditions (Adkins et al. 2011; Goergen and Daehler 2002), and it has been suggested that they do not tolerate more than one month of drought (Rahlao et al. 2010). Interestingly, once the seedlings are established, the amount of water does not seem to affect their performance, even under higher temperature regimes. When favourable conditions occur, *P. setaceum* can quickly thrive (already under sub-optimal ecological conditions) and overcome native coexisting species through rapid exploitation of available resources (such as water and



**Fig. 4** Relationships between *P. setaceum* HSM predictions and temperatures: blue dots, current climatic conditions; orange dots, RCP2.6 climatic conditions; red dots, RCP8.5 climatic conditions

nutrients), thanks to its broad ecological tolerance and high phenotypic plasticity (Williams and Black 1993; Poulin et al. 2007; Rahlao et al. 2010). Furthermore, under drought or limited resources availability, the alteration of biomass allocation patterns in favour of roots seems to be a conservative strategy (Williams et al. 1995; Rahlao et al. 2010). This may explain why *P. setaceum* is an excellent invader of habitats characterized by fluctuating resources, frequently disturbed or with an irregular rainfall pattern, in agreement with Davis' theory of community invasibility (Davis et al. 2000). Compared to native species, the higher nitrogen and water use efficiency (González-Rodríguez et al. 2010) allow *P. setaceum* to compete effectively in warmer, drier areas, conferring it a considerable competitive advantage (Tordoni et al. 2020). Recently, Rodríguez-Caballero et al. (2017, 2018) have shown how *P. setaceum* could establish symbiosis with different mycorrhizal fungi and act as a transformer (Pyšek et al. 2004) modifying the soil bacterial community related to N cycle. As a consequence, it also tends to increase the risk of

intense wildfires (to which it is well adapted), thus posing a further threat to resident native communities (Tunison 1992).

Other studies on *P. setaceum* ecophysiology have confirmed the broad ecological tolerance of the species in other environments. Specifically, Jacobi and Warshauer (1992) reported that in the Hawaiian Islands, *P. setaceum* has a relatively wide elevation range (500–2900 m) but is limited to areas with median annual rainfall less than 1250 mm, hence only dry and mesic habitats are colonized by this species. In southern California, Sweet and Holt (2015) reported that *P. setaceum* might benefit from dampened temperatures and additional moisture through fog drip, becoming competitive in the cooler season due to phenotypic plasticity in response to temperature and water. Finally, since *P. setaceum* is endowed of C4 metabolism, it is likely that climate change may have positive effects on its performance, further promoting its invasion success, as already observed for other C4 invaders (e.g. Chuine et al. 2012 for *Setaria parviflora*). Indeed, C4 plants are intrinsically adapted to

elevated temperatures and to tolerate heat stress (Sage and Kubien 2003) thanks to higher effective quantum yield of CO<sub>2</sub> fixation, which increases photosynthetic rates, especially at higher temperatures (Ehleringer and Björkman 1977; Sage and Kubien 2003).

For the sake of clarity, we would like to highlight that a possible drawback of our predictions is the violation of the equilibrium assumption. In fact, it was not possible to consider the whole climatic range of the species and train the model including also occurrences present in native range: only 52 *P. setaceum* occurrences are available in an area covering mostly the East Africa and the Arabian peninsula (Sudan, Ethiopia, Eritrea, Kenya, Somalia, Saudi Arabia, Yemen and Oman, GBIF accessed on 26th May 2020). Considering, however, the invasion pressure that the delicate ecosystem of Tenerife is undergoing, predicting the areas that are more prone to invasion in the near future is more valuable (from a conservation and management perspective and under the precautionary principle; West et al. 2016) than modelling the full potential invasive range of *P. setaceum* at equilibrium. Nonetheless, considering the possible violation of this assumption, care must be taken when interpreting the model outputs.

## Conclusions

Our model predictions highlight that the distribution of *P. setaceum* in Tenerife will be strongly affected by climate change and particularly by the predicted increase in aridity on the island. Our model (particularly RCP8.5) foresees that the species would move to higher elevation and to the north-western part of the island, likely to benefit both from orographic precipitations and enhanced water availability promoted by trade winds. Moreover, under the current socio-economic perspectives, the RCP2.6 scenario is becoming year after year less feasible (Davis and Socolow 2014), making the RCP8.5 one even more likely. The species may consequently have negative impacts on entire ecosystems, due to its strong competitive capacities and being a driver of other invasive species. Particularly, the predicted invasion of areas at higher elevation poses the question of its ability to penetrate the pine forests (especially the more xeric ones in the southern part of the island). Its establishment could drastically affect the dynamics of this unique

ecosystem (e.g. natural regeneration, fire regime), if proper conservation plans will not be developed to support ecosystem health and diversity. However, even though our predictions are coherent with the auto-ecology of the species, climate change predictions are extremely simplified and should be therefore interpreted with care. More powerful outcomes may be obtained downscaling climatic data from global models such as Global Circulation Model (GCM), but to our knowledge spatially interpolated fine scale GCM projections for the Canary Islands are still lacking. Moreover, it has been showed that whereas global scale products are generally applicable at broad geographical scale, modelling climate surface on islands needs to consider fine scale spatial variation due to spatial variability and steep climatic gradient variation (Garzón-Machado et al. 2014; Khalyani et al. 2016). In a global warming scenario, a modelling approach as the one proposed here could play a key role in alien species monitoring, highlighting the portions of the territory that are more prone to biological invasions.

To our knowledge, Walentowitz et al. (2019) and ours results are the first *P. setaceum* niche model outputs presented. Such cartographic products provide an immediate tool for understanding potential changes in species distributions, and they can be used to disseminate and increase the awareness of biological invasion outside the academia. Considering the dramatic effects that an uncontrolled increase of *P. setaceum* may have on Tenerife, understanding its future potential habitat distribution is therefore crucial from a conservational perspective. Immediate control measures (e.g. manual and chemical species removal efforts) should be carried out from the border of *P. setaceum* current distribution, focusing especially along invasion pathways (Walentowitz et al. 2019). Furthermore, prohibiting the commercial use of *P. setaceum* propagules could be another tool to control the species further spread. In fact, Potgieter et al. (2019) showed that ornamental plants are still perceived as having aesthetic benefits when confined to private gardens; however, shifts in perceptions may occur when they become widespread in the wild, leading to economic and environmental costs. Finally, specific Citizen Science projects have been shown to be effective to raise the awareness about biological invasions and to engage both public administrations and citizens in monitoring and controlling invasive

species and therefore could be a valuable tool to help controlling the species invasion and its deleterious effect on Canary Islands (e.g. Sladonja and Poljuha 2018).

**Acknowledgements** We would like to thank the Teno Rural Park for having shared its IAS database. We are grateful to the anonymous reviewers for the constructive and positive suggestions. D.D.R. was supported by an ERASMUS+ 2015-2016 grant provided by the European Commission.

## References

- Adkins E, Cordell S, Drake DR (2011) Role of fire in the germination ecology of fountain grass (*Pennisetum setaceum*), an invasive african bunchgrass in Hawai'i. *Pac Sci* 65(1):17–26
- Aiello-Lammens ME, Boria RA, Radosavljevic A, Vilela B, Anderson RP (2015) sptin: an R package for spatial thinning of species occurrence records for use in ecological Niche models. *Ecography* 38(5):541–545
- Arechavaleta M, Rodríguez S, Zurita N, García A (2010) Lista de especies silvestres de canarias. Hongos, plantas y animales terrestres. Gobierno de Canarias, Tenerife
- Arévalo J, Otto R, Escudero C, Fernández-Lugo S, Arteaga M, Delgado J, Fernández-Palacios J (2010) Do anthropogenic corridors homogenize plant communities at a local scale? A case studied in Tenerife (Canary Islands). *Plant Ecol* 209(1):23–35
- Arteaga MA, Delgado JD, Otto R, Fernández-Palacios JM, Arévalo JR (2009) How do alien plants distribute along roads on oceanic islands? A case study in Tenerife, Canary Islands. *Biol Invasions* 11(4):1071–1086
- Bacaro G, Maccherini S, Chiarucci A, Jentsch A, Rocchini D, Torri D, Gioria M, Tordoni E, Martellos S, Altobelli A et al (2015) Distributional patterns of endemic, native and alien species along a roadside elevation gradient in Tenerife, Canary Islands. *Commun Ecol* 16(2):223–234
- Barbet-Massin M, Rome Q, Villemant C, Courchamp F (2018) Can species distribution models really predict the expansion of invasive species? *PLoS ONE* 13:3. <https://doi.org/10.1371/journal.pone.0193085>
- Barni E, Bacaro G, Falzoi S, Spanna F, Siniscalco C (2012) Establishing climatic constraints shaping the distribution of alien plant species along the elevation gradient in the Alps. *Plant Ecol* 213(5):757–767
- Bella S, D'Urso V et al (2012) First record in the Mediterranean basin of the alien leafhopper *Balclutha Brevis* living on invasive *Pennisetum setaceum*. *Bull Insectol* 65(2):195–198
- Bellard C, Bertelsmeier C, Leadley P, Thuiller W, Courchamp F (2012) Impacts of climate change on the future of biodiversity. *Ecol Lett* 15(4):365–377
- Bellard C, Leclerc C, Courchamp F (2014) Impact of sea level rise on the 10 insular biodiversity hotspots. *Glob Ecol Biogeogr* 23(2):203–212
- Benedetti Y, Morelli F (2017) Spatial mismatch analysis among hotspots of alien plant species, road and railway networks in Germany and Austria. *PLoS ONE* 12(8):e0183691
- Brundu G, Richardson DM (2016) Planted forests and invasive alien trees in Europe: a code for managing existing and future plantings to mitigate the risk of negative impacts from invasions
- Chuine I, Morin X, Sonié L, Collin C, Fabreguettes J, Degueldre D, Salager J-L, Roy J (2012) Climate change might increase the invasion potential of the alien C4 grass *Setaria parviflora* (Poaceae) in the Mediterranean Basin. *Divers Distrib* 18(7):661–672
- Cordell S, Sandquist D (2008) The impact of an invasive African bunchgrass (*Pennisetum setaceum*) on water availability and productivity of canopy trees within a tropical dry forest in Hawaii. *Funct Ecol* 22(6):1008–1017
- Courchamp F, Chapuis J-L, Pascal M (2003) Mammal invaders on islands: impact, control and control impact. *Biol Rev* 78(3):347–383
- Cropper T (2013) The weather and climate of Macaronesia: past, present and future. *Weather* 68(11):300–307
- Cropper TE, Hanna E (2014) An analysis of the climate of Macaronesia, 1865–2012. *Int J Climatol* 34(3):604–622
- Da Re D, Tordoni E, Pérez ZN, Fernández-Palacios JM, Arévalo JR, Otto R, Rocchini D, Bacaro G (2019) A spatially-explicit model of alien plant richness in Tenerife (Canary Islands). *Ecol Complex* 38:75–82
- D'Antonio CM, Vitousek PM (1992) Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annu Rev Ecol Syst* 23(1):63–87
- Davis SJ, Socolow RH (2014) Commitment accounting of CO<sub>2</sub> emissions. *Environ Res Lett* 9(8):084018
- Davis MA, Grime JP, Thompson K (2000) Fluctuating resources in plant communities: a general theory of invasibility. *J Ecol* 88(3):528–534
- de Paz PLP, Gallo AG, Heene A (1999) Control y erradicación del “Rabo-Gato” (*Pennisetum setaceum*) en la Isla de Palma. Universidad de la Laguna, San Cristóbal de La Laguna
- Delgado JD, Arévalo JR, Fernández-Palacios J (2004) Consecuencias de la fragmentación viaria: efectos de borde de las carreteras en la laurisilva y el pinar de Tenerife. *Ecología Insular/Island Ecology*. Asociación Española de Ecología Terrestre (AEET)-Cabildo Insular de la Palma, pp 181–225
- Devesa Alcaraz JA, Arnelas I et al (2006) *Pennisetum setaceum* (forssk.) Chiov. (Poaceae), nueva localidad para la flora ibérica. *Acta Bot Malac* 31:190–191
- Di Cola V, Broennimann O, Petitpierre B, Breiner FT, Damen M, Randin C, Engler R, Pottier J, Pio D, Dubuis A et al (2017) ecospat: an r package to support spatial analyses and modeling of species niches and distributions. *Ecography* 40(6):774–787
- Dukes JS, Mooney HA (1999) Does global change increase the success of biological invaders? *Trends Ecol Evol* 14(4):135–139

- Dyderski MK, Paž S, Frelich LE, Jagodziński AM (2018) How much does climate change threaten European forest tree species distributions? *Glob Change Biol* 24(3):1150–1163
- Early R, Sax DF (2014) Climatic niche shifts between species' native and naturalized ranges raise concern for ecological forecasts during invasions and climate change. *Glob Ecol Biogeogr* 23(12):1356–1365
- Ehleringer J, Björkman O (1977) m yields for CO<sub>2</sub> uptake in C3 and C4 plants: dependence on temperature, CO<sub>2</sub>, and O<sub>2</sub> concentration. *Plant Physiol* 59(1):86–90
- Elith J, Phillips SJ, Hastie T, Dudík M, Chee YE, Yates CJ (2011) A statistical explanation of maxent for ecologists. *Divers Distrib* 17(1):43–57
- Fernández-Palacios JM, Rijdsdijk KF, Norder SJ, Otto R, de Nascimento L, Fernández-Lugo S, Tjørve E, Whittaker RJ (2016) Towards a glacial-sensitive model of island biogeography. *Glob Ecol Biogeogr* 25(7):817–830
- Follak S, Eberius M, Essl F, Fördös A, Sedlacek N, Trognitz F (2018) Invasive alien plants along roadsides in Europe. *EPPO Bull* 48(2):256–265
- Foxcroft LC, Spear D, van Wilgen NJ, McGeoch MA (2019) Assessing the association between pathways of alien plant invaders and their impacts in protected areas. *NeoBiota* 43:1
- Francisco-Ortega J, Santos-Guerra A, Bacallado JJ (2009) Canary Islands, biology. In: Gillespie R (ed) *Encyclopedia of islands*. California Press, Berkeley, pp 127–133
- Gallien L, Douzet R, Pratte S, Zimmermann NE, Thuiller W (2012) Invasive species distribution models-how violating the equilibrium assumption can create new insights. *Glob Ecol Biogeogr* 21(11):1126–1136
- García-Herrera R, Gallego D, Hernández E, Gimeno L, Ribera P, Calvo N (2003) Precipitation trends in the Canary Islands. *Int J Climatol* 23(2):235–241
- Garzón-Machado V, Otto R, del Arco Aguilar MJ (2014) Bioclimatic and vegetation mapping of a topographically complex oceanic island applying different interpolation techniques. *Int J Biometeorol* 58(5):887–899
- Giorgi F (2006) Climate change hot-spots. *Geophys Res Lett* 33(8):101029
- Gobierno de Canarias (2019). Biodiversity Data Bank of the Canary Islands (<https://www.biodiversidadcanarias.es/biota>) [Date of consultation: 15/04/2019]
- Goergen E, Daehler CC (2002) Factors affecting seedling recruitment in an invasive grass (*Pennisetum setaceum*) and a native grass (*Heteropogon contortus*) in the Hawaiian Islands. *Plant Ecol* 161(2):147–156
- González-Rodríguez AM, Baruch Z, Palomo D, Cruz-Trujillo G, Jiménez MS, Morales D (2010) Ecophysiology of the invader *Pennisetum setaceum* and three native grasses in the Canary Islands. *Acta Oecologica* 36(2):248–254
- Gremmen N, Chown S, Marshall D (1998) Impact of the introduced grass *Agrostis stolonifera* on vegetation and soil fauna communities at Marion Island, sub-Antarctic. *Biol Conserv* 85(3):223–231
- Guevara L, Gerstner BE, Kass JM, Anderson RP (2018) Toward ecologically realistic predictions of species distributions: a cross-time example from tropical montane cloud forests. *Glob Change Biol* 24(4):1511–1522
- Guisan A, Thuiller W (2005) Predicting species distribution: offering more than simple habitat models. *Ecol Lett* 8(9):993–1009
- Guisan A, Thuiller W, Zimmermann NE (2017) *Habitat suitability and distribution models: with applications in R*. Cambridge University Press, Cambridge
- Hansen A et al (1970) Contributions to the flora of the Canary Islands (especially Tenerife). *Cuad Bot Canar* 9:37–59
- Hernandez RR, Sandquist DR (2011) Disturbance of biological soil crust increases emergence of exotic vascular plants in California sage scrub. *Plant Ecol* 212(10):1709
- Hijmans RJ, Phillips S, Leathwick J, Elith J, Hijmans MRJ (2017) Package 'dismo'. *Circles* 9(1):1–68
- Hirzel AH, Le Lay G, Helfer V, Randin C, Guisan A (2006) Evaluating the ability of habitat suitability models to predict species presences. *Ecol Model* 199(2):142–152
- Hobi S (2008) Analyse der Faktoren Klima und Störung als Höhenlimite des Neophyten *Pennisetum setaceum* auf Teneriffa. Semesterarbeit. Lehrstuhl für Landschaftsökologie, Department für Ökologie und Ökosystemmanagement, Technische Universität München
- Hulme PE (2009) *Handbook of alien species in Europe*, vol 569. Springer, New York
- Jacobi JD, Warshauer FR (1992) Distribution of six alien plant species in upland habitats on the island of Hawaii. Alien plant invasions in native ecosystems of Hawaii. Cooperative National Park Resources Studies Unit, University of Hawaii, Honolulu, pp 155–188
- Jiménez-Valverde A, Peterson AT, Soberón J, Overton J, Aragón P, Lobo JM (2011) Use of niche models in invasive species risk assessments. *Biol Invasions* 13(12):2785–2797
- Kalwij JM, Robertson MP, van Rensburg BJ (2015) Annual monitoring reveals rapid upward movement of exotic plants in a montane ecosystem. *Biol Invasions* 17(12):3517–3529
- Kearney M (2006) Habitat, environment and niche: what are we modelling? *Oikos* 115(1):186–191
- Kenis M, Auger-Rozenberg M-A, Roques A, Timms L, Péré C, Cock MJ, Settele J, Augustin S, Lopez-Vaamonde C (2009) Ecological effects of invasive alien insects. *Biol Invasions* 11(1):21–45
- Khalyani AH, Gould WA, Harmsen E, Terando A, Quinones M, Collazo JA (2016) Climate change implications for tropical islands: interpolating and interpreting statistically down-scaled gcm projections for management and planning. *J Appl Meteorol Climatol* 55(2):265–282
- Kleinbauer I, Dullinger S, Peterseil J, Essl F (2010) Climate change might drive the invasive tree *Robinia pseudacacia* into nature reserves and endangered habitats. *Biol Conserv* 143(2):382–390
- Kueffer C, Daehler CC, Torres-Santana CW, Laverne C, Meyer J-Y, Otto R, Silva L (2010) A global comparison of plant invasions on oceanic islands. *Perspect Plant Ecol Evol Syst* 12(2):145–161
- Luque A, Martín JL, Dorta P, Mayer P et al (2014) Temperature trends on Gran Canaria (Canary Islands), an example of global warming over the subtropical Northeastern Atlantic. *Atmos Clim Sci* 4(1):20–28
- Martín Esquivel J, García H, Redondo C, García I, Carralero I (1995) La red canaria de espacios naturales protegidos. Gobierno de Canarias, Viceconsejería de Medio Ambiente, Santa Cruz
- Martín JL, Bethencourt J, Cuevas-Agulló E (2012) Assessment of global warming on the island of Tenerife, Canary Islands

- (Spain), trends in minimum, maximum and mean temperatures since 1944. *Clim Change* 114(2):343–355
- Matías L, Jump AS (2013) Impacts of predicted climate change on recruitment at the geographical limits of Scots pine. *J Exp Bot* 65(1):299–310
- Medley KA (2010) Niche shifts during the global invasion of the Asian tiger mosquito, *Aedes albopictus* skuse (Culicidae), revealed by reciprocal distribution models. *Glob Ecol Biogeogr* 19(1):122–133
- Merow C, Smith MJ, Silander JA (2013) A practical guide to MaxEnt for modeling species' distributions: what it does, and why inputs and settings matter. *Ecography* 36:1058–1069
- Millennium Ecosystem Assessment (2005) Ecosystems and human well-being: synthesis. Millennium Ecosystem Assessment, Washington, DC
- Morales CL, Aizen MA (2002) Does invasion of exotic plants promote invasion of exotic flower visitors? A case study from the temperate forests of the southern andes. *Biol Invasions* 4(1):87–100
- Muscarella R, Galante PJ, Soley-Guardia M, Boria RA, Kass JM, Uriarte M, Anderson RP (2014) Enm eval: an R package for conducting spatially independent evaluations and estimating optimal model complexity for maxent ecological niche models. *Methods Ecol Evol* 5(11):1198–1205
- Myers DE (1984) Co-kriging—new developments. Springer, Dordrecht, pp 295–305
- Pasta S, Badalamenti E, Mantia TL (2010) Tempi e modi di un'invasione incontrastata: *Pennisetum setaceum* (Forssk.) Chiov. (Poaceae) in sicilia. *Naturalista Sicil* 34:487–525
- Pauchard A, Alaback PB (2004) Influence of elevation, land use, and landscape context on patterns of alien plant invasions along roadsides in protected areas of South-Central Chile. *Conserv Biol* 18(1):238–248
- Pereira HM, Leadley PW, Proença V, Alkemade R, Scharlemann JPW, Fernández-Manjarrés JF, Araújo MB, Balvanera P, Biggs R, Cheung WWL, Chini LP, Cooper HA, Gilman EL, Guénette S, Hurtt GC, Huntington HP, Mace GM, Oberdorff T, Revenga C, Rodrigues PP, Scholes RJ, Sumaila UR, Walpole M (2010) Scenarios for global biodiversity in the 21st century. *Science* 330(6010):1496–501
- Peterson AT, Soberón J, Pearson RG, Anderson RP, Martínez-Meyer E, Nakamura M, Araújo MB (2011) Ecological niches and geographic distributions (MPB-49), vol 56. Princeton University Press, Princeton
- Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modeling of species geographic distributions. *Ecol Model* 190(3–4):231–259
- Phillips SJ, Dudík M, Elith J, Graham CH, Lehmann A, Leathwick J, Ferrier S (2009) Sample selection bias and presence-only distribution models: implications for background and pseudo-absence data. *Ecol Appl* 19(1):181–197
- Phillips SJ, Anderson RP, Dudík M, Schapire RE, Blair ME (2017) Opening the black box: an open-source release of maxent. *Ecography* 40(7):887–893
- Potgieter LJ, Gaertner M, O'Farrell PJ, Richardson DM (2019) Perceptions of impact: invasive alien plants in the urban environment. *J Environ Manag* 229:76–87
- Poulin J, Sakai AK, Weller SG, Nguyen T (2007) Phenotypic plasticity, precipitation, and invasiveness in the fire-promoting grass *Pennisetum setaceum* (Poaceae). *Am J Bot* 94(4):533–541
- Pyšek P, Richardson DM, Rejmánek M, Webster GL, Williamson M, Kirschner J (2004) Alien plants in checklists and floras: towards better communication between taxonomists and ecologists, vol 53. Wiley Online Library, Hoboken, pp 131–143
- Pyšek P, Jarošík V, Hulme PE, Kühn I, Wild J, Arianoutsou M, Bacher S, Chiron F, Didžiulis V, Essl F et al (2010) Disentangling the role of environmental and human pressures on biological invasions across europe. *Proc Nat Acad Sci USA* 107(27):12157–12162
- Pyšek P, Pergl J, Essl F, Lenzner B, Dawson W, Kreft H, Weigelt P, Winter M, Kartesz J, Nishino M, Antonova LA, Barcelona JF, Cabelaz FJ, Cárdenas D, Cárdenas-Toro J, Castaño N, Chacón EL, Chatelain C, Dullinger S, Ebel AL, Figueiredo E, Fuentes NJ, Genovesi P, Groom QJ, Henderson LM, Inderjit Kupriyanov A, Masciadri S, Maurel N, Meerman JC, Morozova OV, Moser D, Nickrent DL, Nowak PM, Pagad S, Patzelt A, Pelser PB, Seebens H, Shu W, Thomas JJ, Velayos M, Weber E, Wieringa JJ, Baptiste MP, van Kleunen M (2017) Naturalized alien flora of the world: species diversity, taxonomic and phylogenetic patterns, geographic distribution and global hotspots of plant invasion. *Czech Botl Soc* 89:203–274
- Qian H, Ricklefs RE (2006) The role of exotic species in homogenizing the North American flora. *Ecol Lett* 9(12):1293–8
- Rahla SJ, Esler KJ, Milton SJ, Barnard P (2010) Nutrient addition and moisture promote the invasiveness of crimson fountaingrass (*Pennisetum setaceum*). *Weed Sci* 58(2):154–159
- Ribeiro PJ Jr, Diggle PJ et al (2001) geor: a package for geostatistical analysis. *R news* 1(2):14–18
- Roberts DR, Bahn V, Ciuti S, Boyce MS, Elith J, Guillera-Arroita G, Hauenstein S, Lahoz-Monfort JJ, Schröder B, Thuiller W et al (2017) Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. *Ecography* 40(8):913–929
- Rodríguez-Caballero G, Caravaca F, del Mar Alguacil M, Fernández-López M, Fernández-González AJ, Roldán A (2017) Striking alterations in the soil bacterial community structure and functioning of the biological N cycle induced by *Pennisetum setaceum* invasion in a semiarid environment. *Soil Biol Biochem* 109:176–187
- Rodríguez-Caballero G, Caravaca F, Roldán A (2018) The unspecificity of the relationships between the invasive *Pennisetum setaceum* and mycorrhizal fungi may provide advantages during its establishment at semiarid mediterranean sites. *Sci Total Environ* 630:1464–1471
- Saavedra M, Alcántara C (2017) *Pennisetum setaceum*, planta invasora en expansión. In: Mercedes Royuela Hernando y Ana Zabalza Aznárez (editoras): XVI Congreso de la Sociedad Española de Malherbología: actas. Pamplona-Iruña, 25-27 octubre, 2017. Universidad Pública de Navarra Nafarroako Unibertsitate Publikoa, 2017. Universidad Pública de Navarra/Nafarroako Unibertsitate Publikoa
- Sage RF, Kubien DS (2003) Quo vadis C4? An ecophysiological perspective on global change and the future of C4 plants. *Photosynth Res* 77(2–3):209–225

- Sánchez-Benítez A, García-Herrera R, Vicente-Serrano SM (2017) Revisiting precipitation variability, trends and drivers in the Canary Islands. *Int J Climatol* 37(9):3565–3576
- Scalera R, Genovesi P, Essl F, Rabitsch W (2012) The impacts of invasive alien species in Europe. *Eur Environ Agency Tech Rep* 16:114
- Schuster MJ, Wragg PD, Reich PB (2018) Using revegetation to suppress invasive plants in grasslands and forests. *J Appl Ecol* 55(5):2362–2373
- Sladonja B, Poljuha D (2018) Citizen science as a tool in biological recording: a case study of *Ailanthus altissima* (Mill.) Swingle. *Forests* 9(1):31
- Somot S, Sevault F, Déqué M, Crépon M (2008) 21st century climate change scenario for the Mediterranean using a coupled atmosphere-ocean regional climate model. *Glob Planet Change* 63(2–3):112–126
- Steinbauer MJ, Irl SDH, González-Mancebo JM, Breiner FT, Hernández-Hernández R, Hopfenmüller S, Kidane YM, Jentsch A, Beierkuhnlein C (2017) Plant invasion and speciation along elevational gradients on the oceanic island La Palma, Canary Islands. *Ecol Evol* 7(2):771–779
- Steiner FM, Schlick-Steiner BC, VanDerWal J, Reuther KD, Christian E, Stauffer C, Suarez AV, Williams SE, Crozier RH (2008) Combined modelling of distribution and niche in invasion biology: a case study of two invasive tetramorium ant species. *Divers Distrib* 14(2008):538–545
- Stephenson NL (1990) Climatic control of vegetation distribution: the role of the water balance. *Am Nat* 135(5):649–670
- Stocker T (2014) Climate change 2013: the physical science basis: working group I contribution to the fifth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge
- Sweet LC, Holt JS (2015) Establishment stage competition between exotic *Crimson fountaingrass* (*Pennisetum setaceum*, C4) and native Purple Needlegrass (*Stipa pulchra*, C3). *Invasive Plant Sci Manag* 8(2):139–150
- Syfert MM, Smith MJ, Coomes DA (2013) The effects of sampling bias and model complexity on the predictive performance of maxent species distribution models. *PLoS ONE* 8(2):e55158
- Thuiller W (2007) Biodiversity: climate change and the ecologist. *Nature* 448:550–552
- Toby Kiers E, Palmer TM, Ives AR, Bruno JF, Bronstein JL (2010) Mutualisms in a changing world: an evolutionary perspective. *Ecol Lett* 13(12):1459–1474
- Tordoni E, Napolitano R, Nimis P, Castello M, Altobelli A, Da Re D, Zago S, Chines A, Martellos S, Maccherini S, Bacaro G (2017) Diversity patterns of alien and native plant species in trieste port area: exploring the role of urban habitats in biodiversity conservation. *Urban Ecosyst* 20(5):1151–1160
- Tordoni E, Petruzzellis F, Nardini A, Savi T, Bacaro G (2019) Make it simpler: alien species decrease functional diversity of coastal plant communities. *J Veg Sci* 30(3):498–509
- Tordoni E, Petruzzellis F, Nardini A, Bacaro G (2020) Functional divergence drives invasibility of plant communities at the edges of a resource availability gradient. *Diversity* 12(4):148
- Tunison JT (1992) Fountain grass control in Hawaii Volcanoes National Park: management considerations and strategies. Alien plant invasions in native ecosystems of Hawaii: management and research. Cooperative National Parks Resources Studies Unitn, University of Hawai'i at Manoa, Honolulu, pp 376–393
- Vilà M, López-Darias M (2006) Contrasting biogeography of endemic and alien terrestrial species in the canary islands. *Orsis organismes i sistemes* 21:91–101
- Vilà M, Espinar JAL, Hejda M, Hulme PE, Jarošík V, Maron JL, Pergl J, Schaffner U, Sun Y, Pyšek P (2011) Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecol Lett* 14(7):702–8
- Walentowitz AJ, Irl SD, Acevedo Rodríguez AJ, Palomares-Martínez Á, Vetter V, Zennaro B, Medina FM, Beierkuhnlein C (2019) Graminoid invasion in an insular endemism hotspot and its protected areas. *Diversity* 11(10):192
- Warren DL, Seifert SN (2011) Ecological niche modeling in maxent: the importance of model complexity and the performance of model selection criteria. *Ecol Appl* 21(2):335–342
- West AM, Kumar S, Brown CS, Stohlgren TJ, Bromberg J (2016) Field validation of an invasive species maxent model. *Ecol Inform* 36:126–134
- Whittaker RJ, Fernández-Palacios JM (2007) Island biogeography: ecology, evolution, and conservation. Oxford University Press, Oxford
- Williams D, Black R (1993) Phenotypic variation in contrasting temperature environments: growth and photosynthesis in *Pennisetum setaceum* from different altitudes on Hawaii. *Funct Ecol* 7:623–633
- Williams DG, Mack RN, Black RA (1995) Ecophysiology of introduced *Pennisetum setaceum* on Hawaii: the role of phenotypic plasticity. *Ecology* 76(5):1569–1580
- Wilson AM, Silander JA (2014) Estimating uncertainty in daily weather interpolations: a Bayesian framework for developing climate surfaces. *Int J Climatol* 34(8):2573–2584

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.