



RESEARCH ARTICLE OPEN ACCESS

Environmental and Geographic Conditions on the Breeding Grounds Drive Bergmannian Clines in Nightjars

Aaron A. Skinner^{1,2} | Alicia M. Korpach^{3,4,5} | Susanne Åkesson⁶ | Marja H. Bakermans⁷ | Erin M. Bayne⁸ | Thomas J. Benson⁹ | Giovanni Boano¹⁰ | R. Mark Brigham³ | Simon S. Christiansen¹¹ | Greg J. Conway¹² | Christina M. Davy¹³ | Ruben Evens^{14,15} | Kevin C. Fraser² | Autumn-Lynn Harrison¹⁶ | Anders Hedenström⁴ | Ian G. Henderson¹² | Juha Honkala¹⁷ | Lars B. Jacobsen¹⁸ | Michiel Lathouwers^{19,20} | Peter P. Marra²¹ | Janet W. Ng²² | Gabriel Norevik²³ | Amy L. Scarpignato²⁴ | Kasper Thorup²⁵ | Christopher M. Tonra² | Steven L. Van Wilgenburg²⁶ | Andrew C. Vitz²⁷ | Michael Ward²⁸ | Elly Knight²⁹

Correspondence: Aaron A. Skinner (skinnerayayron93@gmail.com)

Received: 27 November 2024 | **Revised:** 19 May 2025 | **Accepted:** 20 May 2025

Funding: This work was supported by Ohio Agricultural Research and Development Center, British Trust for Ornithology, Environment and Climate Change Canada, EACOM Timber Corporation, Association of Field Ornithologists, Illinois Department of Natural Resources, Natural Sciences and Engineering Research Council of Canada, U.S. Fish and Wildlife Service, Mitacs, Conoco Phillips, Mark Constantine, Weyerhaeuser Company Limited, Svenska Forskningsrådet Formas, University of Manitoba, The British Birds Charitable Trust and Canadian Foundation for Innovation.

Keywords: annual cycle | Bergmann's rule | body size | Caprimulgidae | GPS tracking | mechanism | nightjars | productivity | seasonality | temperature regulation

ABSTRACT

Aim: To evaluate (1) whether three migratory nightjar species (Family Caprimulgidae) adhere to Bergmann's rule, (2) whether environmental factors on the breeding or wintering grounds determine body size, and (3) which mechanistic hypotheses best explain Bergmannian patterns in body size.

Location: North and South America; Europe and Africa.

Taxon: Eastern whip-poor-will (*Antrostomus vociferus*), Common nighthawk (*Chordeiles minor*) and European nightjar (*Caprimulgus europaeus*).

Methods: We used GPS tracking and morphometric data to assess competing hypotheses explaining variation in body size for each species, based on their breeding ($n = 3388$) and wintering ($n = 189$) locations.

Results: All three species exhibited Bergmannian patterns in body size, providing the first evidence that nightjars conform to Bergmann's rule despite adaptations to severe environmental conditions. Environmental and geographic variables at breeding sites were stronger predictors of body size than wintering-site variables. Although we found partial support for Bergmann's temperature regulation hypothesis, geographic variables, rather than specific environmental factors, emerged as the strongest predictors of body size variation.

Main Conclusions: Latitude and longitude correlated strongly with environmental variables and migratory distance; thus, these geographical variables likely encompass many factors that influence body size in nightjars. The present study is among the first to use tracking data from individual birds to understand how environmental pressures across the annual cycle are related to body size. Our findings highlight the critical role of geographic breeding-ground factors in shaping Bergmannian patterns, offering robust evidence to support nearly two centuries of research since Bergmann's rule was first described in 1847.

First, second, and last authors are ordered by their contribution; Authors 3-28 are listed alphabetically.

For affiliations refer to page 13.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2025 The Author(s). *Journal of Biogeography* published by John Wiley & Sons Ltd.

RESUMEN

Objetivo: Evaluar (1) si la regla de Bergmann se observa en tres especies de Caprimúlgidos (Familia Caprimulgidae) migratorios, (2) si los factores ambientales en las áreas de reproducción o invernada influyen más en el tamaño corporal, y (3) qué hipótesis mecanísticas explican mejor los patrones de tamaño corporal asociados a la regla de Bergmann.

Ubicación: América del Norte y del Sur; Europa y África.

Taxón: *Antrostomus vociferus*, *Chordeiles minor* y *Caprimulgus europaeus*.

Métodos: Utilizamos datos GPS para evaluar hipótesis que expliquen las variaciones en el tamaño corporal de cada especie, considerando su ubicación de reproducción e invernada.

Resultados: Las tres especies mostraron patrones Bergmannianos, siendo esta la primera evidencia de la regla Bergman en los Caprimúlgidos, a pesar de sus adaptaciones fisiológicas a condiciones extremas. Los factores ambientales en las áreas de reproducción fueron mejores predictores del tamaño corporal que los de las áreas de invernada. Aunque encontramos cierta base para la hipótesis de regulación térmica de Bergmann, las variables geográficas, más que los factores ambientales específicos, resultaron ser los principales predictores de la variación en el tamaño corporal.

Conclusiones Principales: La latitud y la longitud mostraron una fuerte correlación con las variables ambientales y la distancia migratoria, lo que sugiere que estas variables geográficas probablemente integran múltiples factores que influyen en el tamaño corporal de los Caprimúlgidos. Este estudio es el primero en utilizar datos de seguimiento de individuos de aves para comprender cómo las presiones ambientales a lo largo del ciclo anual moldean el tamaño corporal. Nuestros hallazgos destacan el papel fundamental de los factores geográficos en las áreas de reproducción para dar forma a los patrones Bergmannianos, proporcionando evidencia sólida que respalda casi dos siglos de investigación desde que la regla de Bergmann fue descrita por primera vez en 1847.

1 | Introduction

Bergmann's Rule describes a pattern wherein "body size varies inversely with ambient temperature, so that body size increases with latitude" (translated from the original German by Watt et al. 2010), a rule broadly supported in endotherms at both intraspecific (Ashton 2002; Henry et al. 2023; Meiri and Dayan 2003) and interspecific levels (Fröhlich et al. 2023; He et al. 2023). While thermoregulatory pressures have historically been considered the most likely mechanism driving Bergmann's Rule, other environmental clines (e.g., productivity, seasonality) that are correlated with latitude could also drive clines in body size (Meiri 2011). Despite strong support for Bergmann's Rule, the primary mechanisms causing widespread latitudinal clines in body size remain uncertain. Identifying these mechanisms is critical for predicting the impact of climate change on body size (Teplitsky and Millien 2014).

Several hypotheses have emerged to explain Bergmannian clines (Figure 1; Blackburn et al. 1999; Jones et al. 2005). Hypothesis 1 is phenomenological, because demonstrating a phenomenon (i.e., adherence to Bergmann's Rule) is a prerequisite to testing the mechanism behind the pattern (Meiri 2011). The remaining hypotheses explore mechanisms underlying potential geographic patterns in body size:

1. The geographic pattern hypothesis predicts body size variation with an organism's geographic position on Earth (latitude, longitude or elevation). Latitude and elevation influence temperature, and longitude is related to other potentially important environmental gradients (e.g., precipitation).
2. The temperature regulation hypothesis, originally invoked by Bergmann, suggests that homeothermic organisms closer to the poles experience selective pressure for efficient heat retention and thus have larger bodies to reduce the ratio of body surface area to volume (Salewski and Watt 2017; Watt et al. 2010). This hypothesis has been

expanded to include more efficient heat dissipation in hotter climates (Blackburn et al. 1999).

3. The productivity hypothesis suggests that food availability is the most important determinant of body size, as juvenile growth can be limited by nutrition (Huston and Wolverton 2011; McNab 2010; Rosenzweig 1968). Compelling evidence exists for this hypothesis, particularly where food availability and latitude are negatively related, and organismal body size increases with food availability (i.e., larger at more southerly latitudes; Katti and Price 2003; Meiri et al. 2007).
4. The seasonality hypothesis (or the fasting endurance hypothesis; Ashton 2002; Blackburn et al. 1999) suggests that larger individuals have relatively lower metabolic rates compared to their smaller-bodied conspecifics, making them more resilient to periods of limited resources found in more seasonal environments (Lindstedt and Boyce 1985).
5. The migratory distance hypothesis is not related to environmental clines, but instead suggests that body size variation is determined by selective pressures related to long-distance migration (Alerstam et al. 2003; Vágási et al. 2016).

Latitude-body size relationships are more complex in migratory species (Mainwaring and Street 2021; Meiri and Dayan 2003), although they tend to exhibit Bergmannian clines as well (Ashton 2002; He et al. 2023). Long-distance migratory birds are morphologically adapted for lengthy migratory flights (Hein et al. 2012; Phillips et al. 2018), although selective pressures on traits often come into conflict. For example, being light-weight facilitates energy-efficient long-distance flight, yet large wings, muscle hypertrophy and energy stores provide the power and fuel for a costly journey (Alerstam et al. 2003). Wing size is also under strong selection from competing pressures including migration strategy (e.g., energy- or time-efficient strategies), overall distance




FIGURE 1 | Body size within a species often varies with latitude or other geographic variables (e.g., elevation and longitude), presumably as a function of underlying environmental gradients. Establishing the existence of a pattern (1) is a prerequisite for testing potential mechanisms (Hypotheses 2–5). Each hypothesis except 5 (migration distance) can be tested in Caprimulgids using environmental data from breeding or winter locations.

and route, foraging behaviour (Marchetti et al. 1995) and habitat associations across the annual cycle (Hedenström 2008; Leisler and Winkler 2003; Rayner 1988; Saino et al. 2017). In species that minimise migration time (Hedenström 2008; Hedenström and Hedh 2024), selection is expected to favour individuals with longer wings and lower body mass at greater migration distances, due to the advantages these traits confer for higher flight speeds, aerodynamic efficiency, and faster energy accumulation at stopovers (Alerstam and Hedenström 1998). Because migration imposes diverse physiological and ecological challenges (Piersma et al. 2005), it is important to consider the migration distance hypothesis (distinct from the “migration ability hypothesis” summarised in Blackburn et al. 1999), where body size evolves in response to the demands of long-distance migration (Gibson et al. 2019).

Uncovering the mechanisms driving Bergmann's Rule is complicated in migratory birds, many of which spend more than half of the year on the wintering grounds (La Sorte et al. 2017), and thus are influenced by varying environmental conditions across the annual cycle. Indeed, some studies have found wintering grounds conditions to be important in determining body size (Bosco et al. 2023; Gibson et al. 2019; Weeks et al. 2020). When this is the case, the distribution of body sizes observed across the breeding range will be influenced by a species' migratory connectivity, or the degree to which individuals that breed in close proximity to each other also overwinter in close proximity (Webster and Marra 2005). Species with weak migratory

connectivity may have varied morphology within breeding populations if diverse conditions (e.g., environmental, geographic) exist throughout the wintering grounds.

Adherence to Bergmann's Rule in migratory birds also depends on how body size is measured (Bailey et al. 2020; Teplitsky and Millien 2014). Most studies examining latitudinal clines in body size in birds use wing chord (i.e., the distance from the bend of the wing to the tip of the longest primary feather) to reflect overall body size (Ashton 2002). However, there is no single best metric of body size, particularly for migratory birds. For example, several studies argue that body mass is the best indicator of size (Blackburn et al. 1999; Freeman and Jackson 1990). However, body mass is flexible over time (Gibson et al. 2019), varying with body condition, reproductive status, and migratory phase (e.g., fattening prior to departure), and can fluctuate within and between seasons (Guglielmo 2018; Scott et al. 1994; Gosler et al. 1998). Wing chord, on the other hand, is relatively ‘fixed’ in species that moult flight feathers once a year (feather wear throughout the annual cycle negligibly influences wing length in most species; e.g., Fernández and Lank 2007). Wing chord, therefore, is the most repeatable single measure of body size in birds (Goodenough et al. 2010; Subasinghe et al. 2021; Gosler et al. 1998); however, it can be confounded with selective pressures from migration, such as migration speed (Bennett et al. 2019), distance (Vágási et al. 2016) or strategy (Vincze et al. 2019). Thus, it is important to consider multiple measures of body size as traits may respond differently to

selective pressures (Bailey et al. 2020), resulting in ‘shape shifting’ (i.e., changes in body proportions) as well as changes in body size (Ryding et al. 2021).

Nightjars (Family Caprimulgidae, hereafter ‘Caprimulgids’) are a widespread group of cryptic, nocturnal, aerial insectivores that are notoriously difficult to study (Holyoak 2001). Temperate-breeding Caprimulgids are migratory with high site fidelity (Bakermans et al. 2022; Ng et al. 2018; Norevik et al. 2025), making them ideal candidates for understanding how evolutionary pressures across the annual cycle influence body size. Furthermore, Caprimulgids have evolved behavioural and physiological adaptations to help withstand severe environmental conditions, which otherwise could strongly select for locally-adapted body sizes. For example, Caprimulgids can experience significant heat stress while roosting or nesting (Newberry et al. 2021; O’Connor et al. 2017), and food intake may be restricted during certain times of the month when low moonlight levels inhibit aerial foraging (Evens et al. 2020; Norevik et al. 2019; Souza-Cole 2021). To withstand these temperature extremes and periods of fasting, Caprimulgids leverage low metabolic rates (Lane et al. 2004), torpor (Brigham et al. 2006; Smit et al. 2011), gular fluttering and efficient evaporative heat dissipation. These adaptations help tolerate severe environmental conditions, which could reduce adherence to Bergmann’s rule (He et al. 2023; Ryding et al. 2021). This is consistent with the only available study examining Bergmann’s rule in Caprimulgids, which found that the Common nighthawk (*Chordeiles minor*) did not exhibit a Bergmannian cline on the breeding grounds (James 1970).

We used temperate-breeding migratory Caprimulgids as a focal group to explore intraspecifically how multiple factors drive patterns in body size across the annual cycle. We used cross-continental breeding and winter locations from three species: the Neotropical-Nearctic Eastern whip-poor-will (*Antrostomus vociferus*) and Common nighthawk, and the palaearctic-Afrotropical European nightjar (*Caprimulgus europaeus*). We had the following objectives: (1) determine whether migratory species from the Caprimulgid family adhere to the pattern of Bergmann’s Rule; (2) evaluate whether environmental conditions at the breeding or wintering grounds have a stronger influence on body size; and (3) test various mechanistic hypotheses for any observed geographic variation in body size (Figure 1).

2 | Methods

2.1 | Capture, Tag Deployment and Measurement

In total, 7318 Caprimulgids were captured using mist nets and conspecific playback across North America and Europe (Figure 2). We ringed birds using uniquely numbered bands, aged and sexed individuals, and measured wing chord using a wing ruler (precision = 1 mm) and mass using a digital scale (precision = 0.1 g), or a Pesola scale in rare cases (precision = 1.0 g; Pyle 1997). For individuals that were captured multiple times, we averaged wing chord and mass measurements across all captures where the individual was the same age, prioritising morphometrics from adults (each individual was included only once to avoid pseudoreplication). We examined error in wing chord

measurements but found negligible differences with individuals recaptured within the same season (Figure S1).

A subset of 215 Caprimulgids were tagged with GPS loggers that tracked their geographic location during migration and on the wintering grounds (Figure 2). Whip-poor-wills and European nightjars were tagged with archival GPS loggers $\leq 3.6\%$ of the bird’s mass, and nighthawks were tagged with Argos-GPS satellite loggers $\leq 5.0\%$ of the bird’s mass. Tags took GPS fixes (accuracy 10 m) during migration and the over-winter period, with fix frequency ranging from every few hours to every 10 days. We recaptured whip-poor-wills and European nightjars the following summer using the same methodology. See the following references for additional details on field methods for each species (Knight et al. 2021 for nighthawks; Bakermans et al. 2022; Korpach et al. 2022; Skinner et al. 2022 for whip-poor-wills; Lathouwers et al. 2022; Norevik et al. 2019 for European nightjars). Given that only a small subset of individuals had wintering data, we divided our data into two datasets: one with all breeding locations of individuals that passed filtering (hereafter, ‘breeding dataset’; see *dataset manipulation* section in the [Supporting Information](#)), and one with just individuals with wintering data (hereafter, ‘annual cycle dataset’).

2.2 | Body Size Measures

We used wing chord and body mass as separate measures of body size. We assumed that, while mass is a dynamic trait that fluctuates within and across seasons, that mass on the breeding grounds is correlated with mass on the wintering grounds. We find that this is true when examining the mass of Kirtland’s Warbler (*Setophaga kirtlandii*) individuals measured first on the wintering grounds, and then measured on the breeding grounds a few months later ($\beta_{\text{scaled}} = 0.78$, $p = 2.84 \times 10^{-6}$; data from Nathan Cooper, Figure S2). We believe this is a reasonable assumption for our focal species as well for two additional reasons. First, body mass is largely a function of structural size outside of periods of migratory fattening and egg-carrying (Freeman and Jackson 1990). Nightjars in our study had little fat within the stationary breeding season (62.5% of fat scores were 0), and migratory fattening appeared minimal in Whip-poor-will and Nighthawk (Figure S3). Thus, there is evidence that these two species do not deposit extensive fat reserves, so we believe their mass may be less variable compared with other species of migratory birds (also see Hidalgo-Rodríguez et al. 2021). European nightjars did increase in fat as migration approached, so we subset our data to remove individuals captured during periods of migratory fattening (Figure S3). Second, of repeat captures in our dataset, > 50% of the variation in mass *across years* is attributed to measurements coming from the same individual (Figure S3). The most parsimonious explanation for this finding is that body mass is correlated across the annual cycle.

We also examined the intraspecific allometric scaling relationship between mass and wing chord in our focal species (Santoro and Calzada 2022). Generally, wing chord increases proportionally with body mass to the one-third power (i.e., under isometric scaling), but this is species-specific and can vary substantially depending on a species’ life history. We used




FIGURE 2 | Panels (A–C) depict breeding locations for 3,388 Caprimulgids captured on the breeding grounds in North America and Europe. Each circle depicts the number of individuals captured in a quarter degree grid cell, and sample sizes should be interpreted as ‘less than’ the number in question (e.g., the smallest circle is < 20 captures). Box and whisker plots show the distribution of latitudes and longitudes where outliers are depicted as points, the whiskers extend to 1.5 *the interquartile range, the hinges represent the 25th and 75th percentiles, and the central bar is the median. Panels (D–F) show breeding and wintering connections for 189 Caprimulgids tagged with archival GPS tags in North America and Europe. Lines represent the great-circle paths between the breeding and wintering locations (i.e., not the actual migratory paths). Panel G shows the distribution of migratory distances for each species.

standardised major axis regression in the ‘smatr’ package in R (Warton et al. 2012) to examine scaling of wing and mass in our three species. We found that allometric scaling in European nightjars is consistent with isometry, whereas scaling in night-hawks and whip-poor-wills is hyperallometric (Figure S4).

2.3 | Dataset Manipulation

We filtered both breeding and annual cycle datasets in several ways prior to analysis to minimise extraneous variation in body size due to age, sex, time of day, time of year and the potential influence of climate change (Table S1). Our final breeding and annual cycle datasets consisted of 3388 and 189 Caprimulgids, respectively (Section ‘Dataset manipulation’, Figures S3, S5, S6).

2.4 | Environmental Predictors

We extracted environmental data from the summer capture site on the breeding grounds and the first wintering location (determined via GPS data; Figure 2) using Google Earth Engine with the package ‘rgee’ (Aybar 2021) for each bird. We extracted climate variables from WorldClim 2.1 (historical monthly averages from 1970 to 2000; Fick and Hijmans 2017), and the Enhanced Vegetation Index (EVI) from Landsat 8 Collection Tier 1 8-Day EVI composites (Chander et al. 2009; Huete et al. 2002; see Table 1 for specific environmental variables). We extracted values within a circular buffer corresponding to the mean home range size for each species in each stationary season (whip-poor-will: breeding 500m, winter 100m | Bakermans et al. 2022; Skinner et al. 2023; Tonra et al. 2019; Wilson 2003; nighthawk: breeding 5000m, winter 1000m | Ng et al. 2018; European nightjar: breeding 500m, winter 100m | Mitchell et al. 2020; Sharps et al. 2015). We used migratory dates from tracked individuals and published literature to establish the relevant months that environmental variables influence individuals on both the breeding and wintering grounds (Table S2). We used mean monthly climatic values because (1) our

hypotheses did not necessarily correspond to minimum or maximum values, (2) environments were relatively stable during the months in question and (3) the mean, maximum and minimum values were highly correlated. We then averaged climatic variables across months in examination of the temperature regulation and productivity hypotheses, whereas we calculated the coefficient of variation (standard deviation/mean) in examination of the seasonality hypothesis (Table 1).

2.5 | Migration Distance

Migration distance was calculated as the sum of the great-circle distances between each consecutive GPS fix between the breeding grounds and the first wintering location. We examined the impact of tag sampling resolution on our calculated migration distances and found that fix rate impacted migration distance only in whip-poor-wills. Thus, we rarefied Whip-poor-will migration data from Canada so that sampling resolution resembled that of other data collectors, after which there was no further relationship between sampling resolution and migration distance ($t = -1.51$; $p = 0.13$, $df = 93$; Supporting Information ‘Rarefaction of GPS data’ section).

2.6 | Statistical Analysis

We created 12 sets of models, one for each combination of body size measure (2), species (3) and dataset (2). For the breeding dataset, we used Akaike’s information criterion corrected for small sample size (AIC_c) to examine how the confounding factors of age and sex influenced wing and mass, and how the quadratic variable ‘time since sunset’ (the difference between the capture time and local sunset) influenced just mass (Figures S7, S8), for each species $\times$ body size measure (Bartoń 2023). For the annual cycle dataset, we examined the importance of the quadratic variable ‘time since sunset’ as a confounding variable using a similar AIC_c framework (‘Confounding variables’ section). While we could not control

TABLE 1 | Nine combinations of potential hypotheses for the pattern/mechanism that explains Bergmann’s rule and the season of influence.

Hypothesis	Season	Global model
1 Geographic pattern	Breeding	Latitude (+), longitude (+/–), elevation (+)
1 Geographic pattern	Winter	Latitude (+), Longitude (+/–), Elevation (+)
2 Temperature regulation	Breeding	Solar radiation (–), Temperature (–)
2 Temperature regulation	Winter	Solar radiation (–), Temperature (–)
3 Productivity	Breeding	EVI (+), Precipitation (+)
3 Productivity	Winter	EVI (+), Precipitation (+)
4 Seasonality	Breeding	EVI CV (+), Precipitation CV (+), Temperature CV (+)
4 Seasonality	Winter	EVI CV (+), Precipitation CV (+), Temperature CV (+)
5 Migration distance	NA	Migratory distance (+/–)

Note: Numbers for hypotheses (H) match those used in the text. We show the associated global model, and the predicted direction of effect (on the breeding grounds) for each variable: Positive (+), negative (–) or differing predictions (+/–) by species or dependent variable. For example, for the migration distance hypothesis we predict that wing length will increase, but body mass will decrease, with longer migratory journeys. Note the geographic pattern hypothesis is phenomenological (i.e., examining patterns in body size), where all other hypotheses test the underlying mechanisms. Abbreviations: CV, coefficient of variation; and EVI, Enhanced Vegetation Index.

for age or sex in analyses using the annual cycle datasets due to small and uneven sample sizes, we conducted the analysis twice, once using all ages and sexes ($n = 189$) and once using only adult males ($n = 145$).

For each of the 12 model sets, we organised the potential predictors of body size into unique and mutually exclusive hypotheses (Table 1) and generated a global model for each hypothesis. Thus, model sets consisted of a global model and all subsets of the global model for each hypothesis, as well as a null model to help determine the explanatory ability of the other models. This approach provides a framework for nested model construction from a theoretically justified global model, rather than arbitrary combinations of predictors. For the annual cycle dataset we generated two global models for each hypothesis, one for the breeding grounds and one for the wintering grounds. This allowed for direct competition of breeding versus wintering models (in line with objective 2), and kept the number of models reasonable for each species $\times$ body size measure (Burnham et al. 2011). The number of models differed in each species $\times$ body size measure $\times$ dataset combination due to collinearity between predictor variables (Figure S9). There were 116 models from the breeding dataset (between 17 and 21 models per species $\times$ body size measure) and 226 models from the annual cycle dataset (between 34 and 42 models per species $\times$ body size measure). We scaled and centered all predictor and response variables to allow for the comparison of parameter estimates across species. We modelled the relationship between body size and our predictors using linear regression, and considered models within $\Delta AIC_c \leq 4$ from the top model to be 'important' for addressing our three objectives, only if the null model was not within $\Delta AIC_c \leq 4$ (Burnham et al. 2011; Burnham and Anderson 2004). We then conducted model averaging on parameters from 'important' models to understand the strength and direction of the effects (Bartón 2023). Inclusion of models $2 < \Delta AIC_c < 4$ in model averaging is a more conservative approach than using a cutoff of $\Delta AIC_c < 2$, as it incorporates additional uncertainty in our parameter estimates (Burnham et al. 2011; Richards et al. 2011). See 'Model validation' section in the Supporting Information for details on the inspection of residuals, multicollinearity and spatial autocorrelation (Figures S10–S11).

2.7 | Evaluation of Objectives

To test whether Caprimulgids adhere to Bergmann's rule (Objective 1), we examined model selection results along with model averaged parameter estimates. A species followed Bergmann's rule if: (1) breeding latitude was in at least one 'important' model, (2) birds were larger at higher breeding latitudes and (3) the 85% unconditional confidence interval did not overlap zero.

To test the relative influence of breeding vs. wintering grounds on body size (Objective 2), we examined the number of models from the breeding vs. wintering grounds within $\Delta AIC_c \leq 4$ of the top model. The number of models with breeding vs. wintering data was roughly equivalent in each annual cycle model set, although across all species $\times$ body size measures there were 8 more models with wintering data due to reduced collinearity in predictor variables on the wintering grounds.

To evaluate which hypothesis best explains variation in body size (Objective 3), we compared the strength and direction of parameter coefficient estimates for each species $\times$ body size measure using the dataset we determined most important in Objective 2. For example, if the breeding grounds are determined more important than the wintering grounds in the annual cycle analysis, we will assess support for the varying hypotheses using the breeding dataset.

3 | Results

3.1 | Confounding Variables

In the breeding dataset, the most supported model controlling for confounding variation in both mass and wing included age and sex for all species (except age for nighthawks; see Supporting Information "Confounding variables" section), and the quadratic variable time since sunset in controlling for mass (Table S3). In the annual cycle dataset, the top model included the time since sunset in mass models for European nightjars and whip-poor-wills (Table S4).

3.2 | Objective 1—Adherence to Bergmann's Rule

Breeding latitude appeared in 'important' models for all species $\times$ body size measures (Table 2). Parameter estimates were always positive and 85% unconditional confidence intervals did not overlap zero (Figure 4).

3.3 | Objective 2—Importance of Breeding vs. Wintering Grounds

Using the annual cycle dataset, models from the breeding season received the highest likelihood in four of six species $\times$ body size measure combinations, while the null model appeared within $\Delta AIC_c \leq 4$ for the other two species $\times$ body size measures (Nighthawk and European nightjar mass; Table S5). We found qualitatively similar patterns whether we ran the analysis with all individuals or only adult males (Figures S12–S13). Thus, we examine Objective 3 using the breeding dataset.

3.4 | Objective 3—Testing Hypotheses

When using the breeding dataset, we observed support for the geographic pattern hypothesis in all species $\times$ body size measures (Figure 3), and support for the temperature regulation hypotheses in the whip-poor-will $\times$ mass model set (Table 2). Latitude was a particularly strong predictor of body size (six of the seven largest parameter estimates; Figure 4 and Table S6), and was positively correlated with both mass and wing chord for all three species. Longitude had a relatively weaker effect on size but still appeared in 'important' models in whip-poor-will and nighthawk mass, as well as whip-poor-will wing, and was positive in these three cases. Elevation was also found in 'important' models for all species $\times$ body size measures, although the direction of effect was opposite to that predicted by Bergmann's rule for whip-poor-will wing chord, and the 85% confidence intervals

TABLE 2 | 'Important' ($\Delta AIC_c \leq 4$) models plus the top model from each hypothesis constructed using the breeding data for six species $\times$ body size measures. K is the number of model parameters (the intercept and residual variance parameters are both estimated as well), wt is the model weight, and R^2 is the coefficient of determination.

Nighthawk mass—fixed variables: time since sunset ² + sex					
Model	K	ΔAIC_c	wt	−2 Log-likelihood	Hypothesis R^2
Elevation + Latitude + Longitude	8	0.00	0.79	−371.49	Geographic pattern 0.39
Latitude + Longitude	7	2.72	0.20	−373.90	Geographic pattern 0.38
Temperature	6	17.98	0.00	−382.58	Temperature Regulation 0.34
EVI CV + Temperature CV	7	28.09	0.00	−386.59	Seasonality 0.33
Precipitation	6	41.85	0.00	−394.51	Productivity 0.29
Null	5	45.87	0.00	−397.57	NA 0.28
AICc value of best model = 759.46.					
Nighthawk wing—fixed variables: sex					
Model	K	ΔAIC_c	wt	−2 Log-likelihood	Hypothesis R^2
Latitude	4	0.00	0.36	−431.70	Geographic pattern 0.18
Elevation + Latitude	5	0.26	0.32	−430.80	Geographic pattern 0.19
Latitude + Longitude	5	1.76	0.15	−431.55	Geographic pattern 0.18
Elevation + Latitude + Longitude	6	2.29	0.11	−430.78	Geographic pattern 0.19
Temperature	4	4.33	0.04	−433.86	Temperature Regulation 0.17
Precipitation	4	21.90	0.00	−442.65	Productivity 0.13
Temperature CV	4	25.07	0.00	−444.24	Seasonality 0.12
Null	3	35.77	0.00	−450.61	NA 0.08
AICc value of best model = 871.52.					
European nightjar mass – fixed variables: age + time since sunset ² + sex					
Model	K	ΔAIC_c	wt	−2 Log-likelihood	Hypothesis R^2
Latitude	7	0.00	0.71	−1546.88	Geographic pattern 0.50
Elevation + Latitude	8	1.78	0.29	−1546.76	Geographic pattern 0.50
Solar radiation + Temperature	8	12.93	0.00	−1552.33	Temperature Regulation 0.49
Temperature CV	7	92.70	0.00	−1593.23	Seasonality 0.46
Precipitation	7	116.26	0.00	−1605.01	Productivity 0.45

(Continues)

TABLE 2 | (Continued)

European nightjar mass – fixed variables: age + time since sunset ² + sex						
Model	K	ΔAICc	wt	–2 Log-likelihood	Hypothesis	R ²
Null	6	177.32	0.00	–1636.55	NA	0.43
AICc value of best model = 3107.83.						
European nightjar wing – fixed variables: age + sex						
Model	K	ΔAICc	wt	–2 Log-likelihood	Hypothesis	R ²
Elevation + Latitude	6	0.00	0.52	–2405.48	Geographic pattern	0.35
Latitude	5	0.16	0.48	–2406.57	Geographic pattern	0.34
EVI CV + Temperature CV	6	250.31	0.00	–2530.64	Seasonality	0.26
Precipitation	5	377.20	0.00	–2595.09	Productivity	0.21
Solar radiation + Temperature	6	550.31	0.00	–2680.63	Temperature Regulation	0.14
Null	4	769.07	0.00	–2792.03	NA	0.03
AICc value of best model = 4823.						
Whip-poor-will mass – fixed variables: age + time since sunset ² + sex						
Model	K	ΔAICc	wt	–2 Log-likelihood	Hypothesis	R ²
Temperature	7	0.00	0.42	–611.32	Temperature Regulation	0.35
Latitude + Longitude	8	0.26	0.37	–610.42	Geographic pattern	0.35
Elevation + Latitude + Longitude	9	1.78	0.17	–610.14	Geographic pattern	0.35
Precipitation CV + EVI CV + Temperature CV	9	12.90	0.00	–615.70	Seasonality	0.34
Precipitation	7	19.84	0.00	–621.24	Productivity	0.32
Null	6	121.33	0.00	–673.02	NA	0.17
AICc value of best model = 1236.87.						
Whip-poor-will wing – fixed variables: age + sex						
Model	K	ΔAICc	wt	–2 Log-likelihood	Hypothesis	R ²
Latitude + Longitude	6	0.00	0.45	–589.66	Geographic pattern	0.42
Elevation + Latitude + Longitude	7	0.22	0.40	–588.74	Geographic pattern	0.42

(Continues)

TABLE 2 | (Continued)

Whip-poor-will wing – fixed variables: age + sex						
Model	K	ΔAICc	wt	–2 Log-likelihood	Hypothesis	R ²
Elevation + Latitude	6	3.17	0.09	–591.24	Geographic pattern	0.42
Precipitation CV + Temperature CV	6	6.44	0.02	–592.88	Seasonality	0.41
EVI + Precipitation	6	13.73	0.00	–596.53	Productivity	0.41
Temperature	5	14.89	0.00	–598.13	Temperature Regulation	0.40
Null	4	144.14	0.00	–663.77	NA	0.23
AICc value of best model = 1191.48.						

overlapped zero half of the time (Figure 4). Finally, the univariate model with temperature was the top model for predicting whip-poor-will mass (Table 2), and had a large effect size in the direction consistent with Bergmann’s rule (Figure 4). Table S6 shows parameter estimates and unconditional confidence intervals from ‘important’ models and Table S7 shows full model selection results.

4 | Discussion

Bergmann’s rule is perhaps the oldest biogeographical rule, yet not all warm-blooded vertebrates follow this rule (Åkesson et al. 2020; Ashton et al. 2000; Blackburn and Gaston 1996; Meiri and Dayan 2003; Watt et al. 2010), and a consensus regarding the underlying mechanisms has evaded scientists. The present work is among the first to leverage tracking technology to understand how selective pressures from across the annual cycle are related to body size in migratory birds (but see Shipley et al. 2022). We found that the three Caprimulgid species we examined (Common nighthawk, Eastern whip-poor-will, and European nightjar) adhere to Bergmann’s rule. We also found that explanatory variables associated with the breeding grounds, particularly related to a bird’s geographical location on Earth, were the best predictors of body size. We did not find conclusive evidence for any single mechanistic hypothesis and suggest that geographic variables likely rose to the top in model selection because they combine information across many important environmental gradients. Our results demonstrate that despite being migratory, nocturnal and having adaptations to mitigate severe environmental conditions, Caprimulgids exhibit similar patterns in body size variation as other groups of birds, reinforcing the generalizability of Bergmann’s rule.

4.1 | Influence of Winter vs. Breeding Grounds on Body Size

Variables measured at breeding locations were better predictors of body size than variables at wintering locations in all three species, particularly when wing chord was used as the size metric. The breeding grounds may be more important than wintering grounds in determining body size for three primary reasons. First, thermoregulation in the nestling period, before birds have grown feathers, may be particularly impactful on stress levels, survival and ultimately in determining body size at the population level (Newberry and Swanson 2018). This is likely exacerbated in Caprimulgids, which build scrape nests in open habitats like the forest floor or bare ground, and are more exposed to environmental conditions relative to species that build insulated nests in closed habitats (e.g., trees or bushes; Mainwaring and Street 2021). Experimental and observational evidence suggests that high temperatures during development lead to smaller nestlings, and that these size differences persist as adults (Andrew et al. 2017; Youtz et al. 2020; but also see Nord and Nilsson 2011). While there is strong evidence for the importance of heat dissipation during hot breeding months (Andrew et al. 2018; Speakman and Król 2010), thermoregulatory pressures from a warming climate may shape body size in complex ways, potentially




FIGURE 3 | Isoclines of body mass based off of the top geographic model for three Caprimulgid species (Nighthawk: Latitude + longitude + elevation; European nightjar: Latitude; Whip-poor-will: Latitude + longitude). Mass was predicted for adult males and while holding the time since sunset at the mean for each species. Spatial predictions for wing chord were qualitatively similar.

exerting different selective pressures at different life stages (e.g., nestling vs. adult). Thus, additional research is needed to understand how selection acts across life stages and whether body size trade-offs exist between early-life constraints (e.g., chick survival) and adult thermoregulatory demands (Shipley et al. 2022).

Second, it is possible that the responsibilities associated with chick-rearing (particularly feeding and thermoregulation) during the breeding period may prevent Caprimulgids from using evolved behavioural or physiological mechanisms for thermal tolerance (e.g., torpor), or strongly select for birds that can withstand periods of prolonged fasting (e.g., allowing for continued thermoregulation of chicks). Thus, Caprimulgids may rely more on behavioural approaches to maintain thermoregulatory and metabolic equilibrium in the non-breeding season, while relying on optimal surface-area-to-volume ratios during the breeding season.

Third, all three Caprimulgids studied here had low migratory connectivity, such that birds from across large breeding ranges converge on core wintering areas (Knight et al. 2021; Korpach et al. 2022; Norevik et al. 2020; Skinner et al. 2022). We expect winter environmental conditions to be fairly uniform in species with low migratory connectivity, reducing the amount of environmental heterogeneity experienced on the wintering grounds in our study populations (Gibson et al. 2019; Jones et al. 2005). That being said, the distribution of breeding and wintering latitudes were similar in all three species, and birds could select wintering locations at smaller spatial scales to provide more favourable environmental conditions for their given body size (especially when wintering in mountainous areas). Thus, we conclude that the distribution of breeding and wintering latitudes were comparable, and instead the relative stability and benign conditions experienced on the tropical wintering grounds likely result in weaker selective pressures than the breeding grounds (e.g., winter environments may be entirely in species' thermoneutral zones; Ryding et al. 2021).

Finally, it is important to recognise the limitations of our annual cycle dataset, and the inconclusive results from this analysis when mass was the dependent variable. Indeed, the null model was the most supported in nighthawk and European nightjar mass in the annual cycle dataset. Given that nighthawk and European nightjars exhibited strong geographic gradients in body mass when we used the full breeding dataset, there was likely not enough of a latitudinal gradient on the breeding grounds (particularly in European nightjars), or statistical power to detect an effect in the annual cycle dataset in these two species $\times$ body size measures. Furthermore, we recognise that mass was only measured on the breeding grounds, and mass may change significantly throughout the annual cycle (perhaps even adaptively in response to the winter environment; Gibson et al. 2019). Future work measuring mass during wintering months would greatly improve our understanding of how environmental pressures from across the annual cycle influence body size.

4.2 | Mechanisms Underlying Bergmann's Rule

We were unable to identify a single mechanism that clearly explained the Bergmannian pattern for all three Caprimulgid species. Combinations of geographic variables (latitude, longitude and elevation) were almost always better predictors of body size than environmental variables (as in Gibson et al. 2019), suggesting that multiple environmental variables that covary with geographic clines are responsible for determining body size within Caprimulgid species, and that geographic variables may reflect a composite of mechanisms. In migratory organisms, Bergmann's rule has often been assessed with the most easily accessible data—geographic information from the breeding grounds (Ashton 2002). Our research suggests that these readily available data may indeed represent the most important variables influencing body size, and thus, past conclusions on Bergmann's rule in migratory birds are likely robust. Despite the difficulties in determining mechanisms in highly




FIGURE 4 | Parameter estimates (circles) 85% (thick error bars) and 95% (thin error bars) unconditional confidence intervals of all variables from 'important' models (those within 4 Δ AICc points of the top model), estimated using breeding season data from 3388 Caprimulgids. We included the 85% confidence intervals because they align with an AIC model selection framework and acknowledge the explicit link between AIC scores and p -values (Sutherland et al. 2023). We also include 95% confidence intervals to facilitate comparison with related work. Both predictor and response variables were scaled and centered to facilitate direct comparison between species.

collinear systems (Wigginton and Dobson 1999; Yom-Tov and Geffen 2011), examining model selection results, strength and direction of parameter estimates, and correlations between geographic and environmental variables can still provide insights into the mechanisms at play.

Latitude was the strongest predictor of body size in nearly all species $\times$ body size measures, with larger-bodied individuals occurring at higher latitudes. Generally, at more northerly latitudes conditions were cooler (and with less solar radiation), drier, less productive and more seasonal. Temperature regulation likely has a dominant role in determining body size in these Caprimulgid species (especially whip-poor-wills), given that it was often the

mechanistic model with the most support (as in Fan et al. 2019; Henry et al. 2023; Romano et al. 2021). Seasonality models often received some support as well, and parameter estimates tended to be positive and did not overlap zero (as predicted by Bergmann's rule). Low nighttime temperatures early in the breeding season may limit flying insect activity (Taylor 1963), particularly at the northern extremes of breeding ranges. Thus, larger-bodied individuals would be better-equipped to withstand such variations in prey availability (Ashton 2002; Blackburn et al. 1999; Lindstedt and Boyce 1985). Migration distances were also substantially longer for individuals breeding at more northerly latitudes in whip-poor-wills and nighthawks, consistent with the migration distance hypothesis for wing but not mass. Finally, we conclude that productivity is unlikely to be the driving force behind Bergmannian clines in Caprimulgids, given that more northerly sites were less productive than more southerly sites. Note that this is in direct contrast to previous studies that show body size tracking environmental gradients irrespective of latitude (Katti and Price 2003; Meiri et al. 2007; Nwaogu et al. 2018). Importantly, our data exhibited a negative productivity cline in 'ecologically and evolutionarily available' productivity as well (i.e., during the peak breeding season; unpublished data), which may be more relevant in determining body size in some species (Huston and Wolverton 2011).

The correlation between longitude and body size in whip-poor-wills and nighthawks was counterintuitive. Whip-poor-wills and nighthawks tended to be larger farther east, where it is generally hotter, wetter, more productive and less seasonal, and it seems unlikely that these variables would exert selective pressures on body size differently than they do along latitudinal gradients. An examination of parameter estimates associated with productivity models and migration distance in nighthawks confirmed that these factors were not responsible for variation in size along longitudinal gradients. This suggests that there could be unexplored additive or interactive effects (or functional forms) of environmental variables that were not explored in this study, or alternative pressures influencing body size along longitudinal gradients (e.g., pesticide use; Martínez-Padilla et al. 2017). While longitude has not been included in many Bergmann's rule studies, we argue that it can be related to important environmental gradients, similar to latitude or elevation, and thus should be included in studies aiming to understand environment-morphology relationships (e.g., James 1970).

Elevation can also be an important predictor of body size in birds, where colder temperatures at higher elevations are thought to increase body size for thermoregulatory reasons (Blackburn and Ruggiero 2001; García et al. 2021; Romano et al. 2021). We observed this pattern in nighthawks and European nightjars but not in whip-poor-wills. In European nightjars and whip-poor-wills, effect sizes were relatively small, 85% confidence intervals largely overlapped zero, and elevational gradients were minimal (95% of individuals occupied a gradient < 500 m). Thus, we focus on nighthawk response to elevational gradients. In addition to cooler climates at higher elevations, nighthawks experienced increased solar radiation and more arid climates, which may favour larger individuals as water lost to the environment is reduced in larger birds (James 1970; Jirinec et al. 2021; Nwaogu et al. 2018; Yom-Tov and Geffen 2006). Future work should include models testing the impact of both fluid and temperature regulation in

selecting for avian body size. Finally, through examining parameter coefficients we note there is marginal evidence of increased *relative* wing length at higher elevations in European nightjars (Gutiérrez-Pinto et al. 2014; Youngflesh et al. 2022). Future work should tie wing and mass measurements together at the level of the individual (e.g., allometric-informed indices of relative wing length; Youngflesh et al. 2022) to detect shapeshifting along geographical or environmental gradients.

4.3 | Conclusions for Conservation

Biological response to climate change can be split into three main categories: (1) range distribution shifts, (2) adaptation through phenotypical or phenological change or (3) local extinction (Millien et al. 2006). There is evidence that the northern limit of the nighthawk's breeding range is shifting northward (see eBird [trend map](#)) (LaSorte and Thompson III 2007), and climate change models predict that whip-poor-will ranges have the potential to expand at the north edge and contract at the south edge under a warming scenario (<https://www.audubon.org/field-guide/bird/eastern-whip-poor-will>). However, it is unclear whether Caprimulgid morphologies are also tracking environmental conditions through time, as seen by many species globally (Jirinec et al. 2021; Salewski et al. 2010; Weeks et al. 2020). Species that respond to spatial temperature gradients generally also respond to temperature increases through time (Youngflesh et al. 2022), suggesting that our focal species may be capable of altering morphology in response to increasing temperatures associated with climate change. However, given that Caprimulgids have high site fidelity (Bakermans et al. 2022; Ng et al. 2018; Norevik et al. 2025), long lifespans (> 10 years) and low diversity in migratory strategy (e.g., obligate migrants, small non-breeding ranges), they may be particularly susceptible to rapid changes in climate (Both et al. 2010; Gilroy et al. 2016). Our data show that European nightjars have increased in wing chord and decreased in body mass from 1990 to 2022, suggesting that they have increased in relative wing length in response to increasing temperatures (as in several other recent studies; Ryding et al. 2024; Weeks et al. 2020; Youngflesh et al. 2022). Long-term morphological datasets are needed for nighthawks and whip-poor-wills, and future research should work to better understand whether morphological changes (1) represent a genetic or plastic response to temperature gradients (Millien et al. 2006; Romano et al. 2025; Teplitsky and Millien 2014), and (2) if any observed changes are biologically significant for thermoregulation (Nord et al. 2024).

Affiliations

¹Department of Zoology, The University of British Columbia, Vancouver, British Columbia, Canada | ²School of Environment and Natural Resources, The Ohio State University, Columbus, Ohio, USA | ³Department of Biological Sciences, University of Manitoba, Winnipeg, Manitoba, Canada | ⁴Department of Biology, University of Regina, Regina, Saskatchewan, Canada | ⁵University of Winnipeg, Winnipeg, Manitoba, Canada | ⁶Department of Biology, Lund University, Lund, Sweden | ⁷Department of Integrative and Global Studies, Worcester Polytechnic Institute, Worcester, Massachusetts, USA | ⁸Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada | ⁹Illinois Natural History Survey,

Prairie Research Institute, University of Illinois Urbana-Champaign, Champaign, Illinois, USA | ¹⁰Museo Civico di Storia Naturale, Turin, Italy | ¹¹Skagen Bird Observatory, Birdlife Denmark, Skagen, Denmark | ¹²British Trust for Ornithology, Norfolk, UK | ¹³Department of Biology, Carleton University, Ottawa, Ontario, Canada | ¹⁴Biodiversity Research Centre, Université Catholique de Louvain (UCL), Louvain-la-Neuve, Belgium | ¹⁵Department of Biology, Behavioural Ecology and Ecophysiology Group, University of Antwerp, Wilrijk, Belgium | ¹⁶Autumn-Lynn Harrison, Migratory Bird Center, Smithsonian's National Zoo and Conservation Biology Institute, Washington, District of Columbia, USA | ¹⁷Zoology Unit, Finnish Museum of Natural History, Pb 17, 00014, University of Helsinki, Helsinki, Finland | ¹⁸Center for Macroecology, Evolution and Climate, University of Copenhagen, Copenhagen, Denmark | ¹⁹Research Group: Zoology, Biodiversity and Toxicology, Hasselt University, Centre for Environmental Sciences, Diepenbeek, Belgium | ²⁰University of Namur, Department of Geography, Institute of Life, Earth and Environment (ILEE), Namur, Belgium | ²¹The Earth Commons Institute; Department of Biology, McCourt School of Public Policy, Georgetown University, Washington, District of Columbia, USA | ²²Environment & Climate Change Canada, Canadian Wildlife Service, Regina, Saskatchewan, Canada | ²³Department of Biology, Centre for Animal Movement Research, Lund University, Lund, Sweden | ²⁴Migratory Bird Center, Smithsonian's National Zoo and Conservation Biology Institute, Washington, District of Columbia, USA | ²⁵Center for Macroecology, Evolution and Climate, University of Copenhagen, Copenhagen, Denmark | ²⁶Environment & Climate Change Canada, Canadian Wildlife Service, Saskatoon, Saskatchewan, Canada | ²⁷Massachusetts Division of Fisheries and Wildlife, Westborough, Massachusetts, USA | ²⁸Department of Natural Resources and Environmental Sciences, University of Illinois Urbana-Champaign, Urbana, Illinois, USA | ²⁹Migratory Bird Center, Smithsonian's National Zoo and Conservation Biology Institute, Washington, District of Columbia, USA

Author Contributions

A.A.S. and E.K. conceptualized the article. A.A.S. led the analysis with assistance from E.K. and A.M.K. A.A.S. wrote the initial draft. All authors provided morphological and/or tracking data and contributed comments on manuscript drafts.

Acknowledgements

We would like to thank Sylvia Heredia for the drawings of the nightjars in Figures 2 and 3, as well as the nightjar wing on Figure 4. We would also like to thank all technicians and personnel who participated in the collection of the morphological data used in these analyses, particularly the late Fabrizio Silvano (Stazzano, Alessandria, Italy) at the River Scrivia ringing station. We thank the U.S. Fish & Wildlife Service (USFWS-F18AC00618; W-197-R), The Ohio Agricultural Research and Development Center (Ohio State University SEEDS Grant), Association of Field Ornithologists, Illinois Department of Natural Resources, The University of Manitoba, Natural Sciences and Engineering Research Council of Canada (Discovery Program), Canadian Foundation for Innovation (John R Evans Leaders Fund), the American Ornithological Society, Conoco Phillips Global Signature Programs, Environment and Climate Change Canada, Mitacs (Accelerate Program), Weyerhaeuser Company Limited, EACOM Timber Corporation, Resolute Forest Products, Domtar Inc., Natural England, Forestry England, The British Birds Charitable Trust, Mark Constantine, the Swedish Research Council (2016-03625, 2020-03707), and the British Trust of Ornithology for funding.

Mist netting was conducted under the following permits: J.nr. SN 302-009 (Denmark), ringing permit A, 0045 (Italy), VARELY/1320/2022 (Finland), M470-12, M72-15, M74-20 (Sweden), IACUC protocol #22040 and USGS Bird Band Permit #23959 (Illinois, USA), Banding Permit No. 10876F and Wildlife Scientific Collector's Authorization No. 1095705 (Canada), IACUC protocol no. 20-119 and USGS Bird Banding Laboratory permit no. 21963-S (Massachusetts, USA), AUP00001523 (University

of Alberta), 15-16, 18-66 (National Zoological Park and Conservation Biology), 15032-04 (Texas Tech University), 23-05-13-16C, 15-05-16-19C (University of South Dakota), SFCQ2017-02 (Environment and Climate Change Canada), US Federal Bird Banding Permits 09700, 22199, 22834, and Scientific Permit to Capture and Band Migratory Birds 10277, 10365, 10619H, 10534, 10169, 10887 (Environment and Climate Change Canada).

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The data and code needed to reproduce all analyses, figures, and tables in the main text of the manuscript are available on Dryad (https://datadryad.org/stash/share/3Ic-u0MVyd1TgknPDNcMOZBw2O4lBxdD_4np321u2Yw). The collation of this rich morphological dataset with information across the annual cycle is the most complete for Caprimulgids to date. We would be interested in future collaborations using these data.

References

- Åkesson, S., P. W. Atkinson, A. Bermejo, et al. 2020. "Evolution of Chain Migration in an Aerial Insectivorous Bird, the Common Swift *Apus Apus*." *Evolution* 74, no. 10: 2377–2391. <https://doi.org/10.1111/evo.14093>.
- Alerstam, T., and A. Hedenström. 1998. "The Development of Bird Migration Theory." *Journal of Avian Biology* 29, no. 4: 343–369. <https://doi.org/10.2307/3677155>.
- Alerstam, T., A. Hedenstrom, and S. Åkesson. 2003. "Long-Distance Migration: Evolution and Determinants." *Oikos* 103, no. 2: 247–260. <https://doi.org/10.1034/j.1600-0706.2003.12559.x>.
- Andrew, S. C., M. Awasthy, A. D. Griffith, S. Nakagawa, and S. C. Griffith. 2018. "Clinal Variation in Avian Body Size Is Better Explained by Summer Maximum Temperatures During Development Than by Cold Winter Temperatures." *Auk* 135, no. 2: 206–217. <https://doi.org/10.1642/AUK-17-129.1>.
- Andrew, S. C., L. L. Hurley, M. M. Mariette, and S. C. Griffith. 2017. "Higher Temperatures During Development Reduce Body Size in the Zebra Finch in the Laboratory and in the Wild." *Journal of Evolutionary Biology* 30, no. 12: 2156–2164. <https://doi.org/10.1111/jeb.13181>.
- Ashton, K. G. 2002. "Patterns of Within-Species Body Size Variation of Birds: Strong Evidence for Bergmann's Rule." *Global Ecology and Biogeography* 11, no. 6: 505–523.
- Ashton, K. G., M. C. Tracy, and A. de Queiroz. 2000. "Is Bergmann's Rule Valid for Mammals?" *American Naturalist* 156, no. 4: 390–415. <https://doi.org/10.1086/303400>.
- Aybar, C. 2021. "rgee: R Bindings for Calling the "Earth Engine" API (Version R Package Version 1.1.2) [Computer Software]." <https://CRAN.R-project.org/package=rgee>.
- Bailey, L. D., L. E. B. Kruuk, R. Allen, M. Clayton, J. Stein, and J. L. Gardner. 2020. "Using Different Body Size Measures Can Lead to Different Conclusions About the Effects of Climate Change." *Journal of Biogeography* 47, no. 8: 1687–1697. <https://doi.org/10.1111/jbi.13850>.
- Bakermans, M. H., J. M. Driscoll, and A. C. Vitz. 2022. "Habitat Selection and Site Fidelity on Winter Home Ranges of Eastern Whip-Poor-Wills (*Anurostomus Vociiferus*)." *Avian Conservation and Ecology* 17, no. 2: art17. <https://doi.org/10.5751/ACE-02237-170217>.
- Bartoń, K. 2023. "MuMIn: Multi-Model Inference [Computer Software]." <https://CRAN.R-project.org/package=MuMIn>.
- Bennett, R. E., A. D. Rodewald, K. V. Rosenberg, et al. 2019. "Drivers of Variation in Migration Behavior for a Linked Population of Long-Distance Migratory Passerine." *Auk* 136, no. 4. <https://doi.org/10.1093/auk/ukz051>.
- Blackburn, T. M., and K. J. Gaston. 1996. "Spatial Patterns in the Species Richness of Birds in the New World." *Ecography* 19, no. 4: 369–376. <https://doi.org/10.1111/j.1600-0587.1996.tb00001.x>.
- Blackburn, T. M., K. J. Gaston, and N. Loder. 1999. "Geographic Gradients in Body Size: A Clarification of Bergmann's Rule." *Diversity and Distributions* 5, no. 4: 165–174. <https://doi.org/10.1046/j.1472-4642.1999.00046.x>.
- Blackburn, T. M., and A. Ruggiero. 2001. "Latitude, Elevation and Body Mass Variation in Andean Passerine Birds." *Global Ecology and Biogeography* 10: 245–259. <https://doi.org/10.1046/j.1466-822X.2001.00237.x>.
- Bosco, L., A. Otterbeck, T. Fransson, A. Lindén, M. Piha, and A. Lehikoinen. 2023. "Increasing Winter Temperatures Explain Body Size Decrease in Wintering Bird Populations of Northern Europe—But Response Patterns Vary Along the Spatioclimatic Gradient." *Global Ecology and Biogeography* 32, no. 12: 2100–2110. <https://doi.org/10.1111/geb.13754>.
- Both, C., C. A. M. Van Turnhout, R. G. Bijlsma, H. Siepel, A. J. Van Strien, and R. P. B. Foppen. 2010. "Avian Population Consequences of Climate Change Are Most Severe for Long-Distance Migrants in Seasonal Habitats." *Proceedings of the Royal Society B: Biological Sciences* 277, no. 1685: 1259–1266. <https://doi.org/10.1098/rspb.2009.1525>.
- Brigham, R. M., C. P. Woods, J. E. Lane, Q. E. Fletcher, and F. Geiser. 2006. "Ecological Correlates of Torpor Use Among Five Caprimulgiform Birds." *Acta Zoologica Sinica* 52, no. Supplement: 401–404.
- Burnham, K. P., and D. R. Anderson. 2004. "Multimodel Inference: Understanding AIC and BIC in Model Selection." *Sociological Methods & Research* 33, no. 2: 261–304. <https://doi.org/10.1177/0049124104268644>.
- Burnham, K. P., D. R. Anderson, and K. P. Huyvaert. 2011. "AIC Model Selection and Multimodel Inference in Behavioral Ecology: Some Background, Observations, and Comparisons." *Behavioral Ecology and Sociobiology* 65, no. 1: 23–35. <https://doi.org/10.1007/s00265-010-1029-6>.
- Chander, G., B. L. Markham, and D. L. Helder. 2009. "Summary of Current Radiometric Calibration Coefficients for Landsat MSS, TM, ETM+, and EO-1 ALI Sensors." *Remote Sensing of Environment* 113, no. 5: 893–903. <https://doi.org/10.1016/j.rse.2009.01.007>.
- Evens, R., C. Kowalczyk, G. Norevik, et al. 2020. "Lunar Synchronization of Daily Activity Patterns in a Crepuscular Avian Insectivore." *Ecology and Evolution* 10, no. 14: 7106–7116. <https://doi.org/10.1002/ece3.6412>.
- Fan, L., T. Cai, Y. Xiong, G. Song, and F. Lei. 2019. "Bergmann's Rule and Allen's Rule in Two Passerine Birds in China." *Avian Research* 10, no. 1: 34. <https://doi.org/10.1186/s40657-019-0172-7>.
- Fernández, G., and D. Lank. 2007. "Variation in the Wing Morphology of Western Sandpipers (*Calidris Mauri*) in Relation to Sex, Age Class, and Annual Cycle." *Auk* 124, no. 3: 1037–1046. <https://doi.org/10.1093/auk/124.3.1037>.
- Fick, S. E., and R. J. Hijmans. 2017. "WorldClim 2: New 1-km Spatial Resolution Climate Surfaces for Global Land Areas." *International Journal of Climatology* 37: 4302–4315.
- Freeman, S., and W. M. Jackson. 1990. "Univariate Metrics Are Not Adequate to Measure Avian Body Size." *Ornithology* 107, no. 1: 69–74.
- Fröhlich, A., D. Kotowska, R. Martyka, and M. R. E. Symonds. 2023. "Allometry Reveals Trade-Offs Between Bergmann's and Allen's Rules, and Different Avian Adaptive Strategies for Thermoregulation." *Nature Communications* 14, no. 1: 1101. <https://doi.org/10.1038/s41467-023-36676-w>.
- García, J., J. Arizaga, J. I. Rodríguez, D. Alonso, and S. Suárez-Seoane. 2021. "Morphological Differentiation in a Migratory Bird Across

- Geographic Gradients in Mountains of Southern Europe." *Journal of Biogeography* 48, no. 11: 2828–2838. <https://doi.org/10.1111/jbi.14242>.
- Gibson, D., A. D. Hornsby, M. B. Brown, et al. 2019. "Migratory Shorebird Adheres to Bergmann's Rule by Responding to Environmental Conditions Through the Annual Lifecycle." *Ecography* 42, no. 9: 1482–1493. <https://doi.org/10.1111/ecog.04325>.
- Gilroy, J. J., J. A. Gill, S. H. M. Butchart, V. R. Jones, and A. M. A. Franco. 2016. "Migratory Diversity Predicts Population Declines in Birds." *Ecology Letters* 19, no. 3: 308–317. <https://doi.org/10.1111/ele.12569>.
- Goodenough, A. E., R. Stafford, C. L. Catlin-Groves, A. L. Smith, and A. G. Hart. 2010. "Within- and Among-Observer Variation in Measurements of Animal Biometrics and Their Influence on Accurate Quantification of Common Biometric-Based Condition Indices." *Annales Zoologici Fennici* 47, no. 5: 323–334. <https://doi.org/10.5735/086.047.0503>.
- Gosler, A. G., J. J. D. Greenwood, J. K. Baker, and N. C. Davidson. 1998. "The Field Determination of Body Size and Condition in Passerines: A Report to the British Ringing Committee." *Bird Study* 45, no. 1: 92–103. <https://doi.org/10.1080/00063659809461082>.
- Guglielmo, C. G. 2018. "Obese Super Athletes: Fat-Fueled Migration in Birds and Bats." *Journal of Experimental Biology* 221, no. Suppl_1: jeb165753. <https://doi.org/10.1242/jeb.165753>.
- Gutiérrez-Pinto, N., K. G. McCracken, L. Alza, et al. 2014. "The Validity of Ecogeographical Rules Is Context-Dependent: Testing for Bergmann's and Allen's Rules by Latitude and Elevation in a Widespread Andean Duck: Testing Ecogeographical Rules." *Biological Journal of the Linnean Society* 111, no. 4: 850–862. <https://doi.org/10.1111/bij.12249>.
- He, J., J. Tu, J. Yu, and H. Jiang. 2023. "A Global Assessment of Bergmann's Rule in Mammals and Birds." *Global Change Biology* 29: 5199–5210. <https://doi.org/10.1111/gcb.16860>.
- Hedenström, A. 2008. "Adaptations to Migration in Birds: Behavioural Strategies, Morphology and Scaling Effects." *Philosophical Transactions of the Royal Society, B: Biological Sciences* 363, no. 1490: 287–299. <https://doi.org/10.1098/rstb.2007.2140>.
- Hedenström, A., and L. Hedh. 2024. "Seasonal Patterns and Processes of Migration in a Long-Distance Migratory Bird: Energy or Time Minimization?" *Proceedings of the Royal Society B: Biological Sciences* 291, no. 2024: 20240624. <https://doi.org/10.1098/rspb.2024.0624>.
- Hein, A. M., C. Hou, and J. F. Gillooly. 2012. "Energetic and Biomechanical Constraints on Animal Migration Distance." *Ecology Letters* 15, no. 2: 104–110. <https://doi.org/10.1111/j.1461-0248.2011.01714.x>.
- Henry, E., L. Santini, M. A. J. Huijbregts, and A. Benítez-López. 2023. "Unveiling the Environmental Drivers of Intraspecific Body Size Variation in Terrestrial Vertebrates." *Global Ecology and Biogeography* 32, no. 2: 267–280. <https://doi.org/10.1111/geb.13621>.
- Hidalgo-Rodríguez, P., P. Sáez-Gómez, J. Blas, A. Hedenström, and C. Camacho. 2021. "Body Mass Dynamics of Migratory Nightjars Are Explained by Individual Turnover and Fueling." *Behavioral Ecology* 32, no. 6: 1086–1093. <https://doi.org/10.1093/beheco/arab042>.
- Holyoak, D. T. 2001. *Nightjars and Their Allies: The Caprimulgiformes*. OUP Oxford.
- Huete, A., K. Didan, T. Miura, E. P. Rodriguez, X. Gao, and L. G. Ferreira. 2002. "Overview of the Radiometric and Biophysical Performance of the MODIS Vegetation Indices." *Remote Sensing of Environment* 83, no. 1: 195–213. [https://doi.org/10.1016/S0034-4257\(02\)00096-2](https://doi.org/10.1016/S0034-4257(02)00096-2).
- Huston, M. A., and S. Wolverton. 2011. "Regulation of Animal Size by eNPP, Bergmann's Rule and Related Phenomena." *Ecological Monographs* 81, no. 3: 349–405. <https://doi.org/10.1890/10-1523.1>.
- James, F. C. 1970. "Geographic Size Variation in Birds and Its Relationship to Climate." *Ecology* 51, no. 3: 365–390. <https://doi.org/10.2307/1935374>.
- Jirinec, V., R. C. Burner, B. R. Amaral, et al. 2021. "Morphological Consequences of Climate Change for Resident Birds in Intact Amazonian Rainforest." *Science Advances* 7, no. 46: eabk1743. <https://doi.org/10.1126/sciadv.abk1743>.
- Jones, J., C. E. Gibb, S. C. Millard, et al. 2005. "Multiple Selection Pressures Generate Adherence to Bergmann's Rule in a Neotropical Migratory Songbird." *Journal of Biogeography* 32, no. 10: 1827–1833. <https://doi.org/10.1111/j.1365-2699.2005.01320.x>.
- Katti, M., and T. D. Price. 2003. "Latitudinal Trends in Body Size Among Over-Wintering Leaf Warblers (Genus *Phylloscopus*)." *Ecography* 26, no. 1: 69–79. <https://doi.org/10.1034/j.1600-0587.2003.03264.x>.
- Knight, E. C., A.-L. Harrison, A. L. Scarpignato, et al. 2021. "Comprehensive Estimation of Spatial and Temporal Migratory Connectivity Across the Annual Cycle to Direct Conservation Efforts." *Ecography* 44: 1–15. <https://doi.org/10.1111/ecog.05111>.
- Korpach, A. M., C. M. Davy, A. Mills, and K. C. Fraser. 2022. "Migratory Connectivity and Timing for an At-Risk Canadian Landbird, Eastern Whip-Poor-Will (*Antrostomus Vociferus*), From Two Geographically Distant Breeding Areas." *Canadian Journal of Zoology* 100: 660–668. <https://doi.org/10.1139/cjz-2021-0175>.
- La Sorte, F. A., D. Fink, P. J. Blancher, et al. 2017. "Global Change and the Distributional Dynamics of Migratory Bird Populations Wintering in Central America." *Global Change Biology* 23, no. 12: 5284–5296. <https://doi.org/10.1111/gcb.13794>.
- Lane, J. E., D. L. Swanson, R. M. Brigham, and A. E. McKechnie. 2004. "Physiological Responses to Temperature by Whip-Poor-Will: More Evidence for the Evolution of Low Metabolic Rates in Caprimulgiformes." *Condor* 106, no. 4: 921–925.
- LaSorte, F. A., and F. R. Thompson III. 2007. "Poleward Shifts in Winter Ranges of North American Birds." *Ecology* 88, no. 7: 1803–1812. <https://doi.org/10.1890/06-1072.1>.
- Lathouwers, M., T. Artois, N. Dendoncker, et al. 2022. "Rush or Relax: Migration Tactics of a Nocturnal Insectivore in Response to Ecological Barriers." *Scientific Reports* 12, no. 1: 4964. <https://doi.org/10.1038/s41598-022-09106-y>.
- Leisler, B., and H. Winkler. 2003. "Morphological Consequences of Migration in Passerines." In *Avian Migration*, edited by P. Berthold, E. Gwinner, and E. Sonnenschein, 175–186. Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-662-05957-9_11.
- Lindstedt, S. L., and M. S. Boyce. 1985. "Seasonality, Fasting Endurance, and Body Size in Mammals." *American Naturalist* 125, no. 6: 873–878. <https://doi.org/10.1086/284385>.
- Mainwaring, M. C., and S. E. Street. 2021. "Conformity to Bergmann's Rule in Birds Depends on Nest Design and Migration." *Ecology and Evolution* 11: ece3.8034. <https://doi.org/10.1002/ece3.8034>.
- Marchetti, K., T. Price, and A. Richman. 1995. "Correlates of Wing Morphology With Foraging Behaviour and Migration Distance in the Genus *Phylloscopus*." *Journal of Avian Biology* 26, no. 3: 177. <https://doi.org/10.2307/3677316>.
- Martínez-Padilla, J., D. López-Idiáquez, J. J. López-Perea, R. Mateo, A. Paz, and J. Viñuela. 2017. "A Negative Association Between Bromadiolone Exposure and Nestling Body Condition in Common Kestrels: Management Implications for Vole Outbreaks." *Pest Management Science* 73, no. 2: 364–370. <https://doi.org/10.1002/ps.4435>.
- McNab, B. K. 2010. "Geographic and Temporal Correlations of Mammalian Size Reconsidered: A Resource Rule." *Oecologia* 164, no. 1: 13–23. <https://doi.org/10.1007/s00442-010-1621-5>.
- Meiri, S. 2011. "Bergmann's Rule – What's in a Name?" *Global Ecology and Biogeography* 20, no. 1: 203–207. <https://doi.org/10.1111/j.1466-8238.2010.00577.x>.

- Meiri, S., and T. Dayan. 2003. "On the Validity of Bergmann's Rule." *Journal of Biogeography* 30, no. 3: 331–351. <https://doi.org/10.1046/j.1365-2699.2003.00837.x>.
- Meiri, S., Y. Yom-Tov, and E. Geffen. 2007. "What Determines Conformity to Bergmann's Rule?" *Global Ecology and Biogeography* 16, no. 6: 788–794. <https://doi.org/10.1111/j.1466-8238.2007.00330.x>.
- Millien, V., S. Kathleen Lyons, L. Olson, F. A. Smith, A. B. Wilson, and Y. Yom-Tov. 2006. "Ecotypic Variation in the Context of Global Climate Change: Revisiting the Rules." *Ecology Letters* 9, no. 7: 853–869. <https://doi.org/10.1111/j.1461-0248.2006.00928.x>.
- Mitchell, L. J., T. Kohler, P. C. L. White, and K. E. Arnold. 2020. "High Interindividual Variability in Habitat Selection and Functional Habitat Relationships in European Nightjars Over a Period of Habitat Change." *Ecology and Evolution* 10, no. 12: 5932–5945. <https://doi.org/10.1002/eece3.6331>.
- Newberry, G. N., R. S. O'Connor, and D. L. Swanson. 2021. "Urban Rooftop-Nesting Common Nighthawk Chicks Tolerate High Temperatures by Hyperthermia With Relatively Low Rates of Evaporative Water Loss." *Ornithological Applications* 123, no. 3: duab016. <https://doi.org/10.1093/ornithapp/duab016>.
- Newberry, G. N., and D. L. Swanson. 2018. "Elevated Temperatures Are Associated With Stress in Rooftop-Nesting Common Nighthawk (*Chordeiles Minor*) Chicks." *Conservation Physiology* 6, no. 1. <https://doi.org/10.1093/conphys/coy010>.
- Ng, J. W., E. C. Knight, A. L. Scarpignato, A.-L. Harrison, E. M. Bayne, and P. P. Marra. 2018. "First Full Annual Cycle Tracking of a Declining Aerial Insectivorous Bird, the Common Nighthawk (*Chordeiles Minor*), Identifies Migration Routes, Nonbreeding Habitat, and Breeding Site Fidelity." *Canadian Journal of Zoology* 96, no. 8: 869–875. <https://doi.org/10.1139/cjz-2017-0098>.
- Nord, A., and J.-Å. Nilsson. 2011. "Incubation Temperature Affects Growth and Energy Metabolism in Blue Tit Nestlings." *American Naturalist* 178, no. 5: 639–651. <https://doi.org/10.1086/662172>.
- Nord, A., E. Persson, J. K. R. Tabh, and E. Thorald. 2024. "Shrinking Body Size May Not Provide Meaningful Thermoregulatory Benefits in a Warmer World." *Nature Ecology & Evolution* 8, no. 3: 387–389. <https://doi.org/10.1038/s41559-023-02307-2>.
- Norevik, G., S. Åkesson, A. Andersson, J. Bäckman, and A. Hedenström. 2019. "The Lunar Cycle Drives Migration of a Nocturnal Bird." *PLoS Biology* 17, no. 10: e3000456. <https://doi.org/10.1371/journal.pbio.3000456>.
- Norevik, G., S. Åkesson, T. Artois, et al. 2020. "Wind-Associated Detours Promote Seasonal Migratory Connectivity in a Flapping Flying Long-Distance Avian Migrant." *Journal of Animal Ecology* 89, no. 2: 635–646. <https://doi.org/10.1111/1365-2656.13112>.
- Norevik, G., S. Åkesson, and A. Hedenström. 2025. "The Spatial Consistency and Repeatability of Migratory Flight Routes and Stationary Sites of Individual European Nightjars Based on Multiannual GPS Tracks." *Movement Ecology* 13, no. 1: 8. <https://doi.org/10.1186/s40462-025-00537-6>.
- Nwaogu, C. J., B. I. Tieleman, K. Bitrus, and W. Cresswell. 2018. "Temperature and Aridity Determine Body Size Conformity to Bergmann's Rule Independent of Latitudinal Differences in a Tropical Environment." *Journal of Ornithology* 159, no. 4: 1053–1062. <https://doi.org/10.1007/s10336-018-1574-8>.
- O'Connor, R. S., B. O. Wolf, R. M. Brigham, and A. E. McKechnie. 2017. "Avian Thermoregulation in the Heat: Efficient Evaporative Cooling in Two Southern African Nightjars." *Journal of Comparative Physiology B* 187, no. 3: 477–491. <https://doi.org/10.1007/s00360-016-1047-4>.
- Phillips, A. G., T. Töpfer, K. Böhning-Gaese, and S. A. Fritz. 2018. "Evidence for Distinct Evolutionary Optima in the Morphology of Migratory and Resident Birds." *Journal of Avian Biology* 49, no. 10: e01807. <https://doi.org/10.1111/jav.01807>.
- Piersma, T., J. Pérez-Tris, H. Mouritsen, U. Bauchinger, and F. Bairlein. 2005. "Is There a 'Migratory Syndrome' Common to All Migrant Birds?" *Annals of the New York Academy of Sciences* 1046, no. 1: 282–293. <https://doi.org/10.1196/annals.1343.026>.
- Pyle, P. 1997. *Identification Guide to North American Birds, Part I (Vol. 1)*. Slate Creek Press.
- Rayner, J. M. V. 1988. "Form and Function in Avian Flight." *Current Ornithology* 5: 1–66.
- Richards, S. A., M. J. Whittingham, and P. A. Stephens. 2011. "Model Selection and Model Averaging in Behavioural Ecology: The Utility of the IT-AIC Framework." *Behavioral Ecology and Sociobiology* 65, no. 1: 77–89. <https://doi.org/10.1007/s00265-010-1035-8>.
- Romano, A., R. Ambrosini, M. Caprioli, A. Costanzo, A. Novelli, and D. Rubolini. 2025. "Shrinking Body Size Under Climate Warming Is Not Associated With Selection for Smaller Individuals in a Migratory Bird." *Journal of Animal Ecology* 94, no. 5: 958–970. <https://doi.org/10.1111/1365-2656.70027>.
- Romano, A., R. Séchaud, and A. Roulin. 2021. "Generalized Evidence for Bergmann's Rule: Body Size Variation in a Cosmopolitan Owl Genus." *Journal of Biogeography* 48, no. 1: 51–63. <https://doi.org/10.1111/jbi.13981>.
- Rosenzweig, M. L. 1968. "The Strategy of Body Size in Mammalian Carnivores." *American Midland Naturalist* 80, no. 2: 299–315. <https://doi.org/10.2307/2423529>.
- Ryding, S., M. Klaassen, G. J. Tattersall, J. L. Gardner, and M. R. E. Symonds. 2021. "Shape-Shifting: Changing Animal Morphologies as a Response to Climatic Warming." *Trends in Ecology & Evolution* 36, no. 11: 1036–1048. <https://doi.org/10.1016/j.tree.2021.07.006>.
- Ryding, S., A. McQueen, M. Klaassen, G. J. Tattersall, and M. R. E. Symonds. 2024. "Long- and Short-Term Responses to Climate Change in Body and Appendage Size of Diverse Australian Birds." *Global Change Biology* 30, no. 10: e17517. <https://doi.org/10.1111/gcb.17517>.
- Saino, N., R. Ambrosini, M. Caprioli, et al. 2017. "Wing Morphology, Winter Ecology, and Fecundity Selection: Evidence for Sex-Dependence in Barn Swallows (*Hirundo Rustica*)." *Oecologia* 184, no. 4: 799–812. <https://doi.org/10.1007/s00442-017-3918-0>.
- Salewski, V., W. M. Hochachka, and W. Fiedler. 2010. "Global Warming and Bergmann's Rule: Do Central European Passerines Adjust Their Body Size to Rising Temperatures?" *Oecologia* 162, no. 1: 247–260. <https://doi.org/10.1007/s00442-009-1446-2>.
- Salewski, V., and C. Watt. 2017. "Bergmann's Rule: A Biophysiological Rule Examined in Birds." *Oikos* 126, no. 2: 161–172. <https://doi.org/10.1111/oik.03698>.
- Santoro, S., and J. Calzada. 2022. "Allometry to Evaluate Allen's Rule in Climate Warming." *Trends in Ecology & Evolution* 37: 375–377. <https://doi.org/10.1016/j.tree.2022.02.012>.
- Scott, I., P. I. Mitchell, and P. R. Evans. 1994. "Seasonal Changes in Body Mass, Body Composition and Food Requirements in Wild Migratory Birds." *Proceedings of the Nutrition Society* 53, no. 3: 521–531. <https://doi.org/10.1079/PNS19940062>.
- Sharps, K., I. Henderson, G. Conway, N. Armour-Chelu, and P. M. Dolman. 2015. "Home-Range Size and Habitat Use of European Nightjars *Caprimulgus Europaeus* Nesting in a Complex Plantation-Forest Landscape." *IBIS* 157, no. 2: 260–272. <https://doi.org/10.1111/ibi.12251>.
- Shipley, J. R., C. W. Twining, C. C. Taff, M. N. Vitousek, and D. W. Winkler. 2022. "Selection Counteracts Developmental Plasticity in Body-Size Responses to Climate Change." *Nature Climate Change* 12, no. 9: 863–868. <https://doi.org/10.1038/s41558-022-01457-8>.

- Skinner, A. A., S. N. Matthews, M. P. Ward, et al. 2023. "Eastern Whip-Poor-Will Have Larger Nonbreeding Home Ranges in Areas With More Agriculture and Forest Fragmentation." *Ornithological Applications* 125: 1–15. <https://doi.org/10.1093/ornithapp/duac050>.
- Skinner, A. A., M. P. Ward, I. Souza-Cole, et al. 2022. "High Spatiotemporal Overlap in the Non-Breeding Season Despite Geographically Dispersed Breeding Locations in the Eastern Whip-Poor-Will (*Antrostomus Vociferus*)." *Diversity and Distributions*: 1–15. <https://doi.org/10.1111/ddi.13477>.
- Smit, B., J. G. Boyles, R. M. Brigham, and A. E. McKechnie. 2011. "Torpor in Dark Times: Patterns of Heterothermy Are Associated With the Lunar Cycle in a Nocturnal Bird." *Journal of Biological Rhythms* 26, no. 3: 241–248. <https://doi.org/10.1177/0748730411402632>.
- Souza-Cole, I. F. 2021. *Understanding the Diel Activity Patterns and Determinants of Abundance of the Eastern Whip-Poor-Will*. University of Illinois Urbana-Champaign.
- Speakman, J. R., and E. Król. 2010. "Maximal Heat Dissipation Capacity and Hyperthermia Risk: Neglected Key Factors in the Ecology of Endotherms." *Journal of Animal Ecology* 79, no. 4: 726–746. <https://doi.org/10.1111/j.1365-2656.2010.01689.x>.
- Subasinghe, K., M. R. E. Symonds, M. Vidal-García, et al. 2021. "Repeatability and Validity of Phenotypic Trait Measurements in Birds." *Evolutionary Biology* 48: 100–114.
- Sutherland, C., D. Hare, P. J. Johnson, D. W. Linden, R. A. Montgomery, and E. Droge. 2023. "Practical Advice on Variable Selection and Reporting Using Akaike Information Criterion." *Proceedings of the Royal Society B: Biological Sciences* 290: 20231261.
- Taylor, L. R. 1963. "Analysis of the Effect of Temperature on Insects in Flight." *Journal of Animal Ecology* 32, no. 1: 99–117. <https://doi.org/10.2307/2520>.
- Teplitsky, C., and V. Millien. 2014. "Climate Warming and Bergmann's Rule Through Time: Is There Any Evidence?" *Evolutionary Applications* 7, no. 1: 156–168. <https://doi.org/10.1111/eva.12129>.
- Tonra, C. M., J. R. Wright, and S. N. Matthews. 2019. "Remote Estimation of Overwintering Home Ranges in an Elusive, Migratory Nocturnal Bird." *Ecology and Evolution* 00: 1. <https://doi.org/10.1002/ece3.5723>.
- Vágási, C. I., P. L. Pap, O. Vincze, G. Osváth, J. Erritzøe, and A. P. Møller. 2016. "Morphological Adaptations to Migration in Birds." *Evolutionary Biology* 43, no. 1: 48–59. <https://doi.org/10.1007/s11692-015-9349-0>.
- Vincze, O., C. I. Vágási, P. L. Pap, C. Palmer, and A. P. Møller. 2019. "Wing Morphology, Flight Type and Migration Distance Predict Accumulated Fuel Load in Birds." *Journal of Experimental Biology* 222, no. 1: jeb183517. <https://doi.org/10.1242/jeb.183517>.
- Warton, D. I., R. A. Duursma, D. S. Falster, and S. Taskinen. 2012. "Smatr 3—An R Package for Estimation and Inference About Allometric Lines." *Methods in Ecology and Evolution* 3, no. 2: 257–259. <https://doi.org/10.1111/j.2041-210X.2011.00153.x>.
- Watt, C., S. Mitchell, and V. Salewski. 2010. "Bergmann's Rule; A Concept Cluster?" *Oikos* 119, no. 1: 89–100. <https://doi.org/10.1111/j.1600-0706.2009.17959.x>.
- Webster, M. S., and P. P. Marra. 2005. "The Importance of Understanding Migratory Connectivity and Seasonal Interactions." In *Birds of Two Worlds: The Ecology and Evolution of Temperate-Tropical Migration*, edited by R. Greenberg and P. P. Marra, 199–209. John Hopkins University Press.
- Weeks, B. C., D. E. Willard, M. Zimova, et al. 2020. "Shared Morphological Consequences of Global Warming in North American Migratory Birds." *Ecology Letters* 23, no. 2: 316–325. <https://doi.org/10.1111/ele.13434>.
- Wigginton, J. D., and F. S. Dobson. 1999. "Environmental Influences on Geographic Variation in Body Size of Western Bobcats." *Canadian Journal of Zoology* 77, no. 5: 802–813. <https://doi.org/10.1139/z99-037>.
- Wilson, M. 2003. *Distribution, Abundance, and Home Range of the Whip-Poor-Will (*Caprimulgus Vociferus*) in a Managed Forest Landscape*. College of William & Mary. <http://scholarworks.wm.edu/etd/1539626403>.
- Yom-Tov, Y., and E. Geffen. 2006. "Geographic Variation in Body Size: The Effects of Ambient Temperature and Precipitation." *Oecologia* 148, no. 2: 213–218. <https://doi.org/10.1007/s00442-006-0364-9>.
- Yom-Tov, Y., and E. Geffen. 2011. "Recent Spatial and Temporal Changes in Body Size of Terrestrial Vertebrates: Probable Causes and Pitfalls." *Biological Reviews* 86, no. 2: 531–541. <https://doi.org/10.1111/j.1469-185X.2010.00168.x>.
- Youngflesh, C., J. F. Saracco, R. B. Siegel, and M. W. Tingley. 2022. "Abiotic Conditions Shape Spatial and Temporal Morphological Variation in North American Birds." *Nature Ecology & Evolution* 6: 1860–1870. <https://doi.org/10.1038/s41559-022-01893-x>.
- Youtz, J., K. D. Miller, E. K. Bowers, et al. 2020. "Bergmann's Rule Is Followed at Multiple Stages of Postembryonic Development in a Long-Distance Migratory Songbird." *Ecology and Evolution* 10: 10672–10686. <https://doi.org/10.1002/ece3.6721>.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.